Direct and transgenerational effects of an experimental heat wave on early life

stages in a freshwater snail

Katja Leicht[1,2], Otto Seppälä[1,3]

[1] Eawag, Swiss Federal Institute of Aquatic Science and Technology, 8600 Dübendorf,

Switzerland

[2] Department of Biological and Environmental Science, PO Box 35, 40014 University of

Jyväskylä, Finland

[3] ETH Zürich, Institute of Integrative Biology (IBZ), 8092 Zürich, Switzerland

Corresponding author:

Otto Seppälä

Eawag, Swiss Federal Institute of Aquatic Science and Technology, 8600 Dübendorf,

Switzerland

Tel: +41 58 765 5348

Fax: +41 58 765 5802

E-mail: otto.seppaelae@eawag.ch

Running title: Direct and transgenerational effects of a heat wave

Abstract

Global climate change imposes a serious threat to natural populations of many species. Estimates of the effects of climate change-mediated environmental stresses are, however, often based only on their direct effects on organisms, and neglect the potential transgenerational effects. We investigated whether high temperature (i.e. an experimental heat wave) that is known to reduce performance of adult *Lymnaea stagnalis* snails affects their offspring through maternal effects. Specifically, we tested whether eggs and hatched juveniles are affected by maternal thermal environment, and how strong these effects are compared with direct effects of temperature on offspring. We examined the effect of maternal thermal environment (15°C versus 25°C) on per offspring investment (egg size), and the role of both maternal and offspring thermal environments (15°C versus 25°C) on hatching success and developmental time of eggs, offspring survival after hatching, and hatchling size at the age of five weeks. Exposure of mothers to high temperature increased hatching success of eggs, and also made the onset of hatching earlier. However, high maternal temperature reduced the survival and the final size of hatched juveniles. Direct effects of high temperature on offspring survival were negative (both eggs and hatchlings), but increased the developmental rate and growth of those eggs and hatchlings that survived. Interestingly, the magnitude of transgenerational effects of high temperature on hatching success of eggs and hatchling survival were similar to its direct effects. This indicates that heat waves can affect natural populations through transgenerational effects, and that the magnitude of such effects can be equally strong to the direct effects of temperature, although this depends on the trait considered. Our results highlight the importance of considering transgenerational effects of climate warming when estimating its effects in the wild.

48   Keywords: climate change, environmental stress, global warming, *Lymnaea stagnalis*, maternal

49   effects

Introduction

Owing to global climate change, the average temperatures at the Earth's surface as well as the frequency and severity of extreme weather events such as summer heat waves are increasing (Easterling *et al.* 2000; Karl & Trenberth 2003; Meehl & Tebaldi 2004; Kirtman *et al.* 2013). These changes can have strong effects on organisms that escalate to higher levels of biological organization such as populations and communities (Walther *et al.* 2002; Parmesan & Yohe 2003; Walther 2010). Especially extreme weather events can dramatically influence population dynamics, species abundance, and species interactions (e.g. Easterling *et al.* 2000; Bruno *et al.* 2007; Hance *et al.* 2007). However, environmental conditions can not only influence the fitness of the individuals exposed to them but also the fitness of their offspring through transgenerational maternal and/or paternal effects (reviewed in Bernardo 1996; Mousseau & Fox 1998b). Hence, for understanding the effects of climate change on natural populations, studies examining such transgenerational effects are needed.

Especially transgenerational maternal effects after exposure to environmental stress can significantly alter offspring performance (e.g. Silbermann & Tatar 2000; Mitchell & Read 2005; Janhunen, Piironen & Peuhkuri 2010). Such effects can result from reduced physiological condition of the mother that limits the total amount of resources it invests in reproduction (Tessier *et al.* 1983; Steer *et al.* 2004), and/or exposure of offspring to hormones produced by the mother (McCormick 1999; Groothuis & Schwabl 2008). Maternal effects could also take place via altered resource allocation between reproduction and other traits depending on the environmental conditions the mother experiences. Challenging environmental conditions may, for example, reduce resource allocation to produced offspring to sustain self-maintenance, or increase per offspring investment when the reproductive value of individuals is changed so that investment in current reproduction increases at the expense of future reproduction (Fisher 1930;

Williams 1966). Furthermore, maternal effects can be adaptations to prepare offspring for the

future conditions they are about to encounter (e.g. herbivory, parasitism, pollution; Agrawal

2002; Moret 2006; Marshall 2008).

To understand the consequences of such transgenerational effects in context of climate

change, it is essential to estimate their direction and magnitude compared with the direct effects

of the same environmental factors. Maternal effects are typically strongest in early stages of

organisms' life histories (Mousseau & Dingle 1991; Heath, Fox & Heath 1999; Pettay *et al.*

2008), but such stages are often also highly susceptible to the direct effects of environmental

variation (e.g. Jang 1991; Zhang *et al.* 2015; Klockmann, Günter & Fischer 2017). For example,

temperature determines the development of eggs and juveniles by altering their metabolic and

physiological processes in many species (Gillooly *et al.* 2001; Person-Le Ruyet *et al.* 2004; Zuo

et al. 2012). High temperature in particular can impose a serious challenge by reducing the

hatching success of eggs and early survival of hatched offspring (Janhunen *et al.* 2010; Zhang *et*

al. 2015; Klockmann *et al.* 2017). Despite of high interest on transgenerational effects of climate

change in natural populations (reviewed in Donelson *et al.* 2018), their relative importance

compared with direct effects of the same environmental factors is, however, often overlooked

(but see Burgess & Marshall 2011; Parker *et al.* 2012; Salinas & Munch 2012; Shama *et al.* 2014;

Wadgymar, Mactavish & Anderson 2018).

Here, we tested whether high temperature as it can occur during heat waves has

transgenerational effects on offspring performance, which traits they affect, and how strong they

are compared with direct effects of high temperature in the freshwater snail *Lymnaea stagnalis* L.

(Gastropoda: Pulmonata). In this species, exposure of adult individuals to high temperature ($\geq$

25°C) initially increases growth and reproduction, but prolonged exposure (one week or longer)

ceases reproductive rate and reduces immune function (Seppälä & Jokela 2011; Leicht, Jokela &

Seppälä 2013). This indicates that high temperature is physiologically challenging and has strong

negative effects on adult snails. We estimated the effect of maternal thermal environment (15°C

versus 25°C) on per offspring investment by adult snails (egg size), and the role of both maternal

and offspring thermal environments on offspring performance (hatching success and

developmental time of eggs, survival of hatched offspring, offspring size at the age of five weeks)

using a full-factorial design. We found that high temperature affected offspring performance both

directly and through maternal effects. The relative importance and the direction of these effects

varied among traits, and the magnitude of maternal effects were equally strong to direct effects in

some of the examined traits. This highlights the importance of considering transgenerational

effects when estimating the consequences of climate change in natural populations.

Methods

Experimental animals

The snails used in this study came from a laboratory stock population (F_4 generation) originating

from a pond in Zurich, Switzerland (47°22'05''N, 8°34'41''E). The summer water temperature

in ponds typically remains low (< 16°C) in this region, although it depends on pond hydrology

(T. Salo, unpublished data). However, during heat waves, water temperature can rapidly increase

to 20–30°C and remain high for over two weeks (T. Salo, unpublished data). We started the stock

population using 45 adult snails collected from the pond. Since *L. stagnalis* prefers outcrossing

(Puurtinen *et al.* 2007; Nakadera *et al.* 2017), often engages in multiple matings (Nakadera *et al.*

2017), and can store sperm from those matings for over two months (Nakadera, Blom & Koene

2014), the stock population can be expected to reflect the genetic variation in the source

population well. We maintained the stock population in the approximate size of 400 individuals

at 15 ± 2°C (control temperature used in the experiment; see the section about experimental

design below) for two years before the study (see Leicht, Seppälä & Seppälä 2017).

We haphazardly collected 113 adult snails from the stock population and used them as a

maternal generation in the experiment. We placed the snails individually in 2 dl perforated plastic

cups sunk into a water bath (aged tap water at 15 ± 1°C) that was connected to a biological filter.

We used a water bath to provide maximal water quality for snails. This minimises the growth of

microorganisms in water that activate snail immune function (Seppälä & Leicht 2013). This is

important because immune challenge could potentially alter snail reproductive strategy and/or

affect the quality of produced offspring. We fed the snails with fresh lettuce *ad libitum* and

maintained them under these conditions for three days prior to the experiment to acclimate them

to the maintenance conditions. Because *L. stagnalis* snails can reproduce through self-fertilization

as well as through outcrossing using allosperm they have stored from previous matings (Cain

1956; Nakadera *et al.* 2014), experimental snails did not need a mating partner to oviposit eggs

under the used conditions.

Experimental design

Maternal treatments

At the beginning of the experiment, we randomly assigned the snails used as a maternal

generation (see the previous section) into two temperature treatments [15 ± 1°C (54 snails), 25 ±

1°C (59 snails); Fig. 1]. We used 25°C as a high (i.e. heat wave) temperature as it reduces

immune defence and life history traits in adult snails (Seppälä *et al.* 2011; Leicht *et al.* 2013), lies

above the thermal optimum for development and growth of juvenile snails (Vaughn 1953), and

occurs intermittently in habitats of snails during hot summers (T. Salo, unpublished data). We

chose 15°C as a control temperature as it is close to the thermal optimum of *L. stagnalis* (Vaughn

1953) and common in ponds (T. Salo, unpublished data). Note that we assigned more individuals

into the high temperature treatment because we expected increased mortality in those snails. We

transferred the snails to their treatment temperatures in cups filled with aged tap water at 15°C.

This allowed a slow change (over 10 h) to the target temperature for snails assigned to the high

temperature treatment. We then transferred the snails into perforated plastic cups (2 dl) sunk into

similar water baths as above, and exposed them to their respective temperature treatments for

seven days. At 15°C, 51 of these snails survived, and of those 25 oviposited eggs. At 25°C, 39

snails survived, and 34 of them reproduced. We did not measure the number of oviposited eggs in

this experiment as the effect of temperature on snail fecundity has been described in detail in

earlier studies (Leicht *et al.* 2013; 2017).

Figure 1. Summary of the full-factorial experimental design used to examine direct and

transgenerational effects of exposure of *L. stagnalis* snails to two temperatures (benign: 15°C;

heat wave: 25°C). One egg clutch per adult snail was used in offspring treatments. When

available, 15 hatchlings per egg clutch were used. When fewer individuals were available, they

were all used.

After the initial seven-day exposure to temperature treatments, we removed all egg

clutches oviposited by the snails from the cups and continued maintaining the snails under the

same experimental conditions. During the following ten days, we checked the cups twice a day

for new clutches to be collected for the next step of the experiment (see the next section). This

procedure ensured first, that the snails were exposed to their respective temperature treatments

long enough to induce strong effects on their performance before the clutches were collected (see

Leicht *et al.* 2013), and second, that the collected clutches were exposed to the maternal

temperature treatments only briefly. From each snail that oviposited (same individuals as above

reproduced), we collected the first clutch containing more than 20 eggs, or if only smaller

clutches were produced, the largest clutch. We placed each collected clutch on a millimetre paper

and photographed it from 10 cm above with a Fujifilm FinePix F30 digital camera (scene mode:

close up, focal length: 35 mm, aperture: F/2.8, shutter speed: 1/85, sensitivity: ISO-200, image

size: 2848×2136 pixels, focus mode: auto focus). From the digital images, we counted the eggs

in each clutch. Furthermore, we measured the two-dimensional area (mm^2) of five randomly

chosen eggs in each clutch from the digital images using ImageJ software (ImageJ 1.42q, Wayne

Rasband, National Institute of Health, USA). After photographing, we placed the clutches

individually into plastic cups with 0.4 dl of aged tap water to be transferred to the next step of the

experiment (see the next section). It is important to note that the time different snails needed for

ovipositing after the initial exposure period varied between one and ten days, which may have

affected the thermal challenge imposed to them as well as their offspring. However, to our

knowledge, oviposition cannot be induced artificially in this species.

Offspring treatments

We used a full-factorial design to expose egg clutches produced in both maternal temperature

treatments (see the previous section) to two offspring temperature treatments (15°C, 25°C; Fig.

1). In each maternal temperature treatment, we randomly assigned some of the oviposited egg

clutches to remain at the same temperature where they were produced and transferred the rest of

the clutches to the other temperature. The number of egg clutches per treatment combination

varied between 11-18, which was because of unequal mortality and probability to reproduce in

parental snails in different treatments (see the previous section). We slowly warmed up or cooled

down the clutches that were transferred to a different temperature as described above to avoid a

sudden change between temperatures. After that, we checked the clutches daily, counted the

number of hatched snails, and removed the hatchlings from the cups. When possible, we placed

15 hatchlings from each clutch individually in plastic cups filled with 0.4 dl of aged tap water.

We used all hatchlings when less than 15 individuals were available. We fed the snails with

Spirulina ad libitum, and changed the water in the cups twice a week. We reared the isolated

hatchlings for five weeks, and measured their survival and shell length to the nearest 0.1 mm

using a digital calliper at the end of the experiment.

Statistical analyses

We analysed the effect of temperature on the survival of adult snails during the initial exposure to

experimental treatments (a seven-day period before the collection of egg clutches started) using a

generalized linear model. We used the status of snails (survived, died) as a binomial response

variable (logit link function) and temperature treatment as a fixed factor. Additionally, we

analysed variation in snail reproductive status (oviposited, did not oviposit) using a similar

generalised linear model. We analysed the effect of temperature on the size of produced eggs (ln

transformed to homogenize error variance) using a mixed-model analysis of variance (mixed-

model ANOVA). In the analysis, we used a model with maternal temperature treatment as a fixed

and the clutch each egg originated from as a random factor (nested within maternal temperature).

To estimate the effects of maternal and offspring temperature on offspring performance, we

first analysed the variation in hatching success of eggs using a generalized linear model. In the

analysis, we used the proportion of eggs that hatched from each clutch as a binomial response

variable (logit link function), and maternal temperature treatment and offspring temperature

treatment as fixed factors. We included the interaction term between the factors into the model.

Less than three snails hatched from two clutches that were both produced and maintained at 25°C.

We excluded these individuals from all the further analyses as they would not provide sufficient

replication within those clutches.

After that, we calculated the developmental time until hatching for each egg as the

difference between the date the clutch was oviposited and the hatching date. We then analysed

the effects of temperature on developmental time using a multivariate analysis of variance

(MANOVA, with Pillai's trace test statistic for unequal sample sizes). We used the onset of

hatching (i.e. the first hatching day; square-root transformed to homogenize error variance),

median developmental time (we used the median rather than the mean as the distribution of

hatching time within the clutches was skewed), and the end of hatching (i.e. the last hatching day;

ln transformed to homogenize error variance) for each clutch as response variables. We used

maternal temperature treatment and offspring temperature treatment as fixed factors in the

analysis, and included the interaction term between them into the model. Since the MANOVA

indicated effects of temperature on developmental time (see the results section), we conducted

separate ANOVAs using a similar model as above for the different parameters of developmental

time to investigate whether their responses to temperature were different.

We analysed the variation in the survival of hatched offspring during the experiment

using a generalized linear mixed-effects model with the status of snails (survived, died) as a

binomial response variable (logit link function). We used maternal temperature treatment and

offspring temperature treatment as fixed factors, and egg clutch each offspring originated from as

a random factor (nested within the interaction between maternal temperature and offspring

temperature). We included the interaction term between maternal and offspring temperature into

the model. From the offspring that survived until the end of the experiment, we analysed the

variation in size using a mixed-model ANOVA with shell length (square-root transformed to

homogenize error variance) as a response variable, and a similar model as for survival. Survival

and/or size could not be measured from 30 juvenile snails (3.5% of all individuals) because of

human errors. We excluded these snails from the data. The assumptions of all the above analyses

were fulfilled, and we performed them using IBM SPSS Statistics Version 23.0 software

(Armonk, NY: IBM Corp.).

Results

During the initial exposure of adult snails to different temperature treatments (i.e. a seven-day

period before the collection of egg clutches started), the survival of snails exposed to 25°C was

reduced (estimated marginal mean $\pm$ SE: 66.1 $\pm$ 6.2%) compared with snails exposed to 15°C

(estimated marginal mean $\pm$ SE: 94.4 $\pm$ 3.1%; generalized linear model: Wald Chi-Square =

10.940, d.f. = 1, p = 0.001). In those snails that survived, the probability of reproducing was

higher at 25°C (estimated marginal mean $\pm$ SE: 87.2 $\pm$ 5.4%) than at 15°C (estimated marginal

mean $\pm$ SE: 49.0 $\pm$ 7.0%; generalized linear model: Wald Chi-Square = 12.429, d.f. = 1, p <

0.001). Note that the same individuals that reproduced during this initial exposure period also

produced the eggs that were exposed to offspring temperature treatments.

Eggs oviposited by snails at 25°C were smaller than those oviposited at 15°C (two-

dimensional area; estimated marginal mean ± SE: 25°C: 1.11 ± 0.01 mm^2; 15°C: 1.30 ± 0.01

mm^2; ANOVA: $F_{1,57}$ = 26.275, p < 0.001). Hatching success of eggs was affected by both the

maternal temperature treatment and the offspring temperature treatment, and these effects were

independent of each other (Table 1, Fig. 2). High maternal temperature increased hatching

success by 9.0% whereas high offspring temperature reduced it by 7.5% (Fig. 2). Maternal and

offspring temperature also affected the developmental time of eggs, and these effects were

independent of each other (MANOVA, maternal temperature treatment: Pillai's trace = 0.310,

$F_{3,51}$ = 7.654, p < 0.001; offspring temperature treatment: Pillai's trace = 0.918, $F_{3,51}$ = 191.413, p

< 0.001; maternal temperature treatment × offspring temperature treatment: Pillai's trace = 0.046,

$F_{3,51}$ = 0.827, p = 0.485). The effects of temperature on developmental time were first because

offspring started to hatch 12.3% earlier when mothers had been exposed to 25°C (ANOVA: $F_{1,53}$

= 15.806, p < 0.001; Fig. 3a). Second, the onset, median, and the end of hatching were earlier

when offspring were maintained at 25°C (first day of hatching: 55.1% reduction; ANOVA: $F_{1,53}$

= 571.961, p < 0.001; median developmental time: 43.8% reduction; ANOVA: $F_{1,53}$ = 189.817, p

< 0.001; last day of hatching: 43.8% reduction; ANOVA: $F_{1,53}$ = 62.002, p < 0.001; Fig. 3).

Table 1. Generalized linear model for the hatching success of *L. stagnalis* eggs (proportion of

eggs that hatched per clutch) by maternal temperature treatment (15°C, 25°C), offspring

temperature treatment (15°C, 25°C), and their interaction.

Effect	d.f.	Wald-Chi-Square	p
Maternal temperature treatment (M)	1	29.992	< 0.001
Offspring temperature treatment (O)	1	24.833	< 0.001
M × O	1	0.377	0.539

Figure 2. Hatching success of eggs [proportion of eggs that hatched (%; mean ± SE)] for egg

clutches produced at different maternal temperature treatments (15°C, 25°C) and maintained at

15°C (black circles) or at 25°C (white circles).

Figure 3. Developmental time of eggs presented using the mean $\pm$ SE of (a) onset of hatching,

(b) median developmental time, and (c) end of hatching for egg clutches produced at different

maternal temperature treatments (15°C, 25°C) and maintained at 15°C (black circles) or at 25°C

(white circles).

High maternal temperature reduced offspring survival by 17.8%, and this effect was

independent of the temperature offspring were exposed to (Table 2, Fig. 4). When offspring were

maintained at 25°C, they showed tendency towards lower survival (Table 2, Fig. 4). Temperature

affected the size of offspring so that high maternal temperature and low offspring temperature

reduced shell length at the end of the experiment (Table 3, Fig. 4).

Table 2. Generalized linear mixed-effects model for the survival of juvenile *L. stagnalis* snails

during the experiment (survived/died) by maternal temperature treatment (15°C, 25°C), offspring

temperature treatment (15°C, 25°C), their interaction, and the egg clutch each individual

originated from.

Effect	d.f.	F	Estimate	SD	Z	p
Maternal temperature treatment (M)	1	4.452				0.035
Offspring temperature treatment (O)	1	3.588				0.059
M $\times$ O	1	1.587				0.208
Egg clutch (M $\times$ O)			0.582	0.184	3.165	0.002

Figure 4. (a) Proportion (%; estimated marginal mean ± SE) of offspring that survived until the

end of the experiment, and (b) shell length (mm; estimated marginal mean ± SE) of those snails

at the end of the experiment when produced at different maternal temperature treatments (15°C,

25°C) and maintained at 15°C (black circles) or at 25°C (white circles).

Table 3. Mixed-model analysis of variance for the shell length of juvenile *L. stagnalis* snails at the end of the experiment by maternal temperature treatment (15°C, 25°C), offspring temperature treatment (15°C, 25°C), their interaction, and the clutch each individual originated from.

Effect	d.f.	MS	F	p
Maternal temperature treatment (M)	1	3.926	15.012[a]	< 0.001
Offspring temperature treatment (O)	1	54.520	208.927[a]	< 0.001
M × O	1	0.216	0.827[a]	0.367
Egg clutch (M × O)	53	0.302	3.370	< 0.001
Error	455	0.090		

[a] Egg clutch (M × O) as the error term

Discussion

Exposure to an experimental heat wave affected eggs and hatchlings of *L. stagnalis* snails both directly and via transgenerational maternal effects. Direct effects of high temperature were both negative and positive by reducing the hatching success of eggs, but by increasing their development rate and the subsequent growth of those individuals that hatched. The magnitude and the direction of transgenerational effects when compared with direct effects varied among examined traits. In general, high maternal temperature benefitted offspring at very early life stages (hatching success and the onset of hatching), but reduced performance at later stages (survival and final size of hatched snails). Hence, the potential adaptive value of responding to high temperature by maternal effects may be limited only to very early life stages. Interestingly, the magnitude of transgenerational effects on hatching success and survival of offspring were similar to the direct effects of high temperature, although in the case of hatching success the direction of the effects differed. These findings indicate that heat waves can not only impact

natural populations through transgenerational effects, but that the magnitude of those effects can

be equally strong to the direct within generation effects of high temperature. The relative strength

of the direct and transgenerational effects, however, depends on the considered trait.

It is important to note that because all parental snails that we initially exposed to different

temperatures did not survive or reproduce, potential differences between the temperature

treatments in offspring generation could arise from selection in the parental population. This

could be, for instance, if weak snails could not survive under environmental stress. Similarly, not

all offspring hatched or they died before the end of the experiment. While we cannot exclude

neither estimate the potential role of selection in our experiment, it seems unlikely that selection

could explain our results on transgenerational effects. This is because the longest lasting effects

of high maternal temperature on offspring (i.e. hatchling survival and size at the end of the

experiment) were negative. If only high quality individuals were able to survive and reproduce at

high temperature this should lead to the opposite result. Because high temperature had a direct

negative effect on the hatching of juvenile snails, their increased size at the end of the experiment

could be not only due to phenotypic plasticity but also because of selection.

Effects of temperature on eggs

Mothers exposed to high temperature produced smaller eggs that had higher hatching success

compared with snails that oviposited at benign temperature (14.6% reduction in size, 9.0%

increase in hatching success). Thus, exposure of the maternal generation to high temperature can

increase the quality of produced eggs despite their smaller size. High maternal temperature made

the onset of hatching earlier by 2.5 days (12.3% reduction). The magnitude of the direct effect of

high temperature on hatching success of eggs was similar to the observed transgenerational

effect, but negative (7.5% reduction). The direct effect of high temperature on development of

eggs was strong by shortening the median developmental time by approximately two weeks

(43.8% reductions). Faster development and earlier hatching could reduce the risk of eggs being

exposed to natural enemies such as predators (see Warkentin 1995; Chivers *et al.* 2001), and it is

suggested to increase future survival of offspring (Arcese & Smith 1985; Warner & Shine 2007)

as well as allow earlier maturation and increased reproductive output (Uller & Olsson 2010).

However, early hatching may also bring disadvantages by leading to less well developed

offspring (e.g. Warkentin 1999; Buckley, Michael & Irschick 2005).

Large egg size is often beneficial for offspring by increasing their fitness (Hutchings

1991; Fox 1994; Krist 2011). This is likely to be because large eggs can provide more energy and

nutrients for developing embryos (reviewed in Williams 1994). In aquatic organisms, however,

reduced egg size may be beneficial by ensuring oxygen supply to embryos when oxygen

concentration in water decreases under high temperature (Woods 1999; Moran & Woods 2007).

Therefore, egg size might not be a good indicator of egg quality. The observed transgenerational

effect of high temperature on development of eggs (developmental time and hatching success)

may be due to two non-exclusive mechanisms. First, high temperature can increase metabolic rate

of adults that allow production of high quality eggs (see Jann & Ward 1999; Saino *et al.* 2004).

Second, it may increase resource allocation towards reproduction rather than other traits (e.g.

growth) when the residual reproductive value of individuals decrease, for example, due to

increased mortality (Fisher 1930; Williams 1966). The effect of maternal temperature on

developmental time of eggs was, however, limited to the onset of hatching. This indicates that the

possibly increased investment on oviposited eggs may be rapidly depleted. On the other hand,

smaller eggs are expected to develop more slowly (Levitan 2000). Therefore, it is possible that

high maternal investment and small egg size overrode each other's effects so that no net change

in developmental time of eggs could be detected. The direct effect of temperature on hatching

success of eggs is likely to be due to high sensitivity of mollusc embryos to high temperature

(Vaughn 1953), which leads to mortality in several taxa probably due to denaturation of proteins

(reviewed in Pepin 1991; Noble, Stenhouse & Schwanz 2018). The direct effect of temperature

on developmental time of eggs was possibly because temperature determines the speed of

biochemical processes of the developing embryos (e.g. García-Guerrero, Villarreal & Racotta

2003; Sibert, Ouellet & Brêthes 2004).

Effects of temperature on hatchlings

Exposure of mothers to high temperature reduced the probability of offspring to survive until the

age of five weeks (17.8% reduction compared with 15°C). High maternal temperature also

reduced the size offspring reached (28.9% reduction). High temperature showed tendency to have

a negative direct effect on offspring survival (16.1% reduction). However, offspring that survived

grew larger at high temperature (97.4% increase in size). Hence, high maternal temperature

reduced offspring performance in the examined traits while the direct effects of high temperature

were both positive and negative. Reduced size of offspring due to high maternal temperature may

lead to delayed maturity as well as reduced mating success and fecundity (reviewed in Clutton-

Brock 1988), and also increase susceptibility to predators (e.g. Janzen 1993; Craig *et al.* 2006).

Instead, direct effect of high temperature can benefit those individuals that are able to survive

under such conditions.

Reduced offspring size when mothers experience high temperature is found across a wide

range of animal taxa (reviewed in Atkinson *et al.* 2001). The reason for this is not yet clear and

may either be an adaptation to maximize mother's life time fitness (Yampolsky & Scheiner 1996)

or due to physiological constraints under such conditions (Blanckenhorn 2000). The direct effect

of high temperature on offspring survival may be due to temperature-induced changes in, for

instance, protein structures and/or membrane fluidity (reviewed in Pörtner, Lucassen & Storch

2005), which can lead to body malfunctions and increased mortality. On the other hand, high

temperature fastens metabolic rate and can increase growth of organisms (e.g. Iguchi & Ikeda

2005; Salo *et al.* 2017). This, however, could also lead to faster use of energetic reserves that may

reduce survival.

General conclusions

Our finding that the potential adaptive value of responding to high temperature by maternal

effects was limited to very early life stages (i.e. eggs) is in line with earlier research (Mousseau *et*

al. 1991; Heath *et al.* 1999; Pettay *et al.* 2008). Instead, the result of equally strong direct and

transgenerational effects of exposure to high temperature on some of the examined traits

contradict earlier studies that have examined their relative importance in determining offspring

performance and physiology (e.g. Groeters & Dingle 1988; Huey *et al.* 1995; Steigenga &

Fischer 2007; Burgess *et al.* 2011; Salinas *et al.* 2012; Shama *et al.* 2014). In those studies,

transgenerational effects of temperature are typically reported to be weak compared with its

direct effects. To our knowledge, transgenerational effects of environmental change have been

found to be strong compared with its direct effects in a climate change context only in

germination probability of a perennial forb, *Boechera stricta*, when wintering conditions are

manipulated (Wadgymar *et al.* 2018). Together with that finding, our results indicate that climate

change-mediated environmental changes can affect natural populations through transgenerational

effects, and that these effects may be as strong as or even stronger than the direct effects of

environmental change.

Despite of high interest on transgenerational effects of climate change-mediated

environmental change on organisms (Donelson *et al.* 2018) many studies have not tested their

relative importance compared with direct within generation effects. This is because earlier studies

have focused, for example, on testing whether negative effects of environmental change are

reduced if parents have experienced the same environmental conditions (e.g. Donelson *et al.*

2012; Miller *et al.* 2012). Testing this does not necessarily require a full-factorial design that is

needed for examining the relative importance of direct and transgenerational effects. The

alternative approach is relevant in systems where environment changes gradually and relatively

slowly to a predicted direction (e.g. in oceans). Such studies would, however, not be realistic in

terrestrial and freshwater systems that experience high and rapid fluctuations in several

environmental conditions owing to extreme weather events such as heat waves. In marine

species, exposing parents to altered environmental conditions has been found to reduce the

negative effects of increased temperature and CO_2-level on offspring (Donelson *et al.* 2012;

Miller *et al.* 2012; Shama *et al.* 2014). In our study, none of the observed direct effects of

temperature depended on the maternal environment, which was indicated by the lack of

interactive effects between temperature treatments. This may be due to higher unpredictability of

extreme weather events in freshwater systems compared with marine environments that could

limit the ability of such adaptive maternal effects to evolve (see Mousseau & Fox 1998a).

Acknowledgements

We thank K. Räsänen, J. Jokela and the reviewers of PCI Ecology for constructive comments on

the manuscript. The study was funded by the Biological Interactions Doctoral Program (BIOINT)

to KL, and the Emil Aaltonen Foundation and Swiss National Science Foundation (grant 31003A

140876 and 31003A 169531) to OS.

Conflict of interest disclosure

The authors of this preprint declare that they have no financial conflict of interest with the

content of this article.

References

Agrawal, A.A. (2002) Herbivory and maternal effects: mechanisms and consequences of

transgenerational induced plant resistance. *Ecology,* **83,** 3408-3415.

Arcese, P. & Smith, J.N.M. (1985) Phenotypic correlates and ecological consequences of

dominance in song sparrows. *Journal of Animal Ecology,* **54,** 817-830.

Atkinson, D., Morley, S.A., Weetman, D. & Hughes, R.N. (2001) Offspring size responses to

maternal temperature in ectotherms. *Environment and animal development: genes, life*

histories and plasticity (eds D. Atkinson & M. Thorndyke), pp. 269-285. BIOS Scientific,

Oxford.

Bernardo, J. (1996) Maternal effects in animal ecology. *American Zoologist,* **36,** 83-105.

Blanckenhorn, W.U. (2000) Temperature effects on egg size and their fitness consequences in the

yellow dung fly Scathophaga stercoraria. *Evolutionary Ecology,* **14,** 627-643.

Bruno, J.F., Selig, E.R., Casey, K.S., Page, C.A., Willis, B.L., Harvell, C.D., Sweatman, H. &

Melendy, A.M. (2007) Thermal stress and coral cover as drivers of coral disease

outbreaks. *Plos Biology,* **5,** 1220-1227.

Buckley, C.R., Michael, S.F. & Irschick, D.J. (2005) Early hatching decreases jumping

performance in a direct-developing frog, *Eleutherodactylus coqui*. *Functional Ecology,*

19, 67-72.

Burgess, S.C. & Marshall, D.J. (2011) Temperature-induced maternal effects and environmental

predictability. *Journal of Experimental Biology,* **214,** 2329-2336.

Cain, G.L. (1956) Studies on cross-fertilization and self-fertilization in *Lymnaea stagnalis*

appressa Say. *Biological Bulletin,* **111,** 45-52.

Chivers, D.P., Kiesecker, J.M., Marco, A., DeVito, J., Anderson, M.T. & Blaustein, A.R. (2001)

Predator-induced life history changes in amphibians: egg predation induces hatching.

Oikos, **92,** 135-142.

Clutton-Brock, T.H. (1988) *Reproductive success studies of individual variation in contrasting*

breeding systems. University of Chicago Press, Chicago.

Craig, J.K., Burke, B.J., Crowder, L.B. & Rice, J.A. (2006) Prey growth and size-dependent

predation in juvenile estuarine fishes: experimental and model analyses. *Ecology,* **87,**

2366-2377.

Donelson, J.M., Munday, P.L., McCormick, M.I. & Pitcher, C.R. (2012) Rapid transgenerational

acclimation of a tropical reef fish to climate change. *Nature Climate Change,* **2,** 30-32.

Donelson, J.M., Salinas, S., Munday, P.L. & Shama, L.N.S. (2018) Transgenerational plasticity

and climate change experiments: where do we go from here? *Global Change Biology,* **24,**

13-34.

Easterling, D.R., Meehl, G.A., Parmesan, C., Changnon, S.A., Karl, T.R. & Mearns, L.O. (2000)

Climate extremes: observations, modeling, and impacts. *Science,* **289,** 2068-2074.

Fisher, R.A. (1930) *The genetical theory of natural selection*. Clarendon Press, Oxford.

Fox, C.W. (1994) The influence of egg size on offspring performance in the seed beetle,

Callosobruchus maculatus. Oikos, **71,** 321-325.

García-Guerrero, M., Villarreal, H. & Racotta, I.S. (2003) Effect of temperature on lipids,

proteins, and carbohydrates levels during development from egg extrusion to juvenile

stage of *Cherax quadricarinatus* (Decapoda : Parastacidae). *Comparative Biochemistry*

and Physiology a-Molecular & Integrative Physiology, **135,** 147-154.

Gillooly, J.F., Brown, J.H., West, G.B., Savage, V.M. & Charnov, E.L. (2001) Effects of size and

temperature on metabolic rate. *Science,* **293,** 2248-2251.

Groeters, F.R. & Dingle, H. (1988) Genetic and maternal influences on life-history plasticity in

milkweed bugs (*Oncopeltus*): response to temperature. *Journal of Evolutionary Biology,*

1, 317-333.

Groothuis, T.G.G. & Schwabl, H. (2008) Hormone-mediated maternal effects in birds:

mechanisms matter but what do we know of them? *Philosophical Transactions of the*

Royal Society B, **363,** 1647-1661.

Hance, T., van Baaren, J., Vernon, P. & Boivin, G. (2007) Impact of extreme temperatures on

parasitoids in a climate change perspective. *Annual Review of Entomology,* **52,** 107-126.

Heath, D.D., Fox, C.W. & Heath, J.W. (1999) Maternal effects on offspring size: variation

through early development of chinook salmon. *Evolution,* **53,** 1605-1611.

Huey, R.B., Wakefield, T., Crill, W.D. & Gilchrist, G.W. (1995) Within- and between-generation

effects of temperature on early fecundity of *Drosophila melanogaster*. *Heredity,* **74,** 216-

223.

Hutchings, J.A. (1991) Fitness consequences of variation in egg size and food abundance in

brook trout *Salvelinus fontinalis*. *Evolution,* **45,** 1162-1168.

Iguchi, N. & Ikeda, T. (2005) Effects of temperature on metabolism, growth and growth

efficiency of *Thysanoessa longipes* (Crustacea : Euphausiacea) in the Japan Sea. *Journal*

of Plankton Research, **27,** 1-10.

Jang, E.B. (1991) Thermal death kinetics and heat tolerance in early and late 3rd instars of the

oriental fruit fly (Diptera, Tephritidae). *Journal of Economic Entomology,* **84,** 1298-1303.

Janhunen, M., Piironen, J. & Peuhkuri, N. (2010) Parental effects on embryonic viability and

growth in Arctic charr *Salvelinus alpinus* at two incubation temperatures. *Journal of Fish*

Biology, **76,** 2558-2570.

Jann, P. & Ward, P.I. (1999) Maternal effects and their consequences for offspring fitness in the

Yellow Dung Fly. *Functional Ecology,* **13,** 51-58.

Janzen, F.J. (1993) An experimental analysis of natural selection on body size of hatchling

turtles. *Ecology,* **74,** 332-341.

Karl, T.R. & Trenberth, K.E. (2003) Modern global climate change. *Science,* **302,** 1719-1723.

Kirtman, B., Power, S.B., Adedoyin, J.A., Boer, G.J., Bojariu, R., Camilloni, I., Doblas-Reyes,

F.J., Fiore, A.M., Kimoto, M., Meehl, G.A., Prather, M., Sarr, A., Schär, C., Sutton, R.,

van Oldenborg, G.J., Vecchi, G. & Wang, H.J. (2013) Near-term climate change:

projections and predictability. *Climate change 2013: the physical science basis.*

Contribution of working group I to the fifth assessment report of the Intergovernmental

Panel on Climate Change. Summary for policy makers (eds T.F. Stocker, D. Qin, G.-K.

Plattner, M.M.B. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex & P.M.

Midgley), pp. 953-1028. Campridge University Press, Cambridge and New York.

Klockmann, M., Günter, F. & Fischer, K. (2017) Heat resistance throughout ontogeny: body size

constrains thermal tolerance. *Global Change Biology,* **23,** 686-696.

Krist, M. (2011) Egg size and offspring quality: a meta-analysis in birds. *Biological Reviews,* **86,**

692-716.

Leicht, K., Jokela, J. & Seppälä, O. (2013) An experimental heat wave changes immune defense

and life history traits in a freshwater snail. *Ecology and Evolution,* **3,** 4861-4871.

Leicht, K., Seppälä, K. & Seppälä, O. (2017) Potential for adaptation to climate change: family-

level variation in fitness-related traits and their responses to heat waves in a snail

population. *Bmc Evolutionary Biology,* **17,** 140.

Levitan, D.R. (2000) Optimal egg size in marine invertebrates: theory and phylogenetic analysis

of the critical relationship between egg size and development time in echinoids. *American*

Naturalist, **156,** 175-192.

Marshall, D.J. (2008) Transgenerational plasticity in the sea: context-dependent maternal effects

across the life history. *Ecology,* **89,** 418-427.

McCormick, M.I. (1999) Experimental test of the effect of maternal hormones on larval quality

of a coral reef fish. *Oecologia,* **118,** 412-422.

Meehl, G.A. & Tebaldi, C. (2004) More intense, more frequent, and longer lasting heat waves in

the 21st century. *Science,* **305,** 994-997.

Miller, G.M., Watson, S.-A., Donelson, J.M., McCormick, M.I. & Munday, P.L. (2012) Parental

environment mediates impacts of increased carbon dioxide on a coral reef fish. *Nature*

Climate Change, **2,** 858-861.

Mitchell, S.E. & Read, A.F. (2005) Poor maternal environment enhances offspring disease

resistance in an invertebrate. *Proceedings of the Royal Society B,* **272,** 2601-2607.

Moran, A.L. & Woods, H.A. (2007) Oxygen in egg masses: interactive effects of temperature,

age, and egg-mass morphology on oxygen supply to embryos. *Journal of Experimental*

Biology, **210,** 722-731.

Moret, Y. (2006) 'Trans-generational immune priming': specific enhancement of the

antimicrobial immune response in the mealworm beetle, *Tenebrio molitor*. *Proceedings of*

the Royal Society B, **273,** 1399-1405.

Mousseau, T.A. & Dingle, H. (1991) Maternal effects in insect life histories. *Annual Review of*

Entomology, **36,** 511-534.

Mousseau, T.A. & Fox, C.W. (1998a) The adaptive significance of maternal effects. *Trends in*

Ecology & Evolution, **13,** 403-407.

Mousseau, T.A. & Fox, C.W. (1998b) *Maternal effects as adaptations*. Oxford University Press,

New York.

Nakadera, Y., Blom, C. & Koene, J.M. (2014) Duration of sperm storage in the simultaneous

hermaphrodite *Lymnaea stagnalis*. *Journal of Molluscan Studies,* **80,** 1-7.

Nakadera, Y., Mariën, J., Van Straalen, N.M. & Koene, J.M. (2017) Multiple mating in natural

populations of a simultaneous hermaphrodite, *Lymnaea stagnalis*. *Journal of Molluscan*

Studies, **83,** 56-62.

Noble, D.W.A., Stenhouse, V. & Schwanz, L.E. (2018) Developmental temperatures and

phenotypic plasticity in reptiles: a systematic review and meta-analysis. *Biological*

Reviews, **93,** 72-97.

Parker, L.M., Ross, P.M., O'Connor, W.A., Borysko, L., Raftos, D.A. & Portner, H.-O. (2012)

Adult exposure influences offspring response to ocean acidification in oysters. *Global*

Change Biology, **18,** 82-92.

Parmesan, C. & Yohe, G. (2003) A globally coherent fingerprint of climate change impacts

across natural systems. *Nature,* **421,** 37-42.

Pepin, P. (1991) Effect of temperature and size on development, mortality, and survival rates of

the pelagic early life history stages of marine fish. *Canadian Journal of Fisheries and*

Aquatic Sciences, **48,** 503-518.

620 Person-Le Ruyet, J., Mahé, K., Le Bayon, N. & Le Delliou, H. (2004) Effects of temperature on

621   growth and metabolism in a Mediterranean population of European sea bass,

622   *Dicentrarchus labrax. Aquaculture,* **237,** 269-280.

623 Pettay, J.E., Charmantier, A., Wilson, A.J. & Lummaa, V. (2008) Age-specific genetic and

624   maternal effects in fecundity of preindustrial finnish women. *Evolution,* **62,** 2297-2304.

625 Pörtner, H.O., Lucassen, M. & Storch, D. (2005) Metabolic biochemistry: its role in thermal

626   tolerance and in the capacities of physiological and ecological function. *The Physiology of*

627   *Polar Fishes (Fish Physiology 21)* (eds J.F. Steffensen, A.P. Farrell, W.S. Hoar & D.R.

628   Randall). Elsevier Academic Press, San Diego.

629 Puurtinen, M., Knott, K.E., Suonpää, S., Nissinen, K. & Kaitala, V. (2007) Predominance of

630   outcrossing in *Lymnaea stagnalis* despite low apparent fitness costs of self-fertilization.

631   *Journal of Evolutionary Biology,* **20,** 901-912.

632 Saino, N., Romano, M., Ambrosini, R., Ferrari, R.P. & Møller, A.P. (2004) Timing of

633   reproduction and egg quality covary with temperature in the insectivorous Barn Swallow,

634   *Hirundo rustica. Functional Ecology,* **18,** 50-57.

635 Salinas, S. & Munch, S.B. (2012) Thermal legacies: transgenerational effects of temperature on

636   growth in a vertebrate. *Ecology Letters,* **15,** 159-163.

637 Salo, T., Stamm, C., Burdon, F.J., Räsänen, K. & Seppälä, O. (2017) Resilience to heat waves in

638   the aquatic snail *Lymnaea stagnalis*: additive and interactive effects with micropollutants.

639   *Freshwater Biology,* **62,** 1831-1846.

640 Seppälä, O. & Jokela, J. (2011) Immune defence under extreme ambient temperature. *Biology*

641   *Letters,* **7,** 119-122.

Seppälä, O. & Leicht, K. (2013) Activation of the immune defence of the freshwater snail

Lymnaea stagnalis by different immune elicitors. *Journal of Experimental Biology,* **216,**

2902-2907.

Shama, L.N.S., Strobel, A., Mark, F.C. & Wegner, K.M. (2014) Transgenerational plasticity in

marine sticklebacks: maternal effects mediate impacts of a warming ocean. *Functional*

Ecology, **28,** 1482-1493.

Sibert, V., Ouellet, P. & Brêthes, J.C. (2004) Changes in yolk total proteins and lipid components

and embryonic growth rates during lobster (*Homarus americanus*) egg development

under a simulated seasonal temperature cycle. *Marine Biology,* **144,** 1075-1086.

Silbermann, R. & Tatar, M. (2000) Reproductive costs of heat shock protein in transgenic

Drosophila melanogaster. Evolution, **54,** 2038-2045.

Steer, M.A., Moltschaniwskyj, M.A., Nichols, D.S. & Miller, M. (2004) The role of temperature

and maternal ration in embryo survival: using the dumpling squid *Euprymna tasmanica* as

a model. *Journal of Experimental Marine Biology and Ecology,* **307,** 73-89.

Steigenga, M.J. & Fischer, K. (2007) Within- and between-generation effects of temperature on

life-history traits in a butterfly. *Journal of Thermal Biology,* **32,** 396-405.

Tessier, A.J., Henry, L.L., Goulden, C.E. & Durand, M.W. (1983) Starvation in *Daphnia*: energy

reserves and reproductive allocation. *Limnology and Oceanography,* **28,** 667-676.

Uller, T. & Olsson, M. (2010) Offspring size and timing of hatching determine survival and

reproductive output in a lizard. *Oecologia,* **162,** 663-671.

Vaughn, C.M. (1953) Effects of temperature on hatching and growth of *Lymnaea stagnalis*

appressa Say. *American Midland Naturalist,* **49,** 214-228.

Wadgymar, S.M., Mactavish, R.M. & Anderson, J.T. (2018) Transgenerational and within-

generation plasticity in response to climate change: insights from a manipulative field

experiment across an elevational gradient. *American Naturalist,* **in press**.

Walther, G.-R. (2010) Community and ecosystem responses to recent climate change.

Philosophical Transactions of the Royal Society B, **365,** 2019-2024.

Walther, G.-R., Post, E., Convey, P., Menzel, A., Parmesan, C., Beebee, T.J.C., Fromentin, J.M.,

Hoegh-Guldberg, O. & Bairlein, F. (2002) Ecological responses to recent climate change.

Nature, **416,** 389-395.

Warkentin, K.M. (1995) Adaptive plasticity in hatching age: a response to predation risk trade-

offs. *Proceedings of the National Academy of Sciences of the United States of America,*

92, 3507-3510.

Warkentin, K.M. (1999) Effects of hatching age on development and hatchling morphology in

the red-eyed treefrog, *Agalychnis callidryas. Biological Journal of the Linnean Society,*

68, 443-470.

Warner, D.A. & Shine, R. (2007) Fitness of juvenile lizards depends on seasonal timing of

hatching, not offspring body size. *Oecologia,* **154,** 65-73.

Williams, G.C. (1966) Natural selection costs of reproduction and a refinement of Lack's

principle. *American Naturalist,* **100,** 687-690.

Williams, T.D. (1994) Intraspecific variation in egg size and egg composition in birds: effects on

offspring fitness. *Biological Reviews of the Cambridge Philosophical Society,* **69,** 35-59.

Woods, H.A. (1999) Egg-mass size and cell size: effects of temperature on oxygen distribution.

American Zoologist, **39,** 244-252.

Yampolsky, L.Y. & Scheiner, S.M. (1996) Why larger offspring at lower temperatures? A

demographic approach. *American Naturalist,* **147,** 86-100.

Zhang, W., Chang, X.-Q., Hoffmann, A.A., Zhang, S. & Ma, C.-S. (2015) Impact of hot events at

different developmental stages of a moth: the closer to adult stage, the less reproductive

output. *Scientific Reports,* **5,** 10436.

Zuo, W.Y., Moses, M.E., West, G.B., Hou, C. & Brown, J.H. (2012) A general model for effects

of temperature on ectotherm ontogenetic growth and development. *Proceedings of the*

Royal Society B, **279,** 1840-1846.