Role of the *Campylobacter jejuni cheVAWY* chemotaxis genes in chemotactic

motility and biofilm formation

Mark Reuter[1], Eveline Ultee[1], Yasmin Toseafa[1], Andrew Tan[1], and Arnoud H.M. van Vliet[1,2,*]

1.   Quadram Institute Bioscience, Gut Health and Food Safety Programme, Norwich Research

Park, Norwich, United Kingdom.

2.   University of Surrey, Department of Pathology and Infectious Diseases, School of Veterinary

Medicine, Faculty of Health and Medical Sciences, Guildford, United Kingdom

* Address correspondence to Dr Arnoud van Vliet (a.vanvliet@surrey.ac.uk)

Running title: Chemotaxis and biofilm formation in *C. jejuni*

Keywords: Campylobacter, chemotaxis, biofilm formation, flagellar motility

Total number of words in Summary: 238

Total number of words in Manuscript text: 5,152

Number of Figures: 6, Number of Tables: 1

Supplementary information: 6 Figures, 2 Tables, 2 Movies (via Figshare)

Abbreviations: MCPs, methyl-accepting chemotaxis proteins; CheA-Y, receiver domain of CheA;

M-MuLV, Moloney Murine Leukemia Virus; BAB, Blood Agar Base; PMT, photomultiplier tube.

ABSTRACT

Chemotaxis is the ability of motile bacteria to coordinate swimming behaviour to navigate a

changeable environment, by sensing environmental conditions through receptors and transducing

these signals to the flagellar motor. In the foodborne bacterial pathogen *Campylobacter jejuni*,

flagellar motility and chemotaxis are required for intestinal colonisation and virulence. Here we

present a systematic characterisation of the CheVAWY chemotaxis system of *C. jejuni* and its role

in chemotactic motility and biofilm formation. Inactivation of the core chemotaxis genes (*cheA*,

cheY, *cheV* or *cheW*) impaired chemotactic motility but not flagellar assembly. Inactivation of *cheV*

or *cheW* or presence of two copies of *cheV* or *cheW* resulted in reduced chemotaxis. A *cheY* mutant

swam in clockwise loops and this behaviour was complemented by wildtype *cheY* or a *cheY* gene

lacking the D53 conserved phosphorylation site, but not by *cheY* lacking the D7 metal-binding site.

Deletion of the CheY-like received domain from the *cheA* gene did not impair chemotactic motility,

nor could this domain complement a *cheY* mutant. The *cj0350* gene was identified as encoding a

putative CheX phosphatase, and the presence of two copies of *cj0350* resulted in reduced

chemotactic motility. Finally, inactivation of any of the core chemotaxis genes interfered with the

ability to form a discrete biofilm at an air-media interface. This work shows that interference with

the *Campylobacter* chemotaxis system at any level disrupts optimal chemotactic motility, and also

affects transmission modes such as biofilm formation and dispersal.

INTRODUCTION

Campylobacter jejuni is an important causative agent of bacterial gastroenteritis in humans [1],

and is commonly transmitted via contaminated food, especially poultry meat [2]. Infection with *C.*

jejuni is also associated with neurodegenerative diseases like Miller-Fisher and Guillain-Barre

syndrome [3]. One of the key factors for infection by *C. jejuni* is its flagella-based motility, as

aflagellated *C. jejuni* are unable to cause disease in animal models and show strongly reduced host

cell invasion in cell culture-based assays [4-6]. The flagella are also involved in biofilm formation

[7-9], and are targeted by bacteriophages [10].

Bacteria are able to adapt to changeable environments using a number of different strategies.

Over time, genome plasticity and evolution allows for long-term adaptation to a particular niche.

More immediate stresses are sensed by transcriptional regulators, resulting in changes in gene

expression. By far the most rapid means of evading stress by motile bacteria is by changing

swimming behaviour to escape unfavourable conditions or seek more favourable conditions, using

chemotaxis to sense and move according to chemical gradients or intracellular signals [11]. The

core chemotaxis pathway consists of the CheY-CheA two-component signal transduction system

[12]. Stimulation of signal sensing methyl-accepting chemotaxis proteins (MCPs) results in

autophosphorylation of the histidine kinase CheA, which transfers the phosphate group to the

response regulator CheY. Phosphorylated CheY interacts with the flagellar switch to alter flagellar

motor rotation. Counter-clockwise rotation commonly results in straight running motility while

clockwise rotation results in a tumbling behaviour and directional change [13].

While the core CheA-CheY pathway is conserved in motile bacteria, the bacterial kingdom

contains several variations on this theme [14, 15]. A number of accessory proteins can be involved,

such as the CheW protein which is required for CheA-MCP interaction [16, 17], while a

methyltransferase (CheR) and a methylesterase (CheB) modify the MCP proteins to accommodate

adaptation [18]. Some organisms contain the CheW-like protein CheV, which may help integrate

signals from a specific class of sensors and may also act as a phosphate sink [19]. Phosphatases

such as CheZ or CheX can dephosphorylate CheY to reduce its effect on flagellar rotation [20, 21].

Bacteria may also have multiple non-redundant homologs of the chemotaxis proteins or contain

specific adaptations, adjuncts, and domain-fusions that add new functions to the core chemotaxis

system [22].

Compared to other foodborne pathogens such as *E. coli*, *Salmonella*, and *Listeria*, *C. jejuni* has

a greater number of MCPs involved in sensing amino acids, deoxycholate, dicarboxylic acid TCA

intermediates, formic acid, fucose, redox, iron, phosphate and energy status [23-29], plus a core

chemotaxis system consisting of CheVAWY proteins plus a CheR and a truncated CheB protein [4,

11], and the ChePQ regulatory system controlling the core chemotaxis genes *cheVAW* [30].

Chemotaxis-defective mutants have been shown to be attenuated in disease models [31-34], show

reduced immunopathology [35] and chick colonization [34], and were reported to be crucial for

infection in genome-wide transposon mutant screenings for virulence factors [36, 37].

In this manuscript we have used genetic inactivation, mutation and complementation to

systematically examine the *cheVAWY* genes encoding the core chemotaxis pathway in *C. jejuni*

(Fig. 1). We show that each signalling protein is necessary for chemotaxis in complex media, but

not for motility. Furthermore, we show that the metal-binding site encoded by *cheY* is required for

CheY function, but that the phosphorylation site is not required. Finally, we show that the receiver

domain of CheA (CheA-Y) is dispensable for CheA function, but cannot complement a *cheY*

mutant, and that chemotaxis is required for organized biofilm formation.

MATERIALS AND METHODS

C. jejuni **strains and growth conditions.**

Campylobacter jejuni strain NCTC 11168 and its isogenic mutants (Table 1) were routinely

cultured in a MACS-MG-1000 controlled atmosphere cabinet (Don Whitley Scientific) in

microaerobic conditions (85% N_2, 5% O_2, 10% CO_2) at 37°C. For growth on plates, strains were

either grown on Brucella agar or Blood Agar Base (BAB) with Skirrow supplement (10 μg ml^{-1}

vancomycin, 5 μg ml^{-1} trimethoprim, 2.5 IU polymyxin-B). Broth culture was carried out in

Brucella broth (Becton Dickinson).

Construction of insertional inactivation strains

Insertional inactivation mutants were made as described previously [26] using primers listed in

Table S1, resulting in plasmids listed in Table S2. Plasmids were propagated in *E. coli* strain

TOP10. To insert antibiotic resistance cassettes, *Bam*HI sites were introduced in the target genes by

inverse PCR (Table S1) and ligated to either the kanamycin cassette from pMARKan9 (*ΔcheY*,

ΔcheW, *ΔcheV*) or chloramphenicol cassette from pTopCat. All constructs were sequenced prior to

transformation (Eurofins Genomics, Ebersberg, Germany). Single *C. jejuni* mutant strains were

isolated after transformation of the *C. jejuni* NCTC 11168 wildtype strain with plasmids by

electroporation [26], followed by selection on plates supplemented with either 50 μg ml^{-1}

kanamycin or 10 μg ml^{-1} chloramphenicol. To confirm the position of the antibiotic cassette in

antibiotic resistant clones, genomic DNA was isolated from four ml of overnight culture (DNeasy

kit, QIAGEN). Diluted genomic DNA (50 ng) was used as template for PCR using primers that

anneal outside of the cloned flanking regions in combination with antibiotic cassette-specific

primers (Table S1).

Construction of complementation constructs and site directed mutagenesis

C. jejuni mutants were complemented by inserting the individual chemotaxis genes (*cheA*,

cheY, *cheW*, *cheV*, *cheAY*, *cj0350, fliN*) *in trans* using the *cj0046* pseudogene, as described

previously [26]. In these plasmids, the genes are expressed from the *fdxA* promoter. To make the

CheY^{D53A} substitution, pCASO59 was used as template for inverse PCR using primers

CheYDA01/02 (Table S1). To make the CheY^{D7A} substitution, pCASO59 was used as template for

inverse PCR using primers cheYD7AFwd/Rev (Table S1). To make the CheAΔReceiver domain

construct (CheA^{ΔRec}), pCASO58 was used as template for inverse PCR using primers

cheARec01/02, which changes the codon encoding lysine 648 to a stop codon (see Table S1). The

inverse PCR products were digested with *Dpn*I for 60 minutes at 37°C and then purified (PCR

purification kit, QIAGEN) and transformed into *E. coli*. To identify clones containing altered

sequences, plasmid DNA was purified and sequenced using appropriate primers (Eurofins

Genomics, Ebersberg, Germany).

RNA Extraction and RT-PCR

RNA was extracted using the hot phenol method [38]. RNA concentration was determined

using a Nanodrop 2000 (Thermo Fisher Scientific). A mix of Moloney Murine Leukemia Virus (M-

MuLV) Reverse Transcriptase and murine RNase Inhibitor with an optimized random primer mix

was used to make cDNA (NEB). Two μg of RNA was mixed with random primers (6 μM final

concentration) and water and incubated at 70°C for 5 minutes then on ice for 5 minutes. This mix

was added to 10 μl of M-MuLV reaction mix and 2 μl M-MuLV enzyme mix in a total volume of

20 μl. Control reactions were also prepared containing 2 μl water in place of enzyme to show any

amplification from contaminating DNA. Reactions were incubated at 25°C for 5 minutes, 42°C for

60 minutes, then 80°C for 4 minutes. Reactions were made to total volume of 50 μl by adding 30 μl

water. For PCR, 2 μl of cDNA was mixed with OneTaq HotStart master mix (1× final

concentration, NEB), and gene-specific primers (0.2 μM final concentration) (Table S1) in a total

volume of 25 μl. Extension reactions were carried out at 68°C.

Assessment of growth

A 50 µl single-use glycerol stock, routinely stored at -80°C, was used to inoculate a BAB plate

with Skirrow supplements and these cells were used to inoculate fresh Brucella broth. Cultures were

grown in microaerobic conditions with shaking overnight at 37°C. The overnight culture was

diluted to $A_{600} \approx 0.05$ ($\sim 1 \times 10^7$ CFU ml^{-1}) in 200 µl in a 96-well plate (flat bottom, non-treated,

sterile, polystyrene, (Corning, NY, USA)). Growth in 96-well plates was assessed using an Omega

plate reader (FLUOstar Omega (BMG Labtech, Germany)) linked to an atmospheric control unit in

microaerobic conditions at 37°C. Omega assays were run for 24 hours (double orbital shaking at

400 rpm) and A_{600} data was recorded every 60 minutes.

Light microscopy and flagella staining using the Ryu stain

Typically, 10 µl of an overnight culture was examined using an Eclipse 50i microscope ($\times$100

lens) to monitor swimming phenotypes. When necessary, a Coolpix 4500 digital camera (Nikon)

was used to capture video (15 frames second^{-1}, 320$\times$240 pixels). Video compilations were made by

extracting appropriate frames using ImageJ (Rasband, W.S., ImageJ, U. S. National Institutes of

Health, Bethesda, Maryland, USA, http://imagej.nih.gov/ij/, 1997-2014). To plot swimming

behaviour, individual cells were tracked using the Manual tracking plug-in for ImageJ (Fabrice

Cordeli, http://rsb.info.nih.gov/ij/plugins/track/track.html) then visualized using the Chemotaxis and

Migration Tool (Ibidi, http://ibidi.com/software/chemotaxis_and_migration_tool/). To visualise

flagella, cells were stained using the Ryu stain [39] as described previously [40]. ImageJ was used

to add a scale bar and prepare montage images.

Chemotaxis assays

Chemotactic motility was measured using soft agar motility assays and tube taxis assays [26].

All soft agar motility assays were carried out using Brucella soft agar in square 10 mm^2 petri plates

(Sterilin) inoculated with wildtype and three test strains, as described previously [26]. For each

plate, halo size was expressed as a percentage of the corresponding wildtype and each strain was

tested for significance using a one-sample t-test (alpha = 0.05), compared to a hypothetical value of

100 (GraphPad Prism 6.01). Strain-to-strain comparisons were made using a two-tailed Mann-

Whitney test.

Tube taxis assays were prepared as described previously using Brucella soft agar [26]. The

tubes were incubated at 37°C in air in a waterbath (Grant). Tubes were photographed after 24, 40,

48, 64, and 72 hours and the dye front was measured from the top of the agar using the ImageJ

software and expressed as a percentage of the wildtype strain. Each strain was tested for

significance using a one-sample t-test (alpha = 0.05) compared to a hypothetical value of 100

(GraphPad Prism 6.01) and strain-to-strain comparisons were made using a two-tailed Mann-

Whitney test.

Biofilm assays

A 50 µl single-use glycerol stock, routinely stored at -80°C, was used to inoculate a BAB plate

with Skirrow supplements and these cells were used to inoculate fresh Brucella broth. Cultures were

grown in microaerobic conditions with shaking overnight at 37°C. The overnight culture was

diluted to $A_{600} \approx 0.05$ ($\sim 1 \times 10^7$ CFU ml^{-1}) in 22 ml sterile Brucella in a 50 ml flacon tube. A twin-

frost microscope slide (sterilized in 70% ethanol) was inserted into each falcon tube. Tubes were

incubated at 37°C in microaerobic conditions without shaking. After 48 hours, slides were removed

from the tubes using flamed-sterilized tweezers and briefly washed in water. Slides were dried in air

before staining with 1% crystal violet. Unbound crystal violet was washed off with water and slides

were dried in air. Biofilms were imaged using a GenePixPro microarray scanner (Axon). The

photomultiplier tube (PMT) gain of either the 635 nm or 532 nm laser was adjusted to achieve a

balanced image. To assess biofilm shedding, the A_{600} of the planktonic cultures were measured;

statistically different results were determined using a two-tailed Mann-Whitney test (GraphPad

Prism 6.01).

Bioinformatic analysis.

Pfam (http://pfam.sanger.ac.uk/ and InterProScan (http://www.ebi.ac.uk/Tools/pfa/iprscan/)

were used to search for protein domains and architectures. All BLAST searches were conducted

within BioEdit. EMBOSS Water was used for pairwise protein alignments

(http://www.ebi.ac.uk/Tools/psa/emboss_water/). ClustalX was used for multiple sequence

alignment (http://www.clustal.org/clustal2/). DeepView version 4.1 was used to make the CheY

model (http://spdbv.vital-it.ch/).

RESULTS

Inactivation and complementation of the genes involved in the core chemotaxis system

To assess the roles of the *cheVAWY* genes in *C. jejuni* chemotaxis, each individual gene was

inactivated by a kanamycin or chloramphenical antibiotic resistance cassette in the same orientation

as the gene. To test for polar effects, each mutated gene was complemented by introduction of the

different chemotaxis genes mutant *in trans* in the *cj0046* pseudogene under control of the *fdxA*

promoter [26]. We also used this complementation system to introduce altered versions of the *cheY*

and *cheA* genes in *C. jejuni* mutants, and tested the effect of alteration of expression levels by

introducing a second copy of chemotaxis genes in wildtype *C. jejuni* NCTC11168 (Table 1). All the

mutant strains did not show significant changes in growth (Fig. S1) and expressed flagella at both

poles (Fig. 2). For all mutants generated here, chemotactic motility was assessed using soft agar

motility plates and tube taxis assays, which measures motility in both an energy and redox gradient

[26].

CheV is the dominant adaptor protein in the chemotaxis signalling pathway

In liquid media, motility of the *cheW* and *cheV* mutants was comparable to the wildtype, but

both halo formation and migration were reduced compared to the wildtype (Fig. 3, Fig. S2).

Chemotactic motility of the *cheW* mutant was reduced to less than 50% compared to the wildtype

strain at both 24 and 48 hours, respectively (Fig. 3, Fig. S2). The *cheV* mutant showed greater

reduction in halo formation and migration than the *cheW* mutant when compared to the wildtype

strain at both 24 and 48 hours (Fig. 3, Fig. S2). Complementation of the *cheW* mutant restored

chemotaxis phenotypes to that of the wildtype strain (Fig. 3, Fig. S2). However, complementation

of the *cheV* mutant did not restore chemotactic motility. The *cheV* mutant still expressed the

downstream *cheA* and *cheW* genes as shown by reverse transcriptase PCR (Fig. S3A), suggesting

the phenotype of the *cheV* mutant is not due to polar effects on the transcription of the downstream

cheA and *cheW* genes. Supplementation of the wildtype strain with a second copy of *cheV* (WT

+*cheV*) did not alter swimming in liquid media, but led to a significant reduction in chemotactic

motility (Fig. S4), suggesting that the stoichiometry of CheV and other chemotaxis proteins is of

importance for its function. Finally, complementation of the *cheW* mutant with the *cheV* gene did

not restore chemotactic motility (Fig. 3, Fig. S2).

Increased levels of CheY affect chemotactic motility

The *cheY* mutant showed a 'Catherine Wheel' swimming behaviour [41] in liquid media, where

a cell would appear to get trapped in a clockwise swimming loop (see Supplementary Movie 1 and

Fig. 4B) before resuming a darting motility as observed in the wildtype (Fig. 4A). The *cheY* mutant

displayed the previously described chemotaxis defect [32], with chemotactic motility reduced to

less than 20% of wildtype (Fig. 3, Fig. S2). Chemotactic motility was restored to wildtype levels by

complementation with the *cheY* gene, and even exceeded that observed in the wildtype strain (Fig.

3, Fig. S2), with the wildtype swimming behaviour also restored (Fig. 4C). A wildtype strain

containing an additional copy of *cheY* (WT + *cheY*) displayed increased chemotactic motility

compared to the wildtype strain (Fig. S4).

CheY requires the metal binding pocket but not the phosphorylation site for activity

An alignment of the seed sequences for a canonical Receiver Domain (Pfam, PF00072) with

the *C. jejuni* CheY sequence showed that both the active site aspartic acid residue (D53) and the N-

terminal 'acid-pocket' aspartic acid residues (D7, D8) were well conserved (Fig. S5A). A three-

dimensional model of *C. jejuni* CheY, based on the *Thermotoga maritima* CheY structure [42]

shows that the three aspartic acid residues are predicted to cluster to form the active site (Fig. S5

inset). These predictions were tested experimentally using site-directed mutagenesis of aspartate

residues 7 and 53 to alanine (D7A and D53A) of the *cheY* gene, to eliminate either phosphorylation

or metal binding, respectively, and the mutated versions were introduced in the *cheY* mutant. The

cheY mutant containing *cheY*[D53A] produced flagella (Fig. 2), and showed full restoration of wildtype

phenotype with swimming without the 'Catherine Wheel' phenotype (Fig. 4D), and restoration of

chemotactic motility to wildtype levels (Fig. 3, Fig. S2). In contrast, the *cheY* mutant containing

cheY[D7A] was even more defective in chemotactic motility (Fig. 3, Fig. S2), despite possessing

flagella (Fig. 2). The *cheY*[D7A] strain displayed swimming in very tight repeating spirals (Fig. 4E).

RT-PCR showed that all *cheY*-complemented strains made a *cheY* transcript that was absent in the

cheY mutant (Fig. S3D).

CheA is required for chemotaxis but the C-terminal receiver domain is not

Inactivation of the *cheA* gene strongly reduced chemotactic motility to levels similar to the

cheV and *cheY* mutants (Fig. 3, Fig. S2). However, inactivation of *cheA* did not disrupt transcription

of the downstream *cheW* gene, as determined with RT-PCR (Fig. S3B) Introduction of the *cheA*

gene *in trans* did not complement the *cheA* mutant (Fig. 3, Fig. S2), but introduction of a second

cheA copy in the wildtype strain resulted in significantly reduced chemotactic motility (Fig. S4).

The architecture of *C. jejuni* CheA differs to that seen in *E. coli* and *B. subtilis* (Fig. 1), as it has

a C-terminal receiver domain, CheA-Y, with 58% similarity (33% identity) to CheY (Fig. S5B).

The multiple alignment of CheY sequences including CheA-Y shows the presence of two aspartic

acid residues at the N-terminus and the active site aspartate, suggesting that the acid-pocket active

site is conserved (Fig. S5A). To assess the role of the CheA-Y domain in the function of CheA, the

cheA gene was mutated to contain a stop-codon at position 648 (*cheA*$^{\Delta Rec}$), truncating the CheA

protein at the start of the C-terminal receiver domain. Introduction of the *cheA*$^{\Delta Rec}$ gene in the

wildtype and *cheA* mutant strain resulted in transcription of a *cheA* transcript, which was absent in

the *cheA* mutant (Fig. S3E). The *cheA*$^{\Delta Rec}$ construct complemented the *cheA* mutant better than the

full length *cheA* gene, with restoration of about half of chemotactic motility by the *cheA*$^{\Delta Rec}$ version,

and no complementation by the wildtype *cheA* gene (Fig. 3, Fig. S2). When the *cheA*$^{\Delta Rec}$ gene was

introduced into the wildtype strain, this resulted in increased levels of chemotactic motility

compared to the wildtype strain (Fig. S4). We also investigated the ability of the CheA-Y receiver

domain alone to complement the *cheY* mutant, but this did not restore chemotactic motility (Fig. 3,

Fig. S2, Fig. S3), and the swimming behaviour still showed the characteristic catherine wheels seen

in the *cheY* mutant (Fig. 4). Thus, the sequence of the CheA-Y domain has diverged to the point

where it is functionally distinct from CheY in *C. jejuni*.

The *C. jejuni cj0350* gene encodes a putative CheX protein

A domain search of the NCTC11168 genome identified the *cj0350* gene as encoding a putative

CheX domain. CheX is an ancillary chemotaxis protein known to dephosphorylate CheY in

Borrelia burgdorferi [43] and *Thermotoga maritima* [44]. The *C. jejuni* CheX sequence is well

conserved in the genus *Campylobacter* (but absent from the genus *Helicobacter*), although

divergent from the canonical CheX sequences that comprise the Pfam seed alignment PF13690. We

constructed a *cj0350* mutant, which showed decreased chemotactic motility, although this

phenotype was not rescued by complementation with *cj0350* (Fig. 5). However, complementation

by the gene downstream of *cj0350*, *fliN*, did restore wildtype chemotactic motility (Fig. 5),

suggesting that the *cj0350* (*cheX*) phenotype is probably due to a disruption of *fliN* expression.

Introduction of a second copy of the *cj0350* gene in wildtype *C. jejuni* did not affect flagellar

assembly (Fig. 5D) or swimming in liquid media (see Supplementary Movie 2), but resulted in

reduced chemotactic motility (Fig. 5A-C). Given that phosphatase activity on CheY should result in

decreased chemotactic motility due to less stimulation of the flagella switch, this is consistent with

the possibility that Cj0350 is a divergent CheX ortholog.

Chemotaxis is required for organized biofilm formation at the air-media interface

We investigated the role of chemotaxis in biofilm formation, by developing a glass-slidebased

assay combined with crystal violet staining, with detection using a microarray scanner (employing

both lasers at 532 and 635 nm). The wildtype strain formed a biofilm on the glass slide at the air-

media interface (Fig. 6). Below the air-media interface, a less-intense ~2 mm band of adhered cells

was observed. All of the chemotaxis mutants used in this study were defective in formation of a

biofilm at the air-media interface (Fig. 6). In the *cheW* mutant, the air-media interface biofilm was

present, but a second more intense area of adhered cells was also visible ~5 mm below the air-

media interface. Both the *cheV* mutant and the complemented *cheV* mutant showed defective air-

media interface biofilm. The *cheA* and *cheY* mutants both showed a disorganized air-media

interface biofilm with a greater population of submerged cells. In all cases, except for the *cheV*

mutant and the *cheY^{D7A}*-complemented *cheY* mutant, complementation with the respective wildtype

gene restored the organised interface biofilm phenotype (Fig. 6). To assess the level of shedding of

cells from the biofilm, the A_{600} of planktonic fraction for each assay was recorded. Inactivation of

cheY resulted in fewer cells in the planktonic fraction compared to the wildtype strain, while the

complemented *cheY* mutant and the *cheY*^{D53A} complemented mutant both had more cells in the

planktonic fraction (Fig. S6A). Other *cheY* strains with reduced chemotactic motility also showed a

lower level of cells in the planktonic fraction. Therefore, the biofilm phenotype of at least the *cheY*

mutant could in part be due to a disruption of normal cell shedding/dispersal from the biofilm. To

test the hypothesis that *C. jejuni* may require the fully functional chemotactic motility to seek the

optimal environment for growth as a biofilm, the slide biofilm assay was performed using a

paralysed flagella (*pflA*) mutant [40, 45]. This mutant displayed low levels of biofilm formation at

the air-media interface (Fig. S6B). Thus, while the chemotaxis system is not required for biofilm

formation *per se*, it is necessary for organized biofilm formation at the air-media interface and may

play a role in dispersal/shedding of cells from the biofilm.

DISCUSSION

Flagellar motility plays an important role in colonisation and virulence of many bacterial

pathogens. While the ability to move to or from locations is an important contributor to the

infection process, such movement needs to be directed, and is commonly based on detection of

external (e.g. chemical gradient) and internal stimuli (e.g. metabolic state). These stimuli can be

positive or negative, with examples of positive stimuli being attractants such as nutrients, while

negative stimuli can be repellents such as bile acids. The chemotaxis system integrates the signals

from external and internal sensors through a signal transduction cascade consisting of MCPs,

CheW/CheV, CheA and CheY, while other factors such as CheB, CheR, CheC, CheZ, or CheX may

modulate and fine-tune the signal transduction cascade, with diverse combinations observed

throughout the bacterial kingdom [4, 46].

In this study we have dissected the core chemotaxis signalling pathway of the important

foodborne zoonotic pathogen *C. jejuni*, which is the most common bacterial cause of gastroenteritis

in the developed world. Directed motility is central to the process of intestinal colonisation by *C.*

jejuni, as mutants defective in motility or chemotaxis are either unable to colonise or show

significantly reduced efficiency of colonisation [32, 36]. Although *C. jejuni* shares many

components of the core chemotaxis pathway with model organisms such as *E. coli* and *B. subtilis*,

its chemotaxis system contains several additional components and modifications, and in this study

we have used a combination of gene inactivation and complementation experiments with

phenotypic assays to study the *C. jejuni* chemotaxis system.

Roles of *C. jejuni* CheW and CheV adaptor proteins in chemotaxis

Campylobacter is not exceptional in containing both CheV and CheW proteins. Although

CheW is the most common chemotaxis 'adaptor' protein (found in nearly 100% of genomes

containing the core chemotaxis components [47], CheV was found in almost 40% of chemotaxis-

positive genomes [47] and three genomes (*B. thuringiensis*, *B. weihenstephanensis*, and *L.*

monocytogenes) were found to contain CheV orthologs without CheW orthologs, suggesting that

CheV alone can function in the MCP-CheA complex [47]. In this study, the chemotaxis phenotype

of the *cheW* mutant is the least severe, suggesting that CheV does function to elicit some signal

transduction event. Similar effects were observed in *B. subtilis* [48] and further showed that the

CheW domain of CheV was sufficient for signal transduction. A direct protein-protein interaction

between CheV and Tlp4 (deoxycholate sensor, Cj0262c), Tlp6 (TCA intermediates sensor,

Cj0448c), and Tlp8 (redox sensor CetZ) has been documented [49] and Yeast Two- and Three-

Hybrid analysis and immunoprecipitation were used to show that Tlp1 interacts with both CheW

and CheV, although the interaction with CheV was much stronger and localised to the CheV-W

domain [50]. Both *cheV* and *cheW* mutants had chemotactic motility defects in Brucella media

although chemotaxis towards aspartate was still competent in either mutant. Similarly, the novel

chemotaxis protein Cj0371 of *C. jejuni* was shown to interact with the CheV protein, and influence

ATPase activity of CheA [51], suggesting that multiple systems engage in the chemotaxis pathway

to modulate signal transduction. An evolutionary genomics study showed that the presence of CheV

is correlated with increased numbers of MCPs and proposed that CheV functions to accommodate

signal transduction from a specific group of MCPs, exemplified by the *Salmonella* McpC chemo-

receptor [19]. None of the ten *C. jejuni* MCPs share the architecture of the McpC-type sensor.

Moreover, *H. pylori* genomes encode only four chemotaxis sensors and three CheV proteins [52].

The C-terminal receiver domain on CheV most likely refines the function of this protein. It has

previously been proposed that CheV acts as a phosphate sink, normalizing over-stimulation of

CheA [19, 53, 54]. The observation that expressing two copies of CheV in *C. jejuni* reduces

chemotactic motility (Fig. 3, Fig. S2) supports this theory, as increasing the level of the CheV

protein will divert phosphate flow from CheY to CheV. Interestingly, the chemotaxis phenotype of

the WT +*cheV* strain was more pronounced than that of the WT +*cheX* strain, which suggests that

the phosphate sink is more effective at dampening signal transduction than the possible CheX-

mediated dephosphorylation of CheY. Organisms lacking CheY-phosphatases may also utilize a

phosphate sink mechanism to modulate chemotaxis.

Role of CheY domains

The CheY protein is the final effector protein in the signal transduction cascade, directly

interacting with the flagellar switch to alter rotation between clockwise and counter-clockwise

rotation. Our observation that the *cheY*[D53A] variant could restore both chemotactic motility and

darting swimming in the *cheY* mutant (Fig. 3, Fig. S2) suggests that either a) unphosphorylated

CheY can interact with the flagella switch or b) the conserved aspartate 53 residue is not the only

site of activating phosphorylation. Sequence alignment of *C. jejuni* CheY with the seed receiver

domain alignment unambiguously identifies the putative site of phosphorylation and a structural

model clearly places Aspartate 53 in the acid pocket active site. Expression of CheY[D53A] from the

fdxA promoter may result in expression of the protein at increased levels. Indeed, expression of

wildtype CheY from the *fdxA* promoter significantly stimulated chemotactic motility (Fig. 3, Fig.

S2). Based on the CheY^{D53A} data, the chemotactic motility observed in the complemented *cheY*

mutant may be a mixture of CheY and CheY-P interaction at the flagella switch. If

unphosphorylated CheY can modify the flagella switch operation, the question of resetting the

chemotaxis system in *C. jejuni* becomes all the more intriguing. If one further considers the C-

terminal receiver domain of CheA as a further phosphate sink, the complete model of phosphate-

flow in *C. jejuni* chemotaxis becomes a complex story. The CheA-Y domain has all the hallmarks

of the classical receiver domain and appears to be dispensable for the core signal transduction

function. Moreover, CheA-Y is no substitute for CheY as the *cheA-Y* gene could not complement

the inactivation of *cheY* (Fig. 3, Fig. S2). Residues that are known to be important for binding the

flagella switch (T87, Y106, and K109 (*E. coli* numbering [55])) are conserved in CheA-Y and

amino acid differences between CheY and CheA-Y are spread across the entire sequence, and not

localized to the β4-α4 region, which shows the largest structural changes following phosphorylation

[56, 57]. Sequence divergence resulting in loss of chemotactic function is well documented: *R.*

sphaeroides che genes only partially restore taxis in *E. coli* [58]. Thus, the *C. jejuni* CheY/CheA-Y

proteins provide a good model for studying mutational drift resulting in loss or change of function.

C. jejuni CheA lacks the P2 domain found in *E. coli* and *B. subtilis* CheA (Fig. 1), which raises the

question of how it interacts with CheY. In *C jejuni*, CheA and CheY interact with each other [49].

However, since CheA^{ΔRec} restores signalling, the CheA-Y domain is not the likely interaction

partner for CheY.

Other components of the chemotaxis pathway

In this study we have identified the *cj0350* gene as a potential new chemotaxis gene in *C.*

jejuni. Although inactivation of the *cj0350* gene did affect chemotaxis, this could be explained by a

polar effect on the downstream *fliN* flagellar gene, and hence we cannot be certain about the role of

Cj0350 in chemotaxis yet. Motif analysis suggests the Cj0350 protein may be a novel variant of the

CheX phosphatase, representing a *Campylobacter*-specific chemotaxis adaptation in the

Epsilonproteobacteria, as no CheX ortholog could be detected in any available *Helicobacter*

genome. The primary sequence of the putative CheX protein identified in this study is divergent

from the canonical CheX sequence. Thus, in organisms where no current phosphatase has been

identified, cryptic phosphatase proteins may yet be discovered. In addition to CheX and CheZ, a

number of other CheY phosphatases have been characterized. *B. subtilis* uses CheC, CheD, and a

FliY/N fusion protein [59-61]. However, chemotaxis systems lacking any clear phosphatases are

also known (~20% of chemotaxis positive genomes [47]); for instance, *R. sphaeroides*,

Rhodospirillum centenum, and *Sinorhizobium meliloti* have complex chemotaxis systems but lack

known CheY-phosphatases [22, 58]. *H. pylori* further regulates CheZ activity by complex formation

with the novel chemotaxis protein ChePep [20, 62]. *Campylobacter jejuni* contains a ChePep

homolog (NCTC11168 - Cj1178c), which can complement an *H. pylori chepep* mutant [62]

although the role of this protein in chemotaxis in *C. jejuni* is currently not known.

Chemotaxis and biofilm formation

Previous studies have implicated chemotaxis sensors in biofilm formation. Tlp3 (Cj1564)

mutants showed increased biofilm formation [63], while CetZ (Tlp8) mutants showed decreased

biofilm formation [33]. We show that removing each component of the chemotaxis pathway results

in disruption of biofilm formation at a discrete air-surface interface, and that this phenotype can be

rescued by complementation for most of the chemotaxis genes investigated here. Thus, the

chemotaxis system likely has a role in coordinating biofilm formation, and this coordinating role

may extend to shedding and dispersal of cells from the biofilm. This study is therefore the first to

present a role for signal transduction in the active dispersal of cells from a *C. jejuni* biofilm. In

processing environments, biofilms present a reservoir of cells that can subsequently re-contaminate

the food chain highlighting the need to learn more about biofilm dispersal.

Conclusions

The ability to couple flagella rotation with environmental sensing is an effective adaptive

mechanism allowing bacteria to seek an optimum environment. While chemotaxis systems from

model organisms such as *E. coli* and *B. subtilis* provide an effective model for studying signal

transduction in non-paradigm organisms, species-specific modulations and augmentations abound

and require focussed investigation. Such genus- and species-specific analyses will be required to

better understand chemotaxis in other Epsilon-proteobacteria and other organisms that share

elements of the *Campylobacter* chemotaxis system and provides a further paradigm for chemotaxis

signal transduction.

ACKNOWLEDGMENTS

The authors wish to thank Dr Henri Tapp for assistance with statistical evaluations, and members of

the former IFR *Campylobacter* research group for helpful discussions. We would also like to thank

Rachael Stanley for electron microscopy and Maddy Houchen and Robert Hindmarsh for

microbiology media support.

FUNDING STATEMENT

The authors gratefully acknowledge the support of the Biotechnology and Biological Sciences

Research Council (BBSRC) via the Gut Health and Food Safety Institute Strategic Programme

(BB/J004529/1). The funder had no role in study design, data collection and analysis, decision to

publish, or preparation of the manuscript.

CONFLICTS OF INTEREST

The authors declare that there are no conflicts of interest.

REFERENCES

1. **Nichols GL, Richardson JF, Sheppard SK, Lane C, Sarran C**. Campylobacter epidemiology: a descriptive study reviewing 1 million cases in England and Wales between 1989 and 2011. *BMJ Open* 2012;2:e001179. doi: 10.1136/bmjopen-2012-001179

2. **Tam CC, Higgins CD, Neal KR, Rodrigues LC, Millership SE et al.** Chicken Consumption and Use of Acid-Suppressing Medications as Risk Factors for Campylobacter Enteritis, England. *Emerg Infect Dis* 2009;15:1402-1408. doi: 10.3201/eid1509.080773

3. **Poropatich KO, Walker CL, Black RE**. Quantifying the association between Campylobacter infection and Guillain-Barre syndrome: a systematic review. *J Health Popul Nutr* 2010;28:545-52. doi:

4. **Lertsethtakarn P, Ottemann KM, Hendrixson DR**. Motility and chemotaxis in Campylobacter and Helicobacter. *Annu Rev Microbiol* 2011;65:389-410. doi: 10.1146/annurev-micro-090110-102908

5. **Gao B, Lara-Tejero M, Lefebre M, Goodman AL, Galan JE**. Novel components of the flagellar system in epsilonproteobacteria. *MBio* 2014;5:e01349-14. doi: 10.1128/mBio.01349-14

6. **Asakura H, Hashii N, Uema M, Kawasaki N, Sugita-Konishi Y et al.** Campylobacter jejuni pdxA affects flagellum-mediated motility to alter host colonization. *PLoS One* 2013;8:e70418. doi: 10.1371/journal.pone.0070418

7. **Reuter M, Mallett A, Pearson BM, van Vliet AHM**. Biofilm formation by *Campylobacter jejuni* is increased under aerobic conditions. *Appl Environ Microbiol* 2010;76:2122-8. doi: 10.1128/AEM.01878-09

8. **Svensson SL, Pryjma M, Gaynor EC**. Flagella-mediated adhesion and extracellular DNA release contribute to biofilm formation and stress tolerance of Campylobacter jejuni. *PLoS One* 2014;9:e106063. doi: 10.1371/journal.pone.0106063

9. **Brown HL, Reuter M, Salt LJ, Cross KL, Betts RP et al.** Chicken juice enhances surface attachment and biofilm formation of Campylobacter jejuni. *Appl Environ Microbiol* 2014;80:7053-60. doi: 10.1128/AEM.02614-14

10. **Baldvinsson SB, Sorensen MC, Vegge CS, Clokie MR, Brondsted L**. Campylobacter jejuni motility is required for infection of the flagellotropic bacteriophage F341. *Appl Environ Microbiol* 2014;80:7096-106. doi: 10.1128/AEM.02057-14

11. **Korolik V**. The role of chemotaxis during Campylobacter jejuni colonisation and pathogenesis. *Curr Opin Microbiol* 2018;47:32-37. doi: 10.1016/j.mib.2018.11.001

12. **Hazelbauer GL, Lai WC**. Bacterial chemoreceptors: providing enhanced features to two-component signaling. *Curr Opin Microbiol* 2010;13:124-32. doi: 10.1016/j.mib.2009.12.014

13. **Patteson AE, Gopinath A, Goulian M, Arratia PE**. Running and tumbling with E. coli in polymeric solutions. *Sci Rep* 2015;5:15761. doi: 10.1038/srep15761

14. **Krell T, Lacal J, Munoz-Martinez F, Reyes-Darias JA, Cadirci BH et al.** Diversity at its best: bacterial taxis. *Environ Microbiol* 2011;13:1115-24. doi: 10.1111/j.1462-2920.2010.02383.x

15. **Cluzel P, Surette M, Leibler S**. An ultrasensitive bacterial motor revealed by monitoring signaling proteins in single cells. *Science* 2000;287:1652-5. doi:

16. **Martin AC, Wadhams GH, Armitage JP**. The roles of the multiple CheW and CheA homologues in chemotaxis and in chemoreceptor localization in *Rhodobacter sphaeroides*. *Mol Microbiol* 2001;40:1261-72. doi:

17. **Boukhvalova MS, Dahlquist FW, Stewart RC**. CheW binding interactions with CheA and Tar. Importance for chemotaxis signaling in *Escherichia coli*. *J Biol Chem* 2002;277:22251-9. doi: 10.1074/jbc.M110908200

18. **Parkinson JS, Hazelbauer GL, Falke JJ**. Signaling and sensory adaptation in *Escherichia coli* chemoreceptors: 2015 update. *Trends Microbiol* 2015;23:257-266. doi: 10.1016/j.tim.2015.03.003

19. **Ortega DR, Zhulin IB**. Evolutionary Genomics Suggests That CheV Is an Additional Adaptor for Accommodating Specific Chemoreceptors within the Chemotaxis Signaling Complex. *PLoS computational biology* 2016;12:e1004723. doi: 10.1371/journal.pcbi.1004723

20. **Lertsethtakarn P, Howitt MR, Castellon J, Amieva MR, Ottemann KM**. *Helicobacter pylori* CheZ(HP) and ChePep form a novel chemotaxis-regulatory complex distinct from the core chemotaxis signaling proteins and the flagellar motor. *Mol Microbiol* 2015;97:1063-78. doi: 10.1111/mmi.13086

21. **Muff TJ, Foster RM, Liu PJ, Ordal GW**. CheX in the three-phosphatase system of bacterial chemotaxis. *J Bacteriol* 2007;189:7007-13. doi: 10.1128/JB.00896-07

22. **Porter SL, Wadhams GH, Armitage JP**. Signal processing in complex chemotaxis pathways. *Nat Rev Microbiol* 2011;9:153-65. doi: 10.1038/nrmicro2505

23. **Day CJ, King RM, Shewell LK, Tram G, Najnin T et al.** A direct-sensing galactose chemoreceptor recently evolved in invasive strains of Campylobacter jejuni. *Nat Commun* 2016;7:13206. doi: 10.1038/ncomms13206

24. **Li Z, Lou H, Ojcius DM, Sun A, Sun D et al.** Methyl-accepting chemotaxis proteins 3 and 4 are responsible for *Campylobacter jejuni* chemotaxis and jejuna colonization in mice in response to sodium deoxycholate. *J Med Microbiol* 2014;63:343-54. doi: 10.1099/jmm.0.068023-0

25. **Vegge CS, Brondsted L, Li YP, Bang DD, Ingmer H**. Energy taxis drives *Campylobacter jejuni* toward the most favorable conditions for growth. *Appl Environ Microbiol* 2009;75:5308-14. doi: 10.1128/aem.00287-09

26. **Reuter M, van Vliet AHM**. Signal balancing by the CetABC and CetZ chemoreceptors controls energy taxis in *Campylobacter jejuni*. *PLoS One* 2013;8:e54390. doi: 10.1371/journal.pone.0054390

27. **Dwivedi R, Nothaft H, Garber J, Xin Kin L, Stahl M et al.** L-fucose influences chemotaxis and biofilm formation in *Campylobacter jejuni*. *Mol Microbiol* 2016;101:575-589. doi: 10.1111/mmi.13409

28. **Lubke AL, Minatelli S, Riedel T, Lugert R, Schober I et al.** The transducer-like protein Tlp12 of Campylobacter jejuni is involved in glutamate and pyruvate chemotaxis. *BMC Microbiol* 2018;18:111. doi: 10.1186/s12866-018-1254-0

29. **Chandrashekhar K, Srivastava V, Hwang S, Jeon B, Ryu S et al.** Transducer-Like Protein in Campylobacter jejuni With a Role in Mediating Chemotaxis to Iron and Phosphate. *Front Microbiol* 2018;9:2674. doi: 10.3389/fmicb.2018.02674

30. **Cha G, Chen Z, Mo R, Lu G, Gao B**. The Novel Regulators CheP and CheQ Control the Core Chemotaxis Operon cheVAW in Campylobacter jejuni. *Mol Microbiol* 2018:Epub ahead of print, doi: 10.1111/mmi.14144. doi: 10.1111/mmi.14144

31. **Takata T, Fujimoto S, Amako K**. Isolation of nonchemotactic mutants of *Campylobacter jejuni* and their colonization of the mouse intestinal tract. *Infect Immun* 1992;60:3596-600. doi:

32. **Yao R, Burr DH, Guerry P**. CheY-mediated modulation of *Campylobacter jejuni* virulence. *Mol Microbiol* 1997;23:1021-31. doi:

33. **Chandrashekhar K, Gangaiah D, Pina-Mimbela R, Kassem, II, Jeon BH et al.** Transducer like proteins of *Campylobacter jejuni* 81-176: role in chemotaxis and colonization of the chicken gastrointestinal tract. *Front Cell Infect Microbiol* 2015;5:46. doi: 10.3389/fcimb.2015.00046

34. **Hendrixson DR, DiRita VJ**. Identification of *Campylobacter jejuni* genes involved in commensal colonization of the chick gastrointestinal tract. *Mol Microbiol* 2004;52:471-84. doi: 10.1111/j.1365-2958.2004.03988.x

35. **Bereswill S, Fischer A, Plickert R, Haag LM, Otto B et al.** Novel murine infection models provide deep insights into the "menage a trois" of *Campylobacter jejuni*, microbiota and host innate immunity. *PLoS One* 2011;6:e20953. doi: 10.1371/journal.pone.0020953

36. **de Vries SP, Linn A, Macleod K, MacCallum A, Hardy SP et al.** Analysis of Campylobacter jejuni infection in the gnotobiotic piglet and genome-wide identification of bacterial factors required for infection. *Sci Rep* 2017;7:44283. doi: 10.1038/srep44283

37. **Gao B, Vorwerk H, Huber C, Lara-Tejero M, Mohr J et al.** Metabolic and fitness determinants for in vitro growth and intestinal colonization of the bacterial pathogen Campylobacter jejuni. *PLoS Biol* 2017;15:e2001390. doi: 10.1371/journal.pbio.2001390

38. **Porcelli I, Reuter M, Pearson BM, Wilhelm T, van Vliet AHM**. Parallel evolution of genome structure and transcriptional landscape in the Epsilonproteobacteria. *BMC Genomics* 2013;14:616. doi: 10.1186/1471-2164-14-616

39. **Heimbrook ME, Wang WL, Campbell G**. Staining bacterial flagella easily. *J Clin Microbiol* 1989;27:2612-5. doi:

40. **Reuter M, Periago PM, Mulholland F, Brown HL, van Vliet AHM**. A PAS domain-containing regulator controls flagella-flagella interactions in *Campylobacter jejuni*. *Front Microbiol* 2015;6:770. doi: 10.3389/fmicb.2015.00770

41. **Lauga E, DiLuzio WR, Whitesides GM, Stone HA**. Swimming in circles: motion of bacteria near solid boundaries. *Biophys J* 2006;90:400-12. doi: 10.1529/biophysj.105.069401

42. **Usher KC, de la Cruz AF, Dahlquist FW, Swanson RV, Simon MI et al.** Crystal structures of CheY from Thermotoga maritima do not support conventional explanations for the structural basis of enhanced thermostability. *Protein Sci* 1998;7:403-12. doi: 10.1002/pro.5560070221

43. **Motaleb MA, Miller MR, Li C, Bakker RG, Goldstein SF et al.** CheX is a phosphorylated CheY phosphatase essential for *Borrelia burgdorferi* chemotaxis. *J Bacteriol* 2005;187:7963-9. doi: 10.1128/jb.187.23.7963-7969.2005

44. **Park SY, Chao X, Gonzalez-Bonet G, Beel BD, Bilwes AM et al.** Structure and function of an unusual family of protein phosphatases: the bacterial chemotaxis proteins CheC and CheX. *Mol Cell* 2004;16:563-74. doi: 10.1016/j.molcel.2004.10.018

45. **Yao R, Burr DH, Doig P, Trust TJ, Niu H et al.** Isolation of motile and non-motile insertional mutants of *Campylobacter jejuni*: the role of motility in adherence and invasion of eukaryotic cells. *Mol Microbiol* 1994;14:883-93. doi:

46. **Micali G, Endres RG**. Bacterial chemotaxis: information processing, thermodynamics, and behavior. *Curr Opin Microbiol* 2016;30:8-15. doi: 10.1016/j.mib.2015.12.001

47. **Wuichet K, Zhulin IB**. Origins and Diversification of a Complex Signal Transduction System in Prokaryotes. *Science Signaling* 2010;3:ra50-ra50. doi: 10.1126/scisignal.2000724

48. **Rosario MM, Fredrick KL, Ordal GW, Helmann JD**. Chemotaxis in Bacillus subtilis requires either of two functionally redundant CheW homologs. *J Bacteriol* 1994;176:2736-9. doi:

49. **Parrish JR, Yu J, Liu G, Hines JA, Chan JE et al.** A proteome-wide protein interaction map for *Campylobacter jejuni*. *Genome Biol* 2007;8:R130. doi: 10.1186/gb-2007-8-7-r130

50. **Hartley-Tassell LE, Shewell LK, Day CJ, Wilson JC, Sandhu R et al.** Identification and characterization of the aspartate chemosensory receptor of *Campylobacter jejuni*. *Mol Microbiol* 2010;75:710-30. doi: 10.1111/j.1365-2958.2009.07010.x

51. **Du X, Wang N, Ren F, Tang H, Jiao X et al.** cj0371: A Novel Virulence-Associated Gene of Campylobacter jejuni. *Front Microbiol* 2016;7:1094. doi: 10.3389/fmicb.2016.01094

52. **Jimenez-Pearson MA, Delany I, Scarlato V, Beier D**. Phosphate flow in the chemotactic response system of *Helicobacter pylori*. *Microbiology (Reading, England)* 2005;151:3299-311. doi: 10.1099/mic.0.28217-0

53. **Karatan E, Saulmon MM, Bunn MW, Ordal GW**. Phosphorylation of the response regulator
CheV is required for adaptation to attractants during *Bacillus subtilis* chemotaxis. *J Biol Chem*
2001;276:43618-26. doi: 10.1074/jbc.M104955200
54. **Pittman MS, Goodwin M, Kelly DJ**. Chemotaxis in the human gastric pathogen *Helicobacter*
pylori: different roles for CheW and the three CheV paralogues, and evidence for CheV2
phosphorylation. *Microbiology (Reading, England)* 2001;147:2493-504. doi:
10.1099/00221287-147-9-2493
55. **Schuster M, Silversmith RE, Bourret RB**. Conformational coupling in the chemotaxis
response regulator CheY. *Proc Natl Acad Sci U S A* 2001;98:6003-8. doi:
10.1073/pnas.101571298
56. **Sola M, Lopez-Hernandez E, Cronet P, Lacroix E, Serrano L et al.** Towards understanding
a molecular switch mechanism: thermodynamic and crystallographic studies of the signal
transduction protein CheY. *J Mol Biol* 2000;303:213-25. doi: 10.1006/jmbi.2000.4507
57. **Lee SY, Cho HS, Pelton JG, Yan D, Berry EA et al.** Crystal structure of activated CheY.
Comparison with other activated receiver domains. *J Biol Chem* 2001;276:16425-31. doi:
10.1074/jbc.M101002200
58. **Porter SL, Wadhams GH, Armitage JP**. *Rhodobacter sphaeroides*: complexity in
chemotactic signalling. *Trends Microbiol* 2008;16:251-60. doi: 10.1016/j.tim.2008.02.006
59. **Szurmant H, Bunn MW, Cannistraro VJ, Ordal GW**. Bacillus subtilis hydrolyzes CheY-P
at the location of its action, the flagellar switch. *J Biol Chem* 2003;278:48611-6. doi:
10.1074/jbc.M306180200
60. **Szurmant H, Muff TJ, Ordal GW**. Bacillus subtilis CheC and FliY are members of a novel
class of CheY-P-hydrolyzing proteins in the chemotactic signal transduction cascade. *J Biol*
Chem 2004;279:21787-92. doi: 10.1074/jbc.M311497200
61. **Silversmith RE**. Auxiliary phosphatases in two-component signal transduction. *Curr Opin*
Microbiol 2010;13:177-83. doi: 10.1016/j.mib.2010.01.004
62. **Howitt MR, Lee JY, Lertsethtakarn P, Vogelmann R, Joubert LM et al.** ChePep controls
Helicobacter pylori Infection of the gastric glands and chemotaxis in the Epsilonproteobacteria.
MBio 2011;2. doi: 10.1128/mBio.00098-11
63. **Rahman H, King RM, Shewell LK, Semchenko EA, Hartley-Tassell LE et al.**
Characterisation of a multi-ligand binding chemoreceptor CcmL (Tlp3) of *Campylobacter*
jejuni. *PLoS Pathog* 2014;10:e1003822. doi: 10.1371/journal.ppat.1003822
64. **Parkhill J, Wren BW, Mungall K, Ketley JM, Churcher C et al.** The genome sequence of
the food-borne pathogen *Campylobacter jejuni* reveals hypervariable sequences. *Nature*
2000;403:665-8. doi: 10.1038/35001088
653

654   **Table 1.** *Campylobacter jejuni* strains used in this study.

Strain	Description [a]
C. jejuni strains	
NCTC 11168	Wildtype *C. jejuni* [64]
11168 Δ*cheY*	NCTC 11168 *cj1118c::kan^R*
11168 Δ*cheW*	NCTC 11168 *cj0283c::kan^R*
11168 Δ*cheA*	NCTC 11168 *cj0284c::cat^R*
11168 Δ*cheV*	NCTC 11168 *cj0285c::kan^R*
11168 Δ*cheY +cheY^{fdxApr*}*	NCTC 11168 *cj1118c::kan^R cj0046::cj1118c*^{fdxApr*}*cat^R*
11168 Δ*cheY +cheYD53A^{fdxApr*}*	NCTC 11168 *cj1118c::kan^R cj0046::cj1118c*^{D53AfdxApr*}*cat^R*
11168 Δ*cheY +cheYD7A^{fdxApr*}*	NCTC 11168 *cj1118c::kan^R cj0046::cj1118c*^{D7AfdxApr*}*cat^R*
11168 Δ*cheY +cheA-Y^{fdxApr*}*	NCTC 11168 *cj1118c::kan^R cj0046::cj0284c*^{648-769-fdxApr*}*cat^R*
11168 Δ*cheW +cheW^{fdxApr*}*	NCTC 11168 *cj0283c::kan^R cj0046::cj0283c*^{fdxApr*}*cat^R*
11168 Δ*cheA +cheA^{fdxApr*}*	NCTC 11168 *cj0284c::cat^R cj0046::cj0284c*^{fdxApr*}*kan^R*
11168 Δ*cheA +cheARec^{fdxApr*}*	NCTC 11168 *cj0284c::cat^R cj0046::cj0284c*ΔRec^{fdxApr*}*kan^R*
11168 Δ*cheV +cheV^{fdxApr*}*	NCTC 11168 *cj0285c::kan^R cj0046::cj0285c*^{fdxApr*}*cat^R*
11168 Δ*cheW +cheV^{fdxApr*}*	NCTC 11168 *cj0283c::kan^R cj0046::cj0285c*^{fdxApr*}*cat^R*
11168 *cheY^{fdxApr*}*	NCTC 11168 *cj0046::cj1118c*^{fdxApr*}*cat^R*
11168 *cheYD53A^{fdxApr*}*	NCTC 11168 *cj0046::cj1118c*^{D53AfdxApr*}*cat^R*
11168 *cheW^{fdxApr*}*	NCTC 11168 *cj0046::cj0283c*^{fdxApr*}*cat^R*
11168 *cheA^{fdxApr*}*	NCTC 11168 *cj0046::cj0284c*^{fdxApr*}*kan^R*
11168 Δ*cj0350*	NCTC 11168 *cj0350::cat^R*
11168 *cj0350^{fdxApr*}*	NCTC 11168 *cj0046::cj0350*^{fdxApr*}*kan^R*
11168 Δ*cj0350 +cj0350^{fdxApr*}*	NCTC 11168 *cj0350::cat^R cj0046::cj0350*^{fdxApr*}*kan^R*
11168 Δ*cj0350 +fliN^{fdxApr*}*	NCTC 11168 *cj0350::cat^R cj0046::cj0351*^{fdxApr*}*kan^R*
11168 Δ*pflA*	NCTC11168 *cj1565c::kan^R*

a.   *kan^R*, kanamycin resistance cassette; *cat^R*, chloramphenicol resistance cassette; * denotes

complementation construct: ^{fdxApr}, gene under control of the constitutive *fdxA* promoter.

LEGENDS TO FIGURES

Figure 1. Architecture of the core CheV-A-W-Y chemotaxis system in *C. jejuni* compared to a

model Gram-negative (*E. coli* K-12) and Gram-positive (*B. subtilis* 168) organism.

Figure 2. Disrupting chemotaxis proteins does not impair flagella assembly. Brucella broth cultures

of wild-type NCTC11168 and the chemotaxis gene mutants and complemented strains were grown

overnight at 37°C in microaerobic conditions. Cell were mounted between a microscope slide and

coverslip and freshly prepared Ryu stain added adjacent to the coverslip. After five minutes, areas

where the Ryu stain had penetrated were photographed at ×100 magnification using a Nikon

Coolpix 4500 digital camera. Montage images and scale bars were prepared using ImageJ. Scale bar

= 2 microns. Pictures are representative examples of multiple cultures each examined.

Figure 3. Chemotaxis mutants show a significant reduction in chemotactic motility in soft agar. For

each chemotaxis mutant, the mutant strain was inoculated into 0.4% agar with the wildtype strain.

Halo formation was measured after 24 and 48 hours and halo area expressed as the percentage of

the wildtype halo for the same plate. Raw mean area of WT was 293.2 ± 15.52 and 1230 ± 63.69

mm^2 (24 and 48 hrs). Results are the mean from at least three biological replicates and error bars

show standard deviation. An asterisk denotes statistically significant results, based on a one-sample

t-test (comparison with wildtype) or a two-tailed Mann-Whitney test (*p < 0.05, **p < 0.01).

Figure 4. The *cheY* mutants show a 'catherine wheel' swimming phenotype consisting of repeated

clockwise swimming. Following $6-7$ hours growth at 37°C in microaerobic conditions, movies of

swimming cells (a, wildtype; b, *cheY*; c, *cheY +cheY*; d, *cheY +cheY*^{D53A}; e, *cheY +cheY*^{D7A}; f, *cheY*

+cheA-Y) were recorded at ×100 magnification using a light microscope and attached Nikon

camera. Individual cells were tracked using the Manual Tracking ImageJ plug-in and swimming

trajectories plotted using the Chemotaxis and Migration Tool (Ibidi).

Figure 5. Expressing a second copy of *cj0350* encoding a putative CheX homolog decreases

swarming motility. a) Chemotactic motility. Each strain was inoculated into 0.4% agar with the

wild-type strain. Halo formation was measured after 24 and 48 hours and halo area expressed as the

percentage of the wild-type halo for the same plate. Results are the mean from at least three

biological replicates and error bars show standard deviation. b) Tube Taxis. Assays were monitored

over 72 hours at 37°C and the dye front measured after 24, 48, and 72 hours. Migration was

calculated as a percentage of the wild-type used in the same assay. Results are the mean from at

least three biological replicates and error bars show standard deviation. c. Representative swarm

plate and tube taxis assays showing the WT +*cheX* phenotype. d. WT +*cheX* morphology shown

using scanning electron microscopy and Ryu staining. SEM scale bar = 1 micron, Ryu staining

scale bar = 2 microns.

Figure 6. Chemotaxis is required for biofilm formation at the air-media interface. Static cultures of

each strain were grown in the presence of a sterile glass slide, as shown in the cartoon on the right

(37°C, microaerobic, 48 hrs). Glass slides were removed and stained with 1% crystal violet. CV-

stained biofilm was detected using a GenePix microarray scanner employing both 635 and 532 nm

lasers. An asterisk shows the position of the air-media interface. Three independent biological

repeats are shown for each strain.

SUPPLEMENTARY INFORMATION

Supplementary figures

Figure S1. Chemotaxis mutants do not show impaired growth in Brucella media in microaerobic

conditions.

Figure S2. Chemotaxis mutants show a significant reduction in migration in an energy and redox

gradient.

Figure S3. RT-PCR analysis of chemotaxis gene expression in wildtype and mutant strains.

Figure S4. Strains containing two copies of chemotaxis genes have altered swarming and migration

in taxis tubes.

Figure S5. *C. jejuni* CheY and CheA-Y share 58% similarity and contain the conserved acid pocket

and aspartic acid active site residues.

Figure S6. The biofilm phenotype of the *che* mutants is linked to shedding of cells from the biofilm

and the ability to swim to the air-media interface.

Supplementary Tables

Table S1. Oligonucleotide primers used in this study.

Table S2. Plasmids used in this study.

Other supplementary information

Supplementary Movie 1. Swimming behaviour of the *C. jejuni cheY* mutant.

Available via FigShare, doi: 10.15126/surreydata.7530881

Supplementary Movie 2. Swimming behaviour of the *C. jejuni* WT +*cj0350* (*cheX*) strain.

Available via FigShare, doi: 10.15126/surreydata.7530884

Figure 1. Architecture of the core CheV-A-W-Y chemotaxis system in *C. jejuni* compared to a model Gram-negative (*E. coli* K-12) and Gram-positive (*B. subtilis* 168) organism.

Figure 2. Disrupting chemotaxis proteins does not impair flagella assembly. Brucella broth cultures were grown overnight at 37°C in microaerobic conditions. Cell were mounted between a microscope slide and coverslip and freshly prepared Ryu stain added adjacent to the coverslip. After five minutes, areas where the Ryu stain had penetrated were photographed at ×100 magnification using a Nikon Coolpix 4500 digital camera. Montage images and scale bars were prepared using ImageJ. Scale bar = 2 microns.

Figure 3. Chemotaxis mutants show a significant reduction in chemotactic motility in soft agar. For each chemotaxis mutant, the mutant strain was inoculated into 0.4% agar with the wildtype strain. Halo formation was measured after 24 and 48 hours and halo area expressed as the percentage of the wildtype halo for the same plate. Raw mean area of WT was 293.2 ± 15.52 and 1230 ± 63.69 mm^2 (24 and 48 hrs). Results are the mean from at least three biological replicates and error bars show standard deviation. An asterisk denotes statistically significant results, based on a one-sample *t*-test (comparison with wildtype) or a two-tailed Mann-Whitney test (*p < 0.05, **p < 0.01).

Figure 4. The *cheY* mutants show a 'catherine wheel' swimming phenotype consisting of repeated clockwise swimming. Following 6 – 7 hours growth at 37°C in microaerobic conditions, movies of swimming cells (a, wildtype; b, *cheY*; c, *cheY +cheY*; d, *cheY +cheY*^{D53A}; e, *cheY +cheY*^{D7A}; f, *cheY +cheA-Y*) were recorded at ×100 magnification using a light microscope and attached Nikon camera. Individual cells were tracked using the Manual Tracking ImageJ plug-in and swimming trajectories plotted using the Chemotaxis and Migration Tool (Ibidi).

Figure 5. Expressing a second copy of *cj0350* encoding a putative CheX homolog decreases swarming motility. a) Chemotactic motility. Each strain was inoculated into 0.4% agar with the wild-type strain. Halo formation was measured after 24 and 48 hours and halo area expressed as the percentage of the wild-type halo for the same plate. Results are the mean from at least three biological replicates and error bars show standard deviation. b) Tube Taxis. Assays were monitored over 72 hours at 37°C and the dye front measured after 24, 48, and 72 hours. Migration was calculated as a percentage of the wild-type used in the same assay. Results are the mean from at least three biological replicates and error bars show standard deviation. c. Representative swarm plate and tube taxis assays showing the WT +*cheX* phenotype. d. WT +*cheX* morphology shown using scanning electron microscopy and Ryu staining. SEM scale bar = 1 micron, Ryu staining scale bar = 2 microns.

Figure 6. Chemotaxis is required for robust biofilm formation at the air-media interface. Static cultures of each strain were grown in the presence of a sterile glass slide, as shown in the cartoon on the right (37°C, microaerobic, 48 hrs). Glass slides were removed and stained with 1% crystal violet. CV-stained biofilm was detected using a GenePix microarray scanner employing both 635 and 532 nm lasers. An asterisk shows the position of the air-media interface. Three independent biological repeats are shown for each strain.