Massively parallel fitness profiling reveals multiple novel enzymes in *Pseudomonas putida*

lysine metabolism

Mitchell G. Thompson[1,2,3], Jacquelyn M. Blake-Hedges[1,2,4], Pablo Cruz-Morales[1,2,5], Jesus F.

Barajas[2,6], Samuel C. Curran[1,2,7], Christopher B. Eiben[1,2,8], Nicholas C. Harris[3], Veronica T.

Benites[1,2], Jennifer W. Gin[1,2], William A. Sharpless[1,2,3], Frederick F. Twigg[8], Will Skyrud[4],

Rohith N. Krishna[1,2,4], Jose Henrique Pereira[1,9], Edward E. K. Baidoo[1,2], Christopher J.

Petzold[1,2], Paul D. Adams[1,9,10], Adam P. Arkin[10,11], Adam M. Deutschbauer[3,11], Jay D.

Keasling[1,2,8,10,12]

[1]Joint BioEnergy Institute, 5885 Hollis Street, Emeryville, CA 94608, USA.

[2]Biological Systems & Engineering Division, Lawrence Berkeley National Laboratory, Berkeley,

CA 94720, USA.

[3]Department of Plant and Microbial Biology, University of California, Berkeley, CA 94720,

USA

[4]Department of Chemistry, University of California, Berkeley, CA 94720, USA

[5]Centro de Biotecnología FEMSA, Tecnológico de Monterrey, NL, Mexico

[6]Department of Energy Agile BioFoundry, Emeryville, CA 94608, USA

[7]Comparative Biochemistry Graduate Group, University of California, Berkeley, CA 94720,

USA

[8]Department of Chemical and Biomolecular Engineering, University of California, Berkeley, CA

94720, USA

[9]Molecular Biophysics and Integrated Bioimaging Division, Lawrence Berkeley National

Laboratory, Berkeley, CA 94720, USA.

[10]Joint Program in Bioengineering, University of California, Berkeley/San Francisco, CA 94720,

USA

[11]Environmental Genomics and Systems Biology Division, Lawrence Berkeley National

Laboratory, Berkeley, CA 94720, USA.

[12]The Novo Nordisk Foundation Center for Biosustainability, Technical University of Denmark,

Denmark

Abstract

Despite intensive study for 50 years, the biochemical and genetic links between lysine

metabolism and central metabolism in *Pseudomonas putida* remain unresolved. To establish

these biochemical links, we leveraged Random Barcode Transposon Sequencing (RB-TnSeq), a

genome-wide assay measuring the fitness of thousands of genes in parallel, to identify multiple

novel enzymes in both L- and D-lysine metabolism. We first describe three pathway enzymes

that catabolize L-2-aminoadipate (L-2AA) to 2-ketoglutarate (2KG), connecting D-lysine to the

TCA cycle. One of these enzymes, PP_5260, contains a DUF1338 domain, a family with no

previously described biological function. Our work also identified the recently described CoA

independent route of L-lysine degradation that metabolizes to succinate. We expanded on

previous findings by demonstrating that glutarate hydroxylase CsiD is promiscuous in its 2-

oxoacid selectivity. Proteomics of select pathway enzymes revealed that expression of catabolic

genes is highly sensitive to particular pathway metabolites, implying intensive local and global

regulation. This work demonstrates the utility of RB-TnSeq for discovering novel metabolic

pathways in even well-studied bacteria, as well as a powerful tool for validating previous

research.

Importance

P. putida lysine metabolism can produce multiple commodity chemicals, conferring great

biotechnological value. Despite much research, connecting lysine catabolism to central

metabolism in *P. putida* remained undefined. Herein we use Random Barcode Transposon

Sequencing to fill in the gaps of lysine metabolism in *P. putida*. We describe a route of 2-

oxoadipate (2OA) catabolism in bacteria, which utilizes DUF1338 containing protein PP_5260.

Despite its prevalence in many domains of life, DUF1338 containing proteins had no known

biochemical function. We demonstrate PP_5260 is a metalloenzyme which catalyzes an unusual

2OA to D-2HG decarboxylation. Our screen also identified a recently described novel glutarate

metabolic pathway. We validate previous results, and expand the understanding of glutarate

hydroxylase CsiD by showing can it use either 2OA or 2KG as a cosubstrate. Our work

demonstrates biological novelty can be rapidly identified using unbiased experimental genetics,

and that RB-TnSeq can be used to rapidly validate previous results.

Introduction

Pseudomonas putida is an ubiquitous saprophytic soil bacterium and is a model organism

for bioremediation (1). Interest in utilizing *P. putida* KT2440 as a chassis organism for metabolic

engineering has recently surged due to the existence of well-established genetic tools and its

robust metabolism of aromatic compounds that resemble lignin hydrolysis products (2–4). As

lignin valorization remains essential for the economic feasibility of cellulosic bioproducts, a

nuanced and predictable understanding of *P. putida* metabolism is highly desirable (5).

Although its aromatic metabolism has garnered much attention, the lysine metabolism of

P. putida has also been rigorously studied for over fifty years (6). An understanding of lysine

metabolism has had biotechnological value, as it has been used to produce glutarate, 5-

aminovalerate (5AVA), as well as valerolactam in *P. putida* and in the other bacteria (7–10). Our

current understanding of lysine catabolism however, remains incomplete. In particular, the

connection between D-lysine metabolism and central metabolism in *P. putida* is unclear and has

not been fully characterized.

Figure S1**: Metabolic pathways of lysine catabolism in *P. putida* KT2440**. A) L-lysine metabolic pathway is

shown in blue, while the known steps of D-lysine metabolism are shown in red. B) Proposed route of 2AA

metabolism in *P. putida*.

P. putida employs bifucating pathways to catabolize lysine, separately metabolizing the

L- and D-isomers (Figure S1a) (11). The L-lysine degradation pathway proceeds to glutarate,

which can then be either be degraded to acetyl-CoA via a glutaryl-CoA intermediate, or to

succinate without a CoA bound intermediate (Figure S1a) (9). The final steps of D-lysine

catabolism remain more elusive. The initial steps of D-lysine catabolism are well described, but

the genetic basis stops at 2AA (12). Furthermore, ^{13}C labeling experiments by Revelles et al.

demonstrated a putative metabolic connection between the D- and L-lysine pathways at 2AA

(11). Subsequent steps to central carbon metabolism have never been fully validated. (6, 11–13).

Given the importance of lysine metabolism, and recent availability of high-throughput genetic

tools, we sought to identify the missing steps in D-lysine metabolism that have remained despite

50 years of research.

Random barcode transposon sequencing (RB-TnSeq) is a genome-wide approach that

measures the importance of each gene to growth (or fitness) in a massively parallel assay (14).

RB-TnSeq can identify phenotypes for thousands of previously uncharacterized genes (14, 15),

including the levulinic acid degradation pathway in *P. putida* KT2440 (16). In this study, we

applied RB-TnSeq to uncover multiple novel genes implicated in L- and D-lysine metabolism in

P. putida. We first describe a three enzyme route connecting L-2AA to 2KG (Figure S1B).

Within this pathway, D-lysine metabolism connects to central metabolism through a 2HG

intermediate, which is directly produced from 2OA in a reaction catalyzed by a DUF1338-

containing protein. This protein family, widely distributed across many domains of life,

previously had no known function. Subsequently, we further characterize the glutarate

hydroxylase CsiD, by demonstrating its 2-oxoacid promiscuity during the hydroxylation of

glutarate. Finally, we show the expression of all newly discovered enzymes changes significantly

in response to specific metabolites within the two catabolic pathways.

Results

Identification of lysine catabolism genes via RB-TnSeq

Figure 1: **Results of RB-TnSeq screen**. A) Genes that showed less than -2 log$_2$ fitness on either D-lysine, L-lysine, or 5AVA but showed no less than -0.5 log$_2$ fitness defect when grown on glucose. B) Plot of genome wide fitness values of libraries grown on either L-lysine or D-lysine. Genes encoding for enzymes known to be involved in D-lysine metabolism are shown in red, while those known to be involved in L-lysine metabolism are shown in blue. C) Venn diagram of genes with significant fitness defects when grown on either D-lysine, L-lysine, or 5AVA.

To identify mutants defective in lysine catabolism in *P. putida* KT2440, an RB-TnSeq

library of this bacterium (16) was grown on minimal medium supplemented with either D-lysine

or L-lysine as the sole carbon source. To evaluate whether D-lysine metabolism was required for

the metabolism of other downstream metabolites of L-lysine, the library was also grown on

5AVA. As a control, we also grew the library on glucose. Fitness was calculated as the log$_2$ ratio

of strain and gene abundance at the end of selective growth relative to initial abundance (14).

Fitness profiling revealed 39 genes with significant fitness values of less than -2 for 5AVA, D-

lysine, or L-lysine, and no less than -0.5 fitness for glucose (Figure 1a, Supplementary Table 1).

Within this set, 10 of the 12 known lysine degradation genes were identified, with the exception

of the two enzymes in the CoA-dependent route of glutarate degradation (*gcdH* and *gcdG*),

which both had significant fitness values ($t < -4$) but whose fitness was greater than -2. Instead,

we identified the recently-characterized genes involved in the CoA independent pathway (*csiD*

and *lghO*) (9).

The fitness data corroborated previous work showing a functional D-lysine pathway is

required for L-lysine catabolism (6, 11). None of the known L-lysine catabolic genes showed

fitness defects for growth on D-lysine, but transposon insertions in all previously-identified D-

lysine genes showed negative fitness scores when grown on L-lysine (Figure 1b). No known D-

lysine catabolic enzymes showed fitness defects when grown on 5AVA, suggesting the D-lysine

dependence of L-lysine catabolism may only occur for early catabolic steps (Figure 1c).

In addition to catabolic enzymes, lysine transporters and multiple transcriptional

regulators were identified (Figure 1a). The putative lysine amino acid ABC transporter system

(PP_3593, PP_3394, and PP_3395) showed significant fitness defects when grown with either

isomer of lysine. Some of the transcriptional regulators were located near known catabolic or

transport enzymes (PP_0384, PP_3592, and PP_3603), while others were not clustered with any

obviously related genes (PP_1109, PP_2868, PP_3649, and PP_4482). Two known global

regulators were identified in our screen: *cbrA* (PP_4695), a histidine kinase sensor that showed

fitness defects on both lysine isomers, and the alternative sigma factor *rpoX* (PP_2088) which

only had fitness defects when grown on D-lysine.

Additionally there were 15 genes which, when disrupted, displayed fitness advantages

greater than 2 on 5AVA, D-lysine, or L-lysine and less than 0.5 fitness when grown on glucose.

This positive fitness value indicates these mutations confer a competitive advantage compared to

other strains when grown on these carbon sources. Most striking amongst these genes were the

sigma factor *rpoS* and the LPS export system (PP_1778/9), which when disrupted, both

displayed fitness benefits on all three non-glucose carbon sources (Figure S2).

Figure S2: Results of RB-TnSeq screen. A) Genes that showed great than 2 log2 fitness on either D-lysine, L-lysine, or 5AVA but showed no less than 0.5 log2 fitness defect when grown on glucose.

Only one gene (PP_0787, a quinolinate phosphoribosyltransferase) showed fitness

defects on all three non-glucose carbon sources (Figure 1c). However, disruption of PP_0787

also showed a significant fitness defect when grown on levulinic acid, suggesting it is unlikely to

be uniquely important to lysine metabolism (16). Only 3 genes shared fitness defects between

5AVA and L-lysine (*davT*, *davD*, and *lghO*), all of which have been previously implicated in

5AVA metabolism (Figure 1c) (9).

 <u>PP_4108 is a L-2AA aminotransferase</u>

Figure 2: **Identification of PP_4108 as a L-2AA aminotransferase**. A) Growth of wild-type KT2440 and PP_4108 mutant on 2AA as a sole carbon source. Shaded area represents 95% confidence interval (CI), n=3. B) *In vivo* accumulation of 2AA in wild-type KT2440 and a PP_4108 mutant after 12 hours of growth on minimal medium supplemented with 10 mM glucose and 10 mM D-lysine. Bars represent $\log_{10}$ transformed spectral counts, error bars show 95% CI, n=3. C) *In vitro* transamination reactions of PP_4108 with 2KG as an amino acceptor. Bars represent µM metabolite concentration of either 2OA (black) or 2AA (white) in either boiled or native protein reactions. Error bars show 95% CI, n=3. D) *In vitro* transaminations of PP_4108 incubated with different possible amino donors and 2KG as acceptor. Bars represent relative activity of enzyme standardized to L-2AA after 16 hour incubation. Error bars show standard deviation, n=2.

In humans and other animals, L-lysine degradation proceeds through a 2AA intermediate,

which a transaminase converts to 2OA (9, 11, 17). Yet, no such transaminase has been identified

in *P. putida*. We identified a candidate aminotransferase, PP_4108, for which gene inactivation

showed a significant growth defect on D-lysine (-5.9) and a relatively minor defect on L-lysine (-

1.2). To corroborate our RB-TnSeq fitness data, we constructed a deletion mutant of PP_4108

that failed to grow in a plate reader assay on 10 mM DL-2AA (Figure 2a). The mutant showed a

severe growth defect on 10 mM D-lysine and an increased lag time when grown on 10 mM L-

lysine (Figure S3).

Figure S3**: Growth of PP_4108 mutants on lysine**. A) Growth of wild-type KT2440 and PP_4108 mutant on D-

lysine as a sole carbon source. Shaded area represents 95% CI, n=3. B) Growth of wild-type KT2440 and PP_4108

mutant on L-lysine as a sole carbon source. Shaded area represents 95% CI, n=3. C) Colorimetric glutamate

formation time course of PP_4108. Error bars are standard deviation of n=2.

To further validate this hypothesis, the ΔPP_4108 strain was subjected to metabolomics

analysis to monitor the accumulation of its expected substrate, 2AA, when grown on glucose and

D-lysine. After 12 hours of growth on minimal media supplemented with 10 mM each of glucose

and D-lysine, the PP_4108 deletion strain showed a 6.3-fold increase ($p = 0.00016$) in

normalized intracellular 2AA concentration compared to WT (Figure 2b). Next PP_4108 was

expressed and purified from *E. coli* for biochemical characterization. After purified enzyme

incubation with DL-2AA, 2KG, and pyridoxal phosphate (PLP) for 16 hours, the reaction

mixture was analyzed with LC-TOF. The expected product, 2OA, was detected in the enzymatic

reaction but not in a boiled enzyme control, confirming PP_4108 as a transaminase that converts

2AA to 2OA (Figure 2c). As many transaminases have broad substrate specificity (18), we also

probed the substrate range of PP_4108 using a colorimetric assay for glutamate, a stoichiometric

product of the transamination reaction (Figure 2d). The enzyme was most active on L-2AA, and

only showed 2.8% relative activity ($p = 0.0057$) on its enantiomer, D-2AA. This specificity for

the L-2AA isomer may explain why only 50% of the DL-2AA was transformed in the previous

experiment (Figure 2c). No activity was observed on either lysine isomer; however, the enzyme

had slight activity towards 4-aminobutyrate/γ-aminobutyrate (GABA) (2.8% relative activity, $p =$

0.0057) and moderate activity on 5AVA (30.5% relative activity, $p = 0.0139$). Over shorter time

scales PP_4108 had no activity on any substrate expect L-2AA (Figure S3c). These results

suggest *P. putida* KT2440 metabolizes D-lysine to L-2AA, which is then converted to 2OA by

the transaminase PP_4108.

PP_5260 is a novel DUF1338 family enzyme that catalyzes the conversion of 2OA to 2HG

Early work proposed 2OA is converted to 2KG via a 2HG intermediate (13, 19), while

later results suggested a direct conversion of 2OA to glutarate (11). Either route likely requires a

decarboxylation of 2OA, so we initially searched for decarboxylases within our dataset. Our

fitness data on either lysine isomer revealed no obvious decarboxylases or enzymes likely to

contain a thiamine pyrophosphate (TPP) cofactor, which are commonly employed by

decarboxylases. However, a gene near other D-lysine catabolic genes in the *P. putida* genome,

PP_5260, showed a significant fitness defect. A ΔPP_5260 strain was unable to grow on either

isomer of lysine verifying its importance in lysine degradation (Figure 3a).

Figure 3: **Identification of ydcJ (PP_5260) as a 2OA decarboxylase/hydroxylase.** A) Growth of wild-type

(black) and PP_5260 mutant (red) on D-lysine (line) or L-lysine (dashed line) as a sole carbon source.

Shaded area represents 95% CI, n=3. B) HPLC traces of *in vitro* reactions run with apo PP_5260 with

exogenous metals added at 50 µM. Retention times for 2OA and 2HG are shown by vertical dashed lines.

Metal or EDTA control is indicated to the right of traces. C) *In vitro* assay of 2OA conversion to 2HG by

purified PP_5260 protein analysed via LC-TOF. 2OG in white, 2HG in black. D) *In vitro* assay of purified

PP_5260 protein with 2OA as substrate. Black bar represents concentration of D-2HG measured by enzyme

coupled assay. White bar represents total 2HG concentration as measured by LC-TOF. Error bars represent

95% CI, n=3. E) Initial velocity of reaction catalyzed by PP_5260 as a function of 2OA concentration. Blue

dots represent individual measurements, while the black fit line shows a Michaelis-Menten fit. F) Chemical

reaction catalyzed by PP_5260, 2OA is decarboxylated to D-2HG.

PP_5260 belongs to the DUF1338 protein family (http://pfam.xfam.org/family/PF07063).

Although several unpublished crystal structures of DUF1338 domain containing proteins have

been deposited into the Protein Data Bank, their biological function remains elusive. However,

these structures combined with protein sequence alignments suggest a putative metal binding site

is conserved throughout the DUF1338 family. As we hypothesized PP_5260 serves as the

missing decarboxylase in D-lysine metabolism, we purified the enzyme for biochemical analysis.

Enzymatic activity on 2OA was probed and analyzed via LC-TOF. After incubation of 2OA with

PP_5260, we observed a ~92% (*p*=0.00034) reduction in the abundance of 2OA, whereas no

2OA was consumed in a boiled enzyme control or enzyme treated with EDTA confirming it to

be a metalloenzyme (Figure S4a). Initially we believed the product would be either glutarate or

glutarate semialdehyde, however neither of these was detected in the reaction. Early biochemical

work suggested 2HG as a potential intermediate in pipecolate metabolism (19), and when the

enzymatic product was compared to a racemic 2HG standard they shared the same mass,

retention time and mass-to-charge ratio (Figure S4b), as well identical isotopic distributions of

[M-H] peaks in the mass spectra (Figure S4c).

Figure S4: **Characterization of PP_5260**. A) In vitro reactions of boiled or native PP_5260 incubated with 2OA

with 50 μM EDTA (white), or without EDTA (black). Bars represent $\log_{10}$ transformed spectral counts, errors bars

represent 95% CI, n=3 B) E LC-TOF analysis of a 2HG standard, and products of PP_5260 incubated with 2OA,

and boiled control. C) Mass spectra of 2HG standard and product of PP_5260 in vitro reaction. D) Standard curves

of D-2HG and L-2HG using a D-2HG specific enzymatic detection assay. Shaded areas represent 95% CI, n=3.

218   To identify the metal cofactor, the enzyme was dialyzed against EDTA to remove metals,

and individual divalent metals were added back. Only the addition of Fe(II) restored enzymatic

activity as measured by HPLC (Figure 3b). Subsequent reactions quenched after 5 minutes

showed 200 uM of 2HG formed and 800 uM 2OA remaining, demonstrating 1:1 2OA to 2HG

reaction stoichiometry (Figure 3c). Whether the product of the PP_5260 reaction is either L-2HG

or D-2HG is critical to understanding the eventual fate of D-lysine, as *lghO* is specific for L-

2HG (Figure S1a). An enzyme coupled assay specific for the detection of D-2HG was used to

assess the stereochemistry of the PP_5260 product. Standard curves of D-2HG and L-2HG

showed that the assay was only responsive to D-2HG (Figure S4d). The concentration of *in vitro*

reactions of PP_5260 were then measured by both LC-TOF as well as the enzyme coupled assay,

revealing all 2HG as the D-isomer (Figure 3d).

Kinetic parameters of PP_5260 were determined using an enzyme-coupled assay to

spectrophotometrically measure CO_2 evolution via NADH oxidation (20). PP_5260 displayed a

V_{max} of 0.33 mM/min (+/- 0.08 mM), a K_m of 0.06 mM (+/- 0.03 mM), and a K_{cat} of 330 m^{-1}

using 2OA as a substrate. Taken together these results reveal that PP_5260 is novel Fe(II)

dependent decarboxylase that converts 2OA to D-2HG (Figure 3f), a chemical reaction not

previously observed in nature.

<u>DUF1338 proteins are a widely distributed enzyme family with a putative conserved role in</u>

<u>amino acid catabolism</u>

After functional characterization of PP_5260, we use phylogenomics to propagate the

annotation and further explore the biological role of DUF1338 proteins found in other organisms.

We found that DUF1338 proteins are widely distributed across the tree of life, with homologs of

PP_5260 found in plants and green algae (22), fungi, and bacteria, though they were not found in

animals or archaea (Figure 4a). Homologs are widely distributed amongst bacteria, with the

Firmicutes being a notable exception. PP_5260 homologs within the plant group Streptophyta, as

well as bacterial groups Actinobacteria, Cyanobacteria, and Bacteroidetes formed monophyletic

clades, while homologs from other taxonomic groups were not monophyletic (Figure 4a).

DUF1338 homologs are found in bacteria important to biotechnology (*Corynebacterium*

glutamicum), the environment (*Nostoc puncitforme*), and medicine (*Yersinia pestis*,

Mycobacterium tuberculosis, *Burkholderia pseudomallei*).

Figure 4: **Phylogenomics of the DUF1338 enzyme family**. A) Phylogenetic relationships among DUF1338

homologs and their distribution among major phyla. Branches in the tree are colored by phylum. DUF1338 is

found in most bacterial phyla as well as in plants and fungi. Non-monophyletic clades suggest pervasive

horizontal gene transfer events in the family. B) Phylogenomics of selected DUF1338 homologs in bacteria.

The phylogeny in the left shows the phylogenetic relationships between selected homologs, the branches

have been colored according to their adscription to a given phylum and the support values are shown at the

nodes. The boxes in the right represent the gene neighborhood for each homolog. The genes have been

colored to represent their annotated functions.

Publicly available fitness data show both *Pseudomonas fluorescens* FW300-N2C3 and

Sinorhizobium meliloti PP_5260 homologs have L-lysine specific defects when interrupted (15).

Genomic contexts within other bacteria suggest many DUF1338-containing enzymes may be

involved in lysine or other amino acid metabolism (Figure 4b). Within the Actinobacteria

DUF1338 homologs are often found adjacent to sarcosine oxidases, aldehyde dehydrogenases,

and transaminases implying an additional catabolic amino acid function. In both the oleaginous

bacterium *Rhodococcus opacus* B4 and *M. tuberculosis*, DUF1338 homologs are found next to

predicted L-lysine aminotransferases, suggesting an ancestral homolog functioned in lysine

catabolism. Interestingly, the *R. opacus* B4 genome has three DUF1338 homologs, only one of

which contains genes predicted to be specific to lysine catabolism. The other two gene

neighborhoods are similar in their functional content, mainly differing by containing an

oxidoreductase or glycolate dehydrogenase, either of which may perform the same biochemical

function. In Alphaproteobacteria, Betaproteobacteria, and Cyanobacteria, the presence of

aldehyde dehydrogenases, oxidoreductases, glycolate dehydrogenases, and aminotransferases

implies a metabolic function similar to PP_5260.

<u>PP_4493 putatively oxidizes D-2HG to 2KG and connects D-lysine to central metabolism</u>

In the CoA independent route of glutarate metabolism, LghO oxidizes L-2HG to 2KG,

however this enzyme is highly specific towards the L-2HG isomer and showed no fitness defect

on D-lysine in our RB-TnSeq data (Figure S1a). A putative FAD-dependent and 4Fe-4S cluster-

containing glycolate dehydrogenase, PP_4493, did show fitness defects on both D-lysine and L-

lysine (fitness scores of -5.4 and -2.7 respectively) (Figure 1a). Glycolate dehydrogenases are

members of a larger family of enzymes that oxidize the alcohol group of an alpha-hydroxyacid to

their corresponding alpha-ketoacid (Figure 5a). Therefore, we hypothesized PP_4493 could

potentially oxidize a similar 2-hydroxyacid, 2HG, to the corresponding alpha-ketoacid, 2KG.

Moreover, many PP_5260 homologs were located next to or near putatively annotated glycolate

dehydrogenases in other bacteria, underscoring their potential metabolic link (Figure 4b). To

confirm these hypotheses, we again constructed a deletion strain, *P. putida* ΔPP_4493, which

could not grow on D-lysine as a sole carbon source (Figure 5b), and showed attenuated growth

on L-lysine (Figure S5). Furthermore, when grown on 10 mM glucose and 10 mM D-lysine the

mutant accumulated ~500 uM 2HG normalized to optical density, whereas wild type *P. putida*

did not accumulate any detectable 2HG (Figure 5c). Subsequent analysis of accumulated 2HG

via a D-2HG specific detection kit revealed that this accumulated 2HG was indeed D-2HG

(Figure 5c). These data and the conserved function and genomic context of glycolate

dehydrogenases strongly suggest PP_4493 catalyzes the last step of L-2AA metabolism,

oxidizing D-2HG to 2KG (Figure S1b).

293
294

Figure 5: **Identification of PP_4493 as a putative D-2HG dehydrogenase**. A) General chemical reaction of

a dehydrogenase converting a 2-hydroxyacid to a 2-ketoacid B) Growth of *P. putida* KT2440 and PP_4108

mutant on D-lysine as a sole carbon source. Shaded area represents 95% CI, n=3. C) *In vivo* accumulation of

2HG in wild-type KT2440 and a PP_4108 mutant after 12 hours of growth on minimal medium

supplemented with 10 mM glucose and 10 mM D-lysine. White bar represents concentration of D-2HG

measured by enzyme coupled assay. Black bar represent total 2HG concentration as measured by LC-TOF.

Red line represents limit of detection of enzyme coupled assay for D-2HG. Bars represent $\log_{10}$ transformed

spectral counts, error bars show 95% CI, n=3.

Figure S5: **Growth of PP_4493 mutant on L-lysine.** Growth of wild-type KT2440 and PP_4493 mutant on L-

lysine as a sole carbon source. Shaded area represents 95% CI, n=3

CsiD is highly specific for glutarate hydroxylation but promiscuous in 2-oxoacid selectivity

During the initial preparation of this manuscript, Zhang et al. discovered a novel pathway

of glutarate metabolism in *P. putida (9)*. They describe a cyclic reaction cascade wherein a novel

2KG-dependent non-heme Fe(II) oxygenase, PP_2909 (CsiD), hydroxylates glutarate to form

2HG and succinate using 2KG as a cosubstrate. PP_2910 (LghO), a putative L-2HG oxidase,

then subsequently converts L-2HG to 2KG, regenerating the 2KG consumed in the initial

reaction. These reactions result in the net incorporation of succinate into central metabolism

(Figure S1). Our fitness results of the library grown on 5AVA also identified both *csiD* and

lghO, in addition to the two enzymes from the CoA-dependent glutarate pathway, glutaryl-CoA

ligase (*gcdG*) and glutaryl-CoA dehydrogenase (*gcdH*), mutants of which showed mild fitness

defects when grown on 5AVA (Figure 6a). We also purified *csiD* and confirmed it hydroxylates

glutarate in a 2KG-dependent manner (Figure S6a). HPLC analysis demonstrated that as

glutarate was consumed, equimolar quantities of succinate and L-2HG were produced (Figure

S6b). Additionally, a *csiD* deletion mutant showed increased lag time when grown on either L-

lysine or 5AVA. By deleting the glutaryl-CoA ligase *gcdG*, and disrupting the CoA-dependent

glutarate pathway, we completely prevented growth on 5AVA or L-lysine (Figure S6c). These

results are in agreement with those found with Zhang et al (9).

325
326

Figure 6: **Role of CsiD in *P. putida* lysine metabolism**. A) Plot of genome wide fitness values of libraries

grown on either 5AVA or glucose. CoA-dependent glutarate degradation genes are shown in red, while those

known to be involved succinate producing metabolism are shown in blue. B) Phylogenetic tree of bacterial

CsiD homologs. Homologs used in *in vitro* assays are highlighted in red. C) *in vitro* reactions of CsiD with

different substrates using 2KG as a 2-oxoacid. Bars show peak area of 2-hydroxyacid, error bars show 95%

CI, n=3. D) *In vitro* reactions of CsiD homologs with different 2-oxoacids. Bars represent spectral counts of

L-2HG. Error bars show 95% CI, n=3.

Because non-heme Fe(II) oxidases can be promiscuous with respect to the 2-oxoacid

cosubstrate (21, 22), we evaluated the 2-oxoacid specificity of CsiD. First, we evaluated the

hydroxyl acceptor substrate specificity of CsiD family proteins by purifying two additional

homologs from *E. coli* and a halophilic bacterium, *Halobacillus* sp. BAB-2008 (Figure 6b). We

probed the activity of the homologs against a panel of 3 to 6 carbon fatty acids and diacids in the

presence of 2KG, and found only glutarate served as a hydroxylation substrate (Figure 6c). These

results are consistent with the work recently reported by Zhang *et al (9)* and further suggests the

specificity of CsiD homologs is conserved across phyla. Although extremely specific for the

hydroxylation substrate, all three CsiD homologs could utilize both 2OA and 2KG, but not

oxaloacetate, as a cosubstrate for L-2HG formation (Figure 6d). The coproduct of the reaction

using 2OA as a 2-oxoacid would be glutarate, rather than succinate. This result is particularly

interesting as it provides a possible mechanism for the previously observed metabolic link

between D-lysine and L-lysine catabolism. Growth defects observed in a ΔPP_2909 ΔPP_0158

double mutant grown on D-lysine also support this hypothesis (Figure S7a).

Figure S6: **Characterization of PP_2909 (CsiD) in *P. putida***. A) In vitro reactions of PP_2909. Bars show mM of

either succinate (blue), or 2HG (green) formed by boiled enzyme control, no 2KG control, or native enzyme with

2KG added. Errors bars show 95% CI, n=3. B) Time course in vitro reaction of PP_2909. Plot shows 2HG, 2KG,

succinate, and glutarate overtime. Shaded region shows 95% CI, n=3. C) Growth curves of wild-type KT2440,

PP_0158, PP_2909, PP_2910, or PP_2909/PP_0158 double mutants grown on either L-lysine (left column), or

5AVA (right column). Shaded region shows 95% CI, n=3.

Figure S7: **Growth of PP_2909/PP_0158 and PP_2088 mutants on D-lysine**. A) Growth of wild-type KT2440 and

PP_2909/PP_0158 mutant on D-lysine as a sole carbon source. Shaded area represents 95% CI, n=3. A) Growth of

wild-type KT2440 and PP_2088 mutant on D-lysine as a sole carbon source. Shaded area represents 95% CI, n=3.

<u>Expression of lysine metabolic proteins is responsive to pathway metabolites</u>

Multiple studies have demonstrated the expression of lysine catabolic genes is

upregulated in the presence of pathway metabolites (9, 12, 23). To investigate the regulation of

the newly-discovered lysine catabolic enzymes from this study, wild-type *P. putida* KT2440 was

grown in minimal media on glucose or a single lysine metabolite (e.g. D-lysine, L-lysine, 5AVA,

2AA, or glutarate) as a sole carbon source until cultures reached an OD_{600} of 1.0. We then

quantified the relative abundance of D- and L-lysine catabolic proteins via targeted proteomics

(Figure 7). For each protein, all pairwise statistical comparisons of different carbon sources can

be found in Supplemental Table 2. All five D-lysine pathway proteins measured (AmaA

(PP_5257), AmaB (PP_5258), PP_4108, YdcJ (PP_5260), and YdiJ (PP_4493)) were

upregulated when grown on L-lysine, D-lysine or 2AA compared to the glucose control. Neither

5AVA nor glutarate significantly induced expression of any measured D-lysine proteins. Of all

the targeted proteins, the three identified in this study that directly degrade 2AA were most

strongly induced by 2AA. Somewhat surprisingly, we also found the two enzymes involved in

2AA formation, AmaA and AmaB, were also more highly expressed in the presence of 2AA

suggesting the possible involvement of a global regulator. An interesting finding from our initial

screen showed sigma factor RpoX (PP_2088) to be required for fitness on D-lysine (Figure 1a).

Deletion mutants of *rpoX* were severely attenuated in their ability to grown on D-lysine as a sole

carbon source (Figure S7b). Further work will be necessary to examine complex regulatory

network that controls D-lysine metabolism.

The initial two enzymes from L-lysine metabolism, DavA and DavB, were most highly

expressed in the presence of L-lysine, but also significantly with D-lysine. As previously

observed, DavT and DavD were most strongly upregulated on 5AVA, moderately upregulated on

L-lysine, and to a lesser extent D-lysine. The induction of LhgO and CsiD was highest when

grown on glutarate, although these proteins were also moderately upregulated by 5AVA and L-

lysine. By comparison, PP_0159 (GcdG) expression in the presence of glutarate was stimulated

to a lesser extent than LhgO and CsiD expression; in addition, GcdG was slightly upregulated on

5AVA and L-lysine.

Figure 7: **Expression of lysine degradation pathways in response to different lysine metabolites**.

Relative abundance of selected lysine degradation enzymes expressed in wild-type KT2440 in response to

different carbon sources. Bars show spectral counts of proteins after 36 hours of growth on 10 mM glucose

(black), 5AVA (purple), D-lysine (green), L-lysine (red), glutarate (blue), or 2AA (yellow). Error bars show

95% CI, n=3.

Discussion

Despite intensive study, a complete biochemical and genetic understanding of D-lysine

catabolism in *P. putida* has remained elusive. A 2OA degradation pathway has been extensively

characterized in mammals, because of its implications in human disease (24). In the mammalian

pathway, L-lysine is metabolized to 2OA and eventually converted to acetyl-CoA via a glutaryl-

CoA intermediate (24). However, this pathway has not been observed in bacteria. Previous work

suggested 2OA is either converted via decarboxylation to glutarate or through several enzymatic

steps to 2HG, yet none of these studies conclusively demonstrated a genetic and biochemical

basis for these hypotheses (11, 19, 25). In this work we demonstrate plausible biochemical routes

to account for both of these previously hypothesized pathways.

The first route, catalyzed by the DUF1338-containing metalloenzyme PP_5260, involves

the direct conversion of 2OA to D-2HG. The formation of the D-2HG isomer by PP_5260

maintains stereochemical separation from the L-2HG formed by L-lysine degradation, thus

requiring the dehydrogenase PP_4493 rather than the L-2HG specific oxidase *lghO*. This

transformation seemingly involves two separate reactions: a decarboxylation and a

hydroxylation. Hydroxymandelate synthase has been shown to catalyze a similar enzymatic

reaction, via an intramolecular oxidative decarboxylation similar to 2KG dependent Fe(II)

oxidases (26). PP_5260 is also a Fe(II) dependent decarboxylase, and the two share similar Kcat

values for their given substrates (330 m^{-1} for PP_5260 and 270 m^{-1} for hydroxymandelate

synthase) (27). Though PP_5260 and hydroxymandelate synthase share little sequence

homology, this enzyme may give us insight into the molecular mechanism of DUF1338

enzymes. We have given PP_5260 the tentative title of 2-hydroxyglutarate synthase (*hglS*) until

further mechanistic studies (currently underway in our group) are completed and a more accurate

enzyme name can be assigned.

In bacteria, homologs of PP_5260 appear widely distributed with their genomic contexts

suggesting functions both within and beyond lysine metabolism. Genomic contexts in other

bacteria, particularly *Actinobacteria*, suggest these homologs may be involved in other amino

acid catabolic pathways. Unfortunately, there is scant evidence for homologous function in

model organisms. For example, although DUF1338 proteins are present in other *Ascomycota*,

there is no homolog in *Saccharomyces cerevisiae*. Interestingly, the *E. coli* homolog of PP_5260

is located next to a potential glucans biosynthesis gene: Glucans biosynthesis protein D (28).

Another DUF1338-containing protein from rice has been characterized and was implicated in

starch granule formation (29). These results suggest DUF1338 proteins could play a role in sugar

metabolism.

Recently Zhang *et al*. thoroughly characterized a CoA independent glutarate catabolism

route ending at succinate involving the Fe(II) dependent oxygenase CsiD (9). Our RB-TnSeq

screening convergently uncovered this pathway, and our biochemical and physiological results

fully corroborate their findings. While both works show multiple CsiD homologs from divergent

bacteria are highly specific towards glutarate as a hydroxyl acceptor, all three homologs we

tested showed promiscuous activity toward 2-oxoacid cosubstrates. The ability of the *P. putida*

CsiD to utilize 2OA as a cosubstrate is particularly interesting as it may directly connect L- and

D-lysine metabolism. The promiscuity of CsiD may explain earlier studies which reported

glutarate formation from D-lysine (11). Further studies involving labelled substrates may help

elucidate the potential link between the two pathways. While CsiD plays a clear role in L-lysine

metabolism in *P. putida*, its role in other organisms remains a mystery. In *E. coli*, RpoS controls

the expression of CsiD, but *rpoS* mutants showed fitness benefits on all three lysine metabolites

tested in our RB-TnSeq data (30). Recent work has shown that *E. coli* also uses CsiD to

metabolize lysine, suggesting a possible conserved role for this pathway across bacteria (31).

Work presented here and previous reports have shown expression of both lysine

catabolism pathways are highly responsive to their respective metabolites. While this metabolism

appears highly coordinated, the genes themselves are dispersed across the genome, with both

PP_4018, and PP_4493 found outside of operons, and relatively distant from other lysine

catabolic genes. At least two global regulators appeared to be important to lysine metabolism

based on our Rb-TnSeq data, *cbrA* (PP_4695) and *rpoX* (PP_2088). The two-component system

CbrAB has been implicated in catabolite repression and C/N balance in *P. aeruginosa*, with

mutants unable to grown on multiple amino acids (32). Further work in *P. putida* KT2440

showed the CbrAB system behaved similarly to that in *P. aeruginosa* (33). It would be

unsurprising if this regulator controls the expression of various genes within lysine catabolism;

more work into uncovering the regulon is warranted. RpoX on the other hand has been

implicated in osmotic tolerance in *P. aeruginosa* (34, 35). This is interesting as lysine

metabolism, and specifically pipecolate metabolism, has been associated with osmotic tolerance

across multiple bacteria (36). As a *rpoX* deletion mutant was unable to grow on D-lysine, these

results suggest D-lysine metabolism of *P. putida* may be involved in adaptation to saline or other

osmotically stressful environments.

An interesting question remains as to why *P. putida* maintains separate metabolic

pathways for D- and L-lysine, and why L-lysine metabolism seems dependent on the presence of

an intact D-lysine metabolism. Previously work has proposed that the D-lysine pathway may

provide a way of resolving a C/N imbalance that may occur when L-lysine is metabolized.

However we believe this is unlikely as both lysine degradation pathways contain one

deamination and one transamination reaction (11). Our fitness results indicate D-lysine

metabolism is dispensable for growing on 5AVA. This would suggest only the initial two steps

of L-lysine metabolism, the oxidation of lysine to 5-aminopentanamide by DavB and its

subsequent deamination to 5AVA by DavA are dependent on D-lysine catabolism. We propose

the adjacent AsnC family regulator PP_0384 likely responds to L-lysine as many proteins within

this family respond to amino acids including lysine (37, 38) and expression of these two enzymes

is most responsive to L-lysine. To our knowledge there has been no rigorous characterization of

the regulation of the *davAB* operon, nor of the biochemical activities of these two enzymes *in*

vitro. Future studies to uncover the mechanistic regulation at the transcriptional and post-

translational levels at these two steps may uncover the necessity of D-lysine dependency of the

L-lysine catabolic pathway. Overall our work highlights the utility of global fitness profiling to

discover novel, complex, metabolic pathways in even well-characterized bacteria.

Materials and Methods

<u>Media, chemicals, and culture conditions</u>

Routine bacterial cultures were grown in in Luria-Bertani (LB) Miller medium (BD

Biosciences, USA). *E. coli* was grown at 37 °C, while *P. putida* was grown at 30 °C unless

otherwise noted. When indicated, *P. putida* was grown on modified MOPS minimal medium

(39). Cultures were supplemented with kanamycin (50 mg/L, Sigma Aldrich, USA), gentamicin

(30 mg/L, Fisher Scientific, USA), or carbenicillin (100mg/L, Sigma Aldrich, USA), when

indicated. D-2AA was purchased from Takara Bioscience (USA), all other compounds were

purchased through Sigma Aldrich.

<u>Strains and plasmids</u>

All bacterial strains and plasmids used in this work are listed in Supplemental Table 1.

All strains and plasmids created in this work are available through the public instance of the

JBEI registry. (https://public-registry.jbei.org/folders/391). All plasmids were designed using

Device Editor and Vector Editor software, while all primers used for the construction of

plasmids were designed using j5 software (40–42). Synthetic DNA coding for the *Halobacillus*

sp. BAB-2008 *csiD* homolog was purchased from Integrated DNA Technologies (IDT,

Coralville, IA). Plasmids were assembled via Gibson Assembly using standard protocols (43), or

Golden Gate Assembly using standard protocols (44). Plasmids were routinely isolated using the

Qiaprep Spin Miniprep kit (Qiagen, USA), and all primers were purchased from Integrated DNA

Technologies (IDT, Coralville, IA).

<u>Random barcode TnSeq experiments</u>

P. putida RB-TnSeq library JBEI-1 was created by diluting a 1 mL aliquot of the

previously described *P. putida* RB-TnSeq library (16) in 500 mL of LB media supplemented

with kanamycin which was then grown to an OD_{600} of 0.5 and frozen as 1 mL aliquots after

adding glycerol to a final concentration of 20% v/v. Libraries were stored at −80 °C until used. A

1 mL aliquot of *P. putida* RB-TnSeq library JBEI-1 was thawed on ice and diluted in 25 mL of

LB supplemented with kanamycin. The culture was grown until it reached an OD_{600} of 0.5, at

which point 3 1-mL aliquots were taken, pelleted, decanted, and then stored at −80 °C to use as a

time zero control. The library was then washed once in MOPS minimal medium without any

carbon source, and then diluted 1:50 into 10 mL MOPS minimal medium supplemented with

either 10 mM glucose, 5AVA, D-lysine, or L-lysine. Cells were grown in 50 mL culture tubes

for 48 hours at 30 °C shaking at 200 rpm. After growth 2 ml aliquots from the culture tubes were

pelleted, decanted and frozen at −80 °C for barcode sequencing. We performed DNA barcode

sequencing (BarSeq) as previously described (14, 16). The fitness of a strain is the normalized

log2 ratio of barcode reads in the experimental sample to barcode reads in the time zero sample.

The fitness of a gene is the weighted average of the strain fitness for insertions in the central 10–

90% of the gene. The gene fitness values are normalized so the typical gene has a fitness of zero.

The primary statistic *t*-value is of the form of fitness divided by the estimated variance across

different mutants of the same gene. Statistic *t*-values of >|4| were considered significant. All

experiments described herein pass the quality metrics described previously unless noted

otherwise. All fitness data in this work is publically available at http://fit.genomics.lbl.gov.

Construction of deletion mutants

Deletion mutants in *P. putida* were constructed by homologous recombination and *sacB*

counterselection using the allelic exchange vector pMQ30 (45). Briefly, homology fragments

ranging from 1 kbp to 2 kbp up- and downstream of the target gene, including the start and stop

codons respectively, were cloned into pMQ30. An exception to these design parameters was

plasmid pMQ-PP_5260 which maintained an additional 21 nt at the 5' end of the gene in

addition to the stop codon. Plasmids were then transformed via electroporation into *E. coli* S17

and then mated into *P. putida* via conjugation. Transconjugants were selected for on LB agar

plates supplemented with gentamicin 30 mg/mL, and chloramphenicol 30 mg/mL.

Transconjugants were then grown overnight on LB media also supplemented with 30 mg/mL

gentamicin, and 30 mg/mL chloramphenicol, and then plated on LB agar with no NaCl

supplemented with 10% w/v sucrose. Putative deletions were restreaked on LB agar with no

NaCl supplemented with 10% w/v sucrose, and then were screened via PCR with primers

flanking the target gene to confirm gene deletion.

Plate based growth assays

Growth studies of bacterial strains were conducted a microplate reader kinetic assays.

Overnight cultures were inoculated into 10 mL of LB medium from single colonies, and grown

at 30 °C. These cultures were then washed twice with MOPS minimal media without any added

carbon and diluted 1:100 into 500 uL of MOPS medium with 10 mM of a carbon source in 48-

well plates (Falcon, 353072). Plates were sealed with a gas-permeable microplate adhesive film

(VWR, USA), and then optical density was monitored for 48 hours in an Biotek Synergy 4 plate

reader (BioTek, USA) at 30 ℃ with fast continuous shaking. Optical density was measured at

600 nm and all OD_{600} measurements are reported without pathlength corrections.

<u>Expression and purification of proteins</u>

A 5 mL overnight culture of *E. coli* BL21 (DE3) containing the expression plasmid was

used to inoculate a 500 mL culture of LB. Cells were grown at 37 ℃ to an OD_{600} of 0.6 then

induced with isopropyl β-D-1-thiogalactopyranoside to a final concentration of 1 mM. The

temperature was lowered to 30 ℃ and cells were allowed to express for 18 hours before being

harvested via centrifugation. Cell pellets were stored at -80 ℃ until purification. For purification,

cell pellets were resuspended in lysis buffer (50 mM sodium phosphate, 300 mM sodium

chloride, 10 mM imidazole, 8% glycerol, pH 7.5) and sonicated to lyse cells. Insolubles were

pelleted via centrifugation (30 minutes at 40,000xg). The supernatant was applied to a fritted

column containing Ni-NTA resin (Qiagen, USA) which had been pre-equilibrated with several

column volumes of lysis buffer. The resin was washed with lysis buffer containing 50 mM

imidazole, then the protein was eluted using a step-wise gradient of lysis buffer containing

increasing imidazole concentrations (100 mM, 200 mM, and 400 mM). Fractions were collected

and analyzed via SDS-PAGE. Purified protein was dialyzed overnight at 4 ˚C against 50 mM

HEPES pH 7.5, 5% glycerol.

<u>CsiD *in vitro* assays</u>

The activity of purified CsiD homologs was analyzed in 100 µL reaction mixtures

containing 50 mM HEPES (pH 7), 5 mM glutarate, 5 mM 2KG, 25 µM $FeSO_4$, 0.1 mM

ascorbate, and 0.5 mM dithiothreitol. For negative control reactions, each respective reaction

component was omitted. To initiate reactions, CsiD was added to a final concentration of 7 µM.

For the no enzyme control, CsiD was denatured at 98 °C for 10 minutes prior to addition to the

reaction mix. Reactions were allowed to proceed at 22 °C for 3 hours. Products were analyzed

via LC-TOF method 3 after quenching via the addition of acetonitrile and methanol for a final

ACN:H_2O:MeOH ratio of 6:3:1 To analyze products from substrate range as well 2-oxoacid

specificity experiments, reactions were measured via LC-TOF method 1.

Transamination assays

To determine product formation via PP_4108, assays were conducted in 50 mM HEPES

(pH 7.5), with 5 mM 2KG, 0.1 mM PLP, and 5 mM of substrate, and 10 µM of purified enzyme

or boiled enzyme control in 100µL volumes. Reactions were incubated at 30 °C for 16 hours and

quenched via the addition of ACN and MeOH for a final ACN:H_2O:MeOH ratio of 6:3:1 for LC-

TOF method 3. To determine substrate specificity reactions were set up at 75 µL scale and

carried out at 30°C for up to 16 hours before freezing. For analysis, reactions were diluted 15-

fold in water and assessed by a colorimetric assay for glutamate (Sigma MAK004) in 96-well

format via a SpectraMax M4 plate reader (Molecular Devices, USA).

PP_5260 *in vitro* assays

The activity of PP_5260 was initially assessed in 50 mM HEPES, with 5 mM 2OA as

substrate and 10 µM purified enzyme or boiled enzyme control. Reactions were incubated for 16

hours at 30 °C. To test the necessity of metal cofactors EDTA was added to a final concentration

of 50 µM. Reactions and quenched via the addition of ACN and methanol MeOH for a final

ACN:H_2O:MeOH ratio of 6:3:1 for LC-TOF analysis method 3, or with an equal volume of ice-

cold methanol for HPLC analysis and LC-TOF method 2.

To determine the metal cofactor, after purification over Ni-NTA resin the protein was

concentrated and dialyzed overnight against 50mM HEPES, 100mM NaCl, pH 7.5. To generate

apo-enzyme, the protein was then dialyzed four times at a protein:dialysis buffer ratio of 1:300

against the same buffer containing 5mM EDTA in order to remove any bound metal. The

enzyme was dialyzed once more against buffer without EDTA overnight in order to remove any

remaining chelating reagent. The apo-enzyme was then assayed in the presence of 50 μM of a

variety of potential metal cofactors in 50 mM HEPES with 10 mM 2OA as substrate and 10 μM

purified enzyme. Reactions were incubated for 30 minutes at 30 °C and activity was assessed via

HPLC.

Determination of enzyme stoichiometry was assessed in 50 mM HEPES, 50 μM $FeCl_2$,

with 1 mM 2OA as substrate and 0.1 μM purified enzyme or boiled enzyme control. Reactions

were incubated for 5 minutes at 30 °C and then quenched with an equal volume of ice-cold

methanol then quantified with LC-TOF method 2.

Enzyme coupled decarboxylation assays were carried out as previously described (20).

Reaction mixtures contained 100 mM Tris-HCl (pH 7), 10 mM $MgCl_2$, 0.4 mM NADH, 4 mM,

50 μM $FeCl_2$, phosphoenol pyruvate (PEP), 100U/mL pig heart malate dehydrogenase(Roche),

2U/mL microbial PEP carboxylase (Sigma), and 10 mM 2OA. Reactions were initiated by the

addition of purified PP_5260 or boiled enzyme controls, and absorbance at 340 nm was

measured via a SpectraMax M4 plate reader (Molecular Devices, USA). Michaelis-Menten

behavior was formulated as previously described (46). Least-squares minimization was used to

derive K_m and K_{cat}. Determination of D-2HG concentration was assayed with a D-2-

Hydroxyglutarate (D-2HG) Assay Kit (Sigma MAK320).

Acknowledgements

This manuscript is dedicated to the memory of Cornell Professor Dr. Eugene Madsen. The

authors would like to thank Morgan Price, Dr. John Hangasky, Dr. Jamie Meadows, Dr. Robert

Haushalter, Dr. Bo Pang, Dr. Nick Weathersby, Mary Thompson, and Catharine Adams for their

helpful discussions in preparing this manuscript. We would also like to thank the UC Berkeley

SMART program for providing support for R.K to conduct summer research. This work was part

of the DOE Joint BioEnergy Institute (https://www.jbei.org) supported by the U.S. Department

of Energy, Office of Science, Office of Biological and Environmental Research, and protein

purification and homology modelling components were part of the Agile BioFoundry

(http://agilebiofoundry.org) supported by the U.S. Department of Energy, Energy Efficiency and

Renewable Energy, Bioenergy Technologies Office, through contract DE-AC02-05CH11231

between Lawrence Berkeley National Laboratory and the U.S. Department of Energy. The views

and opinions of the authors expressed herein do not necessarily state or reflect those of the

United States Government or any agency thereof. Neither the United States Government nor any

agency thereof, nor any of their employees, makes any warranty, expressed or implied, or

assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any

information, apparatus, product, or process disclosed, or represents that its use would not

infringe privately owned rights.

Contributions

Conceptualization, M.G.T., and J.M.B.; Methodology, M.G.T., J.M.B, J.F.B., P.C.M., S.C.C.,

N.C.H, C.B.E, E.E.K.B, C.J.P., and A.M.D.; Investigation, M.G.T., J.M.B, W.N.S., R.A.K,

J.F.B., V.T.B, P.C.M., J.W.G, C.J.P, N.C.H., F.F.T., J.H.P W.S., E.E.K.B.; Writing – Original

Draft, M.G.T.; Writing – Review and Editing, All authors.; Resources and supervision, P.D.A.,

A.P.A., A.M.D., and J.D.K.

Competing Interests

J.D.K. has financial interests in Amyris, Lygos, Constructive Biology, Demetrix, Napigen and

Maple Bio.

Table 1

Strain	JBEI Part ID	Reference	Genotype
E. coli DH10B		ThermoFisher	
E. coli S17		ATCC 47055	
E. coli BL21(DE3)		Novagen	
P. putida KT2440		ATCC 47054	
P. putida ΔPP_0158	JPUB_010967	This work	
P. putida ΔPP_2088	JPUB_013224	This work	
P. putida ΔPP_2909	JPUB_010968	This work	
P. putida ΔPP_2910	JPUB_010969	This work	
P. putida ΔPP_0158 ΔPP_2909	JPUB_010970	This work	
P. putida ΔPP_4108	JPUB_010971	This work	
P. putida ΔPP_4493	JPUB_010972	This work	
P. putida ΔPP_5260	JPUB_010973	This work	
Plasmids			
pMQ30		45	Gm, SacB
pET28		Novagen	Kan
pET21b		Novagen	Amp
pMQ30-PP_0158	JPUB_010989	This work	Gm, SacB
pMQ30-PP_2088	JPUB_013222	This work	Gm, SacB
pMQ30-PP_2909	JPUB_010991	This work	Gm, SacB
pMQ30-PP_2910	JPUB_010995	This work	Gm, SacB
pMQ30-PP_4108	JPUB_010981	This work	Gm, SacB
pMQ30-PP_4493	JPUB_010979	This work	Gm, SacB
pMQ30-PP_5260	JPUB_010977	This work	Gm, SacB

pET28-CsiD_Halo	JPUB_010987	This work	Kan
pET28-CsiD_Ecoli	JPUB_010993	This work	Kan
pET28-CsiD_Pput	JPUB_010975	This work	Kan
pET21b-PP_4108	JPUB_010983	This work	Amp
pET21b-PP_5260	JPUB_010985	This work	Amp

References

1. Nelson KE, Weinel C, Paulsen IT, Dodson RJ, Hilbert H, Martins dos Santos VAP, Fouts

DE, Gill SR, Pop M, Holmes M, Brinkac L, Beanan M, DeBoy RT, Daugherty S, Kolonay

J, Madupu R, Nelson W, White O, Peterson J, Khouri H, Fraser CM. 2002. Complete

genome sequence and comparative analysis of the metabolically versatile Pseudomonas

putida KT2440. Environ Microbiol 4:799–808.

2. Nikel PI, Chavarría M, Danchin A, de Lorenzo V. 2016. From dirt to industrial

applications: Pseudomonas putida as a Synthetic Biology chassis for hosting harsh

biochemical reactions. Curr Opin Chem Biol 34:20–29.

3. Loeschcke A, Thies S. 2015. Pseudomonas putida-a versatile host for the production of

natural products. Appl Microbiol Biotechnol 99:6197–6214.

4. Jiménez JI, Miñambres B, García JL, Díaz E. 2002. Genomic analysis of the aromatic

catabolic pathways from Pseudomonas putida KT2440. Environ Microbiol 4:824–841.

5. Ragauskas AJ, Beckham GT, Biddy MJ, Chandra R, Chen F, Davis MF, Davison BH,

Dixon RA, Gilna P, Keller M, Langan P, Naskar AK, Saddler JN, Tschaplinski TJ, Tuskan

GA, Wyman CE. 2014. Lignin valorization: improving lignin processing in the

biorefinery. Science 344:1246843.

6.   Chang YF, Adams E. 1971. Induction of separate catabolic pathways for L- and D-lysine

in Pseudomonas putida. Biochem Biophys Res Commun 45:570–577.

7.   Chae TU, Ko Y-S, Hwang K-S, Lee SY. 2017. Metabolic engineering of Escherichia coli

for the production of four-, five- and six-carbon lactams. Metab Eng 41:82–91.

8.   Zhang J, Barajas JF, Burdu M, Wang G, Baidoo EE, Keasling JD. 2017. Application of an

Acyl-CoA Ligase from Streptomyces aizunensis for Lactam Biosynthesis. ACS Synth Biol

6:884–890.

9.   Zhang M, Gao C, Guo X, Guo S, Kang Z, Xiao D, Yan J, Tao F, Zhang W, Dong W, Liu

P, Yang C, Ma C, Xu P. 2018. Increased glutarate production by blocking the glutaryl-

CoA dehydrogenation pathway and a catabolic pathway involving L-2-hydroxyglutarate.

Nat Commun 9:2114.

10. Kim HT, Khang TU, Baritugo K-A, Hyun SM, Kang KH, Jung SH, Song BK, Park K, Oh

M-K, Kim GB, Kim HU, Lee SY, Park SJ, Joo JC. 2018. Metabolic engineering of

Corynebacterium glutamicum for the production of glutaric acid, a C5 dicarboxylic acid

platform chemical. Metab Eng 51:99–109.

11. Revelles O, Espinosa-Urgel M, Fuhrer T, Sauer U, Ramos JL. 2005. Multiple and

interconnected pathways for L-lysine catabolism in Pseudomonas putida KT2440. J

Bacteriol 187:7500–7510.

12. Revelles O, Wittich R-M, Ramos JL. 2007. Identification of the initial steps in D-lysine

catabolism in Pseudomonas putida. J Bacteriol 189:2787–2792.

13.   Perfetti R, Campbell RJ, Titus J, Hartline RA. 1972. Catabolism of pipecolate to glutamate

676   in Pseudomonas putida. J Biol Chem 247:4089–4095.

677   14.   Wetmore KM, Price MN, Waters RJ, Lamson JS, He J, Hoover CA, Blow MJ, Bristow J,

678   Butland G, Arkin AP, Deutschbauer A. 2015. Rapid quantification of mutant fitness in

679   diverse bacteria by sequencing randomly bar-coded transposons. MBio 6:e00306-15.

680   15.   Price MN, Wetmore KM, Waters RJ, Callaghan M, Ray J, Liu H, Kuehl JV, Melnyk RA,

681   Lamson JS, Suh Y, Carlson HK, Esquivel Z, Sadeeshkumar H, Chakraborty R, Zane GM,

682   Rubin BE, Wall JD, Visel A, Bristow J, Blow MJ, Deutschbauer AM. 2018. Mutant

683   phenotypes for thousands of bacterial genes of unknown function. Nature 557:503–509.

684   16.   Rand JM, Pisithkul T, Clark RL, Thiede JM, Mehrer CR, Agnew DE, Campbell CE,

685   Markley AL, Price MN, Ray J, Wetmore KM, Suh Y, Arkin AP, Deutschbauer AM,

686   Amador-Noguez D, Pfleger BF. 2017. A metabolic pathway for catabolizing levulinic acid

687   in bacteria. Nat Microbiol 2:1624–1634.

688   17.   Okuno E, Tsujimoto M, Nakamura M, Kido R. 1993. 2-Aminoadipate-2-oxoglutarate

689   aminotransferase isoenzymes in human liver: a plausible physiological role in lysine and

690   tryptophan metabolism. Enzyme Protein 47:136–148.

691   18.   Wilding M, Peat TS, Newman J, Scott C. 2016. A β-Alanine Catabolism Pathway

692   Containing a Highly Promiscuous ω-Transaminase in the 12-Aminododecanate-Degrading

693   Pseudomonas sp. Strain AAC. Appl Environ Microbiol 82:3846–3856.

694   19.   Kopchick JJ, Hartline RA. 1979. alpha-Hydroxyglutarate as an intermediate in the

695   catabolism of alpha-aminoadipate by Pseudomonas putida. J Biol Chem 254:3259–3263.

696   20.   Witkowski A, Joshi AK, Smith S. 2002. Mechanism of the β-Ketoacyl Synthase Reaction

697   Catalyzed by the Animal Fatty Acid Synthase[†]. Biochemistry 41:10877–10887.

698   21.   Martinez S, Hausinger RP. 2016. Biochemical and Spectroscopic Characterization of the

699   Non-Heme Fe(II)- and 2-Oxoglutarate-Dependent Ethylene-Forming Enzyme from

700   Pseudomonas syringae pv. phaseolicola PK2. Biochemistry 55:5989–5999.

701   22.   Martinez S, Hausinger RP. 2015. Catalytic Mechanisms of Fe(II)- and 2-Oxoglutarate-

702   dependent Oxygenases. J Biol Chem 290:20702–20711.

703   23.   Revelles O, Espinosa-Urgel M, Molin S, Ramos JL. 2004. The davDT operon of

704   Pseudomonas putida, involved in lysine catabolism, is induced in response to the pathway

705   intermediate delta-aminovaleric acid. J Bacteriol 186:3439–3446.

706   24.   Danhauser K, Sauer SW, Haack TB, Wieland T, Staufner C, Graf E, Zschocke J, Strom

707   TM, Traub T, Okun JG, Meitinger T, Hoffmann GF, Prokisch H, Kölker S. 2012.

708   DHTKD1 mutations cause 2-aminoadipic and 2-oxoadipic aciduria. Am J Hum Genet

709   91:1082–1087.

710   25.   REITZ M, RODWELL V. 1969. a-Hydroxyglutarate Oxidoreductase of Pseudomonas

711   putida. J Bacteriol.

712   26.   Wójcik A, Broclawik E, Siegbahn PEM, Borowski T. 2012. Mechanism of benzylic

713   hydroxylation by 4-hydroxymandelate synthase. A computational study. Biochemistry

714   51:9570–9580.

715   27.   Di Giuro CML, Konstantinovics C, Rinner U, Nowikow C, Leitner E, Straganz GD. 2013.

716   Chiral hydroxylation at the mononuclear nonheme Fe(II) center of 4-(S) hydroxymandelate

717   synthase--a structure-activity relationship analysis. PLoS ONE 8:e68932.

28.   Lequette Y, Odberg-Ferragut C, Bohin J-P, Lacroix J-M. 2004. Identification of mdoD, an

mdoG paralog which encodes a twin-arginine-dependent periplasmic protein that controls

osmoregulated periplasmic glucan backbone structures. J Bacteriol 186:3695–3702.

29.   Zhang L, Ren Y, Lu B, Yang C, Feng Z, Liu Z, Chen J, Ma W, Wang Y, Yu X, Wang Y,

Zhang W, Wang Y, Liu S, Wu F, Zhang X, Guo X, Bao Y, Jiang L, Wan J. 2016.

FLOURY ENDOSPERM7 encodes a regulator of starch synthesis and amyloplast

development essential for peripheral endosperm development in rice. J Exp Bot 67:633–

647.

30.   Vijayakumar SRV, Kirchhof MG, Patten CL, Schellhorn HE. 2004. RpoS-regulated genes

of Escherichia coli identified by random lacZ fusion mutagenesis. J Bacteriol 186:8499–

8507.

31.   Knorr S, Sinn M, Galetskiy D, Williams RM, Wang C, Müller N, Mayans O, Schleheck D,

Hartig JS. 2018. Widespread bacterial lysine degradation proceeding via glutarate and L-2-

hydroxyglutarate. Nat Commun 9:5071.

32.   Nishijyo T, Haas D, Itoh Y. 2001. The CbrA-CbrB two-component regulatory system

controls the utilization of multiple carbon and nitrogen sources in Pseudomonas

aeruginosa. Mol Microbiol 40:917–931.

33.   Valentini M, García-Mauriño SM, Pérez-Martínez I, Santero E, Canosa I, Lapouge K.

2014. Hierarchical management of carbon sources is regulated similarly by the CbrA/B

systems in Pseudomonas aeruginosa and Pseudomonas putida. Microbiology (Reading,

Engl) 160:2243–2252.

34.   Bouffartigues E, Gicquel G, Bazire A, Bains M, Maillot O, Vieillard J, Feuilloley MGJ,

Orange N, Hancock REW, Dufour A, Chevalier S. 2012. Transcription of the oprF gene of

Pseudomonas aeruginosa is dependent mainly on the SigX sigma factor and is sucrose

induced. J Bacteriol 194:4301–4311.

35.   Blanka A, Schulz S, Eckweiler D, Franke R, Bielecka A, Nicolai T, Casilag F, Düvel J,

Abraham W-R, Kaever V, Häussler S. 2014. Identification of the alternative sigma factor

SigX regulon and its implications for Pseudomonas aeruginosa pathogenicity. J Bacteriol

196:345–356.

36.   Neshich IAP, Kiyota E, Arruda P. 2013. Genome-wide analysis of lysine catabolism in

bacteria reveals new connections with osmotic stress resistance. ISME J 7:2400–2410.

37.   Brinkman AB, Bell SD, Lebbink RJ, de Vos WM, van der Oost J. 2002. The Sulfolobus

solfataricus Lrp-like protein LysM regulates lysine biosynthesis in response to lysine

availability. J Biol Chem 277:29537–29549.

38.   Thaw P, Sedelnikova SE, Muranova T, Wiese S, Ayora S, Alonso JC, Brinkman AB,

Akerboom J, van der Oost J, Rafferty JB. 2006. Structural insight into gene transcriptional

regulation and effector binding by the Lrp/AsnC family. Nucleic Acids Res 34:1439–1449.

39.   LaBauve AE, Wargo MJ. 2012. Growth and laboratory maintenance of Pseudomonas

aeruginosa. Curr Protoc Microbiol Chapter 6:Unit 6E.1.

40.   Ham TS, Dmytriv Z, Plahar H, Chen J, Hillson NJ, Keasling JD. 2012. Design,

implementation and practice of JBEI-ICE: an open source biological part registry platform

and tools. Nucleic Acids Res 40:e141.

41.   Chen J, Densmore D, Ham TS, Keasling JD, Hillson NJ. 2012. DeviceEditor visual

biological CAD canvas. J Biol Eng 6:1.

42. Hillson NJ, Rosengarten RD, Keasling JD. 2012. j5 DNA assembly design automation

software. ACS Synth Biol 1:14–21.

43. Gibson DG, Young L, Chuang R-Y, Venter JC, Hutchison CA, Smith HO. 2009.

Enzymatic assembly of DNA molecules up to several hundred kilobases. Nat Methods

6:343–345.

44. Engler C, Kandzia R, Marillonnet S. 2008. A one pot, one step, precision cloning method

with high throughput capability. PLoS ONE 3:e3647.

45. Shanks RMQ, Kadouri DE, MacEachran DP, O'Toole GA. 2009. New yeast

recombineering tools for bacteria. Plasmid 62:88–97.

46. Cornish-Bowden A. 2012. Fundamentals of Enzyme Kinetics, 4th ed. Wiley, John & Sons,

Incorporated.

Massively parallel fitness profiling reveals multiple novel enzymes in *Pseudomonas putida* lysine metabolism

Mitchell G. Thompson[1,2,3], Jacquelyn M. Blake-Hedges[1,2,4], Pablo Cruz-Morales[1,2,5], Jesus F. Barajas[2,6], Samuel C. Curran[1,2,7], Christopher B. Eiben[1,2,8], Nicholas C. Harris[3], Veronica T. Benites[1,2], Jennifer W. Gin[1,2], William A. Sharpless[1,2,3], Frederick F. Twigg[8], Will Skyrud[4], Rohith N. Krishna[1,2,4], Jose Henrique Pereira[1,9], Edward E. K. Baidoo[1,2], Christopher J. Petzold[1,2], Paul D. Adams[1,9,10], Adam P. Arkin[10,11], Adam M. Deutschbauer[3,11], Jay D. Keasling[1,2,8,10,12]

[1]Joint BioEnergy Institute, 5885 Hollis Street, Emeryville, CA 94608, USA.

[2]Biological Systems & Engineering Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA.

[3]Department of Plant and Microbial Biology, University of California, Berkeley, CA 94720, USA

[4]Department of Chemistry, University of California, Berkeley, CA 94720, USA

[5]Centro de Biotecnología FEMSA, Tecnológico de Monterrey, NL, Mexico

[6]Department of Energy Agile BioFoundry, Emeryville, CA 94608, USA

[7]Comparative Biochemistry Graduate Group, University of California, Berkeley, CA 94720, USA

[8]Department of Chemical and Biomolecular Engineering, University of California, Berkeley, CA 94720, USA

[9]Molecular Biophysics and Integrated Bioimaging Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA.

[10]Joint Program in Bioengineering, University of California, Berkeley/San Francisco, CA 94720, USA

[11]Environmental Genomics and Systems Biology Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA.

[12]The Novo Nordisk Foundation Center for Biosustainability, Technical University of Denmark, Denmark

Supplementary Materials and Methods

HPLC analysis

HPLC analysis was performed on an Agilent Technologies 1200 series liquid chromatography instrument coupled to a refractive index detector (35°C, Agilent Technologies, Santa Clara, CA). Samples were injected onto an Aminex HPX-87H Ion Exclusion Column (300 x 7.8 mm, 60°C, Bio-Rad, Hercules, CA) and eluted isocratically with 4 mM H_2SO_4 at 600 uL/min for 20 minutes. Compounds were quantified via comparison to a calibration curve prepared with authentic standards and normalized to injection volume.

Proteomics analysis

$P. putida$ KT2440 wild type was grown on MOPS minimal media with 10 mM of either glucose, L-lysine, D-lysine, 5AVA, 2AA, or glutarate. Cells were harvested when cultures reached an OD_{600} of 1.0 with a 1 cm pathlength. Cell lysis and protein precipitation were achieved by using a chloroform-methanol extraction as previously described (1). Thawed pellets were loosened from 14 mL falcon tubes and transferred to PCR 8-well tube strip, followed by the

addition of 80 µL of methanol, 20 µL of chloroform, and 60 µL of water, with vortexing. The samples were centrifuged at ~20,000 x g for 1 minute for phase separation. The methanol and water (top) layer was removed, then 100 µL of methanol was added and the sample was vortexed briefly. The samples were centrifuged at ~20,000 x g for 1 minute to isolate the protein pellet. The protein pellet was air-dried for 10 minutes and resuspended in 100 mM ammonium bicarbonate with 20% methanol. The protein concentration was measured using the DC Protein Assay Kit (Bio-Rad, Hercules, CA) with bovine serum albumin for the standard curve. A total of 100 µg of protein from each sample was digested with trypsin for targeted proteomic analysis. The protein was reduced by adding tris 2-(carboxyethyl) phosphine (TCEP) at a final concentration of 5 mM, alkylated by adding iodoacetamide at a final concentration of 10 mM, and digested overnight at 37 ℃ with trypsin at a ratio of 1:50 (w/w) trypsin:total protein. As previously described (2), peptides were analyzed using an Agilent 1290 liquid chromatography system coupled to an Agilent 6460QQQ mass spectrometer (Agilent Technologies, Santa Clara, CA). Peptide samples (10 µg) were separated on an Ascentis Express Peptide ES-C18 column (2.7 µm particle size, 160 Å pore size, 50 mm length x 2.1 mm i.d., 60 ℃; Sigma-Aldrich, St. Louis, MO) by using a chromatographic gradient (400 µL/min flow rate) with an initial condition of 95% buffer A (99.9% water, 0.1% formic acid) and 5% buffer B (99.9% acetonitrile, 0.1% formic acid) then increasing linearly to 65% buffer A/35% buffer B over 5.5 minutes. Buffer B was then increased to 80% over 0.3 minutes and held at 80% for two minutes followed by ramping back down to 5% buffer B over 0.5 minutes where it was held for 1.5 minutes to re-equilibrate the column for the next sample. The peptides were ionized by an Agilent Jet Stream ESI source operating in positive-ion mode with the following source parameters: gas Temperature: 250 ℃, gas Flow: 13 L/min, nebulizer pressure: 35 psi, sheath gas temperature:

250 ℃, sheath gas flow: 11 L/min, nozzle voltage: 0 V, chamber voltage: 3,500 V. The data were acquired using Agilent MassHunter, version B.08.02, processed using Skyline (3) version 4.1, and peak quantification was refined with mProphet (4) in Skyline. Data are available at Panorama Public via this link:

https://panoramaweb.org/massive_fitness_profiling_Pseudomonas_putida.url. All pairwise combinations of spectral counts from carbon sources for each protein were compared via Student's t-test followed by a Bonferroni correction.

Detection of metabolites

Sampling of intracellular metabolites was conducted as described previously (5). Multiple methods were used to detect compounds in this work. Method (1) HILIC-HRMS analysis was performed using an Agilent Technologies 6510 Accurate-Mass Q-TOF LC-MS instrument using positive mode and an Atlantis HILIC Silica 5 µm column (150 x 4.6 mm) with a linear of 95 to 50% acetonitrile (v/v) over 8 minutes in water with 40 mM ammonium formate, pH 4.5, at a flow rate of 1 mL minute^{-1}. Method (2) HILIC-HRMS analysis was performed using an Agilent Technologies 6510 Accurate-Mass Q-TOF LC-MS instrument using negative mode and an Atlantis HILIC Silica 5 µm column (150 x 4.6 mm) with an isocratic mobile phase (80% acetonitrile (v/v) with 40 mM ammonium formate, pH 4.5) for 20 minute at a flow rate of 1 mL minute^{-1}. Method (3) is described in George et al (5). Briefly, samples were separated via a SeQuantZIC-pHILIC guard column (20-mm length, 2.1-mm internal diameter, and 5-µm particle size; from EMD Millipore, Billerica, MA, USA), then with a short SeQuantZIC-pHILIC column (50-mm length, 2.1-mm internal diameter, and 5-µm particle size) followed by a long SeQuantZIC-pHILIC column (150-mm length, 2.1-mm internal diameter, and 5-µm particle size) using an Agilent Technologies 1200 Series Rapid Resolution HPLC system (Agilent

Technologies, Santa Clara, CA, USA). The mobile phase was composed of 10 mM ammonium carbonate and 118.4 mM ammonium hydroxide in acetonitrile/water (60.2:39.8, v/v). Metabolites were eluted isocratically via a flow rate of 0.18 mL/min from 0 to 5.4 minutes, which was increased to 0.27 mL/min from 5.4 to 5.7 minutes, and held at this flow rate for an additional 5.4 minutes. The HPLC system was coupled to an Agilent Technologies 6210 TOF-MS system in negative mode. Determination of D-2HG concentration was assayed with a D-2-Hydroxyglutarate (D2HG) Assay Kit (Sigma MAK320).

Phylogenomic analyses

Amino acid sequences of CsiD homologs were downloaded from the pFAM database and aligned with MAFFT-linsi (13). Phylogenetic trees of CsiD alignments were constructed with FastTree 2, and trees were visualized on iTOL (14, 15).

Representative DUF1338 sequences were obtained from pFAM (https://pfam.xfam.org/family/PF07063#tabview=tab3). All genomes analyzed were downloaded from the NCBI FTP site and annotated using RAST (16). Amino acid sequences of DUF1338 proteins from these genomes were retrieved using BlastP with a bit score cutoff of 150 and an E-value of 0.000001. All sequences alignments were performed using Muscle v3.8 (17) and the alignments were manually curated using Jalview V2 (18).

For the phylogenetic reconstructions, the best amino acid substitution model was selected using ModelFinder implemented on IQ-tree (19) the phylogenies were obtained using IQ-tree v 1.6.7 (20), with 10,000 bootstrap replicates. The final trees were visualized and annotated using FigTree v1.4.3 (http://tree.bio.ed.ac.uk/software/figtree/). Genome neighborhoods of DUF1338 were obtained using CORASON-BGC (21) and manually colored and annotated.

Statistical analyses and data presentation

All numerical data were analyzed using custom Python scripts. All graphs were visualized using either Seaborn or Matplotlib. Calculation of 95% confidence intervals, standard deviations, and T-test statistics were conducted via the Scipy library. Bonferroni corrections were calculated using the MNE python library (22).

Bibliography

1.   González Fernández-Niño SM, Smith-Moritz AM, Chan LJG, Adams PD, Heazlewood JL, Petzold CJ. 2015. Standard flow liquid chromatography for shotgun proteomics in bioenergy research. Front Bioeng Biotechnol 3:44.

2.   Batth TS, Singh P, Ramakrishnan VR, Sousa MML, Chan LJG, Tran HM, Luning EG, Pan EHY, Vuu KM, Keasling JD, Adams PD, Petzold CJ. 2014. A targeted proteomics toolkit for high-throughput absolute quantification of Escherichia coli proteins. Metab Eng 26:48–56.

3.   MacLean B, Tomazela DM, Shulman N, Chambers M, Finney GL, Frewen B, Kern R, Tabb DL, Liebler DC, MacCoss MJ. 2010. Skyline: an open source document editor for creating and analyzing targeted proteomics experiments. Bioinformatics 26:966–968.

4.   Reiter L, Rinner O, Picotti P, Hüttenhain R, Beck M, Brusniak M-Y, Hengartner MO, Aebersold R. 2011. mProphet: automated data processing and statistical validation for large-scale SRM experiments. Nat Methods 8:430–435.

5.   George KW, Thompson MG, Kim J, Baidoo EEK, Wang G, Benites VT, Petzold CJ, Chan LJG, Yilmaz S, Turhanen P, Adams PD, Keasling JD, Lee TS. 2018. Integrated analysis of isopentenyl pyrophosphate (IPP) toxicity in isoprenoid-producing Escherichia coli. Metab Eng 47:60–72.

6.   Zhang Y. 2008. I-TASSER server for protein 3D structure prediction. BMC Bioinformatics 9:40.

7.   Roy A, Kucukural A, Zhang Y. 2010. I-TASSER: a unified platform for automated protein

structure and function prediction. Nat Protoc 5:725–738.

8. Yang J, Yan R, Roy A, Xu D, Poisson J, Zhang Y. 2015. The I-TASSER Suite: protein structure and function prediction. Nat Methods 12:7–8.

9. Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, Ferrin TE. 2004. UCSF Chimera—a visualization system for exploratory research and analysis. J Comput Chem 25:1605–1612.

10. Adams PD, Afonine PV, Bunkóczi G, Chen VB, Davis IW, Echols N, Headd JJ, Hung L-W, Kapral GJ, Grosse-Kunstleve RW, McCoy AJ, Moriarty NW, Oeffner R, Read RJ, Richardson DC, Richardson JS, Terwilliger TC, Zwart PH. 2010. PHENIX: a comprehensive Python-based system for macromolecular structure solution. Acta Crystallogr D Biol Crystallogr 66:213–221.

11. Grosdidier A, Zoete V, Michielin O. 2011. SwissDock, a protein-small molecule docking web service based on EADock DSS. Nucleic Acids Res 39:W270-7.

12. Grosdidier A, Zoete V, Michielin O. 2011. Fast docking using the CHARMM force field with EADock DSS. J Comput Chem 32:2149–2159.

13. Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. Mol Biol Evol 30:772–780.

14. Price MN, Dehal PS, Arkin AP. 2010. FastTree 2--approximately maximum-likelihood trees for large alignments. PLoS ONE 5:e9490.

15. Letunic I, Bork P. 2016. Interactive tree of life (iTOL) v3: an online tool for the display and annotation of phylogenetic and other trees. Nucleic Acids Res 44:W242-5.

16. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Zagnitko O. 2008. The RAST Server: rapid annotations using subsystems technology. BMC Genomics 9:75.

17. Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high

throughput. Nucleic Acids Res 32:1792–1797.

18. Waterhouse AM, Procter JB, Martin DMA, Clamp M, Barton GJ. 2009. Jalview Version 2--a multiple sequence alignment editor and analysis workbench. Bioinformatics 25:1189–1191.

19. Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. 2017. ModelFinder: fast model selection for accurate phylogenetic estimates. Nat Methods 14:587–589.

20. Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol 32:268–274.

21. Cruz-Morales P, Ramos-Aboites HE, Licona-Cassani C, Selem-Mójica N, Mejía-Ponce PM, Souza-Saldívar V, Barona-Gómez F. 2017. Actinobacteria phylogenomics, selective isolation from an iron oligotrophic environment and siderophore functional characterization, unveil new desferrioxamine traits. FEMS Microbiol Ecol 93.

22. Gramfort A, Luessi M, Larson E, Engemann DA, Strohmeier D, Brodbeck C, Parkkonen L, Hämäläinen MS. 2014. MNE software for processing MEG and EEG data. Neuroimage 86:446–460.

A

gene	D-Lysine	L-Lysine	5-AVA	Glucose	
davD	-1.3	-2.5	-3	0	succinate-semialdehyde dehydrogenase (NADP+)
davT	-0.77	-2.7	-2.6	0.28	4-aminobutyrate aminotransferase
glcB	0.04	-2.8	-1.9	0.07	malate synthase G
davA	-1.2	-3.2	-0.01	0.16	5-aminopentanamidase
davB	-1.5	-2.5	0.08	-0.15	Lysine 2-monooxygenase
PP_0384	-0.54	-2.5	-0.02	-0.09	Transcriptional regulator, AsnC family
PP_0787	-3.2	-3.6	-3.3	-0.46	quinolinate phosphoribosyltransferase (decarboxylating) monomer
PP_0949	-1.4	-2.9	0.5	-0.18	putative ATP-binding protein UPF0042
PP_1109	-2.2	-0.35	-1.1	-0.15	Transcriptional regulator, GntR family
PP_1673	-0.24	-2.1	0.06	-0.38	cobyrinate a,c-diamide synthase
PP_1941	-0.8	-2	-0.36	-0.07	conserved protein of unknown function
rpoX	-4.4	0.13	-0.5	-0.02	RNA polymerase sigma factor SigX
PP_2326	-0.25	-2.7	-0.48	-0.1	Universal stress protein
PP_2868	0.05	-2.2	-0.36	-0.18	Transcriptional regulator, Cro/CI family
csiD	-0.28	-0.21	-2.2	-0.04	carbon starvation induced protein
lghO	-0.09	-4.4	-3.5	-0.06	L-2-hydroxyglutarate oxidase
dpkA	-6.3	-5.7	0.48	-0.36	Delta 1-piperideine-2-carboxylate reductase
PP_3592	-4.1	-4.4	0.46	-0.01	Transcriptional regulator, RpiR family
PP_3593	-4.9	-5	-0.09	-0.15	Amino acid ABC transporter, periplasmic binding protein
PP_3594	-5.5	-4.7	-0.11	-0.08	Amino acid ABC transporter, membrane protein
PP_3595	-4	-3.5	-0.2	0.1	Amino acid ABC transporter, membrane protein
amaD	-3.7	-3	0.14	-0.4	D-lysine oxidase
PP_3597	-4.3	-6.6	0.25	0.12	Amino-acid ABC transporter, ATP-binding protein
PP_3603	-1.6	-3.1	2.4	-0.45	Transcriptional regulator, GntR family
PP_3649	-0.56	-4	-0.65	0.28	Transcriptional regulator, GntR family
PP_3790	0.14	0.13	-2	-0.38	diaminopimelate epimerase
PP_3821	0.81	0.83	-2.1	-0.11	UTP-glucose-1-phosphate uridylyltransferase
PP_3924	-0.38	-3.7	0.11	-0.15	conserved protein of unknown function
PP_4108	-5.9	-1.2	-0.18	0.21	putative 4-aminobutyrate aminotransferase
aceA	-0.12	-2.1	-1.1	0.19	isocitrate lyase
PP_4482	1.8	-3.4	-0.54	-0.17	Transcriptional regulator, AraC family
PP_4493	-5.4	-2.7	-0.37	-0.07	putative oxidoreductase
PP_4695	-2.7	-3.4	-0.49	-0.49	Sensory box histidine kinase
PP_4829	-0.55	-2.4	-0.34	0.02	putative precorrin-3B synthase
PP_4937	-1.4	-2.8	-0.25	-0.24	toluene tolerance protein
maeB	-4.7	-1.6	2.2	0.22	malic enzyme B
amaA	-4.7	-4	0.14	-0.08	L-pipecolate oxidase
amaB	-4.4	-4.5	-0.55	0.04	L-piperidine-6-carboxylate dehydrogenase
PP_5260	-2.4	-6.1	-0.25	-0.03	metalloprotein, putative enzyme

B

C

A

B

	D-Lysine	L-Lysine	5AVA	Glucose	
PP_0138	3	1.7	2.8	0.11	conserved membrane protein of unknown function
PP_0406	1	0.76	2.3	-0.61	Nucleotidyltransferase family protein
mpl	3.8	-0.37	3.1	0.079	UDP-N-acetylmuramate:L-alanyl-gamma-D-glutamyl- meso-diaminopimelate ligase
PP_0725	0.64	3.2	-0.57	0.18	conserved hypothetical TPR repeat protein
dapA-I	2.5	3.1	0.39	-0.92	4-hydroxy-tetrahydrodipicolinate synthase
rpoS	4	5.4	3.2	0.43	RNA polymerase, sigma S (sigma 38) factor
PP_1778	4.1	2.1	2.8	0.14	Lipopolysaccharide ABC export system, permease protein
PP_1779	4	1.6	2.5	0.24	Lipopolysaccharide ABC export system, ATP-binding protein
PP_2144	0.99	2.4	1.7	-0.12	Transcriptional regulator, TetR family
PP_3603	-1.6	-3.1	2.4	-0.45	Transcriptional regulator, GntR family
PP_3731	-0.77	2.6	-0.11	0.11	Transcriptional regulator, TetR family
astA-II	-0.16	4	-0.15	-0.14	arginine N-succinyltransferase, subunit alpha
PP_5058	0.91	3.1	0.54	0.36	Carboxy-terminal-processing protease
maeB	-4.7	-1.6	2.2	0.22	malic enzyme B
PP_5331	2.1	1.3	-0.16	-0.59	Long-chain acyl-CoA thioester hydrolase

Figure S2: Results of RB-TnSeq screen. A) Genes that showed great than 2 log2 fitness on either D-lysine, L-lysine, or 5AVA but showed no less than 0.5 log2 fitness defect when grown on glucose.