Prediction of Algal Blooms in the Great Lakes through a Convolution Neural Network of Remote Sensing Data

Daniel Hess[1], Vikram Duvvur[1], Karthik Srinivasan[1]

1: Affiliation: New Trier High School, Winnetka, Illinois, United States of America

Author Emails:

Daniel Hess: Daniel.o.hess@gmail.com,

Vikram Duvvur: vikduvvur@gmail.com

Karthik Srinivasan: karthik02@outlook.com

Manuscript Elements: Abstract (150 words), Main Text (X words), Figures X-X

Manuscript Type: Research Article

Abstract:

Harmful Algal Blooms (HABs) are the proliferation of algae due to eutrophication, and have severe repercussions to the ecological balance in many water bodies, attributable to the toxins the algae produce. Additionally, the identification and prediction of these HABs has been a recurring problem in the scientific community due to the interactions between both biological and physical processes that cause the HABs. Here we used remote sensing data to bypass these issues; remote sensing data provides significant information about the coverage of Chlorophyll which can be used to locate HABs. Using this indicator of HABs, we trained a Convolution Neural Network (CNN) to identify 9 types of Algal Blooms, using 25 epochs of 397 images, which can predict Algal Bloom shapes with over a 90% accuracy. This approach of HAB identification can easily be applied to other aquatic ecosystems, and thus can be used on other coastal and lake ecosystems.

Introduction

Algal blooms are a type of algal growth that can be dangerous to aquatic ecosystems in a variety of ways. Some blooms produce toxins capable of killing fish, mammals, and birds, and those toxins may also be harmful to humans (Landsberg, 2010). The death of the algae in the blooms can allow aerobic bacteria to consume all of the oxygen in their environment, leading to the formation of a dead zone. Eutrophication in a water body occurs when the level of nutrients, primarily phosphorus and nitrogen, increases significantly. The increased nutrient load causes a cascade of effects in which algal populations grow out of control and lead to a reduction in water quality and a destruction of the aqueous environment (Yang et al, 2008).

The criteria of surface water quality for lakes or reservoir (CNEPA, 2002)

Items	Surface water quality classification				
	Class I	Class II	Class III	Class IV	Class V
Water temperature (°C)	Maximum week increase≤1; maximum week decrease≤2				
pH	6~9				
DO (mg/L)	Saturation≥90%	≥6	≥5	≥3	≥2
COD_{Mn} (mg/L)	≤2	≤4	≤6	≤10	≤15
COD_{Cr} (mg/L)	≤15	≤15	≤20	≤30	≤40
BOD_5 (mg/L)	≤3	≤3	≤4	≤6	≤10
TN (mg/L)	≤0.2	≤0.5	≤1.0	≤1.5	≤2.0
NH_3-N (mg/L)	≤0.15	≤0.5	≤1.0	≤1.5	≤2.0
NO_2-N (mg/L)	≤0.06	≤0.1	≤0.15	≤1.0	≤1.0
TP (mg/L)	≤0.01	≤0.025	≤0.05	≤0.1	≤0.2
Chlorophyll a	≤0.001	≤0.004	≤0.01	≤0.03	≤0.065
Transparency (m)	≥15	≥4	≥2.5	≥1.5	≥0.5
Escherichia coli (L^{-1})	≤200	≤2000	≤10000	≤20000	≤40000

Table 1 shows a water quality classification table.

DO: dissolved-oxygen; CODMn: Chemical oxygen demand by K2MnO4 oxidation method;

CODCr: Chemical oxygen demand by chromium oxidation method; BOD5: Biological oxygen

demand; TN: Total nitrogen; TP: Total phosphorus

Table 2

The burden values of N and P in various eutrophicated water

Eutrophic status	TP (µg/L)	TN (µg/L)	Primary productivity	TNI	References
Oligotropic water	5~10	250~600	5~300 mg C/m^2	0~30	Likens et al.(1977); Cheng and Li (2006); Richardson et al.(2007)
Moderately eutrophic	10~30	500~1100	1000 mg C/(m^2·d)	31~60	
Eutrophic	30~100	1000~2000	–	61~100	
Hypereutropic	>100	>2000	–	>100	

TN: Total nitrogen; TP: Total phosphorus, TNI: Total nutrient status index

Table 2 shows a comparison of different types of aquatic ecosystems and their water quality.

A variety of conditions may cause harmful algal blooms, or HABs. The main factors are temperature, turbidity, nutrient levels, wind, water currents, and light levels. Of these factors nutrients - most notably phosphorus and nitrogen - are the single most significant cause of HABs. Nutrient buildup is often caused by runoff from fertilizers used on lawns and in agriculture. When it rains, these fertilizers are washed into lakes, seas, rivers, and oceans, where they wreak havoc. The increase in nutrient levels, especially limiting nutrients, allow the algal population to significantly proliferate.

The same factors that precipitate algal blooms may be used to predict their occurrence, shape, and severity, if those blooms occur between May and September. But by using a different set of tools - specifically, supervised machine learning and satellite imagery - it is possible to

predict the likely occurrence and distribution of algal blooms a month in advance. Supervised machine learning is the use of an artificial intelligence that creates an algorithm based on classified, or organized and labeled, data to, in this case, predict algal blooms. The software learns from the example, and builds its own algorithm based solely on the inputs and outputs. The satellite imagery showed the micrograms per liter of chlorophyll in the Great Lakes.

Chlorophyll monitoring is essential for the prediction of algal. Satellite remote sensing is the best method for identifying HABs in a large area. Remote sensing is more advantageous than any other form of HAB identification (Wei et al). Because remote sensing of chlorophyll enables the large scale detection of HAB's we utilized it for our predictor.

If researchers could use satellite images to measure nitrate and phosphate levels, algal blooms could be forecasted earlier and with greater precision than they are today. But using current sensing technology, these ions are not visible from space. The amount of chlorophyll in water, however, can be used as a proxy. Photosynthetic activity is easily detected by satellite imaging. We chose to use chlorophyll levels for two reasons: they are easy to measure from space, and they provide a good, early indication of potentially problematic algal blooms. Research has shown a strong positive empirical relationship between the number of bacteria and the amount of chlorophyll in both freshwater and marine environments. The difference between freshwater and marine environments is negligible as the linear regression equations are statistically indistinguishable (Bird and Kalff). Due to this relationship, high levels of chlorophyll correlate to HABs. We therefore can use the chlorophyll levels to predict the HABs. Our accuracy for identifying HABs is 85 percent.

A comparison of Chlorophyll and HAB's in Lake Erie on August 23rd 2018.

A comparison of Chlorophyll and HAB's in Lake Erie on July 4th 2018.

Methods

Machine learning is a type of artificial intelligence (AI) in which a training set is used to teach the AI without the need for it to be explicitly programmed. We needed a back-propagation artificial neural network capable of identifying several types of common algal bloom shapes in the form of images, so we decided to create a Convolution Neural Network. A neural network is made up of multiple "layers," each of which serves a different purpose and is composed of nodes that contain the logic of the network. We chose to make our neural network a back-propagation network, because the network can evaluate the error and propagate it back through earlier layers to get more accurate classifications of algal bloom shapes. The results are evaluated by loss functions, which reward higher accuracies with lower loss values. Ultimately we want to place an image into one of 11 categories: nine shapes of algal blooms, water or land. These categories are known as classes, hence the name multiclass identifier. We coded the project in Python 3.4 using the Keras and TensorFlow libraries, but the methods will be described in pseudocode

Neural Network Structure

We decided to use a specific type of machine learning model called a Convolution Neural Network because it is frequently used for multiclass classifiers for visual images. To prevent overfitting of the data, we started by setting up a convolution layer that applies 32, 3x3 filters to each image. An example of overfitting is if we trained our model to look for flowers, but only fed it images of red roses; The model would overfit to our training set, and think if an object is not red, then it is not a flower. To reduce the difficulty of classifying images, we simplify the upcoming layers by setting up a 2x2 pooling layer. This pooling layer aggregates information from the previous layer. After the pooling layer, we flatten the data into a one-dimensional single vector to ensure the next layer analyzes all the data from the previous layers. We then create a layer with 128 leaky rectifier linear unit nodes, because this fully connected layer is an easy way to find non-linear combinations of features extrapolated from the previous layers. Leaky rectifier nodes ensure that the value of the nodes do not get stuck at zero and cripple the neural network's signature back propagation. The final output layer is an 11-unit sigmoid node layer, which gives 11 values from zero to one depending on how closely it matches it matches each class, with "1" being an exact match and "0" being not close at all. The 11 classes are nine types of algal blooms, one of land, and one of water. The purpose of the sigmoid function allows the 11 classes to be easily compared, to see which shape the image most likely is. Then we compiled the network with the Adam loss function, which is used to evaluate how effective the weights of the current iteration of the model are. We chose the the Adam optimizer because it is considered to be computationally efficient (Kingma and Lei Ba, 2015).

Pictured above is the graph of the loss of our model versus its epoch through the data on a

logarithmic scale.

Training Images and Labeling

We used 900 images from the Great Lakes Remote Sensing Chlorophyll-A map, created

by the Michigan Tech Research Institute, to select the training examples for our Convolutional

Neural Network. The images were taken from 2010-2015. We used 50 to 10 images for each of

the 11 categories. Because we had a convolution layer, we could go over the same 912 images 25

times, effectively increasing our total training set to over 16,000 images.

100 example training set pictures

Prediction

Given a Chlorophyll-A map, the program first cuts it up into 32 by 32 pixel squares. It then runs the identification algorithm on all of the 33 by 32 squares. It then predicts the next stage of the Algal Bloom according to the flowchart below, and replaces the square if it has a progression on the chart. Finally, the image is stitched back together into one output image.

This is a flow chart of how the prediction algorithm predicts the algal bloom growth based on the previous bloom.

Discussion

Our algal bloom predictor identification accuracy was 85 percent across the nine types of algal blooms, compared to 9.1 percent identification rate from the random assignment of classes. The model will be able to provide an image predicting where and how algal blooms are likely to grow in the future. The predictive power of our software can be best leveraged to mitigate the the damages to local businesses and wildlife from harmful algal blooms. That is because algal

control methods taken in advance of a bloom are the most effective way of preventing and minimizing their effects.

Algal blooms have become a greater threat to natural environments in recent years, and predicting and monitoring them is essential to mitigating the damage they may cause. This approach was advocated in 2001 in a report to Congress by the National Oceanic and Atmospheric Administration and the National Sea Grant College Program that was created by the Woods Hole Oceanographic Institution. The monitoring method would need to allow local, state, and federal groups to work together. It would have to have early warning capabilities and accurate forecasting. It would have to predict bloom occurrence, development, and location. This type of prediction would enable the deployment of realistic mitigation strategies, such as reducing nutrient runoff from yards or farms. By doing this, human health and economic impacts could be reduced. Our software provides the exact kind of service that was called for in 2001. It is real time, predictive, accurate, and would enable both preventative and mitigatory steps to be taken to combat algal blooms.

Selected samples of water eutrophication occurrence in lake, reservoir, estuary and river in the world

Water body	Nutrient (N, P) concentrations	Phytoplankton biomass	History of eutrophication	References
Erie Lake, USA	TP: 115 μg/L	Chlorophyll a: 58 μg/L	Blu-green algae bloom between 1965 and 1979; Declined quality from 1995	Reutter, 1989; Maggie, 2004
Okeechobee Lake, Florida, USA	TP: 50~100 μg/L	N and P are limitors for phytoplankton	Eutrophic at present	Schelske, 1989
City Park Lake, Louisiana, USA	TN: 682 μg/L, TP: 330 μg/L	Chlorophyll a: 35.1 μg/L	Eutrophic (2000~2001)	Ruley and Rusch, 2002
Lugano Lake	TP: 140 μg/L	Phytoplankton dry weight: 7~16 g/m^2	Strongly eutrophic from 1960s	Barbieri and Simona, 2001
Danish lakes	TN: 2.4 mg/L, TP: 0.37 mg/L	Chlorophyll a: 73 μg/L	Highly eutrophic	Jeppesen et al., 1999
Pamvotis Lake, Northwest Greece	TP: 0.11 μg/L, NH$_4^+$: 0.25 mg N/L, NO$_3^-$: 0.56 mg N/L	–	Cultural eutrophication over the past 40 years; Eutrophication at present	Romero et al., 2002
Berg River, South Africa	TN: 217 mg/L, TP: 70 mg/L	–	Hypertrophic	de Villiers, 2007
Chivero Lake, Zimbabwe	TN: 0.3~8.4 mg/L, TP: 1.01~5.01 mg/L	Chlorophyll a: 18.02~22.48 μg/L	Highly eutrophic since 1980s	Nhapi, 2004; Ndebele and Magadza, 2006
Taihu Lake, China	TN: 2.56~4.5 mg/L, TP: 0.25~0.35 g/L	Algae biomass: 2.7~6.4 mg/L	Class I/II in the early 1960s; Class IV in the mid-1990s; Now, inferior Class V	Ye et al., 2007; Jin et al., 2006
Songhua Lake, China	TN: 1.14~1.98 mg/L, TP: 0.038~0.102 mg/L	Algae density: 210.84×10^4~432.68×10^4 cell/L		Wang et al., 2004
Dianchi Lake, China	TN: 2.13~8.27 mg/L, TP: 0.33~0.59 mg/L	–	Class III in the early 1970s; Now, inferior Class V	Guo and Sun, 2002

Table 3 shows selected samples of water eutrophication occurrence in lake, reservoir, estuary and river in the world.

Algal blooms are based on the water conditions that are most often tracked in real time through buoys in the water or with water sample testing. (Yang et al, 2008).) This is very constraining on predicting the size, scope and nature of algal blooms, because such an approach requires that scientists have direct access to a specific area of a body of water. This limits the area that you can effectively forecast for because you need to have access to that water area or place sensors in those areas. On top of this, it greatly increases cost because you have to collect water samples or place sensors. By utilizing data that is already being collected and employing machine learning, we sidestep the normal costs that are associated with forecasting algal blooms. We do not need to be at the site of the algal bloom, and we do not need to collect water samples, which are costly factors. All the work done with our prediction system happens on a computer and can be done by anyone.

Methods for combatting algal blooms are grouped into three main categories: mechanical, physical or chemical, and biological. Mechanical control methods involve filters, pumps, and barriers. These are used to remove blooms or contain them in an area so they do not spread. These can also be used to deal with dead fish and other bloom by products. Physical or chemical control relies on the use of chemicals or mineral compounds to combat algal blooms. The chemical or physical methods usually kill, inhibit, or remove HABs. Biological controls are often the most complicated. They rely on the use of organisms or pathogens, such as viruses, bacteria, parasites, zooplankton, and shellfish, that are able to kill lyse or HAB cells (Wei et al)

HABs are not only dangerous to the aquatic systems that they happen in but to humans that use those systems, as well. Because algae are eaten in large amounts by many aquatic animals and help to form the basis of the trophic pyramid, dangerous algae can cause a build up of toxins in animals that humans consume. HABs have a significant impact on the aquatic resources and can affect how resources are used. The best example of the human impact of harmful algal bloom is the death of humans caused by the consumption of shellfish. Shellfish consume the toxic HABs, and the toxins build up inside of the shellfish. When humans consume these tainted shellfish, they can then become sick and die (Wei et al). The direct human impact of HABs make it essential that our software is utilized to reduce and combat algal blooms.

Algal blooms do not just present a health risk for humans; fish die-offs are a common side effect of algal blooms. When algae flourish in an aquatic environment, they rapidly use up all the dissolved oxygen in whatever area they are growing in. Once they consume oxygen, very little oxygen remains available for other organisms, and mass die-offs may follow. In addition, algal blooms almost always occur during times of warmer temperatures. When water is warmer,

it holds less dissolved oxygen naturally, meaning that there are often two factors depleting levels of dissolved oxygen. Due to the shortages of oxygen caused by these two factors, fish can often suffer die-offs (Yang et al, 2008).

On top of the health risks caused by algal blooms, they also may impose significant economic costs. HABS have a massive economic impact both directly and directly in the areas they occur in. An outbreak of paralytic in the Northeast, That was caused by HABs, had an estimated cost of 6 million dollars (Shumway 1988). In North Carolina between 1987 and 1988 a four to six month period of red tide had an estimated cost on the community of 25 million dollars (Tester and Fowler 1990). A 1991 domoic acid and amnesic shellfish poisoning outbreak in Washington State had a huge impact on the entire community. It affected tourism, fisheries (mostly oyster), and had losses between 15 and 20 million dollars. An outbreak in North Carolin of Pfiesteria piscicida and Pfiesteria between 1995 and 1996 caused massive damage. Millions of fish, including the commercially fished menhaden, died to toxins and or secondary infections Burkholder and Glasgow 1997). Another outbreak of Pfiesteria or Pfiesteria like organisms caused an estimated 43 million dollars in damages to the Chesapeake bay region. It resulted in a public outcry and several reported cases of illness (Sieling and Lipton 1998). The economic impact of HABs is substantial and affects many parts of the world. Because of this it is essential that we create new prediction methods like our algorithm so that control methods can be better utilized (Bushaw-Newton, K.L. and Sellner, K.G).

Though marine coasts experience numerous algal blooms, the issue also is significant in the Great Lakes. Federal agencies have begun to take note of the problem in the Great Lakes, especially in Lake Erie, where the blooms are particularly common. Recent research has shown a

link between health problems in residents of the Great Lakes area and blooms of blue-green

algae, otherwise known as Cyanobacteria. Blue-green algae blooms occur frequently in the U.S.

They are often associated with eutrophication and the nutrient enrichment caused by runoff that

causes the blooms. Blue-green algal blooms usually form on the surface of water and occur most

commonly during the warmest times of the year. They are frequently a source of annoyance for

people who participate in boating, fishing, and swimming. They also often cause flavor and odor

problems in water and at water treatment plants (Wei et al).

Results

Prediction of only HABs:

The cyclical nature of the HABs throughout the seasons of the year hints to the fact that

HABs are in fact predictable. The correlation between a predictive algorithm for HABs and the

HABs that occur in the future is quite pronounced (Muttil et al, 2005). The satellite imaging data

captured the amounts of photosynthesis occurring. We were able to use this data to predict HABs

as opposed to merely photosynthetic activity due to the correlation aforementioned. The

reasoning behind this logic is sound. Although the images are marked as measuring all

photosynthetic activity, the power of the satellite images would not be nearly enough to reach the

activity of photosynthetic plants in the benthic region of deep, oligotrophic lakes such as those in

the Great Lakes. As a result, the photosynthetic activity being depicted in the images must be in

the epilimnion. Apart from those that are able to grow in the littoral zone, the only

photosynthetic activity that would be occurring in the epilimnion would be that of algae, and any

algal colony large enough to be significant in the satellite images suggests that a bloom is occurring.

Other Methods:

HABs are a major problem all over the world. In the Great Lakes, Lake Erie, in particular, has suffered from blooms. Our prediction method enables prediction of algal blooms not just in the Great Lakes, but in all bodies of water. A widespread literature review was performed to understand the HAB problem in Lake Erie. The current methods to forecast HABs all over the world and specifically in Lake Erie were examined. An extensive literature review and analysis was performed on the possible variables for forecasting HABs. Two forecasting methods - CART, classification and regression tree, and ANN, artificial neural network - as well as two training periods and two input variable selection methods, nutrient loading period and Spearman's rank correlation were used. The Spearman rank correlation describes how well two variables' relationship can be explained with a monotonic function. A monotonic function is a function that is always either entirely non-decreasing or entirely non-increasing. For the nutrient loading period selection method, only one set of input variables is used for forecasting. The Spearman selection method, on the other hand, examines more variables than the nutrient loading period, considering up to 28 different averaging periods and lag times for each considered variable. First, the CART forecasting models were tested with both classification methods, a 3-class and 5- class system resulting in the 3-class system being selected. The CART models were then created for both methods and training periods.

Initially, when using the first training period of 2002 to 2011, the loading period method showed better precision in forecasting HABs when compared to the Spearman selection method. When the training period was increased to 2013, both methods showed an improvement in the overall accuracy, with Spearman having an 8.9 percent improvement, and loading period a 5.4 percent improvement. However, with the extended training period, the loading period decision trees for August and September showed a slight increase in precision over the Spearman method and the Spearman method being slightly more precise in October. After the CART models, the ANN models were created and analyzed for both selection methods and training periods. For both selection methods in the first training period, the models often underpredicted the higher magnitude blooms.

Many of the forecasts did not predict the exact same CI, or cyanobacterial index, as observed. However in most cases, both methods predicted the same class of bloom as observed. In most cases after increasing the training period to 2013, both ANN models improved their accuracy for predicting the higher magnitude of blooms. The correlation coefficient increased from 0.70 to 0.77 for the loading period selection method and from 0.79 to 0.83 for the Spearman selection method when extending the training period. Both input selection methods had some difficulty in predicting the 2015 HAB because the 2015 bloom was a special case in terms of nutrient loading, as well as bloom time. There was a large amount of loading in June and July, which is atypical due to how early it was happening. The 2015 bloom in July was 382 percent larger than any bloom recorded from 2002 to 2014. The monthly discharge for June was the highest recorded and third highest on record since the United States Geological Survey

started collecting data in 1939. Similar to the CART method, the ANN model showed an increase in accuracy when forecasting HABs with an extended training period.

Our method, however does not rely on nutrient variables and instead looks at the growth of algal blooms themselves through measured chlorophyll levels. Our method enables the prediction of severity, location, and distribution of algal blooms, just like both the CART and ANN methods. Because we utilize satellite imagery, however, we are able to simplify the prediction process and increase success. Our algorithm successfully identifies algal bloom components 85 percent of the time, due not only to the neural network but also due to the aforementioned near-certainty with which our satellite images highlight algal blooms.

Conclusion:

Understanding the nature of HABs can lead to their prediction through the data gathered by remote sensing images. We found that not only did the data accurately convey where the HABs would occur, but that the bloom events were predictable. Using a Neural Network to classify the HABs and predict where future HABs would occur, our algorithm was able to render success rates of 85%. The future impact of HABs in the Great Lakes can be feasibly lowered by our algorithm, as early response to the HABs can stop their successive propagation through the ecosystem. This is can redeem us from our anthropogenic effect on the prevalence of HABs in the Great Lakes, especially Lake Eerie, and can help us keep the Great Lakes a healthy, biodiversity, and oligotrophic ecosystem.

References

Intro References:

Landsberg, Jan H. (2010). The Effects of Harmful Algal Blooms on Aquatic Organisms. *Reviews in Fisheries Science Vol 10:* 133-390

Yang, Xiao-e et al. "Mechanisms and Assessment of Water Eutrophication ." *Journal of Zhejiang University. Science. B* 9.3 (2008): 197–209.

Wei, Bin, Sugiura, Norio, Maekawa, & Maekawa, Takaaki. (2001). Use of artificial neural network in the prediction of algal blooms. *Water Research Vol 35: 2022-2028*

Bird, D.F., & Kalff, J. (1989). Phagotrophic sustenance of a metalimnetic phytoplankton peak. *Limnology and Oceanography Vol 34, Iss. 1*

Methods References:

Ba, Jimmy Lei, and Diederik P Kingma. "ADAM: A METHOD FOR STOCHASTIC OPTIMIZATION." *Arxiv*, arxiv.org/pdf/1412.6980.pdf.

Discussion References:

"Woods Hole Oceanographic Institution." *Woods Hole Oceanographic Institution*, National Oceanic and Atmospheric Administration, Sept. 2001,

Yang, Xiao-e et al. "Mechanisms and Assessment of Water Eutrophication ." *Journal of Zhejiang University. Science. B* 9.3 (2008): 197–209. *PMC*. Web. 14 Oct. 2018.

Shumway, S.E. 1988. A review of the effects of algal blooms on shellfish and aquaculture. Journal of the World Aquaculture Society 21:65-104.

Tester, P.A. and P.K. Fowler. 1990. Brevetoxin contamination of Mercenaria mercenaria and Crassostrea virginica: A management issue. In: Graneli, E., B. Sundstrom, L. Edler, and D.M. Anderson (eds.), Toxic marine phytoplankton. New York: Elsevier Press. pp. 499-503.

Burkholder, J.M. and H.B. Glasgow Jr. 1997. Pfiesteria piscicida and other Pfiesteria- like dinoflagellates: Behavior, impacts, and environmental controls. Limnology and Oceanography 42:1052-1075.

Sieling, W. and D. Lipton. 1998. Pfiesteria's impact on seafood industry sales.

Bushaw-Newton, K.L. and Sellner, K.G. 1999. Harmful Algal Blooms. In: NOAA's State of the Coast Report. Silver Spring, MD: National Oceanic and Atmospheric Administration. http://state-of-coast.noaa.gov/ bulletins/html/hab_14/hab.html

Muttil, Nitin, and Joseph H.W. Lee. "Genetic Programming For Analysis And Real-Time Prediction Of Coastal Algal Blooms." *Ecological Modelling* 189.3-4 (2005): 363-376. Web. 23 Oct. 2018.

Results References:

Muttil, N, & Chau, KW. (2006). Neural network and genetic programming for modelling coastal algal blooms. *International Journal of Environment and Pollution Vol 28:* 223-238

"Distribution of chlorophyll and harmful algal blooms (HABs): A review on space based studies in the coastal environments of Chinese marginal seas". *Advances in Space Research,* vol. 41, no. 1, 2007, pp. 12-19. doi:10.1016/j.asr.2007.01.037.

Jan H. Landsberg (2002) The Effects of Harmful Algal Blooms on Aquatic Organisms, Reviews in Fisheries Science, 10:2, 113-390

Reutter JM. Lake Erie: Phosphorus and Eutrophication. Columbus: 1989. Ohio Sea Grant College Program.

Maggie BM. Total Maximum Daily Load Effectiveness Monitoring Study: Lakes Erie and Campbell.2004. Quality Assurance Project Plan, No. 04-03-206.

Schelske PC. Assessment of nutrient effects and nutrient limitation in Lake Okeechobee. Journal of the American Water Resources Association. 1989;25(6):1119–1130

Ruley JE, Rusch KA. An assessment of long-term post-restoration water quality trends in a shallow, subtropical, urban hypereutrophic lake. Ecological Engineering. 2002;19(4):265–280.

Barbieri A, Simona M. Trophic evolution of Lake Lugano related to external load reduction: Changes in phosphorus and nitrogen as well as oxygen balance and biological parameters. Lakes and Reservoirs Research and Management. 2001;6(1):37–47.

Jeppesen E, Søndergaard M, Kronvang B, Jensen JP, Svendsen LM, Lauridsen TL. Lake and catchment management in Denmark. Hydrobiologia. 1999;395-396:419–432.

Romero JR, Kagalou I, Imberger J, Hela D, Kotti M, Bartzokas A, Albanis T, Evmirides N, Karkabounas S, Papagiannis J, et al. Seasonal water quality of shallow and eutrophic Lake Pamvotis, Greece: Implications for restoration. Hydrobiologia. 2002;474(1-3):91–105.

de Villiers S. The deteriorating nutrient status of the Berg River, South Africa. Water SA. 2007;33(5):659–664.

Ndebele MR, Magadza CHD. The occurrence of microcystin-LR in Lake Chivero, Zimbabwe. Lakes and Reservoirs Research and Management. 2006;11(1):57–62.

Nhapi I. Options for Wastewater Management in Harare, Zimbabwe. Wageningen, the Netherlands: Wageningen University; 2004. (Ph.D Thesis)

Ye C, Xu QJ, Kong HN, Shen ZM, Yan CZ. Eutrophication conditions and ecological status in typical bays of Lake Taihu in China. Environmental Monitoring and Assessment. 2007;135(1-3):217–225.

Wang X, Lu XG, Zhang XL, Zhang ZQ. Eutrophication aspects and effective factors analysis in Songhua Lake. Wetland Science. 2004;2(4):273–278.

Guo HC, Sun YF. Characteristic analysis and control strategies for the eutrophicated problem of the Lake Dianchi. Progress in Geography. 2002;21(5):500–506.

Michalak, A., et al. (2013). Record-setting algal bloom in Lake Erie caused by agricultural and meteorological trends consistent with expected future conditions. Proceedings of the National Academy of Sciences of the United States of America.

Likens GE, Bormann FH, Pierce RS. (1977). Biogeochemistry of a Forested Ecosystem. New York: Springer-Verlag

Cheng XY, Li SJ. An analysis on the evolvement processes of lake eutrophication and their characteristics of the typical lakes in the middle and lower reaches of Yangtze River. Chinese Science Bulletin. 2006;51(13):1603–1613.

Richardson CJ, King RS, Qian SS, Vaithiyanathan P, Qualls RG, Stow CA. Estimating ecological thresholds for phosphorus in the Everglades. Environmental Science and Technology. 2007;41(23):8084–8091.

CNEPA (Environmental Protection Agency of China) Environmental Quality Standard for Surface Water. GB 3838-2002. 2002.

The criteria of surface water quality for lakes or reservoir (CNEPA, 2002)

Item	Surface water quality cover bodies				
	Class I	Class II	Class III	Class IV	Class V
DO mg/L	Saturate≥90%	6	5	3	2
CODMn mg/L	≤2	4	6	10	15
COD mg/L	≤15	15	20	30	40
BOD5 mg/L	≤3	3	4	6	10
NH3-N mg/L	≤0.15	0.5	1.0	1.5	2.0
TP mg/L	≤0.01	0.025	0.05	0.1	0.2
TN mg/L	≤0.2	0.5	1.0	1.5	2.0
Chloride	≤0.01	0.01	0.05	0.2	1.0
Fluoride(mg/L)	≤1.0	1.0	1.0	1.5	1.5

Table 2

The haematic indices of M and P in various experimental sets.

	Hb (g/L)	ERY (×10⁶/μL)	Platelet probability	DD	Reference
Oligemic state	7–8	115–140	—	—	
Moderately anaemic	9–10	60–120	—	12–20	
Severe anaemia	14	120	—	24–40	
Hyperaemia	24	97–98	—	0.60	

(1) Hemoglobin (2) Erythrogenesis (3a) Antithrombae materials.

Loss vs Epoch

Nose Sheen Nose Sheen Minor Bloom	Nose Sheen Nose Sheen Medium Bloom	Nose Sheen Nose Sheen Major Bloom
Nose Sheen Tail and Wide Minor Bloom	Wide Tail and Wide Medium Bloom	Nose Sheen Tail and Wide Major Bloom
Far From Sheen Minor Bloom	Far From Sheen Medium Bloom	Far From Sheen Major Bloom

Selected samples of water column/data on occurrence in lake, reservoir, estuary and river in the world

Water body	Nature of Pb concentration	Phytoremediation	History of contamination	Reference
Ganga Lake, USA	BDL-0.8 µg/L	Chlorophyll < 5 µg/L		
Pontchartrain Lake, Florida, USA	BDL - 40 µg/L	Water Pycnocephalus filter		Sreehari, 1980
Kep Park Lake	Low ranges	Chlorophyll < 5 µg/L		Bailey and Stevens, 1983
Argenta Lake	BDL ranges	Precipitation dry weight < 5 µg/L		
Barent lakes	BDL-74 µg/L, BDL-7.6 µg/L	Chlorophyll < 5 µg/L	Highly eutrophic	
Victoria Lake, Northeast	BDL-2 µg/L, BDL<2 µg/L, Nil, 0.5 µg/L, 0.5, Nil	Chlorophyll < 5 µg/L	Criteria proportion over to per 5 year	Barrow et al, 2001
Zeekoe Lake, South Africa	BDL<2 µg/L	Chlorophyll < 5 µg/L	Eutrophic	de Villiers, 2007
Chioumri Lake, Cameroon	BDL-12.4 µg/L, BDL-0.9 µg/L	Chlorophyll - BDL < 5 µg/L	Nil	Nkoue Ndondo and Ngoupayou, 2009
Toba Lake, China	BDL-1.6 µg/L, BDL<2 µg/L	Chlorophyll < 5 µg/L	Class III in the early Holocene	
Yangtze Lake, China	BDL-1.6 µg/L, BDL<2 µg/L	Algae zone Chlorophyll < 5 µg/L	Class III in the early Holocene	Wang et al., 2008
Shuvrili Lake, China	BDL-1.6 µg/L	Chlorophyll < 5 µg/L	Class III in the early Holocene	