BRASS: permutation methods for binary traits in genetic association studies with structured samples

Joelle Mbatchou[1,2], Mark Abney[3], Mary Sara McPeek[2,3*]

1 Regeneron Genetics Center, Tarrytown, NY, USA

2 Department of Statistics, The University of Chicago, Chicago, IL, USA

3 Department of Human Genetics, The University of Chicago, Chicago, IL, USA

* mcpeek@uchicago.edu

Abstract

In genetic association analysis of complex traits, permutation testing can be a valuable tool for assessing significance when the distribution of the test statistic is unknown or not well-approximated. This commonly arises, e.g, in tests of gene-set, pathway or genome-wide significance, or when the statistic is formed by machine learning or data adaptive methods. Existing applications include eQTL mapping, association testing with rare variants, inclusion of admixed individuals in genetic association analysis, and epistasis detection among many others. For genetic association testing in samples with population structure and/or relatedness, use of naive permutation can lead to inflated type 1 error. To address this in quantitative traits, the MVNpermute method was developed. However, for association mapping of a binary trait, the relationship between the mean and variance makes both naive permutation and the MVNpermute method invalid. We propose BRASS, a permutation method for binary traits, for use in association mapping in structured samples. In addition to modeling structure in the sample, BRASS allows for covariates, ascertainment and simultaneous testing of multiple markers, and it accommodates a wide range of test statistics. In simulation studies, we compare BRASS to other permutation and resampling-based methods in a range of scenarios that include population structure, familial relatedness, ascertainment

and phenotype model misspecification. In these settings, we demonstrate the superior control of type 1 error by BRASS compared to the other 7 methods considered. We apply BRASS to assess genome-wide significance for association analyses in domestic dog for elbow dysplasia (ED) and idiopathic epilepsy (IE). For both traits we detect previously identified associations, and in addition, for ED, we detect significant association with a SNP on chromosome 35 that was not detected by previous analyses, demonstrating the potential of the method.

Author summary

To determine whether genetic association with a trait is significant, permutation methods are an attractive and popular approach when analytic methods based on distributional assumptions are not available, e.g., when applying machine learning or data adaptive methods, or when performing a multiple testing correction, e.g., to assess region-wide or genome-wide significance in association mapping studies. Existing applications include eQTL mapping, association testing with rare variants, inclusion of admixed individuals in genetic association analysis, and detection of genetic interaction among many others. However, when there is population structure in the sample, naive permutation of the data can lead to inflated significance of the association results. For continuous traits, linear mixed-model based approaches have been proposed for permutation-based tests that can also adjust for sample structure; however, these do not remain valid when applied to binary traits, as key features of binary data are not well accounted for. We propose BRASS, a permutation-based testing method for binary data that incorporates important characteristics of binary data in the trait model, can accommodate relevant covariates and ascertainment, and adjusts for the presence of structure in the sample. In simulations, we demonstrate the superior control of type 1 error by BRASS compared to other methods, and we apply BRASS in the context of correcting for multiple testing in two genome-wide association studies in domestic dog: one for elbow dysplasia and one for idiopathic epilepsy.

Introduction

In genome-wide association studies (GWAS), the primary objective is generally to identify associations between a phenotype of interest and genetic variants. This involves assessing the p-values of certain test statistics by deriving the appropriate null distribution or an asymptotic approximation to it. However, this is not always feasible as the distribution may be intractable. This can arise when the test statistics result from black-box machine learning methods [1,2], or in region-based tests where association signals over multiple sites are combined, e.g., rare variant tests [3], and/or when the test statistic involves data-adaptive weights [4]. A further limitation can arise when, even if the distribution (or the asymptotic distribution) of the test statistic is known for single tests, significance needs to be assessed for the maximum of many correlated tests. This occurs in genome scans to establish a genome-wide significance threshold, where the linkage disequilibrium present between the markers induces correlation between the association tests [5–7].

To overcome these limitations, a common approach is to perform permutation testing so as to obtain replicates of the data under the null hypothesis from which an empirical distribution can be derived [6,7]. A fundamental assumption underlying permutation testing is exchangeability of the subjects in the sample, which is often approximately satisfied in population-based samples. However, this assumption can be violated in the presence of genetic structure in the sample (e.g., kinship, cryptic relatedness, and/or population structure), as it will introduce correlation in the sample, including in the phenotype values through polygenic effects [8,9]. Hence, naive application of permutation testing will usually not preserve the correlation structure and can result in inflated type 1 error rates [10] (though this can be avoided if, for example, all subjects in the sample are equally related [11]). We consider here the problem of permutation testing for a binary trait in the presence of polygenic effects in a sample with population structure, cryptic and/or family relatedness. While an exact permutation test may not be feasible, a permutation-based test that adjusts for the correlation structure is feasible and has been proposed for quantitative traits [9,11]. The approach, called MVNpermute, is based on a linear mixed model (LMM) and is applicable for multivariate-normal data. It incorporates genetic relatedness through the

inclusion of random effects in the LMM and can also adjust for covariates. However, as MVNpermute is primarily designed for quantitative traits that are multivariate-normal, there will be more model misspecification when applied to binary traits as the binary nature of the data is not incorporated into the LMM.

We propose BRASS (for "binary trait resampling method adjusting for sample structure"), a permutation procedure for a binary trait which incorporates both covariates and the correlation structure present in the sample. In contrast to the LMM-based approach, it accommodates the binary nature of the trait through a quasi-likelihood framework that considers the effect of covariates on a logit scale in the mean structure as well as the relationship between the trait mean and its variance, both of which are important features of binary data. Hence, BRASS benefits from less model misspecification in the presence of important covariates relative to the LMM-based approach. Here, we show that the permutation-based replicates obtained with BRASS are able to maintain correct type 1 error control, and we compare it with several alternative approaches. We demonstrate the use of our method in a context of assessing genome-wide significance in domestic dog GWAS. In animal studies, permutation testing is commonly used when there is not a consensus threshold for genome-wide significance [12–15], and it is also used in some human studies when the consensus threshold might not be applicable [6]. We apply BRASS to perform association mapping in two GWAS of domestic dogs, one for elbow dysplasia (ED [OMIA: 000330-9615]) and and one for idiopathic epilepsy (IE [OMIA: 000344-9615]).

Description of the Methods

Overview of BRASS method

We consider the problem of resampling a binary trait variable that is correlated across individuals, where the correlation can arise from, e.g., population structure or related individuals. Our aim is to derive replicates of the trait under the null hypothesis of no association while accounting for the correlation that is present in the sample but whose structure is unknown. To obtain these replicates, we start by modeling the response using a previously described quasi-likelihood framework for correlated binary

traits [16, 17],

$$\boldsymbol{\mu} := \mathbb{E}(\mathbf{Y}|\mathbf{X}, \mathbf{G}) = \text{logit}^{-1}(\mathbf{X}\boldsymbol{\beta} + \mathbf{G}\gamma), \text{ and} \tag{1}$$

$$\boldsymbol{\Omega} := \text{Var}(\mathbf{Y}|\mathbf{X}, \mathbf{G}) = \boldsymbol{\Gamma}^{1/2}\,\boldsymbol{\Sigma}\,\boldsymbol{\Gamma}^{1/2}. \tag{2}$$

where $\mathbf{Y} = (Y_1,...,Y_n)^T$ denotes the phenotype vector for n subjects, $\mathbf{X}$ is the $n \times k$
matrix of k covariates (including an intercept term), which is assumed to be of rank k,
$\mathbf{G} = (G_1,...,G_n)^T$ is the vector of genotypes at the marker of interest, where G_i denotes
the minor allele count for the ith individual, $\boldsymbol{\beta}$ is a k-length vector representing the
unknown effects of covariates, γ represents the unknown effect of the genetic marker, $\boldsymbol{\Gamma}$
is an $n \times n$ diagonal matrix with ith diagonal entry $\mu_i(1 - \mu_i)$ where μ_i is the ith
element of $\boldsymbol{\mu}$, the $n \times n$ matrix $\boldsymbol{\Sigma}$ is given by

$$\boldsymbol{\Sigma} = \xi\,\boldsymbol{\Phi} + (1 - \xi)\,\mathbf{I}_n, \tag{3}$$

where the unknown scalar parameter $\xi \in [0, 1]$ can be viewed as a heritability parameter,
and the $n \times n$ matrix $\boldsymbol{\Phi}$ is generally a kinship or genetic relatedness matrix (GRM).

The framework characterized by (1) and (2) allows for modeling important
covariates, such as biological or ancestry informative covariates, in the mean structure,
while the residual correlation between subjects is captured directly within the variance
structure by the inclusion of $\boldsymbol{\Phi}$ in (3). Furthermore, by including the matrix $\boldsymbol{\Gamma}$ in (2),
this framework models the dependence of the phenotypic variance on the mean, a key
feature of binary data.

Since we wish to simulate replicates of the binary trait under the null hypothesis of
no association, $H_0 : \gamma = 0$, the remaining unknown parameters are $\boldsymbol{\beta}$ and ξ, which need
to be estimated from the data. The null estimate $(\widehat{\boldsymbol{\beta}}, \hat{\xi})$ is easily obtained by iteratively
solving the following system of estimating equations [17] under the constraint $\gamma = 0$,

$$D_{\boldsymbol{\beta}}^T \boldsymbol{\Omega}^{-1}(\mathbf{Y} - \boldsymbol{\mu}) = \mathbf{0}, \tag{4}$$

$$(\mathbf{Y} - \boldsymbol{\mu})^T \boldsymbol{\Gamma}^{-1/2} \boldsymbol{\Sigma}^{-1}(\boldsymbol{\Phi} - \mathbf{I})\boldsymbol{\Sigma}^{-1}\boldsymbol{\Gamma}^{-1/2}(\mathbf{Y} - \boldsymbol{\mu}) = \text{trace}(\boldsymbol{\Sigma}^{-1}(\boldsymbol{\Phi} - \mathbf{I})), \tag{5}$$

where $\boldsymbol{D_\beta} = \frac{\partial \boldsymbol{\mu}}{\partial \boldsymbol{\beta}}$ is a Jacobian of the conditional mean of the trait with respect to $\boldsymbol{\beta}$. In our model with a logit link function, $\boldsymbol{D_\beta} = \boldsymbol{\Gamma X}$.

The justification underlying an ordinary permutation test is an assumption of exchangeability under the null hypothesis, which does not generally hold in most situations of correlated data. Instead, we develop and apply an approximate permutation test based on second-order exchangeability, in which we permute a vector, $\hat{\boldsymbol{\zeta}}$, whose entries each have the same mean and variance and are uncorrelated with each other. Our approach builds on previous work [9, 11] that was limited to quantitative traits, and we extend the ideas to binary traits.

We start by fitting the null model given by Equations 1 and 2 with $\gamma = 0$, and we let $\mathbf{Y} - \hat{\boldsymbol{\mu}}$ denote the resulting residual vector. We aim to map $\mathbf{Y} - \hat{\boldsymbol{\mu}}$ to a lower-dimensional orthonormal space in which its entries will be second-order exchangeable, with the result of this map being $\hat{\boldsymbol{\zeta}}$. To form $\hat{\boldsymbol{\zeta}}$, we need to first obtain the approximate covariance matrix of $\mathbf{Y} - \hat{\boldsymbol{\mu}}$ under the null hypothesis, which is challenging because $\mathbf{Y} - \hat{\boldsymbol{\mu}}$ has no closed-form expression as a function of the data.

To obtain the approximate covariance matrix, we first fix ξ and let $U(\boldsymbol{\beta}) = \boldsymbol{D_\beta^T} \boldsymbol{\Omega}^{-1} (\mathbf{Y} - \boldsymbol{\mu})$, which corresponds to the quasi-score function for $\boldsymbol{\beta}$. Under the null hypothesis of $\gamma = 0$, let $\boldsymbol{\beta}_0$ denote the true value of $\boldsymbol{\beta}$, and let $\boldsymbol{\mu}_0, \boldsymbol{\Gamma}_0, \mathbf{D}_0$ and $\boldsymbol{\Omega}_0$ correspond to $\boldsymbol{\mu}, \boldsymbol{\Gamma}, \mathbf{D}_\beta$ and $\boldsymbol{\Omega}$, respectively, evaluated at $\boldsymbol{\beta}_0$ and $\gamma = 0$. Similarly, let $\hat{\boldsymbol{\mu}}, \hat{\boldsymbol{\Gamma}}, \hat{\mathbf{D}}$ and $\hat{\boldsymbol{\Omega}}$ be the same quantities evaluated at $\hat{\boldsymbol{\beta}}$ and $\gamma = 0$. We apply a Taylor series expansion to $U(\boldsymbol{\beta})$ around $\boldsymbol{\beta}_0$, evaluated at $\hat{\boldsymbol{\beta}}$:

$$U(\hat{\boldsymbol{\beta}}) \approx U(\boldsymbol{\beta}_0) + \frac{\partial U(\boldsymbol{\beta})}{\partial \boldsymbol{\beta}} \bigg|_{\boldsymbol{\beta} = \boldsymbol{\beta_0}} \cdot (\hat{\boldsymbol{\beta}} - \boldsymbol{\beta}_0). \tag{6}$$

Using the fact that $U(\hat{\boldsymbol{\beta}}) = \mathbf{0}$, replacing the Jacobian in (6) by its expectation (similar to Fisher scoring), and solving for $\hat{\boldsymbol{\beta}} - \boldsymbol{\beta}_0$ we get,

$$\hat{\boldsymbol{\beta}} - \boldsymbol{\beta}_0 \approx - \left\{ \mathbb{E} \left[\frac{\partial U(\boldsymbol{\beta})}{\partial \boldsymbol{\beta}} \right] \bigg|_{\boldsymbol{\beta} = \boldsymbol{\beta_0}} \right\}^{-1} U(\boldsymbol{\beta}_0),$$

$$= (\mathbf{D}_0^T \, \boldsymbol{\Omega}_0^{-1} \, \mathbf{D}_0)^{-1} \mathbf{D}_0^T \, \boldsymbol{\Omega}_0^{-1} (\mathbf{Y} - \boldsymbol{\mu}_0). \tag{7}$$

As $\hat{\boldsymbol{\mu}}$ is a non-linear function of $\hat{\boldsymbol{\beta}}$, a Taylor series expansion is performed on $\boldsymbol{\mu}$

around $\boldsymbol{\beta}_0$, and evaluated at $\widehat{\boldsymbol{\beta}}$,

$$\widehat{\boldsymbol{\mu}} \approx \boldsymbol{\mu}_0 + \mathbf{D}_0 \, (\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}_0), \tag{8}$$

Combining (7) and (8), we get for the residual vector

$$\begin{aligned} \mathbf{Y} - \widehat{\boldsymbol{\mu}} &= \mathbf{Y} - \boldsymbol{\mu}_0 - (\widehat{\boldsymbol{\mu}} - \boldsymbol{\mu}_0), \\ &\approx \left[\mathbf{I} - \mathbf{D}_0 \, (\mathbf{D}_0^T \, \boldsymbol{\Omega}_0^{-1} \, \mathbf{D}_0)^{-1} \mathbf{D}_0^T \, \boldsymbol{\Omega}_0^{-1} \right] (\mathbf{Y} - \boldsymbol{\mu}_0). \end{aligned} \tag{9}$$

As a result, we can approximate the covariance matrix of the residuals as

$$\text{Var}(\mathbf{Y} - \widehat{\boldsymbol{\mu}}) \approx \boldsymbol{\Omega}_0 - \boldsymbol{\Gamma}_0 \mathbf{X} \, (\mathbf{X}^T \boldsymbol{\Gamma}_0 \, \boldsymbol{\Omega}_0^{-1} \, \boldsymbol{\Gamma}_0 \mathbf{X})^{-1} \mathbf{X}^T \boldsymbol{\Gamma}_0. \tag{10}$$

The correlation present in $(\mathbf{Y} - \widehat{\boldsymbol{\mu}})$ arises from two sources: that introduced by $\boldsymbol{\Phi}$ through $\boldsymbol{\Omega}_0$ and that introduced from using the estimated mean $\widehat{\boldsymbol{\mu}}$ instead of the true unknown mean $\boldsymbol{\mu}_0$. We use a factorization $\mathbf{C}$ of $\boldsymbol{\Omega}_0$, with $\boldsymbol{\Omega}_0 = \mathbf{C}^T \mathbf{C}$, to remove the correlation due to $\boldsymbol{\Phi}$ and obtain,

$$\text{Var}[\mathbf{C}^{-T}(\mathbf{Y} - \widehat{\boldsymbol{\mu}})] \approx \mathbf{I} - \mathbf{W}(\mathbf{W}^T\mathbf{W})^{-1}\mathbf{W}^T = \boldsymbol{\Psi}_0, \tag{11}$$

where $\mathbf{W} = \mathbf{C}^{-T}\boldsymbol{\Gamma}_0\mathbf{X}$. The matrix $\boldsymbol{\Psi}_0$ in (11) is symmetric, idempotent and of rank $n - k$, so it can be expressed as $\boldsymbol{\Psi}_0 = \mathbf{V}\mathbf{V}^T$ where the columns of $\mathbf{V}$ contain the eigenvectors of $\boldsymbol{\Psi}_0$ corresponding to eigenvalue 1, and $\mathbf{V}^T\mathbf{V} = \mathbf{I}_{n-k}$. We use $\mathbf{V}^T$ to remove the remaining correlation that is driven by parameter estimation.

Applying these two linear transformations, we obtain $\boldsymbol{\zeta} = \mathbf{V}^T\mathbf{C}^{-T}(\mathbf{Y} - \widehat{\boldsymbol{\mu}})$, which has covariance matrix

$$\text{Var}(\boldsymbol{\zeta}) \approx \mathbf{V}^T\boldsymbol{\Psi}\mathbf{V} = \mathbf{I}_{n-k}. \tag{12}$$

Thus, we obtain a transformation of the residuals where the entries have approximately the same variance and are uncorrelated.

In practice ξ is also unknown, and the matrices $\boldsymbol{V}$ and $\boldsymbol{C}$ are functions of both the unknown parameters ξ and $\boldsymbol{\beta}_0$, so we plug in the values $\hat{\xi}$ and $\widehat{\boldsymbol{\beta}}$ estimated under the null hypothesis and call the resulting matrices $\widehat{\boldsymbol{V}}$ and $\widehat{\boldsymbol{C}}$. Finally we obtain

$\widehat{\boldsymbol{\zeta}} = \widehat{\boldsymbol{V}}^T \widehat{\boldsymbol{C}}^{-T}(Y - \widehat{\boldsymbol{\mu}})$. For each permutation replicate, we draw a random permutation matrix $\boldsymbol{\Pi}$ and obtain $\boldsymbol{\Pi}\widehat{\boldsymbol{\zeta}}$, a permuted version of $\widehat{\boldsymbol{\zeta}}$, which we then back-transform to the residual space by applying $\widehat{\boldsymbol{C}}^T \widehat{\boldsymbol{V}}$. Finally, we add on $\widehat{\boldsymbol{\mu}}$ in order to obtain the trait replicate:

$$\mathbf{Y}_\pi = \widehat{\boldsymbol{\mu}} + \widehat{\mathbf{C}}^T \widehat{\mathbf{V}} \boldsymbol{\Pi} \widehat{\mathbf{V}}^T \widehat{\mathbf{C}}^{-T}(\mathbf{Y} - \widehat{\boldsymbol{\mu}}). \tag{13}$$

Under the identity permutation, meaning that $\boldsymbol{\Pi} = \mathbf{I}$, we would recover the original response vector. We note that because $\widehat{\boldsymbol{\Gamma}}$ is just a diagonal matrix, $\widehat{\mathbf{C}}$ can be obtained from a factorization (e.g., cholesky or eigen decomposition) of $\widehat{\boldsymbol{\Sigma}} = \hat{\xi}\,\boldsymbol{\Phi} + (1 - \hat{\xi})\,\mathbf{I}$, which in turn could be obtained from a factorization of $\boldsymbol{\Phi}$ if desired, as was done in [9].

Examining the form of (13), there are three main steps used to obtain a transformation of the residuals with second-order exchangeable entries. The first is to center the response by using the estimated phenotypic mean. The second is to remove the correlation present due to polygenic effects, which is done by the pre-multiplication by $\widehat{\mathbf{C}}^{-T}$. The last step is to remove the correlation that is generated from using parameter estimates instead of the true values when centering, and is represented by the pre-multiplication by $\widehat{\mathbf{V}}^T$. It is possible that the last step might have only a minor impact on the method for large sample sizes and low number of parameters, though we did not investigate this. The replicates generated from this approach are quantitative, i.e., the binary nature of the response is not preserved during resampling. In subsection **Additional resampling methods considered**, we describe how the replicates can be transformed to be binary, and we consider both approaches in simulations. The fixed effects and covariance structure estimated under the null are preserved in the trait replicates, making our method applicable to general scenarios where there are important covariates and/or population structure.

LogMM-PQL

A logistic mixed model (LogMM) is a natural modeling choice for correlated binary data, and an alternative approach to BRASS for generating replicates of correlated binary data could be to fit a LogMM under the null hypothesis and then generate binary replicates from the fitted model. We have developed such an approach, which we call LogMM-PQL. The model underlying LogMM-PQL is that, conditional on $\mathbf{X}$, $\mathbf{G}$

and $\mathbf{u}$, Y_i is Bernoulli(m_i), independently across $i = 1, \ldots, n$, where m_i is the ith component of the vector

$$\mathbf{m} := E(\mathbf{Y}|\mathbf{X}, \mathbf{G}) = \mathrm{logit}^{-1}(\mathbf{X}\boldsymbol{\beta} + \mathbf{G}\gamma + \mathbf{u}), \tag{14}$$

where $\boldsymbol{\Phi}$, $\mathbf{Y}$, $\mathbf{X}$, $\mathbf{G}$, $\boldsymbol{\beta}$ and γ are as before; $\mathbf{u}$ is a multivariate normal random effects vector of length n with $\mathbf{u} \sim \mathrm{MVN}(0, \sigma^2\boldsymbol{\Phi})$ where σ^2 is an unknown scalar, and where γ will be set to zero because the model will be fitted under the null hypothesis of no effect.

The major difficulty with use of LogMMs in the GWAS context is that they are computationally challenging to fit as they involve high-dimensional integrals. Therefore, their use can involve a trade-off between computational feasibility and accuracy of the results. For the parameter estimation part of LogMM-PQL, we use a fast penalized quasi-likelihood (PQL) approach, as implemented in GMMAT [18]. With γ set to zero, we use GMMAT to obtain estimates $\widehat{\boldsymbol{\beta}}$ and $\hat{\sigma}^2$, and then generate trait replicates by drawing random vectors $\mathbf{Y}$ from the LogMM specified by equation (14) with $(\boldsymbol{\beta}, \gamma, \sigma^2)$ set equal to $(\widehat{\boldsymbol{\beta}}, 0, \hat{\sigma}^2)$. LogMM-PQL renders binary replicates that incorporate the mean and variance structure specified by the fitted logistic model.

Additional resampling methods considered

In addition to BRASS and LogMM-PQL, the other resampling methods we compare in simulations include the previously proposed MVNpermute [9,11], as well as a method we call Naive, which involves fitting a LMM whose form under the null is $\mathbf{Y} = \mathbf{X}\boldsymbol{\beta} + \mathbf{e}$, with $\mathbf{e} \sim \mathrm{MVN}(0, \sigma_1^2\boldsymbol{\Phi} + \sigma_2^2\mathbf{I})$. After obtaining the estimated coefficients $\widehat{\boldsymbol{\beta}}$, the Naive method replicates are obtained by permuting the residuals $(\mathbf{Y} - \mathbf{X}\widehat{\boldsymbol{\beta}})$, and adding back $\mathbf{X}\widehat{\boldsymbol{\beta}}$.

While the LogMM-PQL replicates are binary, those from BRASS, MVNpermute and Naive are not. However, we consider alternative approaches based on converting these replicates to binary, referred to as BRASS$_{mod}$, MVNpermute$_{mod}$ and Naive$_{mod}$, respectively, where the subscript "mod" indicates that the trait replicate has been modified to convert it to binary. To convert a continuous trait replicate $\mathbf{Y}_\pi$ to binary, a threshold is set and the values of $\mathbf{Y}_\pi$ above and below that threshold are converted to 1 and 0, respectively, with the threshold chosen so that the original trait $\mathbf{Y}$ and the

binary replicate $\mathbf{Y}_\pi$ have the same case/control proportions.

The features of the various methods are summarized and compared in S1 Table.

Modeling sample structure in the replicates

All 7 resampling strategies described above involve fitting a trait model that
incorporates the structure present in the sample. A common approach to modeling trait
correlation due to additive polygenic effects is to include a GRM in a LMM for the trait
so as to model the structure using random effects [19]. Alternatively, the top principal
components of a given GRM can be used as covariates to model the structure present
using fixed effects [20]. Here, we allow either of the two approaches or they can be
combined (i.e., both fixed and random effects can be used) to capture the sample
structure in the null model. To combine the two approaches, we use the top PCs from a
GRM as fixed effects in the null model and then build a new GRM that is adjusted to
exclude the effects of these top PCs in order to capture the leftover structure as random
effects. In the simulations and data analysis, we form the new GRM using equation 4 of
a previous work [21].

When generating replicates, it is of course important to accurately model the
structure present in the sample, but in addition, it is also important to avoid creating
new structure (not present in the data) as an artifact of the replication procedure. To
do this, we recommend the use of leave-one-chromosome-out (LOCO) GRMs [22, 23] in
the replication procedure. In that case, replicates would be generated separately for
each chromosome, with the SNPs for the given chromosome left out of the GRM. This
avoids introducing new trait-SNP association in the replicates.

Methods for simulation studies

We perform simulation studies to compare the effectiveness of the 7 proposed
permutation-based methods for generating binary trait replicates in the context of
correcting for multiple testing, with both population structure and pedigree structure
present in the sample. We consider a setting in which a binary trait is tested for
association with each of m markers and the aim is to assess the significance of the
smallest p-value out of the m association tests. This context would arise in testing

significance of a genomic region or in assessing genome-wide significance. We simulate non-causal markers that are tested for association with a binary trait and estimate the significance threshold for the top signal amongst them using the empirical distribution from trait replicates generated under the null hypothesis of no association. Genotype, trait and covariates are generated under multiple simulation settings. In real data applications, the true trait model is usually not known a priori, so we consider the effects of model misspecification due to (1) assuming a logistic model when the true model is a liability threshold model and (2) exclusion of an important covariate from the model. In addition, we investigate the impact of ascertainment in the sample as case-control studies often involve phenotype-based ascertainment which can introduce misspecification in the model as well.

For each setting, we perform a simulation study consisting of 4 stages: (1) data set creation, i.e., simulation of genotype and trait data under the null hypothesis of no association, with 20,000 data sets per setting; (2) generation of permutation replicates, in which we consider every possible pair (i, j) of simulated data set i and permutation method j, for $1 \leq i \leq 20,000$ and $1 \leq j \leq 7$, and for each of these, 10,000 permutation replicates are drawn for data set i using permutation method j; (3) analysis of permutation replicates, where for each trio (i, j, k) of simulated data set i, permutation method j, and permutation replicate k, $1 \leq k \leq 10,000$, the trait in permutation replicate k is tested for association with each of the m non-causal markers in data set i, the smallest p-value is recorded in each case, and then the resulting 10,000 p-values for each pair (i, j) are used to estimate a genomewide significance threshold (at level $\alpha = 10^{-3}$) for data set i based on permutation method j; and (4) evaluation of the type 1 error rate of the genomewide threshold for each permutation method j by comparing the smallest p-value for each data set i to the threshold determined in (3) for pair (i, j) and assessing whether the overall proportion of rejections among the 20,000 data sets is significantly different from α for method j. We now describe each of these stages.

For stage (1) data set creation, we simulate genotypes using a 2 sub-population Balding-Nichols model [24] with $F = .01$ and ancestral allele frequencies for SNPs drawn independently and uniformly on $(.2, .8)$. We simulate multiple pedigrees of the same configuration (S1 Fig), with equal numbers of pedigrees assigned to each of the two sub-populations. Founder alleles for a pedigree are assumed to be drawn from the

sub-population to which the pedigree is assigned, and gene dropping is used to 241
determine genotypes for other pedigree members. We simulate phenotypes according to 242
two different generative models for the trait. In each case, $M_c = 10^4$ causal SNPs are 243
simulated as well as 3 covariates: age, sex and an i.i.d. standard normal covariate. The 244
first generative model we use is logistic: 245

$$Y_i | \mathbf{X}_i, \mathbf{W}_i, \boldsymbol{\alpha} \sim \text{Bernoulli}(p_i), \text{ independently, with logit}(p_i) = \mathbf{X}_i \boldsymbol{\beta} + \mathbf{W}_i \boldsymbol{\alpha}, \quad (15)$$

where $\mathbf{X}_i$ is the covariate row vector for the ith individual; $\boldsymbol{\beta}$ are the fixed effects for 246
the covariates; $\mathbf{W}_i$ is a row vector representing the SNP information for the ith 247
individual corresponding to the M_c causal markers standardized to have mean 0 and 248
variance 1; $\boldsymbol{\alpha} = (\alpha_1, \cdots, \alpha_{M_c})^T \sim MVN(0, \sigma^2 \mathbf{I})$ represent the random effects of the M_c 249
causal markers with resulting total additive polygenic variance $\sigma_a^2 = \sigma^2 M_c$. The values 250
for $\boldsymbol{\beta}$ and σ_a^2 are chosen so that, considering the variance of logit(p_i) due to covariate 251
effects and additive polygenic random effects, the fraction of this variance due to 252
covariates is fixed at 20%, 40%, 60% or 80% (while the remaining fraction is due to 253
additive polygenic random effects), and so that Bernoulli error explains on average 254
about 20% of the total phenotypic variability. In addition, the effect size of the 255
standard normal covariate is set so that the p-value for its significance in a LMM Wald 256
test is on average 0.05. (This is the covariate that is set to be missing in the settings in 257
which an important covariate is missing from the observed data set.) The second 258
generative model we use is a liability threshold model: 259

$$Y_i = \mathbb{1}_{\{L_i > 0\}}, \text{ with } L_i = \mathbf{X}_i \boldsymbol{\beta} + \mathbf{W}_i \boldsymbol{\alpha} + \epsilon_i, \quad (16)$$

where L_i is the latent liability for individual i, and $\boldsymbol{\epsilon} = (\epsilon_1, \ldots, \epsilon_n)^T \sim MVN(0, \sigma_e^2 \mathbf{I})$. 260
The values of $\boldsymbol{\beta}$, σ_a^2, and σ_e^2 are chosen using analogous constraints as for the logistic 261
model, where the phenotypic variability due to $\boldsymbol{\epsilon}$ is constrained to be about 20% of the 262
total phenotypic variability. In both 15 and 16, larger values of σ_a^2 correspond to more 263
severe confounding effects of population/family structure on phenotype-genotype 264
association. For the settings in which ascertainment is used, we simulate data under the 265
model in (15) and select 500 cases and 500 controls at random to be retained in the 266

sample. In all settings, in addition to the M_c causal SNPs, $m = 100$ non-causal SNPs are generated as above and used for testing under the null hypothesis of no association. (Further details on the stage (1) simulations can be found in S1 Text.)

We now describe stage (2) generation of permutation replicates for each data set and for each of the 7 permutation methods (BRASS, LogMM-PQL, MVNpermute, Naive, BRASS_{mod}, MVNpermute_{mod}, and Naive_{mod}). For a given data set, the same covariate and GRM information is input to each of the 7 permutation methods. The GRM $\boldsymbol{\Phi}$ is calculated using the M_c causal markers, and then $D = 1$ top PC is removed from $\boldsymbol{\Phi}$ using equation 4 of a previous work [21], resulting in adjusted GRM $\tilde{\boldsymbol{\Phi}}$, which is the GRM input to each permutation method. In most settings, the 4 covariates input to each permutation method are the top PC plus the 3 covariates used to generate the trait. In the setting in which an important covariate is assumed to be missing from the data set, the 3 covariates input to each permutation method are the top PC, age and sex, with the normal covariate left out.

In stage (3), for each pair (i, j) of data set and permutation method, the 10^4 permutation replicates from method j are each tested for association against the panel of m SNPs in data set i, and the smallest p-value among the m SNPs is recorded for each permutation replicate, resulting in 10^4 p-values for pair (i, j). The genome-wide threshold value, corresponding to significance level α, for pair (i, j) is estimated by the 100αth percentile of the p-values observed for that pair. In principle, any association testing method could be used for the stage (3) association tests, provided that the same method is also used in stage (4). For the stage (3) and (4) association tests, we use a slightly modified version of CARAT [17], a binary trait mapping method, where the modification is in the estimation of σ_g^2, the genotypic variance parameter for CARAT. In the original CARAT paper [17] σ_g^2 is estimated by $\hat{\sigma}_g^2 = 2\hat{f}(1 - \hat{f})$ with $\hat{f} = .5\bar{G}$, where $\bar{G}$ is the sample average of $\boldsymbol{G}$ in the data. Our modification is to replace this with $\tilde{\sigma}_g^2 = \mathbf{G}^T \mathbf{P} \mathbf{G}/(n - D - 1)$ where D is the number of PCs removed from the GRM, and $\mathbf{P} = \tilde{\boldsymbol{\Phi}}^{-1} - \tilde{\boldsymbol{\Phi}}^{-1} \mathbf{Z}(\mathbf{Z}^T \tilde{\boldsymbol{\Phi}}^{-1} \mathbf{Z})^{-1} \mathbf{Z}^T \tilde{\boldsymbol{\Phi}}^{-1}$, where $\mathbf{Z}$ is the $n \times (D + 1)$ matrix whose columns are the D eigenvectors that are removed as well as a columns of ones. (The rationale for this modification is detailed in S1 Text.) When we apply CARAT, we use $\tilde{\boldsymbol{\Phi}}$ as the GRM, and we include as fixed effects the top PC plus the 3 covariates that were used to generate the trait. For the setting when an important covariate is assumed

to be missing from the data set, the 3 covariates used in the analysis of replicates are ²⁹⁹
the top PC, age and sex, with the normal covariate left out. ³⁰⁰

Finally, in stage (4), we evaluate the type 1 error of the genomewide threshold for ³⁰¹
each permutation method j by by comparing the p-value for each data set i to the ³⁰²
threshold determined in stage (3) for the pair (i, j) and assessing whether the overall ³⁰³
proportion of rejections out of 20,000 data sets is significantly different from the ³⁰⁴
nominal level α for method j. To make our massive simulation project more ³⁰⁵
computationally efficient, we actually used an adaptive procedure so that, e.g., if a ³⁰⁶
dataset is sufficiently far from the significance threshold based on analysis of the first ³⁰⁷
10^3 permutation replicates, it would not be necessary to analyze the remainder of the ³⁰⁸
10^4 permutation replicates (see S1 Text for details). ³⁰⁹

Verification and Comparison ³¹⁰

Assessment of type 1 error ³¹¹

We assess and compare the performance of the 7 proposed resampling methods in the ³¹²
context of correcting for multiple testing, with both population and pedigree structure ³¹³
present in the sample in addition to important covariates. Fig 1 shows the empirical ³¹⁴
type 1 error rates of all methods, compared to the nominal, when the true trait model is ³¹⁵
logistic. The type 1 error of BRASS is well-controlled in all settings. For LogMM-PQL, ³¹⁶
we observe significant inflation in the type 1 error rate, with the amount of inflation ³¹⁷
increasing as the confounding due to structure present in the sample increases. This ³¹⁸
likely occurs because the PQL model fitting approach tends to underestimate the ³¹⁹
variance component and hence, results in less correlation present in the simulated data ³²⁰
compared to that in the original data. We observe that permuting the residuals from a ³²¹
linear mixed model (Naive) leads in all settings to lack of control of the type 1 error. ³²²
More precisely, when covariates explain most of the variation on the logit scale, the ³²³
method is overly conservative and when instead polygenic effects explain most of the ³²⁴
variability, the type 1 error rate is significantly inflated. Moreover, we also find that ³²⁵
adjusting for sample correlation prior to permutation based on LMM residuals, as done ³²⁶
in MVNpermute, still leads to a very conservative test when covariates highly influence ³²⁷

Fig 1. Empirical Type 1 Error Rates with Logistic Model Including all Covariates at Nominal Level 0.01.
The error rate is based on 20,000 simulated replicates. The solid horizontal line represents the nominal level and the dashed lines represent rejection bounds outside of which the z-test comparing the estimated type 1 error to the nominal level is rejected at level .05. Estimates inside the rejections bounds are represented by circles and those outside the bounds are represented by triangles. Red, black and blue symbols represent inflated, well-controlled and conservative type 1 error rates, respectively. The proportion of variability on the logit scale attributable to polygenic effects vs. covariates is varied from 20 to 80% in increments of 20%.

the variability on the logit scale. This is most likely because a linear mixed model does not allow any dependence between the trait variance and the mean, which is influenced by the covariates. As covariates become less important, we do see somewhat better control of the type 1 error. For Naive$_{mod}$ and MVNpermute$_{mod}$, control of the type 1 error is improved compared to the corresponding original methods when covariates strongly influence the phenotypic mean, and for BRASS$_{mod}$, type 1 error control is

worse compared to BRASS. All of the methods except BRASS show inflated type 1 ₃₃₄
error in at least one setting, whereas BRASS maintains the correct type 1 error rate in ₃₃₅
all settings. ₃₃₆

Robustness to model misspecification: liability threshold vs. ₃₃₇ logistic model ₃₃₈

As the true model for the trait is generally unknown in an association study, we are ₃₃₉
interested in assessing robustness of the type 1 error results to model misspecification. ₃₄₀
Thus, we also perform simulations in which the phenotype is simulated from a liability ₃₄₁
threshold model, which is more dissimilar to the model fit by both BRASS and ₃₄₂
LogMM-PQL than logistic is. The type 1 error results for the liability threshold model ₃₄₃
are shown in Fig 2. The primary effect of the model misspecification in this case seems ₃₄₄
to be to make all the methods except Naive slightly more conservative. ₃₄₅

Robustness to missing an important covariate ₃₄₆

We assess the robustness of the type 1 error results when an important covariate is ₃₄₇
omitted from both the fitted model and the model used for generating replicates. As it ₃₄₈
is usually not known a priori which variables should be kept in the analysis, one would ₃₄₉
commonly try including different combinations of covariates to finally determine the ₃₅₀
ones to include in the final model. It may occur that one of the covariates has a ₃₅₁
moderate effect on the trait and leads to a p-value close to the significance threshold ₃₅₂
(e.g. 0.05). Hence, a judgment call would be required for whether to keep the covariate ₃₅₃
in the model and one may decide to exclude it. It is thus of interest to see how the ₃₅₄
proposed methods would fare in such a scenario as the replicates generated would come ₃₅₅
from a more misspecified model. ₃₅₆

The simulation results are displayed in Fig S1. Similarly to the previous model ₃₅₇
misspecification results, the general effect is to make the results more conservative for ₃₅₈
all permutation methods except Naive. ₃₅₉

Fig 2. Empirical Type 1 Error Rates with Liability Threshold Model at Nominal Level 0.01.
The error rate is based on 20,000 simulated replicates. The solid horizontal line represents the nominal level and the dashed lines represent rejection bounds outside of which the z-test comparing the estimated type 1 error to the nominal level is rejected at level .05. Estimates inside the rejections bounds are represented by circles and those outside the bounds are represented by triangles. Red, black and blue symbols represent inflated, well-controlled and conservative type 1 error rates, respectively. The proportion of variability on the liability scale attributable to polygenic effects vs. covariates is varied from 20 to 80% in increments of 20%.

Robustness to ascertainment

We determine the robustness of the proposed methods when trait-based ascertainment has been applied to the sample. This is commonly used in case-control studies where individuals are included in the sample based on their disease affection status, such as if the prevalence of the trait in the population is too low to obtain sufficient power.

The results are shown in Fig 3, where it can be seen that BRASS is the only method

Fig 3. Empirical Type 1 Error Rates with Ascertainment Present at Nominal Level 0.01.
The error rate is based on 20,000 simulated replicates. The solid horizontal line represents the nominal level and the dashed lines represent rejection bounds outside of which the z-test comparing the estimated type 1 error to the nominal level is rejected at level .05. Estimates inside the rejections bounds are represented by circles and those outside the bounds are represented by triangles. Red, black and blue symbols represent inflated, well-controlled and conservative type 1 error rates, respectively. The proportion of variability on the logit scale attributable to polygenic effects vs. covariates is varied from 20 to 80% in increments of 20%.

that retains good control of the type 1 error rate in all of the ascertainment settings ₃₆₆

considered. In contrast, both logMM-PQL and Naive$_{mod}$ show significantly inflated ₃₆₇

type 1 error in every setting, while both MVNpermute and Naive give excessively ₃₆₈

conservative results when covariates have a major impact on the trait and tend to give ₃₆₉

significantly inflated results when the effect of covariates is low. ₃₇₀

From the results across all the simulation settings, with and without ascertainment ₃₇₁

or model misspecification, it can be seen that BRASS provides the best type 1 error control and is the only method that does not have significantly inflated type 1 error in any setting.

Applications

We apply BRASS to the problem of determining genome-wide significance for GWAS of two different traits in domestic dog. Unlike with humans, a genome-wide threshold for significance in domestic dog GWAS has not been well established. We analyze data from a large domestic dog study [25], with 4,224 dogs representing over 150 breeds genotyped at 185,805 SNPs with multiple phenotypes recorded. We analyze two binary traits from this study, ED, for which the data contain 113 cases and 633 controls among 82 breeds, and IE, for which there are 34 cases and 168 controls from the Irish Wolfhound breed. For both traits, we compute a GRM after LD pruning and filtering for MAF $< .05$. For ED, we take the top 10 PCs out of the GRM and include them as covariates, and sex is included as a covariate for both traits. The data are then analyzed with CARAT. To generate trait replicates using BRASS, we create LOCO GRMs for each chromosome, where for ED we also take out the top 10 PCs in each case and include them as covariates, and sex is included as a covariate for both traits. For genome-wide significance assessment for each trait, we used BRASS to generate 10,000 replicates under the null hypothesis of no association. We analyzed each replicate in the same way that we analyzed the data and used the results of these analyses to form an empirical distribution of the test statistic under the null hypothesis. Genome-wide p-values were then estimated using the empirical distribution for each trait.

Manhattan plots of the p-values of the single-SNP tests for the observed data are presented in Fig 4 for both ED and IE phenotypes, along with the estimated genome-wide significance thresholds, which are estimated at nominal level 0.05 from the empirical distribution of the top association signal based on the trait replicates. The genomic control inflation factors for ED and IE are 1.02 and 0.98, respectively. For IE, we detect a previously-identified [25] 12Mb region on chromosome 4 (position 7.5-19.3 Mb) that reaches the BRASS genome-wide significance threshold, with nominal p-value 2.1×10^{-8} corresponding to a genomewide p-value of .0008 obtained from BRASS. For

A. ED

B. IE

Fig 4. Genome screen results in the domestic dog data.
Manhattan plots of the single-SNP association p-values using CARAT (on $-\log_{10}$ scale) for (A) ED and (B) IE, with genomic position on the x-axis. The horizontal lines represent estimated genome-wide thresholds at nominal level 0.05 using BRASS.

ED, we detect 3 loci, summarized in Table 1. Compared to a previous reference [25], we 402

detected one additional significant locus (rs23910667 on chromosome 35) that is not 403

detected there. 404

Table 1. Top association signals for ED

SNP	Chr	Position	MAF	Single SNP p-value	Genomewide p-value
rs9000666	26	16554631	.18	1.0×10^{-7}	.010
rs21895578	1	77938330	.17	1.5×10^{-7}	.014
rs23910667	35	7272763	.08	3.6×10^{-7}	.033

Computation time

In order to generate replicates from BRASS, the null prospective model of CARAT needs to be fitted, which includes specifying covariates and a GRM $\mathbf{\Phi}$. If not provided, obtaining the eigen decomposition of $\mathbf{\Phi}$ is the main computationally intensive step involved in fitting the null model [17]. However, this step (or one of equivalent computational complexity) is also typically performed when analyzing the data or when using permutation methods based on a linear or logistic mixed model. Once parameter null estimates are obtained, the transformed residuals $\hat{\boldsymbol{\zeta}}$ as well as the linear map $\widehat{\mathbf{C}}^T \widehat{\mathbf{V}}$ in (13) need to be computed; however, once obtained, they can be re-used when generating trait replicates. As the eigen decomposition of $\mathbf{\Phi}$, and hence $\widehat{\mathbf{\Sigma}}$, is obtained prior to fitting the null model, we use it to compute $\widehat{\mathbf{C}}$. Therefore, the most computationally intensive step is to obtain the eigenvectors of $\widehat{\mathbf{\Psi}}$ (the estimate for $\mathbf{\Psi}_0$ in (11)), which involves a time complexity of $O(n^3)$. By using the singular value decomposition of the matrix $\mathbf{W}$ in (11), which involves a time complexity of $O(nk^2)$ (assuming the number of covariates $k < n$), we are able to obtain the eigenvectors of $\widehat{\mathbf{\Psi}}$ without having to construct it and compute its eigen decomposition. Past this step, the additional cost for generating L replicates is $O(n^2 L)$; this computation can easily be parallelized to generate sets of traits replicates independently.

BRASS is implemented in a freely downloadable software package at `https://github.com/joellesophya/BRASS`. We report run times for BRASS in simulated data. Using a single processor on a machine with 6 core Intel Xeon 3.50 GHz CPUs and 32 GB RAM, on simulated data with 1,000, 5,000 and 10,000 individuals, it takes 1.6 s , 1 min and 4.6 min (275 s), respectively, to generate 1,000 trait replicates. As the use of BRASS will vary based on the context (e.g. multiple testing correction, region-based association testing), the computational burden involved in comparing the statistic between the observed trait and its replicates will vary based on what statistic is being considered.

Discussion

In genetic studies of binary traits, it may often be of interest to assess significance of a test statistic whose distribution is unknown or not well-approximated, or to assess

significance of the maximum of many correlated tests. Examples could include test statistics based on machine learning methods [1], rare variant tests [3], test statistics that involve data-adaptive weights [4], and assessment of genomewide significance [5] in a variety of situations such as inclusion of admixed individuals in GWAS [6]. Permutation tests are an attractive and popular approach, but in the presence of population structure, cryptic relatedness and/or family structure, which are all common sources of confounding in genetic association studies, ordinary permutation tests are usually not appropriate as they fail to retain the structure present in the data.

We propose BRASS, a novel permutation-based resampling procedure for generating binary trait replicates in samples with population structure, cryptic relatedness and/or family structure. BRASS allows for covariates and ascertainment, and it accommodates a wide range of test statistics. It uses an estimating equation approach that can be viewed as a hybrid of logistic regression and linear mixed-effects model methods, which allows for use of a GRM and/or principal components or other ancestry-informative covariates to account for sample structure. BRASS relies on obtaining an invertible transformation of phenotypic residuals to achieve approximate second-order exchangeability. After permutation, new trait replicates are obtained by inverting the transformation, in order to preserve structure present in the original sample. BRASS differs from existing methods such as Naive and MVNpermute [11] in that it incorporates key features of a binary trait such as nonlinear effects of covariates and dependence of the variance on the mean, which we show improve the performance of BRASS on binary traits.

In simulations, we demonstrate the superior type 1 error control of BRASS compared to other methods. Across simulation settings, we find that BRASS maintains control of the type 1 error under varying amounts of population structure, familial relatedness, ascertainment and sources of trait model misspecification. In contrast, all 6 of the other methods we consider show significant inflation of type 1 error in multiple settings.

In addition to showing significant inflation of type 1 error in some settings, the Naive and MVNpermute methods, which are two of the 4 methods that are based on an LMM, are excessively conservative when covariate effects are substantial. Taken together, the results show that simply accounting for the sample structure with an LMM, without incorporating key features associated with the binary nature of the trait

such as nonlinear scale for covariate effects and the dependence of the variance on the ⁠467

mean, is not sufficient to obtain replicates that correctly estimate the null distribution ⁠468

of the statistic. Secondarily, we find that when polygenic effects are important, ⁠469

adjusting for the correlation in the residuals prior to permutation, as is done in BRASS ⁠470

and MVNpermute leads to better control of the type 1 error compared to ignoring it, as ⁠471

is done when permuting the raw residuals from a LMM (Naive method). Furthermore, ⁠472

we find that the additional step of binarizing the quantitative replicates results in ⁠473

improvement in the accuracy of the type 1 error rate for the LMM-based methods when ⁠474

the polygenic component has a low impact on the trait distribution relative to covariates, ⁠475

but does not control the type 1 error when more structure is present in the sample. ⁠476

In principle, use of a logistic mixed model would seem to be a natural and promising ⁠477

alternative to BRASS for fitting the binary trait data and generating appropriate ⁠478

replicates that preserve the correlation structure. However, for LogMM-PQL, we ⁠479

observe significant inflation in the type 1 error rate, with the amount of inflation ⁠480

increasing as the confounding due to structure present in the sample increases. This is ⁠481

likely explained by the use of PQL to fit the LogMM, as PQL is known to be fast but to ⁠482

give biased estimates in binary data [26]. However, fitting a LogMM is computationally ⁠483

intensive and requires a trade-off between computational time and accuracy of the ⁠484

approximation of the high-dimensional integral involved. Algorithms with higher ⁠485

accuracy than PQL (e.g. the Laplace approximation [27] or Gauss-Hermite ⁠486

quadrature [28]) lead to increased computational burden and do not scale well for even ⁠487

moderately large GWAS. In contrast, BRASS is seen to be extremely fast ⁠488

computationally. ⁠489

We applied BRASS in the context of multiple testing correction in association ⁠490

mapping studies of ED and IE in domestic dogs. In the analysis of ED, we detected ⁠491

genomewide significant association with 3 loci, 2 of which were previously reported [25], ⁠492

with the 3rd locus also reaching significance based on the BRASS threshold. In the ⁠493

analysis of IE, we detected a 12 Mb region on chromosome 4 that reached the BRASS ⁠494

genome-wide significance threshold and has been previously associated with IE [25]. For ⁠495

the ED data analysis, breed information was available, but we chose to instead use PCs ⁠496

as ancestry informative covariates, due to apparent inaccuracies in the breed labeling ⁠497

information as well as the presence of dogs of mixed or unknown breed in the data set. ⁠498

Interestingly, the genome-wide p-value threshold estimated using replicates from BRASS for the ED trait in a sample of 82 breeds was about half that for the IE trait in a single breed. This is expected given the extremely different population structure between the two samples.

BRASS is designed as a very flexible tool that can be used in a range of situations in which replicates of correlated binary data are needed. Depending on the application, one can consider a different approach to capture the structure present in the sample, as the model used by BRASS can accommodate both covariates and random effects. In the simulation studies and data analysis, we chose to partition the relatedness into random and fixed effects and include both in our model. A previously proposed method, PC-Air [29], obtains PCs from a derived subset of mutually unrelated subjects that are representative of the ancestral diversity present in the sample, so as to ensure that the top PCs will only capture distant genetic relatedness (i.e. population structure). Such methods could easily be used to obtain the population structure information input to BRASS, if desired.

Software availability

Software implementing BRASS is available at
`https://github.com/joellesophya/BRASS`.

Acknowledgments

This work was supported by NIH grant R01 HG001645 (to M.S.M.). This study makes use of data generated from a previous study. [25] The data are available from the Dryad online repository at https://datadryad.org/resource/doi:10.5061/dryad.266k4.

References

1. Li J, Malley JD, Andrew AS, Karagas MR, Moore JH. Detecting gene-gene interactions using a permutation-based random forest method. BioData Mining. 2016;9:14.

2. Cui T, Mekkaoui KE, Reinvall J, Havulinna AS, Marttinen P, Kaski S. Gene–gene interaction detection with deep learning. Communications Biology. 2022; p. 1238.

3. Richter F, Morton S, Kim S, Kitaygorodsky A, Wasson L, KM C, et al. Genomic analyses implicate noncoding de novo variants in congenital heart disease. Nature Genetics. 2020;52:769–777.

4. Lee S, Abecasis GR, Boehnke M, Lin X. Rare-variant association analysis: Study designs and statistical tests. American Journal of Human Genetics. 2014;95(1):5–23. doi:10.1016/j.ajhg.2014.06.009.

5. Dudbridge F, Gusnanto A. Estimation of significance thresholds for genomewide association scans. Genetic Epidemiology. 2008;32(3):227–234. doi:10.1002/gepi.20297.

6. Atkinson EG, Maihofer AX, Kanai M, Martin AR, Karczewski K, Santoro ML, et al. Tractor uses local ancestry to enable the inclusion of admixed individuals in GWAS and to boost power. Nature Genetics. 2021;53:195–204.

7. Sheng X, Guan Y, Ma Z, Wu J, Liu H, Qiu C, et al. Mapping the genetic architecture of human traits to cell types in the kidney identifies mechanisms of disease and potential treatments. Nature Genetics. 2021;53:1322–1333.

8. Astle W, Balding DJ. Population structure and cryptic relatedness in genetic association studies. Statistical Science. 2009; p. 451–471.

9. Abney M, Ober C, McPeek MS. Quantitative-Trait Homozygosity and Association Mapping and Empirical Genomewide Significance in Large, Complex Pedigrees: Fasting Serum-Insulin Level in the Hutterites. The American Journal of Human Genetics. 2002;70(4):920–934. doi:10.1086/339705.

10. Churchill GA, Doerge RW. Naive application of permutation testing leads to inflated type I error rates. Genetics. 2008;178(1):609–610. doi:10.1534/genetics.107.074609.

11. Abney M. Permutation testing in the presence of polygenic variation. Genetic Epidemiology. 2015;39(4):249–258. doi:10.1002/gepi.21893.

12. Bianchi M, Dahlgren S, Massey J, Dietschi E, Kierczak M, Lund-Ziener M, et al. A multi-breed genome-wide association analysis for canine Hypothyroidism identifies a shared major risk locus on CFA12. PLoS ONE. 2015;10(8):e0134720. doi:10.1371/journal.pone.0134720.

13. Melin M, Rivera P, Arendt M, Elvers I, Murén E, Gustafson U, et al. Genome-Wide Analysis Identifies Germ-Line Risk Factors Associated with Canine Mammary Tumours. PLoS Genetics. 2016;12(5):e1006029. doi:10.1371/journal.pgen.1006029.

14. Safra N, Bassuk AG, Ferguson PJ, Aguilar M, Coulson RL, Thomas N, et al. Genome-Wide Association Mapping in Dogs Enables Identification of the Homeobox Gene, NKX2-8, as a Genetic Component of Neural Tube Defects in Humans. PLoS Genetics. 2013;9(7):e1003646. doi:10.1371/journal.pgen.1003646.

15. Tengvall K, Kierczak M, Bergvall K, Olsson M, Frankowiack M, Farias FHG, et al. Genome-Wide Analysis in German Shepherd Dogs Reveals Association of a Locus on CFA 27 with Atopic Dermatitis. PLoS Genetics. 2013;9(5):e1003475. doi:10.1371/journal.pgen.1003475.

16. Zhong S, Jiang D, McPeek MS. CERAMIC: Case-Control Association Testing in Samples with Related Individuals, Based on Retrospective Mixed Model Analysis with Adjustment for Covariates. PLoS Genetics. 2016;12(10):e1006329. doi:10.1371/journal.pgen.1006329.

17. Jiang D, Zhong S, McPeek MS. Retrospective Binary-Trait Association Test Elucidates Genetic Architecture of Crohn Disease. American Journal of Human Genetics. 2016;98(2):243–255. doi:10.1016/j.ajhg.2015.12.012.

18. Chen H, Wang C, Conomos MP, Stilp AM, Li Z, Sofer T, et al. Control for Population Structure and Relatedness for Binary Traits in Genetic Association Studies via Logistic Mixed Models. American Journal of Human Genetics. 2016;98(4):653–666. doi:10.1016/j.ajhg.2016.02.012.

19. Thornton TA. Statistical methods for genome-wide and sequencing association studies of complex traits in related samples. Current Protocols in Human Genetics. 2015;2015(January):1.28.1–1.28.9. doi:10.1002/0471142905.hg0128s84.

20. Price AL, Patterson NJ, Plenge RM, Weinblatt ME, Shadick NA, Reich D. Principal components analysis corrects for stratification in genome-wide association studies. Nature Genetics. 2006;38(8):904–909. doi:10.1038/ng1847.

21. Conomos MP, Reiner AP, Weir BS, Thornton TA. Model-free Estimation of Recent Genetic Relatedness. American Journal of Human Genetics. 2016;98(1):127–148. doi:10.1016/j.ajhg.2015.11.022.

22. Listgarten J, Lippert C, Kadie CM, Davidson RI, Eskin E, D H. Improved linear mixed models for genome-wide association studies. Nature Methods. 2012;9:525–526.

23. Yang J, Zaitlen NA, Goddard ME, Visscher PM, Price AL. Advantages and pitfalls in the application of mixed-model association methods. Nature Genetics. 2014;46:100–106.

24. Balding DJ, Nichols RA. A method for quantifying differentiation between populations at multi-allelic loci and its implications for investigating identity and paternity. Genetica. 1995;96(1-2):3–12. doi:10.1007/BF01441146.

25. Hayward JJ, Castelhano MG, Oliveira KC, Corey E, Balkman C, Baxter TL, et al. Complex disease and phenotype mapping in the domestic dog. Nature Communications. 2016;7. doi:10.1038/ncomms10460.

26. Rodriguez G, Goldman N. Improved estimation procedures for multilevel models with binary response: a case-study. Journal of the Royal Statistical Society: Series A (Statistics in Society). 2001;164(2):339–355. doi:10.1111/1467-985X.00206.

27. Raudenbush SW, Yang ML, Yosef M. Maximum Likelihood for Generalized Linear Models with Nested Random Effects via High-Order, Multivariate Laplace Approximation. Journal of Computational and Graphical Statistics. 2000;9(1):141–157. doi:10.1080/10618600.2000.10474870.

28. Rabe-hesketh S, Skrondal A, Pickles A. Reliable estimation of generalized linear mixed models using adaptive quadrature. Stata Journal. 2002;2(1):1–21.

29. Conomos MP, Miller MB, Thornton TA. Robust inference of population structure for ancestry prediction and correction of stratification in the presence of relatedness. Genetic Epidemiology. 2015;39(4):276–293. doi:10.1002/gepi.21896.

Supporting information

S1 Text.

S1 Fig. Three-generation pedigree used in the simulation studies.

S2 Fig. Empirical Type 1 Error Rates Omitting a Relevant Covariate at Nominal Level 0.01.

S1 Table. Differences between the seven resampling methods compared in the simulation studies.