BRASS: permutation methods for binary traits in genetic association studies with structured samples

Joelle Mbatchou[1,2], Mark Abney[3], Mary Sara McPeek[2,3*]

1 Regeneron Genetics Center, Tarrytown, NY, USA

2 Department of Statistics, The University of Chicago, Chicago, IL, USA

3 Department of Human Genetics, The University of Chicago, Chicago, IL, USA

* mcpeek@uchicago.edu

Abstract

In genetic association analysis of complex traits, permutation testing can be a valuable tool for assessing significance when the distribution of the test statistic is unknown or not well-approximated. This commonly arises, e.g, in tests of gene-set, pathway or genome-wide significance, or when the statistic is formed by machine learning or data adaptive methods. Existing applications include eQTL mapping, association testing with rare variants, inclusion of admixed individuals in genetic association analysis, and epistasis detection among many others. For genetic association testing in samples with population structure and/or relatedness, use of naive permutation can lead to inflated type 1 error. To address this in quantitative traits, the MVNpermute method was developed. However, for association mapping of a binary trait, the relationship between the mean and variance makes both naive permutation and the MVNpermute method invalid. We propose BRASS, a permutation method for binary traits, for use in association mapping in structured samples. In addition to modeling structure in the sample, BRASS allows for covariates, ascertainment and simultaneous testing of multiple markers, and it accommodates a wide range of test statistics. In simulation studies, we compare BRASS to other permutation and resampling-based methods in a range of scenarios that include population structure, familial relatedness, ascertainment

and phenotype model misspecification. In these settings, we demonstrate the superior control of type 1 error by BRASS compared to the other 6 methods considered. We apply BRASS to assess genome-wide significance for association analyses in domestic dog for elbow dysplasia (ED) and idiopathic epilepsy (IE). For both traits we detect previously identified associations, and in addition, for ED, we detect significant association with a SNP on chromosome 35 that was not detected by previous analyses, demonstrating the potential of the method.

Author summary

To determine whether genetic association with a trait is significant, permutation methods are an attractive and popular approach when analytic methods based on distributional assumptions are not available, e.g., when applying machine learning or data adaptive methods, or when performing a multiple testing correction, e.g., to assess region-wide or genome-wide significance in association mapping studies. Existing applications include eQTL mapping, association testing with rare variants, inclusion of admixed individuals in genetic association analysis, and detection of genetic interaction among many others. However, when there is population structure in the sample, naive permutation of the data can lead to inflated significance of the association results. For continuous traits, linear mixed-model based approaches have been proposed for permutation-based tests that can also adjust for sample structure; however, these do not remain valid when applied to binary traits, as key features of binary data are not well accounted for. We propose BRASS, a permutation-based testing method for binary data that incorporates important characteristics of binary data in the trait model, can accommodate relevant covariates and ascertainment, and adjusts for the presence of structure in the sample. In simulations, we demonstrate the superior control of type 1 error by BRASS compared to other methods, and we apply BRASS in the context of correcting for multiple testing in two genome-wide association studies in domestic dog: one for elbow dysplasia and one for idiopathic epilepsy.

Introduction

In genome-wide association studies (GWAS), the primary objective is generally to identify associations between a phenotype of interest and genetic variants. This involves assessing the p-values of certain test statistics by deriving the appropriate null distribution or an asymptotic approximation to it. However, this is not always feasible as the distribution may be intractable. This can arise when the test statistics result from black-box machine learning methods [1,2], or in region-based tests where association signals over multiple sites are combined, e.g., rare variant tests [3], and/or when the test statistic involves data-adaptive weights [4]. A further limitation can arise when, even if the distribution (or the asymptotic distribution) of the test statistic is known for single tests, significance needs to be assessed for the maximum of many correlated tests. This occurs in genome scans to establish a genome-wide significance threshold, where the linkage disequilibrium present between the markers induces correlation between the association tests [5–7].

To overcome these limitations, a common approach is to perform permutation testing so as to obtain replicates of the data under the null hypothesis from which an empirical distribution can be derived [6,7]. A fundamental assumption underlying permutation testing is exchangeability of the subjects in the sample, which is often approximately satisfied in population-based samples. However, this assumption can be violated in the presence of genetic structure in the sample (e.g., kinship, cryptic relatedness, and/or population structure), as it will introduce correlation in the sample, including in the phenotype values through polygenic effects [8,9]. Hence, naive application of permutation testing will usually not preserve the correlation structure and can result in inflated type 1 error rates [10] (though this can be avoided if, for example, all subjects in the sample are equally related [11]). We consider here the problem of permutation testing for a binary trait in the presence of polygenic effects in a sample with population structure, cryptic and/or family relatedness. For quantitative traits, a permutation-based test that adjusts for the correlation structure has previously been proposed [9,11]. The approach, called MVNpermute, is based on a linear mixed model (LMM) and is applicable for multivariate-normal data. It incorporates genetic relatedness through the inclusion of random effects in the LMM and can also adjust for

covariates. However, as MVNpermute is primarily designed for quantitative traits that ₃₂
are multivariate-normal, there will be more model misspecification when applied to ₃₃
binary traits as the binary nature of the data is not incorporated into the LMM. ₃₄

We propose BRASS (for "binary trait resampling method adjusting for sample ₃₅
structure"), a permutation procedure for a binary trait which incorporates both ₃₆
covariates and the correlation structure present in the sample. In contrast to the ₃₇
LMM-based approach, it accommodates the binary nature of the trait through a ₃₈
quasi-likelihood framework that considers the effect of covariates on a logit scale in the ₃₉
mean structure as well as the relationship between the trait mean and its variance, both ₄₀
of which are important features of binary data (see, e.g., [12] for importance of modeling ₄₁
covariates in binary data). Hence, BRASS benefits from less model misspecification in ₄₂
the presence of important covariates relative to the LMM-based approach. Here, we ₄₃
show that the permutation-based replicates obtained with BRASS are able to maintain ₄₄
correct type 1 error control, and we compare it with several alternative approaches. We ₄₅
demonstrate the use of our method in a context of assessing genome-wide significance in ₄₆
domestic dog GWAS. In animal studies, permutation testing is commonly used when ₄₇
there is not a consensus threshold for genome-wide significance [13–16], and it is also ₄₈
used in some human studies when the consensus threshold might not be applicable [6]. ₄₉
We apply BRASS to perform association mapping in two GWAS of domestic dogs, one ₅₀
for elbow dysplasia (ED [OMIA: 000330-9615]) and and one for idiopathic epilepsy (IE ₅₁
[OMIA: 000344-9615]). ₅₂

Description of the Methods ₅₃

Overview of BRASS method ₅₄

We consider the problem of resampling a binary trait variable that is correlated across ₅₅
individuals, where the correlation can arise from, e.g., population structure or related ₅₆
individuals. Our aim is to derive replicates of the trait under the null hypothesis of no ₅₇
association while accounting for the correlation that is present in the sample but whose ₅₈
structure is unknown. To obtain these replicates, we start by modeling the response ₅₉
using a previously described quasi-likelihood framework for correlated binary ₆₀

traits [17, 18],

$$\boldsymbol{\mu} := \mathbb{E}(\mathbf{Y}|\mathbf{X}, \mathbf{G}) = \text{logit}^{-1}(\mathbf{X}\boldsymbol{\beta} + \mathbf{G}\gamma), \text{ and} \tag{1}$$

$$\boldsymbol{\Omega} := \text{Var}(\mathbf{Y}|\mathbf{X}, \mathbf{G}) = \boldsymbol{\Gamma}^{1/2}\boldsymbol{\Sigma}\boldsymbol{\Gamma}^{1/2}. \tag{2}$$

where $\mathbf{Y} = (Y_1,...,Y_n)^T$ denotes the phenotype vector for n subjects, $\mathbf{X}$ is the $n \times k$
matrix of k covariates (including an intercept term), which is assumed to be of rank k,
$\mathbf{G} = (G_1,...,G_n)^T$ is the vector of genotypes at the marker of interest, where G_i denotes
the minor allele count for the ith individual, $\boldsymbol{\beta}$ is a k-length vector representing the
unknown effects of covariates, γ represents the unknown effect of the genetic marker, $\boldsymbol{\Gamma}$
is an $n \times n$ diagonal matrix with ith diagonal entry $\mu_i(1 - \mu_i)$ where μ_i is the ith
element of $\boldsymbol{\mu}$, the $n \times n$ matrix $\boldsymbol{\Sigma}$ is given by

$$\boldsymbol{\Sigma} = \xi\boldsymbol{\Phi} + (1 - \xi)\mathbf{I}_n, \tag{3}$$

where the unknown scalar parameter $\xi \in [0, 1]$ can be viewed as a heritability parameter,
and the $n \times n$ matrix $\boldsymbol{\Phi}$ is generally a kinship or genetic relatedness matrix (GRM).

The framework characterized by (1) and (2) allows for modeling important
covariates, such as biological or ancestry informative covariates, in the mean structure,
while the residual correlation between subjects is captured directly within the variance
structure by the inclusion of $\boldsymbol{\Phi}$ in (3). Furthermore, by including the matrix $\boldsymbol{\Gamma}$ in (2),
this framework models the dependence of the phenotypic variance on the mean, a key
feature of binary data.

Since we wish to simulate replicates of the binary trait under the null hypothesis of
no association, $H_0 : \gamma = 0$, the remaining unknown parameters are $\boldsymbol{\beta}$ and ξ, which need
to be estimated from the data. The null estimate $(\widehat{\boldsymbol{\beta}}, \hat{\xi})$ is easily obtained by iteratively
solving the following system of estimating equations [18] under the constraint $\gamma = 0$,

$$\boldsymbol{D}_{\boldsymbol{\beta}}^T\boldsymbol{\Omega}^{-1}(\mathbf{Y} - \boldsymbol{\mu}) = \mathbf{0}, \tag{4}$$

$$(\mathbf{Y} - \boldsymbol{\mu})^T\boldsymbol{\Gamma}^{-1/2}\boldsymbol{\Sigma}^{-1}(\boldsymbol{\Phi} - \mathbf{I})\boldsymbol{\Sigma}^{-1}\boldsymbol{\Gamma}^{-1/2}(\mathbf{Y} - \boldsymbol{\mu}) = \text{trace}(\boldsymbol{\Sigma}^{-1}(\boldsymbol{\Phi} - \mathbf{I})), \tag{5}$$

where $\boldsymbol{D_\beta} = \frac{\partial \boldsymbol{\mu}}{\partial \boldsymbol{\beta}}$ is a Jacobian of the conditional mean of the trait with respect to $\boldsymbol{\beta}$. In our model with a logit link function, $\boldsymbol{D_\beta} = \boldsymbol{\Gamma X}$.

The justification underlying an ordinary permutation test is an assumption of exchangeability under the null hypothesis, which does not generally hold in most situations of correlated data. Instead, we develop and apply an approximate permutation test based on second-order exchangeability, in which we permute a vector, $\hat{\boldsymbol{\zeta}}$, whose entries each have the same mean and variance and are uncorrelated with each other. Our approach builds on previous work [9, 11] that was limited to quantitative traits, and we extend the ideas to binary traits.

We start by fitting the null model given by Eq 1 and 2 with $\gamma = 0$, and we let $\mathbf{Y} - \hat{\boldsymbol{\mu}}$ denote the resulting residual vector. We aim to map $\mathbf{Y} - \hat{\boldsymbol{\mu}}$ to a lower-dimensional orthonormal space in which its entries will be second-order exchangeable, with the result of this map being $\hat{\boldsymbol{\zeta}}$. To form $\hat{\boldsymbol{\zeta}}$, we need to first obtain the approximate covariance matrix of $\mathbf{Y} - \hat{\boldsymbol{\mu}}$ under the null hypothesis, which is challenging because $\mathbf{Y} - \hat{\boldsymbol{\mu}}$ has no closed-form expression as a function of the data.

To obtain the approximate covariance matrix, we first fix ξ and let $U(\boldsymbol{\beta}) = \boldsymbol{D_\beta}^T \boldsymbol{\Omega}^{-1}(\mathbf{Y} - \boldsymbol{\mu})$, which corresponds to the quasi-score function for $\boldsymbol{\beta}$. Under the null hypothesis of $\gamma = 0$, let $\boldsymbol{\beta}_0$ denote the true value of $\boldsymbol{\beta}$, and let $\boldsymbol{\mu}_0, \boldsymbol{\Gamma}_0, \mathbf{D}_0$ and $\boldsymbol{\Omega}_0$ correspond to $\boldsymbol{\mu}, \boldsymbol{\Gamma}, \mathbf{D}_\beta$ and $\boldsymbol{\Omega}$, respectively, evaluated at $\boldsymbol{\beta}_0$ and $\gamma = 0$. Similarly, let $\hat{\boldsymbol{\mu}}, \hat{\boldsymbol{\Gamma}}, \hat{\mathbf{D}}$ and $\hat{\boldsymbol{\Omega}}$ be the same quantities evaluated at $\hat{\boldsymbol{\beta}}$ and $\gamma = 0$. We apply a Taylor series expansion to $U(\boldsymbol{\beta})$ around $\boldsymbol{\beta}_0$, evaluated at $\hat{\boldsymbol{\beta}}$:

$$U(\hat{\boldsymbol{\beta}}) \approx U(\boldsymbol{\beta}_0) + \frac{\partial U(\boldsymbol{\beta})}{\partial \boldsymbol{\beta}}\bigg|_{\boldsymbol{\beta}=\boldsymbol{\beta_0}} \cdot (\hat{\boldsymbol{\beta}} - \boldsymbol{\beta}_0). \tag{6}$$

Using the fact that $U(\hat{\boldsymbol{\beta}}) = \mathbf{0}$, replacing the Jacobian in (6) by its expectation (similar to Fisher scoring), and solving for $\hat{\boldsymbol{\beta}} - \boldsymbol{\beta}_0$ we get,

$$\hat{\boldsymbol{\beta}} - \boldsymbol{\beta}_0 \approx - \left\{ \mathbb{E}\left[\frac{\partial U(\boldsymbol{\beta})}{\partial \boldsymbol{\beta}}\right]\bigg|_{\boldsymbol{\beta}=\boldsymbol{\beta_0}} \right\}^{-1} U(\boldsymbol{\beta}_0),$$
$$= (\mathbf{D}_0^T \, \boldsymbol{\Omega}_0^{-1} \, \mathbf{D}_0)^{-1} \mathbf{D}_0^T \, \boldsymbol{\Omega}_0^{-1}(\mathbf{Y} - \boldsymbol{\mu}_0). \tag{7}$$

As $\hat{\boldsymbol{\mu}}$ is a non-linear function of $\hat{\boldsymbol{\beta}}$, a Taylor series expansion is performed on $\boldsymbol{\mu}$

around $\boldsymbol{\beta}_0$, and evaluated at $\widehat{\boldsymbol{\beta}}$, 105

$$\widehat{\boldsymbol{\mu}} \approx \boldsymbol{\mu}_0 + \mathbf{D}_0 \, (\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}_0), \tag{8}$$

Combining (7) and (8), we get for the residual vector 106

$$\begin{aligned}
\mathbf{Y} - \widehat{\boldsymbol{\mu}} &= \mathbf{Y} - \boldsymbol{\mu}_0 - (\widehat{\boldsymbol{\mu}} - \boldsymbol{\mu}_0), \\
&\approx \left[\mathbf{I} - \mathbf{D}_0 \, (\mathbf{D}_0^T \, \boldsymbol{\Omega}_0^{-1} \, \mathbf{D}_0)^{-1} \mathbf{D}_0^T \, \boldsymbol{\Omega}_0^{-1} \right] (\mathbf{Y} - \boldsymbol{\mu}_0). \tag{9}
\end{aligned}$$

As a result, we can approximate the covariance matrix of the residuals as 107

$$\mathrm{Var}(\mathbf{Y} - \widehat{\boldsymbol{\mu}}) \approx \boldsymbol{\Omega}_0 - \boldsymbol{\Gamma}_0 \mathbf{X} \, (\mathbf{X}^T \boldsymbol{\Gamma}_0 \, \boldsymbol{\Omega}_0^{-1} \, \boldsymbol{\Gamma}_0 \mathbf{X})^{-1} \mathbf{X}^T \boldsymbol{\Gamma}_0. \tag{10}$$

The correlation present in $(\mathbf{Y} - \widehat{\boldsymbol{\mu}})$ arises from two sources: that introduced by $\boldsymbol{\Phi}$ 108
through $\boldsymbol{\Omega}_0$ and that introduced from using the estimated mean $\widehat{\boldsymbol{\mu}}$ instead of the true 109
unknown mean $\boldsymbol{\mu}_0$. We use a factorization $\mathbf{C}$ of $\boldsymbol{\Omega}_0$, with $\boldsymbol{\Omega}_0 = \mathbf{C}^T \mathbf{C}$, to remove the 110
correlation due to $\boldsymbol{\Phi}$ and obtain, 111

$$\mathrm{Var}[\mathbf{C}^{-T}(\mathbf{Y} - \widehat{\boldsymbol{\mu}})] \approx \mathbf{I} - \mathbf{W}(\mathbf{W}^T \mathbf{W})^{-1} \mathbf{W}^T = \boldsymbol{\Psi}_0, \tag{11}$$

where $\mathbf{W} = \mathbf{C}^{-T} \boldsymbol{\Gamma}_0 \mathbf{X}$. The matrix $\boldsymbol{\Psi}_0$ in (11) is symmetric, idempotent and of rank 112
$n - k$, so it can be expressed as $\boldsymbol{\Psi}_0 = \mathbf{V}\mathbf{V}^T$ where the columns of $\mathbf{V}$ contain the 113
eigenvectors of $\boldsymbol{\Psi}_0$ corresponding to eigenvalue 1, and $\mathbf{V}^T \mathbf{V} = \mathbf{I}_{n-k}$. We use $\mathbf{V}^T$ to 114
remove the remaining correlation that is driven by parameter estimation. 115

Applying these two linear transformations, we obtain $\boldsymbol{\zeta} = \mathbf{V}^T \mathbf{C}^{-T}(\mathbf{Y} - \widehat{\boldsymbol{\mu}})$, which 116
has covariance matrix 117

$$\mathrm{Var}(\boldsymbol{\zeta}) \approx \mathbf{V}^T \boldsymbol{\Psi} \mathbf{V} = \mathbf{I}_{n-k}. \tag{12}$$

Thus, we obtain a transformation of the residuals where the entries have approximately 118
the same variance and are uncorrelated. 119

In practice ξ is also unknown, and the matrices $\boldsymbol{V}$ and $\boldsymbol{C}$ are functions of both the 120
unknown parameters ξ and $\boldsymbol{\beta}_0$, so we plug in the values $\hat{\xi}$ and $\widehat{\boldsymbol{\beta}}$ estimated under the 121
null hypothesis and call the resulting matrices $\widehat{\boldsymbol{V}}$ and $\widehat{\boldsymbol{C}}$. Finally we obtain 122

$\widehat{\boldsymbol{\zeta}} = \widehat{\boldsymbol{V}}^T \widehat{\boldsymbol{C}}^{-T} (Y - \widehat{\boldsymbol{\mu}})$. For each permutation replicate, we draw a random permutation ₁₂₃

matrix $\boldsymbol{\Pi}$ and obtain $\boldsymbol{\Pi}\widehat{\boldsymbol{\zeta}}$, a permuted version of $\widehat{\boldsymbol{\zeta}}$, which we then back-transform to the ₁₂₄

residual space by applying $\widehat{\boldsymbol{C}}^T \widehat{\boldsymbol{V}}$. Finally, we add on $\widehat{\boldsymbol{\mu}}$ in order to obtain the trait ₁₂₅

replicate: ₁₂₆

$$\mathbf{Y}_\pi = \widehat{\boldsymbol{\mu}} + \widehat{\mathbf{C}}^T \widehat{\mathbf{V}} \boldsymbol{\Pi} \widehat{\mathbf{V}}^T \widehat{\mathbf{C}}^{-T} (\mathbf{Y} - \widehat{\boldsymbol{\mu}}). \tag{13}$$

Under the identity permutation, meaning that $\boldsymbol{\Pi} = \mathbf{I}$, we would recover the original ₁₂₇

response vector. We note that because $\widehat{\boldsymbol{\Gamma}}$ is just a diagonal matrix, $\widehat{\mathbf{C}}$ can be obtained ₁₂₈

from a factorization (e.g., cholesky or eigen decomposition) of $\widehat{\boldsymbol{\Sigma}} = \hat{\xi} \, \boldsymbol{\Phi} + (1 - \hat{\xi}) \, \mathbf{I}$, ₁₂₉

which in turn could be obtained from a factorization of $\boldsymbol{\Phi}$ if desired, as was done in [9]. ₁₃₀

Examining the form of (13), there are three main steps used to obtain a ₁₃₁

transformation of the residuals with second-order exchangeable entries. The first is to ₁₃₂

center the response by using the estimated phenotypic mean. The second is to remove ₁₃₃

the correlation present due to polygenic effects, which is done by the pre-multiplication ₁₃₄

by $\widehat{\mathbf{C}}^{-T}$. The last step is to remove the correlation that is generated from using ₁₃₅

parameter estimates instead of the true values when centering, and is represented by the ₁₃₆

pre-multiplication by $\widehat{\mathbf{V}}^T$. It is possible that the last step might have only a minor ₁₃₇

impact on the method for large sample sizes and low number of parameters, though we ₁₃₈

did not investigate this. The replicates generated from this approach are quantitative, ₁₃₉

i.e., the binary nature of the response is not preserved during resampling. In subsection ₁₄₀

Additional resampling methods considered, we describe how the replicates can be ₁₄₁

transformed to be binary, and we consider both approaches in simulations. The fixed ₁₄₂

effects and covariance structure estimated under the null are preserved in the trait ₁₄₃

replicates, making our method applicable to general scenarios where there are important ₁₄₄

covariates and/or population structure. ₁₄₅

LogMM-PQL ₁₄₆

A logistic mixed model (LogMM) is a natural modeling choice for correlated binary ₁₄₇

data, and an alternative approach to BRASS for generating replicates of correlated ₁₄₈

binary data could be to fit a LogMM under the null hypothesis and then generate ₁₄₉

binary replicates from the fitted model. We have developed such an approach, which we ₁₅₀

call LogMM-PQL. The model underlying LogMM-PQL is that, conditional on $\mathbf{X}$, $\mathbf{G}$ ₁₅₁

and $\mathbf{u}$, Y_i is Bernoulli(m_i), independently across $i = 1, \ldots, n$, where m_i is the ith   152

component of the vector   153

$$\mathbf{m} := E(\mathbf{Y}|\mathbf{X}, \mathbf{G}) = \text{logit}^{-1}(\mathbf{X}\boldsymbol{\beta} + \mathbf{G}\gamma + \mathbf{u}), \tag{14}$$

where $\boldsymbol{\Phi}$, $\mathbf{Y}$, $\mathbf{X}$, $\mathbf{G}$, $\boldsymbol{\beta}$ and γ are as before; $\mathbf{u}$ is a multivariate normal random effects   154

vector of length n with $\mathbf{u} \sim \text{MVN}(0, \sigma^2\boldsymbol{\Phi})$ where σ^2 is an unknown scalar, and where γ 155

will be set to zero because the model will be fitted under the null hypothesis of no effect.   156

The major difficulty with use of LogMMs in the GWAS context is that they are   157

computationally challenging to fit as they involve high-dimensional integrals. Therefore,   158

their use can involve a trade-off between computational feasibility and accuracy of the   159

results. For the parameter estimation part of LogMM-PQL, we use a fast penalized   160

quasi-likelihood (PQL) approach, as implemented in GMMAT [19]. With γ set to zero,   161

we use GMMAT to obtain estimates $\widehat{\boldsymbol{\beta}}$ and $\hat{\sigma}^2$, and then generate trait replicates by   162

drawing random vectors $\mathbf{Y}$ from the LogMM specified by Eq (14) with $(\boldsymbol{\beta}, \gamma, \sigma^2)$ set   163

equal to $(\widehat{\boldsymbol{\beta}}, 0, \hat{\sigma}^2)$. LogMM-PQL renders binary replicates that incorporate the mean   164

and variance structure specified by the fitted logistic model.   165

Additional resampling methods considered   166

In addition to BRASS and LogMM-PQL, the other resampling methods we compare in   167

simulations include the previously proposed MVNpermute [9,11], as well as a method   168

we call Naive, which involves fitting a LMM whose form under the null is $\mathbf{Y} = \mathbf{X}\boldsymbol{\beta} + \mathbf{e}$,   169

with $\mathbf{e} \sim \text{MVN}(0, \sigma_1^2\boldsymbol{\Phi} + \sigma_2^2\mathbf{I})$. After obtaining the estimated coefficients $\widehat{\boldsymbol{\beta}}$, the Naive   170

method replicates are obtained by permuting the residuals $(\mathbf{Y} - \mathbf{X}\widehat{\boldsymbol{\beta}})$, and adding back   171

$\mathbf{X}\widehat{\boldsymbol{\beta}}$.   172

While the LogMM-PQL replicates are binary, those from BRASS, MVNpermute and   173

Naive are not. However, we consider alternative approaches based on converting these   174

replicates to binary, referred to as BRASS$_{mod}$, MVNpermute$_{mod}$ and Naive$_{mod}$,   175

respectively, where the subscript "mod" indicates that the trait replicate has been   176

modified to convert it to binary. To convert a continuous trait replicate $\mathbf{Y}_\pi$ to binary, a   177

threshold is set and the values of $\mathbf{Y}_\pi$ above and below that threshold are converted to 1   178

and 0, respectively, with the threshold chosen so that the original trait $\mathbf{Y}$ and the   179

binary replicate $\mathbf{Y}_\pi$ have the same case/control proportions.

The features of the various methods are summarized and compared in S1 Table.

Modeling sample structure in the replicates

All 7 resampling strategies described above involve fitting a trait model that
incorporates the structure present in the sample. A common approach to modeling trait
correlation due to additive polygenic effects is to include a GRM in a LMM for the trait
so as to model the structure using random effects [20]. Alternatively, the top principal
components of a given GRM can be used as covariates to model the structure present
using fixed effects [21]. Here, we allow either of the two approaches or they can be
combined (i.e., both fixed and random effects can be used) to capture the sample
structure in the null model. To combine the two approaches, we use the top PCs from a
GRM as fixed effects in the null model and then build a new GRM that is adjusted to
exclude the effects of these top PCs in order to capture the leftover structure as random
effects. In the simulations and data analysis, we form the new GRM using equation 4 of
a previous work [22].

When generating replicates, it is of course important to accurately model the
structure present in the sample, but in addition, it is also important to avoid creating
new structure (not present in the data) as an artifact of the replication procedure. To
do this, we recommend the use of leave-one-chromosome-out (LOCO) GRMs [23, 24] in
the replication procedure. In that case, replicates would be generated separately for
each chromosome, with the SNPs for the given chromosome left out of the GRM. This
avoids introducing new trait-SNP association in the replicates.

Methods for simulation studies

We perform simulation studies to compare the effectiveness of the 7 proposed
permutation-based methods for generating binary trait replicates in the context of
correcting for multiple testing, with both population structure and pedigree structure
present in the sample. We consider a setting in which a binary trait is tested for
association with each of m markers and the aim is to assess the significance of the
smallest p-value out of the m association tests. This context would arise in testing

significance of a genomic region or in assessing genome-wide significance. We simulate 209
non-causal markers that are tested for association with a binary trait and estimate the 210
significance threshold for the top signal amongst them using the empirical distribution 211
from trait replicates generated under the null hypothesis of no association. Genotype, 212
trait and covariates are generated under multiple simulation settings. In real data 213
applications, the true trait model is usually not known a priori, and it can be important 214
to assess the impact of model misspecification on type 1 error [25]. We consider the 215
effects of model misspecification due to (1) assuming a logistic link function when the 216
true model is a liability threshold model and (2) exclusion of an important covariate 217
from the model. In addition, we investigate the impact of ascertainment in the sample 218
as case-control studies often involve phenotype-based ascertainment which can introduce 219
misspecification in the model as well. 220

For each setting, we perform a simulation study consisting of 4 stages: (1) data set 221
creation, i.e., simulation of genotype and trait data under the null hypothesis of no 222
association, with 20,000 data sets per setting; (2) generation of permutation replicates, 223
in which we consider every possible pair (i, j) of simulated data set i and permutation 224
method j, for $1 \leq i \leq 20,000$ and $1 \leq j \leq 7$, and for each of these, 10,000 permutation 225
replicates are drawn for data set i using permutation method j; (3) analysis of 226
permutation replicates, where for each trio (i, j, k) of simulated data set i, permutation 227
method j, and permutation replicate k, $1 \leq k \leq 10,000$, the trait in permutation 228
replicate k is tested for association with each of the m non-causal markers in data set i, 229
the smallest p-value is recorded in each case, and then the resulting 10,000 p-values for 230
each pair (i, j) are used to estimate a genomewide significance threshold (at level 231
$\alpha = 10^{-3}$) for data set i based on permutation method j; and (4) evaluation of the type 232
1 error rate of the genomewide threshold for each permutation method j by comparing 233
the smallest p-value for each data set i to the threshold determined in (3) for pair (i, j) 234
and assessing whether the overall proportion of rejections among the 20,000 data sets is 235
significantly different from α for method j. We now describe each of these stages. 236

For stage (1) data set creation, we simulate genotypes using a 2 sub-population 237
Balding-Nichols model [26] with $F = .01$ and ancestral allele frequencies for SNPs 238
drawn independently and uniformly on $(.2, .8)$. We simulate multiple pedigrees of the 239
same configuration (S1 Fig), with equal numbers of pedigrees assigned to each of the 240

two sub-populations. Founder alleles for a pedigree are assumed to be drawn from the sub-population to which the pedigree is assigned, and gene dropping is used to determine genotypes for other pedigree members. We simulate phenotypes according to two different generative models for the trait. In each case, $M_c = 10^4$ causal SNPs are simulated as well as 3 covariates: age, sex and an i.i.d. standard normal covariate. The first generative model we use is logistic:

$$Y_i | \mathbf{X}_i, \mathbf{W}_i, \boldsymbol{\alpha} \sim \text{Bernoulli}(p_i), \text{ independently, with logit}(p_i) = \mathbf{X}_i \boldsymbol{\beta} + \mathbf{W}_i \boldsymbol{\alpha}, \qquad (15)$$

where $\mathbf{X}_i$ is the covariate row vector for the ith individual; $\boldsymbol{\beta}$ are the fixed effects for the covariates; $\mathbf{W}_i$ is a row vector representing the SNP information for the ith individual corresponding to the M_c causal markers standardized to have mean 0 and variance 1; $\boldsymbol{\alpha} = (\alpha_1, \cdots, \alpha_{M_c})^T \sim MVN(0, \sigma^2 \mathbf{I})$ represent the random effects of the M_c causal markers with resulting total additive polygenic variance $\sigma_a^2 = \sigma^2 M_c$. The values for $\boldsymbol{\beta}$ and σ_a^2 are chosen so that, considering the variance of $\text{logit}(p_i)$ due to covariate effects and additive polygenic random effects, the fraction of this variance due to covariates is fixed at 20%, 40%, 60% or 80% (while the remaining fraction is due to additive polygenic random effects), and so that Bernoulli error explains on average either (a) about 20% of the total phenotypic variability, resulting in a prevalence of 30% or (b) about 55% of the total phenotypic variability, resulting in a prevalence of 5%. In addition, the effect size of the standard normal covariate is set so that the p-value for its significance in a LMM Wald test is on average 0.05. (This is the covariate that is set to be missing in the settings in which an important covariate is missing from the observed data set.) The second generative model we use is a liability threshold model:

$$Y_i = \mathbb{1}_{\{L_i > 0\}}, \text{ with } L_i = \mathbf{X}_i \boldsymbol{\beta} + \mathbf{W}_i \boldsymbol{\alpha} + \epsilon_i, \qquad (16)$$

where L_i is the latent liability for individual i, and $\boldsymbol{\epsilon} = (\epsilon_1, \ldots, \epsilon_n)^T \sim MVN(0, \sigma_e^2 \mathbf{I})$. The values of $\boldsymbol{\beta}$, σ_a^2, and σ_e^2 are chosen using analogous constraints as for the logistic model, where the phenotypic variability due to $\boldsymbol{\epsilon}$ is constrained to be about 20% of the total phenotypic variability, and the prevalence is set to either 30% or 5%. In both 15 and 16, larger values of σ_a^2 correspond to more severe confounding effects of

population/family structure on phenotype-genotype association. For the settings in ²⁶⁷
which ascertainment is used, we simulate data under either the model in (15) or that in ²⁶⁸
(16) and select 1,000 individuals at random to be retained in the sample using either (a) ²⁶⁹
a balanced case-control ratio or (b) a 3:7 case-control ratio. In all settings, in addition ²⁷⁰
to the M_c causal SNPs, $m = 100$ non-causal SNPs are generated as above and used for ²⁷¹
testing under the null hypothesis of no association. (Further details on the stage (1) ²⁷²
simulations can be found in S1 Text.) ²⁷³

We now describe stage (2) generation of permutation replicates for each data set and ²⁷⁴
for each of the 7 permutation methods (BRASS, LogMM-PQL, MVNpermute, Naive, ²⁷⁵
BRASS$_{mod}$, MVNpermute$_{mod}$, and Naive$_{mod}$). For a given data set, the same covariate ²⁷⁶
and GRM information is input to each of the 7 permutation methods. The GRM $\boldsymbol{\Phi}$ is ²⁷⁷
calculated using the M_c causal markers, and then $D = 1$ top PC is removed from $\boldsymbol{\Phi}$ ²⁷⁸
using equation 4 of a previous work [22], resulting in adjusted GRM $\tilde{\boldsymbol{\Phi}}$, which is the ²⁷⁹
GRM input to each permutation method. In most settings, the 4 covariates input to ²⁸⁰
each permutation method are the top PC plus the 3 covariates used to generate the ²⁸¹
trait. In the settings in which an important covariate is assumed to be missing from the ²⁸²
data set, the 3 covariates input to each permutation method are the top PC, age and ²⁸³
sex, with the normal covariate left out. ²⁸⁴

In stage (3), for each pair (i, j) of data set and permutation method, the 10^4 ²⁸⁵
permutation replicates from method j are each tested for association against the panel ²⁸⁶
of m SNPs in data set i, and the smallest p-value among the m SNPs is recorded for ²⁸⁷
each permutation replicate, resulting in 10^4 p-values for pair (i, j). The genome-wide ²⁸⁸
threshold value, corresponding to significance level α, for pair (i, j) is estimated by the ²⁸⁹
100αth percentile of the p-values observed for that pair. In principle, any association ²⁹⁰
testing method could be used for the stage (3) association tests, provided that the same ²⁹¹
method is also used in stage (4). One consideration is that only four of the permutation ²⁹²
methods we examine create binary replicates, while the other three create non-binary, ²⁹³
real-valued replicates. Whether having non-binary replicates is problematic or not ²⁹⁴
depends on the method of analysis. For example, it is not a problem if the data are ²⁹⁵
analyzed using a linear mixed model or using the CARAT [18] or CERAMIC [17] binary ²⁹⁶
trait methods, because these methods are based on solving certain estimating equations ²⁹⁷
and work well provided that the first and second moments of the data are appropriately ²⁹⁸

modeled. However, analysis by ordinary logistic regression would require binary data. 299
To be able to test all 7 methods, we therefore use a (slightly modified) version of 300
CARAT for the stage (3) and (4) association tests, where the modification is in the 301
estimation of σ_g^2, the genotypic variance parameter for CARAT. In the original CARAT 302
paper [18] σ_g^2 is estimated by $\hat{\sigma}_g^2 = 2\hat{f}(1 - \hat{f})$ with $\hat{f} = .5\bar{G}$, where $\bar{G}$ is the sample 303
average of $\boldsymbol{G}$ in the data. Our modification is to replace this with 304
$\widetilde{\sigma}_g^2 = \mathbf{G}^T \mathbf{P} \mathbf{G}/(n - D - 1)$ where D is the number of PCs removed from the GRM, and 305
$\mathbf{P} = \tilde{\boldsymbol{\Phi}}^{-1} - \tilde{\boldsymbol{\Phi}}^{-1}\mathbf{Z}(\mathbf{Z}^T\tilde{\boldsymbol{\Phi}}^{-1}\mathbf{Z})^{-1}\mathbf{Z}^T\tilde{\boldsymbol{\Phi}}^{-1}$, where $\mathbf{Z}$ is the $n \times (D + 1)$ matrix whose 306
columns are the D eigenvectors that are removed as well as a columns of ones. (The 307
rationale for this modification is detailed in S1 Text.) When we apply CARAT, we use 308
$\tilde{\boldsymbol{\Phi}}$ as the GRM, and we include as fixed effects the top PC plus the 3 covariates that 309
were used to generate the trait. For the settings when an important covariate is 310
assumed to be missing from the data set, the 3 covariates used in the analysis of 311
replicates are the top PC, age and sex, with the normal covariate left out. 312

Finally, in stage (4), we evaluate the type 1 error of the genomewide threshold for 313
each permutation method j by by comparing the p-value for each data set i to the 314
threshold determined in stage (3) for the pair (i, j) and assessing whether the overall 315
proportion of rejections out of 20,000 data sets is significantly different from the 316
nominal level α for method j. To make our massive simulation project more 317
computationally efficient, we actually used an adaptive procedure so that, e.g., if a 318
dataset is sufficiently far from the significance threshold based on analysis of the first 319
10^3 permutation replicates, it would not be necessary to analyze the remainder of the 320
10^4 permutation replicates (see S1 Text for details). The amount of downward bias 321
introduced in the empirical type 1 error estimate by this procedure is very small 322
compared to the Monte Carlo sampling variability. For example, when level .01 is tested 323
under the null hypothesis with 20,000 Monte Carlo replicates using the adaptive 324
procedure, the squared bias in the type 1 error estimate due to the adaptive procedure 325
is less than 5e-10, which is several orders of magnitude smaller than the Monte Carlo 326
error variance of 5e-07. 327

Verification and Comparison

Assessment of type 1 error

We assess and compare the performance of the 7 proposed resampling methods in the

context of correcting for multiple testing, with both population and pedigree structure

present in the sample in addition to important covariates. Fig 1 shows the empirical

Fig 1. Empirical Type 1 Error Rates with Logistic Model Including all Covariates at Nominal Level 0.01.
The error rate is based on 20,000 simulated replicates. The solid horizontal line represents the nominal level and the dashed lines represent rejection bounds outside of which the z-test comparing the estimated type 1 error to the nominal level is rejected at level .05. Estimates inside the rejections bounds are represented by circles and those outside the bounds are represented by triangles. Red, black and blue symbols represent inflated, well-controlled and conservative type 1 error rates, respectively. The proportion of variability on the logit scale attributable to polygenic effects vs. covariates is varied from 20 to 80% in increments of 20%. Prevalence is 30%.

type 1 error rates of all methods, compared to the nominal, when the true trait model is

logistic. The type 1 error of BRASS is well-controlled in all settings. For LogMM-PQL,

we observe significant inflation in the type 1 error rate, with the amount of inflation

increasing as the confounding due to structure present in the sample increases. This

likely occurs because the PQL model fitting approach tends to underestimate the

variance component and hence, results in less correlation present in the simulated data

compared to that in the original data. We observe that permuting the residuals from a

linear mixed model (Naive) leads in all settings to lack of control of the type 1 error.

More precisely, when covariates explain most of the variation on the logit scale, the

method is overly conservative and when instead polygenic effects explain most of the

variability, the type 1 error rate is significantly inflated. Moreover, we also find that

adjusting for sample correlation prior to permutation based on LMM residuals, as done

in MVNpermute, still leads to a very conservative test when covariates highly influence

the variability on the logit scale. This is most likely because a linear mixed model does

not allow any dependence between the trait variance and the mean, which is influenced

by the covariates. As covariates become less important, we do see somewhat better

control of the type 1 error. For Naive$_{mod}$ and MVNpermute$_{mod}$, control of the type 1

error is improved compared to the corresponding original methods when covariates

strongly influence the phenotypic mean, and for BRASS$_{mod}$, type 1 error control is

worse compared to BRASS. All of the methods except BRASS show inflated type 1 ⟨352⟩
error in at least one setting, whereas BRASS maintains the correct type 1 error rate in ⟨353⟩
all settings. ⟨354⟩

Robustness to model misspecification: liability threshold vs. ⟨355⟩ logistic model ⟨356⟩

As the true model for the trait is generally unknown in an association study, we are ⟨357⟩
interested in assessing robustness of the type 1 error results to model misspecification. ⟨358⟩
Thus, we also perform simulations in which the phenotype is simulated from a liability ⟨359⟩
threshold model, which is more dissimilar to the models fit by BRASS and ⟨360⟩
LogMM-PQL than logistic is. The type 1 error results for the liability threshold model ⟨361⟩
are shown in Fig 2. The primary effect of the model misspecification in this case seems

Fig 2. Empirical Type 1 Error Rates with Liability Threshold Model at Nominal Level 0.01.
The model is misspecified with the true model being a liability threshold model. The error rate is based on 20,000 simulated replicates. The solid horizontal line represents the nominal level and the dashed lines represent rejection bounds outside of which the z-test comparing the estimated type 1 error to the nominal level is rejected at level .05. Estimates inside the rejections bounds are represented by circles and those outside the bounds are represented by triangles. Red, black and blue symbols represent inflated, well-controlled and conservative type 1 error rates, respectively. The proportion of variability on the liability scale attributable to polygenic effects vs. covariates is varied from 20 to 80% in increments of 20%. Prevalence is 30%.

⟨362⟩

to be to make all the methods except Naive slightly more conservative. ⟨363⟩

Robustness to missing an important covariate ⟨364⟩

We assess the robustness of the type 1 error results when an important covariate is ⟨365⟩
omitted from both the fitted model and the model used for generating replicates. As it ⟨366⟩
is usually not known a priori which variables should be kept in the analysis, one would ⟨367⟩
commonly try including different combinations of covariates to finally determine the ⟨368⟩
ones to include in the final model. It may occur that one of the covariates has a ⟨369⟩
moderate effect on the trait and leads to a p-value close to the significance threshold ⟨370⟩
(e.g. 0.05). Hence, a judgment call would be required for whether to keep the covariate ⟨371⟩
in the model and one may decide to exclude it. It is thus of interest to see how the ⟨372⟩

proposed methods would fare in such a scenario as the replicates generated would come   ₃₇₃
from a more misspecified model.   ₃₇₄

The simulation results are displayed in Fig S2. Similarly to the previous model   ₃₇₅
misspecification results, the general effect is to make the results more conservative for   ₃₇₆
all permutation methods except Naive.   ₃₇₇

Robustness to ascertainment   ₃₇₈

We determine the robustness of the proposed methods when trait-based ascertainment   ₃₇₉
has been applied to the sample. This is commonly used in case-control studies where   ₃₈₀
individuals are included in the sample based on their disease affection status, such as if   ₃₈₁
the prevalence of the trait in the population is too low to obtain sufficient power.   ₃₈₂

The results are shown in Figs 3 and 4, where it can be seen that BRASS is the only

**Fig 3. Empirical Type 1 Error Rates with Ascertainment Scenario 1 at
Nominal Level 0.01.**
The error rate is based on 20,000 simulated replicates. The solid horizontal line
represents the nominal level and the dashed lines represent rejection bounds outside of
which the z-test comparing the estimated type 1 error to the nominal level is rejected at
level .05. Estimates inside the rejections bounds are represented by circles and those
outside the bounds are represented by triangles. Red, black and blue symbols represent
inflated, well-controlled and conservative type 1 error rates, respectively. The
proportion of variability on the logit scale attributable to polygenic effects vs. covariates
is varied from 20 to 80% in increments of 20%. Prevalence is 30%, and ascertainment
results in a case-control ratio of 1:1.

**Fig 4. Empirical Type 1 Error Rates with Ascertainment Scenario 2 at
Nominal Level 0.01.**
The error rate is based on 20,000 simulated replicates. The solid horizontal line
represents the nominal level and the dashed lines represent rejection bounds outside of
which the z-test comparing the estimated type 1 error to the nominal level is rejected at
level .05. Estimates inside the rejections bounds are represented by circles and those
outside the bounds are represented by triangles. Red, black and blue symbols represent
inflated, well-controlled and conservative type 1 error rates, respectively. The
proportion of variability on the logit scale attributable to polygenic effects vs. covariates
is varied from 20 to 80% in increments of 20%. Prevalence is 5%, and ascertainment
results in a case-control ratio of 3:7.

₃₈₃

method that retains good control of the type 1 error rate in all of the ascertainment   ₃₈₄
settings considered. In contrast, both logMM-PQL and Naive$_{mod}$ show significantly   ₃₈₅
inflated type 1 error in every setting, while both MVNpermute and Naive give   ₃₈₆
excessively conservative results when covariates have a major impact on the trait and   ₃₈₇

tend to give significantly inflated results when the effect of covariates is low. ₃₈₈

Robustness to ascertainment, model misspecification and missing covariate, combined

Finally, we consider a scenario that simultaneously combines multiple features of the previous settings: (1) ascertainment from a prevalence of 5% to a case-control ratio of 3:7; (2) model misspecification in which the true model is a liability-threshold model, which is more dissimilar to the models fit by BRASS and LogMM-PQL than logistic is; and (3) model misspecification due to an important covariate being missing. The results are shown in Fig 5. As before, the model misspecification due to the liability threshold

Fig 5. Empirical Type 1 Error Rates with Ascertainment, Model Misspecification and Missing Covariate at Nominal Level 0.01.
The model is misspecified with the true model being a liability threshold model. The error rate is based on 20,000 simulated replicates. The solid horizontal line represents the nominal level and the dashed lines represent rejection bounds outside of which the z-test comparing the estimated type 1 error to the nominal level is rejected at level .05. Estimates inside the rejections bounds are represented by circles and those outside the bounds are represented by triangles. Red, black and blue symbols represent inflated, well-controlled and conservative type 1 error rates, respectively. The proportion of variability on the liability scale attributable to polygenic effects vs. covariates is varied from 20 to 80% in increments of 20%. Prevalence is 5%, and ascertainment results in a case-control ratio of 3:7. The effect of the omitted covariate on the trait correspond to a Wald test p-value of .05 using a linear mixed model.

model and the omission of an important covariate have the effect of making most of the methods more conservative, where this effect seems most pronounced when covariate effects are much more important than polygenic effects (left-most case corresponding to "20/80" in each plot).

From the results across all the simulation settings, with and without ascertainment or model misspecification, it can be seen that BRASS provides the best type 1 error control and is the only method that does not have significantly inflated type 1 error in any setting.

Applications

We apply BRASS to the problem of determining genome-wide significance for GWAS of two different traits in domestic dog. Unlike with humans, a genome-wide threshold for

significance in domestic dog GWAS has not been well established. We analyze data ₄₀₈
from a large domestic dog study [27], with 4,224 dogs representing over 150 breeds ₄₀₉
genotyped at 185,805 SNPs with multiple phenotypes recorded. We analyze two binary ₄₁₀
traits from this study, ED, for which the data contain 113 cases and 633 controls among ₄₁₁
82 breeds, and IE, for which there are 34 cases and 168 controls from the Irish ₄₁₂
Wolfhound breed. For each trait, we compute a GRM after LD pruning and filtering for ₄₁₃
MAF $< .05$. For ED, we take the top 10 PCs out of the GRM and include them as ₄₁₄
covariates, and sex is included as a covariate for both traits. The data are then analyzed ₄₁₅
with CARAT. To generate trait replicates using BRASS, we create LOCO GRMs for ₄₁₆
each chromosome, where for ED we also take out the top 10 PCs in each case and ₄₁₇
include them as covariates, and sex is included as a covariate for both traits. For ₄₁₈
genome-wide significance assessment for each trait, we used BRASS to generate 10,000 ₄₁₉
replicates under the null hypothesis of no association. We analyzed each replicate in the ₄₂₀
same way that we analyzed the data and used the results of these analyses to form an ₄₂₁
empirical distribution of the test statistic under the null hypothesis. Genome-wide ₄₂₂
p-values were then estimated using the empirical distribution for each trait. ₄₂₃

Manhattan plots of the p-values of the single-SNP tests for the observed data are ₄₂₄
presented in Fig 6 for both ED and IE phenotypes, along with the estimated

Fig 6. Genome screen results in the domestic dog data.
Manhattan plots of the single-SNP association p-values using CARAT (on $-\log_{10}$ scale)
for (A) ED and (B) IE, with genomic position on the x-axis. The horizontal lines
represent estimated genome-wide thresholds at nominal level 0.05 using BRASS. SNPS
from the 38 autosomes of the domestic dog genome are depicted.

₄₂₅

genome-wide significance thresholds, which are estimated at nominal level 0.05 from the ₄₂₆
empirical distribution of the top association signal based on the trait replicates. The ₄₂₇
genomic control inflation factors for ED and IE are 1.02 and 0.98, respectively. For IE, ₄₂₈
we detect a previously-identified [27] 12Mb region on chromosome 4 (position 7.5-19.3 ₄₂₉
Mb) that reaches the BRASS genome-wide significance threshold, with nominal p-value ₄₃₀
2.1×10^{-8} corresponding to a genomewide p-value of .0008 obtained from BRASS. For ₄₃₁
ED, we detect 3 loci, summarized in Table 1. Compared to a previous reference [27], we ₄₃₂
detected one additional significant locus (rs23910667 on chromosome 35) that is not ₄₃₃
detected there. The genomewide p-values by the other 6 methods for these same SNPs ₄₃₄
(detailed in S1 text) are somewhat smaller than those from BRASS, which is in line ₄₃₅

with the simulation studies showing that the other methods tend to be anti-conservative overall and in particular for models in which polygenic effects are more important than covariates.

Table 1. Top association signals for ED

SNP	Chr	Position	MAF	Single SNP p-value	Genomewide p-value
rs9000666	26	16554631	.18	1.0×10^{-7}	.010
rs21895578	1	77938330	.17	1.5×10^{-7}	.014
rs23910667	35	7272763	.08	3.6×10^{-7}	.033

Computation time

In order to generate replicates from BRASS, the null model given by Eq 1 and 2 with $\gamma = 0$ needs to be fitted, which includes specifying covariates and a GRM $\boldsymbol{\Phi}$. If not provided, obtaining the eigen decomposition of $\boldsymbol{\Phi}$ is the main computationally intensive step involved in fitting the null model [18]. However, this step (or one of equivalent computational complexity) is also typically performed when analyzing the data or when using permutation methods based on a linear or logistic mixed model. Once parameter null estimates are obtained, the transformed residuals $\hat{\boldsymbol{\zeta}}$ as well as the linear map $\widehat{\mathbf{C}}^T \widehat{\mathbf{V}}$ in (13) need to be computed; however, once obtained, they can be re-used when generating trait replicates. As the eigen decomposition of $\boldsymbol{\Phi}$, and hence $\widehat{\boldsymbol{\Sigma}}$, is obtained prior to fitting the null model, we use it to compute $\widehat{\mathbf{C}}$. Therefore, the most computationally intensive step is to obtain the eigenvectors of $\widehat{\boldsymbol{\Psi}}$ (the estimate for $\boldsymbol{\Psi}_0$ in (11)), which involves a time complexity of $O(n^3)$. By using the singular value decomposition of the matrix $\mathbf{W}$ in (11), which involves a time complexity of $O(nk^2)$ (assuming the number of covariates $k < n$), we are able to obtain the eigenvectors of $\widehat{\boldsymbol{\Psi}}$ without having to construct it and compute its eigen decomposition. Past this step, the additional cost for generating L replicates is $O(n^2 L)$; this computation can easily be parallelized to generate sets of traits replicates independently.

BRASS is implemented in a freely downloadable software package at `https://github.com/joellesophya/BRASS`. We report run times for BRASS in simulated data. Using a single processor on a machine with 6 core Intel Xeon 3.50 GHz CPUs and 32 GB RAM, on simulated data with 1,000, 5,000 and 10,000 individuals, it

takes 1.6 s , 1 min and 4.6 min (275 s), respectively, to generate 1,000 trait replicates. 461
As the use of BRASS will vary based on the context (e.g. multiple testing correction, 462
region-based association testing), the computational burden involved in comparing the 463
statistic between the observed trait and its replicates will vary based on what statistic is 464
being considered. 465

Discussion 466

In genetic studies of binary traits, it may often be of interest to assess significance of a 467
test statistic whose distribution is unknown or not well-approximated, or to assess 468
significance of the maximum of many correlated tests. Examples could include test 469
statistics based on machine learning methods [1], rare variant tests [3], test statistics 470
that involve data-adaptive weights [4], and assessment of genomewide significance [5] in 471
a variety of situations such as inclusion of admixed individuals in GWAS [6]. 472
Permutation tests are an attractive and popular approach, but in the presence of 473
population structure, cryptic relatedness and/or family structure, which are all common 474
sources of confounding in genetic association studies, ordinary permutation tests are 475
usually not appropriate as they fail to retain the structure present in the data. 476

We propose BRASS, a novel permutation-based resampling procedure for generating 477
binary trait replicates in samples with population structure, cryptic relatedness and/or 478
family structure. BRASS allows for covariates and ascertainment, and it accommodates 479
a wide range of test statistics. It uses an estimating equation approach that can be 480
viewed as a hybrid of logistic regression and linear mixed-effects model methods, which 481
allows for use of a GRM and/or principal components or other ancestry-informative 482
covariates to account for sample structure. BRASS relies on obtaining an invertible 483
transformation of phenotypic residuals to achieve approximate second-order 484
exchangeability. After permutation, new trait replicates are obtained by inverting the 485
transformation, in order to preserve structure present in the original sample. BRASS 486
differs from existing methods such as Naive and MVNpermute [11] in that it 487
incorporates key features of a binary trait such as nonlinear effects of covariates and 488
dependence of the variance on the mean, which we show improve the performance of 489
BRASS on binary traits. 490

In simulations, we demonstrate the superior type 1 error control of BRASS   491
compared to other methods. Across simulation settings, we find that BRASS maintains   492
control of the type 1 error under varying amounts of population structure, familial   493
relatedness, ascertainment and sources of trait model misspecification. In contrast, all 6   494
of the other methods we consider show significant inflation of type 1 error in multiple   495
settings. In addition, the Naive and MVNpermute methods, which are two of the 4   496
methods that are based on an LMM, are excessively conservative when covariate effects   497
are substantial. Taken together, the results show that simply accounting for the sample   498
structure with an LMM, without incorporating key features associated with the binary   499
nature of the trait such as nonlinear scale for covariate effects and the dependence of   500
the variance on the mean, is not sufficient to obtain replicates that correctly estimate   501
the null distribution of the statistic. Secondarily, we find that when polygenic effects   502
are important, adjusting for the correlation in the residuals prior to permutation, as is   503
done in BRASS and MVNpermute leads to better control of the type 1 error compared   504
to ignoring it, as is done when permuting the raw residuals from a LMM (Naive   505
method). Furthermore, we find that the additional step of binarizing the quantitative   506
replicates results in improvement in the accuracy of the type 1 error rate for the   507
LMM-based methods when the polygenic component has a low impact on the trait   508
distribution relative to covariates, but does not control the type 1 error when more   509
structure is present in the sample.   510

In principle, use of a logistic mixed model would seem to be a natural and promising   511
alternative to BRASS for fitting the binary trait data and generating appropriate   512
replicates that preserve the correlation structure. However, for LogMM-PQL, we   513
observe significant inflation in the type 1 error rate, with the amount of inflation   514
increasing as the confounding due to structure present in the sample increases. This is   515
likely explained by the use of PQL to fit the LogMM, as PQL is known to be fast but to   516
give biased estimates in binary data [28]. However, fitting a LogMM is computationally   517
intensive and requires a trade-off between computational time and accuracy of the   518
approximation of the high-dimensional integral involved. Algorithms with higher   519
accuracy than PQL (e.g. the Laplace approximation [29] or Gauss-Hermite   520
quadrature [30]) lead to increased computational burden and do not scale well for even   521
moderately large GWAS. In contrast, BRASS is seen to be extremely fast   522

computationally.   523

We applied BRASS in the context of multiple testing correction in association   524
mapping studies of ED and IE in domestic dogs. In the analysis of ED, we detected   525
genomewide significant association with 3 loci, 2 of which were previously reported [27],   526
with the 3rd locus also reaching significance based on the BRASS threshold. In the   527
analysis of IE, we detected a 12 Mb region on chromosome 4 that reached the BRASS   528
genome-wide significance threshold and has been previously associated with IE [27]. For   529
the ED data analysis, breed information was available, but we chose to instead use PCs   530
as ancestry informative covariates, due to apparent inaccuracies in the breed labeling   531
information as well as the presence of dogs of mixed or unknown breed in the data set.   532
Interestingly, the genome-wide p-value threshold estimated using replicates from   533
BRASS for the ED trait in a sample of 82 breeds was about half that for the IE trait in   534
a single breed. This is expected given the extremely different population structure   535
between the two samples.   536

BRASS is designed as a very flexible tool that can be used in a range of situations in   537
which replicates of correlated binary data are needed. Because BRASS generates   538
replicates of the phenotype, it can be used with a variety of genetic predictors. For   539
example, it could be directly applied to association testing of haplotypes, to association   540
testing of variants on sex chromosomes, or to tests of GxE or GxG interaction.   541
Although imbalanced case-control ratio can have an impact on some association testing   542
methods, it would not be expected to have any particular impact on generation of   543
BRASS replicates, because (1) the appropriate modeling of the connection between the   544
mean and the variance for binary data means the method does not break down for small   545
or large probabilities, in contrast to an LMM, and (2) the asymptotic approximations   546
used for inference in logistic regression are not needed in BRASS. In our simulation   547
results, for the settings with imbalanced case-control ratio (Figs 1, 2, 4, 5), there was no   548
noticeable impact. In our simulations, we have considered various types of model   549
misspecification and demonstrated robustness of BRASS for association testing. We   550
note that certain applications such as testing of interaction have been shown to be more   551
sensitive to model misspecification than association testing is [25], so additional   552
sensitivity analyses may be needed when performing interaction analyses, regardless of   553
the method used to assess significance.   554

Depending on the application, one can consider a different approach to capture the structure present in the sample, as the model used by BRASS can accommodate both covariates and random effects. In the simulation studies and data analysis, we chose to partition the relatedness into random and fixed effects and include both in our model. A previously proposed method, PC-Air [31], obtains PCs from a derived subset of mutually unrelated subjects that are representative of the ancestral diversity present in the sample, so as to ensure that the top PCs will only capture distant genetic relatedness (i.e. population structure). Such methods could easily be used to obtain the population structure information input to BRASS, if desired.

BRASS generates non-binary replicates, which can be used directly in some types of analyses, e.g., when the data are analyzed using a linear mixed model or using the CARAT [18] or CERAMIC [17] binary trait methods, because these methods are based on solving certain estimating equations and work well provided that the first and second moments of the data are appropriately modeled. However, analysis by ordinary logistic regression would require binary data. If binary replicates are needed and if the additive polygenic component is estimated to have a low impact on the trait distribution relative to the covariates, then we would recommend either $BRASS_{mod}$ or $MVNpermute_{mod}$ which both provided good type 1 error control in our simulations in those cases.

Software availability

Software implementing BRASS is available at
`https://github.com/joellesophya/BRASS`.

Acknowledgments

This work was supported by NIH grant R01 HG001645 (to M.S.M.). This study makes use of data generated from a previous study. [27] The data are available from the Dryad online repository at https://datadryad.org/resource/doi:10.5061/dryad.266k4.

References

1. Li J, Malley JD, Andrew AS, Karagas MR, Moore JH. Detecting gene-gene interactions using a permutation-based random forest method. BioData Mining. 2016;9:14.

2. Cui T, Mekkaoui KE, Reinvall J, Havulinna AS, Marttinen P, Kaski S. Gene–gene interaction detection with deep learning. Communications Biology. 2022;5:1238.

3. Richter F, Morton S, Kim S, Kitaygorodsky A, Wasson L, KM C, et al. Genomic analyses implicate noncoding de novo variants in congenital heart disease. Nature Genetics. 2020;52:769–777.

4. Lee S, Abecasis GR, Boehnke M, Lin X. Rare-variant association analysis: Study designs and statistical tests. American Journal of Human Genetics. 2014;95(1):5–23. doi:10.1016/j.ajhg.2014.06.009.

5. Dudbridge F, Gusnanto A. Estimation of significance thresholds for genomewide association scans. Genetic Epidemiology. 2008;32(3):227–234. doi:10.1002/gepi.20297.

6. Atkinson EG, Maihofer AX, Kanai M, Martin AR, Karczewski K, Santoro ML, et al. Tractor uses local ancestry to enable the inclusion of admixed individuals in GWAS and to boost power. Nature Genetics. 2021;53:195–204.

7. Sheng X, Guan Y, Ma Z, Wu J, Liu H, Qiu C, et al. Mapping the genetic architecture of human traits to cell types in the kidney identifies mechanisms of disease and potential treatments. Nature Genetics. 2021;53:1322–1333.

8. Astle W, Balding DJ. Population structure and cryptic relatedness in genetic association studies. Statistical Science. 2009; p. 451–471.

9. Abney M, Ober C, McPeek MS. Quantitative-Trait Homozygosity and Association Mapping and Empirical Genomewide Significance in Large, Complex Pedigrees: Fasting Serum-Insulin Level in the Hutterites. The American Journal of Human Genetics. 2002;70(4):920–934. doi:10.1086/339705.

10. Churchill GA, Doerge RW. Naive application of permutation testing leads to inflated type I error rates. Genetics. 2008;178(1):609–610. doi:10.1534/genetics.107.074609.

11. Abney M. Permutation testing in the presence of polygenic variation. Genetic Epidemiology. 2015;39(4):249–258. doi:10.1002/gepi.21893.

12. Zhang T, Sun L. The hidden factor: accounting for covariate effects in power and sample size computation for a binary trait. Bioinformatics. 2023;39(4):btad139. doi:10.1093/bioinformatics/btad139.

13. Bianchi M, Dahlgren S, Massey J, Dietschi E, Kierczak M, Lund-Ziener M, et al. A multi-breed genome-wide association analysis for canine Hypothyroidism identifies a shared major risk locus on CFA12. PLoS ONE. 2015;10(8):e0134720. doi:10.1371/journal.pone.0134720.

14. Melin M, Rivera P, Arendt M, Elvers I, Murén E, Gustafson U, et al. Genome-Wide Analysis Identifies Germ-Line Risk Factors Associated with Canine Mammary Tumours. PLoS Genetics. 2016;12(5):e1006029. doi:10.1371/journal.pgen.1006029.

15. Safra N, Bassuk AG, Ferguson PJ, Aguilar M, Coulson RL, Thomas N, et al. Genome-Wide Association Mapping in Dogs Enables Identification of the Homeobox Gene, NKX2-8, as a Genetic Component of Neural Tube Defects in Humans. PLoS Genetics. 2013;9(7):e1003646. doi:10.1371/journal.pgen.1003646.

16. Tengvall K, Kierczak M, Bergvall K, Olsson M, Frankowiack M, Farias FHG, et al. Genome-Wide Analysis in German Shepherd Dogs Reveals Association of a Locus on CFA 27 with Atopic Dermatitis. PLoS Genetics. 2013;9(5):e1003475. doi:10.1371/journal.pgen.1003475.

17. Zhong S, Jiang D, McPeek MS. CERAMIC: Case-Control Association Testing in Samples with Related Individuals, Based on Retrospective Mixed Model Analysis with Adjustment for Covariates. PLoS Genetics. 2016;12(10):e1006329. doi:10.1371/journal.pgen.1006329.

18. Jiang D, Zhong S, McPeek MS. Retrospective Binary-Trait Association Test Elucidates Genetic Architecture of Crohn Disease. American Journal of Human Genetics. 2016;98(2):243–255. doi:10.1016/j.ajhg.2015.12.012.

19. Chen H, Wang C, Conomos MP, Stilp AM, Li Z, Sofer T, et al. Control for Population Structure and Relatedness for Binary Traits in Genetic Association Studies via Logistic Mixed Models. American Journal of Human Genetics. 2016;98(4):653–666. doi:10.1016/j.ajhg.2016.02.012.

20. Thornton TA. Statistical methods for genome-wide and sequencing association studies of complex traits in related samples. Current Protocols in Human Genetics. 2015;2015(January):1.28.1–1.28.9. doi:10.1002/0471142905.hg0128s84.

21. Price AL, Patterson NJ, Plenge RM, Weinblatt ME, Shadick NA, Reich D. Principal components analysis corrects for stratification in genome-wide association studies. Nature Genetics. 2006;38(8):904–909. doi:10.1038/ng1847.

22. Conomos MP, Reiner AP, Weir BS, Thornton TA. Model-free Estimation of Recent Genetic Relatedness. American Journal of Human Genetics. 2016;98(1):127–148. doi:10.1016/j.ajhg.2015.11.022.

23. Listgarten J, Lippert C, Kadie CM, Davidson RI, Eskin E, D H. Improved linear mixed models for genome-wide association studies. Nature Methods. 2012;9:525–526.

24. Yang J, Zaitlen NA, Goddard ME, Visscher PM, Price AL. Advantages and pitfalls in the application of mixed-model association methods. Nature Genetics. 2014;46:100–106.

25. Zhang T, Sun L. Beyond the traditional simulation design for evaluating type 1 error control: from the "theoretical" null to "empirical" null. Genetic Epidemiology. 2019;43:166–179.

26. Balding DJ, Nichols RA. A method for quantifying differentiation between populations at multi-allelic loci and its implications for investigating identity and paternity. Genetica. 1995;96(1-2):3–12. doi:10.1007/BF01441146.

27. Hayward JJ, Castelhano MG, Oliveira KC, Corey E, Balkman C, Baxter TL, et al. Complex disease and phenotype mapping in the domestic dog. Nature Communications. 2016;7. doi:10.1038/ncomms10460.

28. Rodriguez G, Goldman N. Improved estimation procedures for multilevel models with binary response: a case-study. Journal of the Royal Statistical Society: Series A (Statistics in Society). 2001;164(2):339–355. doi:10.1111/1467-985X.00206.

29. Raudenbush SW, Yang ML, Yosef M. Maximum Likelihood for Generalized Linear Models with Nested Random Effects via High-Order, Multivariate Laplace Approximation. Journal of Computational and Graphical Statistics. 2000;9(1):141–157. doi:10.1080/10618600.2000.10474870.

30. Rabe-hesketh S, Skrondal A, Pickles A. Reliable estimation of generalized linear mixed models using adaptive quadrature. Stata Journal. 2002;2(1):1–21.

31. Conomos MP, Miller MB, Thornton TA. Robust inference of population structure for ancestry prediction and correction of stratification in the presence of relatedness. Genetic Epidemiology. 2015;39(4):276–293. doi:10.1002/gepi.21896.

Supporting information

S1 Text.
S1 Fig. Three-generation pedigree used in the simulation studies.
S2 Fig. Empirical Type 1 Error Rates Omitting a Relevant Covariate at Nominal Level 0.01.
S1 Table. Differences between the seven resampling methods compared in the simulation studies.

A. ED

B. IE