Sperm gatekeeping: 3D imaging reveals a constricted entrance to zebra finch

sperm storage tubules

Authors: Tania Mendonca[1,2], Ashley J Cadby[2] and Nicola Hemmings[1]

[1]Department of Animal and Plant Sciences, University of Sheffield, Western Bank,

Sheffield S10 2TN, UK

[2]Department of Physics and Astronomy, University of Sheffield, Western Bank, Sheffield

S10 2TN, UK

Corresponding author: Tania Mendonca (t.mendonca@sheffield.ac.uk)

Short Title: Visualising female sperm storage

Abstract:

Females across many internally fertilising taxa store sperm, often in specialised storage

organs in their reproductive tracts. In birds, several hundred sperm storage tubules exist

in the utero-vaginal junction of the oviduct and there is growing evidence that sperm

storage in these tubules is selective. The nature of the mechanisms underlying female

sperm storage in birds remains unknown due to our limited ability to make three

dimensional, live observations inside the large, muscular avian oviduct. Here, we describe

a new application of fluorescence selective plane illumination microscopy to optically

section oviduct tissue from zebra finch *Taeniopygia guttata* females label-free, by

harnessing tissue autofluorescence. Our data provide the first description of the 3D

structure of sperm storage tubules in any bird, and reveal the presence of constricted

openings to tubules, suggesting active entrapment and release of sperm.

Introduction:

Across many internal fertilisers, females have evolved the capacity to maintain viable sperm in specialised sperm storage organs in their reproductive tract as a strategy to maximise fertility. Sperm storage ensures the female has sufficient sperm for fertilisation when copulation and ovulation are not synchronised [1]. Since female promiscuity is common across taxa (e.g. birds [2], mammals [3–5], reptiles [6], fishes [7] and insects [8–10]), storage also provides the opportunity for females to exert control over post-copulatory processes [11–13]. Post-copulatory sexual selection has driven the diversification of sperm storage organs, which vary from single bean-shaped structures in many damselfly species [14,15], to one or more sac-like spermathecae in certain fly species (e.g. three spermathecae in *Scathophaga stercoraria* [16]; two spermathecae along with a seminal receptacle in many Drosophila species [17]), and multiple epithelial crypts in snakes [18], lizards [19], turtles [20] and birds [21,22].

In birds, epithelial sperm storage crypts are called sperm storage tubules (SSTs) and are located in the utero-vaginal junction (UVJ) of the oviduct [23]. The number of SSTs possessed by a single female ranges from around 500 SSTs in the UVJ of the budgerigar *Melopsittam undulatus* to 20,000 in the turkey *Meleagris gallopavo* [24]. A growing body of evidence suggests that avian SSTs may be an important site of sperm selection. Steele and Wishart [25] demonstrated experimentally that when chicken sperm were treated to remove surface proteins from their membranes, they could not enter the SSTs after normal intra-vaginal artificial insemination, even though sperm treated in the same way were capable of fertilising the ovum when inseminated further up the oviduct, past the vagina and UVJ. Bobr *et al.* [23] also noted a lack of abnormal sperm in the SSTs of domestic fowl *Gallus domesticus*, suggesting that abnormal sperm are unable to reach or

enter sperm storage sites. The large number of SSTs present in the avian oviduct may also

allow spatio-temporal segregation of sperm from competing ejaculates within the

oviduct [26–28]. However, the mechanisms by which sperm are selected at the time of

storage remain poorly understood, and how sperm enter and exit the SSTs is unknown.

The role of the SSTs in sperm storage and selection may be relatively passive. Froman

[29] proposed a model where sperm motility, rather than SST function, is pivotal in sperm

retention in SSTs. According to this model, sperm must maintain an optimum swimming

velocity to maintain their position and counter a fluid current within the SST. This model

was supported by evidence that faster sperm emerged out of SSTs later than slower

sperm [30], and that passive loss of sperm from storage might be sufficient to explain last

male precedence in the domestic fowl, turkeys, and zebra finches *Taeniopygia guttata*

[31,32; but see 27]. However, there have been no observations of sperm swimming inside

the SSTs, and other studies have detected the presence of sperm motility suppressors

such as lactic acid in Japanese quail *Coturnix japonica* SSTs [33], calcium and zinc in the

SSTs of chicken, turkeys and Japanese quail [34,35], and carbonic anhydrase in the SSTs

of turkeys, common quail *Coturnix coturnix* and ostriches *Struthio camelus* [36–38]. The

neurotransmitter acetylcholine, released by nerve endings detected in the vicinity of SSTs

[39], has been shown to enhance sperm motility [40], implying a nervous control on

sperm mobilisation at ejection from SSTs. Additionally, Hiyama *et al.* [41] presented

evidence for the potential role of heat shock protein 70 (HSP70) [42] in enhancing sperm

motility at the point of sperm release. HSP70 proteins were found to be localised on the

surface epithelium of the UVJ and their levels were upregulated ~20h post oviposition in

Japanese quail, coincidental with when sperm have been shown to be released from

storage in these birds [41]. The presence of such sperm motility suppressors and

activators within or near the SSTs suggests storage may not be as passive as Froman [29]

suggested.

It is possible that rather than acting as passive refugia, SSTs may be dynamic structures,

capable of active constriction and dilation to mediate the entrance and exit of sperm.

Although numerous studies have failed to find smooth muscle fibres or myoepithelial

cells [39,43,44] around SSTs, Freedman *et al.* [39] detected fibroblast-like cells and an F-

actin rich cytoskeletal mesh called the "terminal web" in turkey SST epithelia. The

terminal web is composed of contractile proteins (actin and myosin) and has been shown

to contribute to contractility in other tissues, such as intestinal brush border cells [45,46]

and embryonic pigmented epithelia in chicken [47]. Freedman *et al.* [39] also found

terminal innervations in the turkey UVJ, suggesting there may be some degree of nervous

control over SST function. Recent evidence also suggests the possibility of SST

contraction, influenced hormonally by the action of progesterone [48,49]. It is therefore

possible that the passage of sperm into and out of storage is controlled, to some degree,

by the physical structure of SSTs themselves.

Our understanding of how SST structure influences sperm storage is limited by our

relatively basic knowledge of SST morphology. The avian oviduct is convoluted, with

opaque, muscular walls, creating numerous practical limitations for making observations

of tubules in living epithelial tissue using conventional microscopy techniques. Empirical

studies of SST morphology have so far used histology [23,50,51] and electron microscopy

[34,43,52] on fixed tissue sections, but these approaches not only remove functional

information, but typically provide two-dimensional information only. Moreover, serial

sectioning is laborious, and loss of material can be difficult to avoid. Commonly used light

microscopy techniques rely on thin sections and squash preparations [26,53], which are

largely inappropriate for large tissue samples since they distort structures of interest and

allow only limited imaging depths.

In this study we used selective plane illumination microscopy (SPIM), to develop a novel

method for live, *ex vivo* imaging and examining the 3D structure of avian SSTs for the first

time. SPIM has been demonstrated to be highly suitable for imaging large samples at

cellular resolution with lower phototoxicity levels than with other optical sectioning

methods [54,55]. Using SPIM, we were able to optically section tissue up to depths of 100

μm, so that entire folds of UVJ mucosal tissue could be imaged live without distorting or

damaging their structure. We provide the first true estimates of the 3D structure of avian

SSTs in living tissue, including the relationship between SST length and diameter, and

report the existence of a previously unreported physical constriction at the entrance to

tubules that may act to regulate sperm transport into and out of storage. We also find no

evidence of sperm motility inside or on release from sperm storage tubules, providing

support for the idea that sperm motility is suppressed inside the SSTs.

Materials and Methods:

Animals

Zebra finches were from a captive population kept at the University of Sheffield

[56,57]. Females (all between one to three years old) were placed in unisex housing

for at least two weeks before being paired with males, in cages (dimensions of each:

0.6m x 0.5m x 0.4m) placed side-by-side and separated by a wire-mesh with the male

and female on either side. A modified nest-box with a wire-mesh partition was

supplied to each pair. This set up ensured that the male and female established a

normal breeding pair bond, leading the female to come into breeding condition, while

ensuring that she had no sperm in her SSTs (sperm can be stored for up to 12 days

after mating in zebra finches [24]). Only females from this no-copulation set-up were

used for SST structural imaging, and all imaging methods described below, unless

stated otherwise, refer to this part of the study. For a separate study, a small number

of females were allowed to copulate with males, and from these females we collected

data on sperm motility inside and on release from the SSTs using different methods

(see "Sperm motility" below). Females were only used once they had started to lay

eggs, to ensure their oviduct was in full reproductive condition. After the second egg

in the ovulatory cycle, females were euthanized (in accordance with Schedule 1

[Animals (Scientific Procedures) Act 1986]] and dissected immediately.

Sample preparation

The oviduct, including the cloaca, was immediately removed from the female and the

connective tissue surrounding it was cleared to uncoil and straighten the vagina and

the UVJ. The lower end of the oviduct was cut through the middle of the uterus to

obtain a segment that included the UVJ, vagina and cloaca. This piece of the oviduct

was then cut open lengthwise and pinned flat on a petri-dish filled with silicone

elastomer (SYLGARD® 184; Dow Corning). A sufficient quantity of Ham's F10 Nutrient

Mix (Invitrogen, UK) was added to keep the tissue moist but not submerged. For SPIM

imaging, UVJ folds were cut individually with iris scissors and mounted one at a time,

on a custom-made sample holder (supplementary materials) using fine insect needles.

The sample holder, with the UVJ fold mounted, was immersed in phenol-free

DMEM/F12 media, at 37º C during imaging.

SPIM imaging

Live UVJ tissue samples, prepared as above, were imaged using a custom-built SPIM

microscope (at the University of Sheffield) with laser excitation at 473 nm and a 520

nm long pass (LP) fluorescence emission filter (Semrock, Inc.). The microscope

hardware and optical components are based on the OpenSPIM platform [58] but with

modifications detailed in Mendonca *et al*. [59 supplementary materials]. The camera,

detection and illumination objectives, and magnification are fixed for the system,

ensuring that the imaging results are reproduceable. The autofluorescence image

stacks were acquired using 500 ms exposure, starting at the outer surface and moving

up to 100 μm deep into the tissue fold.

Characterisation of autofluorescence

SSTs were clearly identifiable in live UVJ tissue during fluorescence imaging on the

SPIM and had a punctate appearance on account of autofluorescent granules (Figure

1, (B and D)) which appeared to be mostly confined to SST epithelial cells and were

present along the entire length of the SST from orifice to blind end. No other cell

structure or organelle was visible in these autofluorescence images.

To determine the organisation of the autofluorescent granules in the SST epithelium,

label-free images from live UVJ folds (n = 10 birds) were acquired on the SPIM were

compared to those of fixed UVJ tissue folds, imaged on the SPIM after staining for

nucleic acids (n = 3 birds), and on a bright-field microscope after histological

sectioning and general histochemical staining (n = 3 birds) (Supplementary

materials).

Figure 1: (A) Schematic of SST transverse section showing cellular polarisation with nuclei (N) towards the basement membrane (BM) and microvilli (MV) at the apical end of the epithelium. The punctate autofluorescence (PA) detected by the SPIM is present proximal to the nucleus but not at the apical end of the epithelium near the lumen (L). (B) Max intensity projection of a UVJ fold with multiple detectable SSTs imaged on the SPIM. (C)-(E) Cross section of SST from histology, autofluorescence imaged on the SPIM, and SYTO-13 labelled nuclei imaged on the SPIM respectively.

Image Analysis

The image stacks acquired using the SPIM were used to reconstruct 480 μm x 480 μm x 100 μm tissue sections containing 3D information on SST structure (n = 10 females, one SST per female). SST shape information was extracted by measuring the diameter

enclosed by the autofluorescence from images of live tissue, at ten equidistant points

along the length of the SST.

UVJ tissue image stacks were first pre-processed in Fiji [60]. Individual unbranched

SSTs were selected from each female such that the entire SST structure was included

in the 3D image stacks. The SSTs follow convoluted paths through the UVJ fold tissue,

so in order to measure cross-sectional diameter at multiple points, it was necessary

to slice the image volume at arbitrary angles to ensure the measurement planes were

perpendicular to the direction of the SST structure. This was accomplished by first

tracing the direction of the SST structure using a dilated version of the SST image

(generated using the 'MorphoLibJ' plugin [61] followed by the application of Gaussian

blur), to smooth the punctate autofluorescence. Two outlines for each SST were then

semi-automatically traced in 3D, from the orifice to the blind end and along opposite

sides of the SST lumen, using the 'Simple Neurite Tracer' [62] plugin in Fiji.

The next stage of image analysis was performed using MATLAB® (2015b, version 8.6,

MathWorks, Natick, MA). An average trace which passed through the SST lumen was

computed from the two traces for each SST. SST lengths were measured from these

average traces. For each SST (n = 10), the average trace was interpolated at ten

equidistant points (the first at the orifice and the tenth point before the blind end of

the tubule) and at each interpolated point a vector describing the direction of the SST

at that point was computed using its nearest neighbouring points on the trace. By

using these vectors and the interpolated points, slicing planes normal to the vectors

were defined. The indices for these slicing planes were used to extract 2D image

sections from the un-dilated original image stacks using the 'ExtractSlice.m' [63]

function. For every extracted slice, its distance from the orifice along the luminal trace

of the SST was computed using the 'Arclength.m' [64] function, and the major axis

diameter (d1) and the minor axis diameter (d2) of the SST (enclosed by

autofluorecence) were measured.

Sperm motility

To test the hypothesis that sperm are motile in the SSTs [29], we used tissue from a

subset of 13 females from a separate study, that had copulated with males prior to

dissection (and therefore had sperm inside their SSTs). These females were dissected

as described above, but instead of mounting UVJ folds for SPIM imaging, folds were

dissected on a microscope slide under a Nikon SMZ25 stereomicroscope to isolate and

open individual SSTs and release sperm. We identified one SST per fold (46 SSTs in

total) with high numbers of stored sperm (mean = 33 per SST), isolated the SST using

methods described in [28] and submerged it in warmed nutrient media (Ham's F10:

Invitrogen, UK; typically used for avian sperm motility analysis) on a microscope slide.

The slide was then placed on a heated stage (set at 38º C) and observed under a 200X

phase contrast microscope for 5 mins to screen for sperm motility inside the tubule. After

this initial observation period, the SST was pulled open to release sperm using fine

dissection needles, as described in [28], and sperm were gently shaken free of the tubule.

The slide was then observed under a 200X phase contrast microscope for another 5 mins

to screen for sperm motility after release from the tubule.

Statistical Analysis:

Data analysis was performed using the statistical package R, version 3.2.3 [65]. We

tested whether SST diameter varied with SST length using a mixed effects model

('lmer' function from the 'lme4' package [66] along with the 'lmerTest' package [67])

with average SST diameter [(d1+ d2) / 2] at the sampled point as the dependent

variable, the distance of sampled point from SST orifice and the SST total length as

fixed effects, and the bird ID as a random effect to account for repeated measures from

each female.

We also assessed if the SST was elliptical or circular in cross-section (the former

providing greater epithelial apical surface area for increased contact with sperm) and

whether any such ellipticity changed in response to SST length. A circularity index was

first calculated by dividing the major axis diameter (d1) by the minor axis diameter

(d2), where a circularity index of one indicates a circular SST cross-section. Data were

then analysed via a mixed effects model using the 'lmer' function [66], with the

circularity index as the dependent variable and the total length of the SST as a fixed

effect. The sum of the major and minor axis diameters (d1+ d2) was also incorporated

as a fixed effect to account for magnitude of change in diameter along each axis, as

well as the distance of sampled point from SST orifice, with an interaction term

between them. As before, bird ID was included as a random effect to account for

repeated measures from each female.

Results:

The diameter of SSTs was found to be notably constricted at their orifice, suggestive

of a structural 'barrier' for entry and exit (Figure 2 (A)). SST diameter increased

slightly increase along the SST's length up to a mid-point, after which it decreased

slightly towards the blind end of the SST, resulting in a significant quadratic

relationship between SST diameter and the distance from the SST orifice (estimated

effect = -16.761, t = -3.085, p = 0.003; Figure 2 (B)). There was no relationship between

SST length and either mean, maximum, or minimum diameter, so long SSTs were

neither wider nor thinner than short SSTs (estimated effect = 0.038, t = 1.058, p =

0.319; Figure 4). The relationship between SST diameter and distance from orifice

also held true with data from labelled tissue (Supplementary Figure S4), confirming

that the shape measured from autofluorescence images was real and not an artefact

resulting from the distribution of the autofluorescence granules.

Figure 2: (A) 3D rendering of an SST showing its constricted orifice (arrowhead). (B) SST diameter

has a quadratic relationship with distance from the SST opening suggesting a constriction at the

orifice and a slight increase in diameter along its length up to the middle of the SST. Each colour

on the plot represents measurements from the same SST (n = 10). Scale on red bounding box is in

microns.

SSTs were found to be slightly elliptical in cross-section, with the major axis diameter

being 1.642 ± 0.168 times larger than the minor axis diameter. The circularity of the

SST in cross-section did not vary significantly with SST diameter (estimated effect =

0.014, t = 0.979, p = 0.330), distance from orifice (estimated effect = 0.003, t = 0.587,

p = 0.558), or the interaction between these two variables (estimated effect = -

0.00009, t = -0.587, p = 0.559). Circularity was also not related to SST total length

(estimated effect = -0.004, t = -1.14, p = 0.282).

Comparisons between the SST measurements from histology and SPIM images

indicate that the autofluorescent granules are present in the supranuclear region of

the SST epithelium (Figure 1). The size of the lumen diameter scaled linearly with the

width of the SST (Supplementary Figure S4), indicating that epithelial cells remained

the same thickness in cross-section with increasing SST diameter. This allowed us to

extrapolate shape information from the above analyses to the SST lumen, and using

this method, we estimated the diameter of the SST orifice to be 3.278 ± 1.115 μm

(mean ± SD; Table 1).

Finally, we found no evidence of sperm motility either inside, or on release from, the

SSTs. Of 1520 sperm across 46 SSTs (from 13 females), no sperm were observed

actively swimming before or after release from the tubules.

Table 1: SST dimensions at orifice and widest section

	Diameter (mean ± SD)	
	at orifice (μm)	**at widest section (μm)**
inter-nuclear diameter[1]	12.060 ± 1.419	30.830 ± 11.055
autofluorescence[1]	10.065 ± 4.288	16.374 ± 6.610
lumen diameter[2]	3.278 ± 1.115	9.128 ± 1.377

SST diameter is the smallest at its orifice and widest at the relative slice position 'slice 6', where

slice 1 is at the orifice and slice 10 is nearest to the blind end of the SST.

[1]Measurements acquired from SPIM image z-stacks

[2]SST lumen diameter values were predicted from the model describing the relationship between

lumen and inter-nuclear diameter.

Discussion:

Using novel 3D imaging methods, we have demonstrated for the first time the

existence of a constricted orifice at the entrance/exit of avian SSTs. Although the

precise function of this constriction remains to be identified, it is likely that it plays

an important role in sperm selection and storage. Consistent with previous findings

of an F-actin rich terminal web in the SSTs in turkeys [39], this constricted opening is

suggestive of contractile activity for active entrapment and release of sperm.

However, even without contractility, the narrow SST orifice (mean diameter of about

3 μm), with the added obstruction of microvilli (1-2 μm in length, [52]) could restrict

the rate of sperm (mean diameter at mid-piece - ~0.6 μm [59]) entering and exiting

the SST, thereby providing a mechanism by which sperm storage and release can be

regulated. This, combined with the fact that sperm do not appear to actively swim

inside or on immediate release from the tubules, supports the idea that avian SSTs

play an active and selective role in sperm storage, regulating sperm uptake and

release and suppressing motility [33–38]. The constricted orifice, together with its

microvilli, may act as a valve, allowing unidirectional movement of sperm only, and

preventing them from being flushed back out. The small luminal diameter along the

SST (mean: 9 μm, Table 1) may also limit the ability of sperm to turn around inside

the SST and swim out.

SSTs were also found to be slightly elliptical in cross-section, with the major axis

diameter being about 1.6 times larger than the minor axis diameter. This ellipticity

was independent of SST radius, the distance along the SST from orifice, or total SST

length. Cross-sectional ellipticity increases the surface area of the SST epithelial apical

surface, allowing for a greater number of microvilli (as compared to a circular lumen

with the same volume) for increased contact with sperm and optimum exchange of

nutrients and waste.

SST diameter varied widely across SSTs. Birkhead *et al.* [68] suggested that some SSTs

might remain inactive in the zebra finch UVJ even in its fully developed state. It is

possible that some of the variation in SST shape that we observed can be explained by

the presence of functional and non-functional SSTs, but it is unclear whether thinner,

more uniform SSTs, or more distended morphs would represent active SSTs. Mero and

Ogasawara [69], and Burke [70] described 'swollen' tubules in chicken and suggested

swelling to be associated with sperm release. Such swellings might help explain the

outliers in our data (Figure 2). It is possible that conformational changes in SST shape

from functional to non-functional may be caused by neural stimulation [39,71] and/or

hormonal effects [48,49], potentially aiding sperm release. Variation in SST shape

might also be explained by factors not tested in this study, including age, hormone

levels and location of the SST in the UVJ.

Figure 3: Sperm storage tubules (SSTs) imaged label-free using the SPIM. SST shape showed a
constricted opening (arrowheads) but varied between more elliptical shapes (left) and thinner,
more uniform shapes (right). Scale on red bounding box is in microns.

About 4 – 27% of all the sperm storage tubules in the zebra finch UVJ are branched

[28,72]. These were not included in the present study, but individual branches are

expected to show similar shapes as unbranched tubules. Hemmings and Birkhead [28]

described sperm from different males differentially stored in separate branches of a

branched SST (albeit a single observation, since in most cases sperm from different

males were stored in different SSTs). Further study of the 3D structure of branched

SSTs could shed light on the mechanisms that might prevent sperm mixing in

branched tubules.

Our novel 3D data on SST structure was made possible by the presence of

punctate/granular autofluorescence, confined to the SST epithelial cells and

uniformly distributed throughout the SST's entire length. These granules were found

to have a supranuclear localisation in the SST epithelial cells (Figure 1). Although

identifying the exact source of the autofluorescence was beyond the scope of this

study, autofluorescence in a similar range has been noted in the ewe *Ovis aries*

endometrium ($\lambda_{ex}/\lambda_{em}$ = 488/525-575 nm) [73] and in human colonic crypts ($\lambda_{ex}/\lambda_{em}$

= 488/580 nm) [74]. While such autofluorescence has been attributed to NADH

metabolism in mitochondria [73,75], another likely source might be lipofuscin in

lysosomes [74]. Mitochondria are not confined to the apical cytoplasm of SST

epithelium as observed in turkeys [76] and domestic fowl [77] so it is unlikely that

these granules represent mitochondria. Lysosomes on the other hand, are globular

vesicles similar to the autofluorescent granules observed here in size (< 1 μm) [78],

and have been detected in the apical cytoplasm of turkey SST epithelia [76], and less

abundantly in domestic fowl [77] and passerine alpine accentor *Prunella collaris* [78].

Multiple studies have also detected the presence of acid phosphatase, an enzyme

found in lysosomes, in the supranuclear cytoplasm of SST epithelia in turkeys [21],

quail [79], domestic fowl [80] and ducks *Anas sp.* [81], but not in the SST lumen, which

corresponds with the autofluorescence pattern we observed here. Acid phosphatase

has been implicated in autolysis associated with oviduct regression [82] as well as

with sperm release [81]. If this is true, the label-free imaging methods developed here

may provide exciting new means for investigating SST functional development

throughout the reproductive cycle. Identifying the chemical nature of the

autofluorescent substance present in SST granules therefore represents an important

avenue for future research.

In summary, we have demonstrated that SSTs in living avian oviductal tissue can be

imaged label-free using SPIM microscopy, and this novel 3D imaging technique has

enabled us to produce the most detailed account of avian SST structure to date,

including the discovery of a previously unknown structural constriction at the

entrance/exit of tubules that may play a key role in the regulation of sperm storage

and selection. The imaging methods described here hold immense potential for

studying SST function and interactions with sperm in real time.

Ethics:

This study was approved by the University of Sheffield, UK. All procedures performed conform to the

legal requirements for animal research in the UK and were conducted under a project licence (PPL

40/3481) issued by the Home Office. Animals were humanely killed under Schedule 1 (Animals

(Scientific Procedures) Act 1986).

Competing interests:

The authors declare no conflict of interest.

Authors' contributions:

T.M. coordinated the study, conducted the experiments and analyses, and wrote the manuscript; N.H.

conceived the study and advised on data analyses. All authors participated in the design of the study,

helped draft the manuscript and gave final approval for publication.

Acknowledgements:

The authors are grateful to Prof Tim Birkhead and Prof Simon Jones from the University of Sheffield

for support. Thanks to Mark Kinch from Skelet.AL, Phil Young, Lynsey Gregory, Emily Glendenning

and Jamie Thompson from the University of Sheffield for technical assistance.

Funding:

TM was funded by an Imagine: Imaging Life studentship granted by the University of Sheffield. NH

was funded by ERC grant no. 268688 and a Patrick and Irwin-Packington Independent Research

Fellowship from the Department of Animal & Plant Sciences, University of Sheffield.

References:

1. Birkhead TR. 1992 Sperm Storage and the Fertile Period in the Bengalese Finch.

Auk **109**, 620–625. (doi:10.2307/4088376)

2. Birkhead TR, Møller AP. 1995 Extra-pair copulation and extra-pair paternity in

birds. *Anim. Behav.* **49**, 843–848. (doi:10.1016/0003-3472(95)80217-7)

3. Hanken J, Sherman PW. 1981 Multiple paternity in Belding's ground squirrel

litters. *Science (80-.).* **212**, 351–353. (doi:10.1126/science.7209536)

4. Tegelström H, Searle J, Brookfield J, Mercer S. 1991 Multiple paternity in wild

common shrews (Sorex araneus) is confirmed by DNA-fingerprinting. *Heredity*

(Edinb). **66 (Pt 3)**, 373–379. (doi:10.1038/hdy.1991.47)

5. Amos W, Twiss S, Pomeroy PP, Anderson SS. 1993 Male mating success and

paternity in the grey seal, Halichoerus grypus: a study using DNA fingerprinting.

Proc. R. Soc. B **252**, 199–207. (doi:10.1098/rspb.1993.0066)

6. Uller T, Olsson M. 2008 Multiple paternity in reptiles: Patterns and processes.

Mol. Ecol. **17**, 2566–2580. (doi:10.1111/j.1365-294X.2008.03772.x)

7. Avise JC, Jones AG, Walker D, DeWoody JA. 2002 Genetic mating systems and

reproductive natural histories of fishes: lessons for ecology and evolution. *Annu.*

Rev. Genet. **36**, 19–45. (doi:10.1146/annurev.genet.36.030602.090831)

8. Gromko MH, Gilbert DG, Richmond RC. 1984 Sperm transfer and use in the

multiple mating system of Drosophila. *Sperm Compet. Evol. Anim. mating Syst.* ,

371–426.

9. Moritz RFA, Kryger P, Koeniger G, Koeniger N, Estoup A, Tingek S. 1995 High

degree of polyandry in Apis dorsata queens detected by DNA microsatellite

variability. *Behav. Ecol. Sociobiol.* **37**, 357–363. (doi:10.1007/BF00174141)

10. Crozier RH, Fjerdingstad EJ. 2001 Polyandry in social Hymenoptera - disunity in

445   diversity? *Ann. Zool. Fennici* **38**, 267–285. (doi:10.2307/23735845)

446 11. Birkhead TR, Møller AP. 1998 *Sperm competition and sexual selection*. Academic

447   Press.

448 12. Simmons LW. 2001 *Sperm competition and its evolutionary consequences in the*

449   *insects*. Princeton University Press.

450 13. Eberhard WG. 1996 *Female Control: Sexual Selection by Cryptic Female Choice*.

451   Princeton University Press.

452 14. Córdoba-Aguilar A, Uhía E, Rivera AC. 2003 Sperm competition in Odonata

453   (Insecta): the evolution of female sperm storage and rivals' sperm displacement. *J.*

454   *Zool.* **261**, 381–398. (doi:10.1017/S0952836903004357)

455 15. Siva-Jothy MT. 1987 The structure and function of the female sperm-storage

456   organs in libellulid dragonflins. *J. Insect Physiol.* **33**, 559–567. (doi:10.1016/0022-

457   1910(87)90070-9)

458 16. Ward PI. 1993 Females influence sperm storage and use in the yellow dung fly

459   Scathophaga stercoraria (L.). *Behav. Ecol. Sociobiol.* **32**.

460   (doi:10.1007/BF00183786)

461 17. Pitnick S, Markow T, Spicer GS. 1999 Evolution of Multiple Kinds of Female

462   Sperm-Storage Organs in Drosophila. *Evolution (N. Y).* **53**, 1804.

463   (doi:10.2307/2640442)

464 18. Fox W. 1956 Seminal receptacles of snakes. *Anat. Rec.* **124**, 519–539.

465   (doi:10.1002/ar.1091240303)

466 19. Conner J, Crews D. 1980 Sperm transfer and storage in the lizard Anolis

467   carolinensis. *J. Morphol.* **163**, 331–348.

468 20. Gist DH, Jones JM. 1989 Sperm storage within the oviduct of turtles. *J. Morphol.*

469   **199**, 379–384. (doi:10.1002/jmor.1051990311)

470   21.   Zavaleta D, Ogasawara F. 1973 A review of the mechanism of the release of

471   spermatozoa from storage tubules in the fowl and turkey oviduct. *Worlds. Poult.*

472   *Sci. J.* **43**, 132–139. (doi:10.1079/WPS19870010)

473   22.   Bakst MR. 1987 Anatomical basis of sperm-storage in the avian oviduct. *Scanning*

474   *Microsc.* **1**, 1257.

475   23.   Bobr LW, Lorenz FW, Ogasawara FX. 1964 Distribution of Spermatozoa in the

476   Oviduct and Fertility in Domestic Birds. I. Residence Sites of Spermatozoa in Fowl

477   Oviducts. *J. Reprod. Fertil.* **8**, 39–47. (doi:10.1530/JRF.0.0110257)

478   24.   Birkhead TR, Møller AP. 1992 Numbers and size of sperm storage tubules and the

479   duration of sperm storage in birds: a comparative study. *Biol. J. Linn. Soc.* **45**, 363–

480   372. (doi:10.1111/j.1095-8312.1992.tb00649.x)

481   25.   Steele MG, Wishart GJ. 1996 Demonstration that the removal of sialic acid from

482   the surface of chicken spermatozoa impedes their transvaginal migration.

483   *Theriogenology* **46**, 1037–1044. (doi:http://dx.doi.org/10.1016/S0093-

484   691X(96)00268-3)

485   26.   King LM, Brillard JP, Garrett WM, Bakst MR, Donoghue AM. 2002 Segregation of

486   spermatozoa within sperm storage tubules of fowl and turkey hens. *Reproduction*

487   **123**, 79–86. (doi:10.1530/rep.0.1230079)

488   27.   Briskie J V. 1996 Spatiotemporal patterns of sperm storage and last-male sperm

489   precedence in birds. *Funct. Ecol.* **10**, 375–383. (doi:10.2307/2390286)

490   28.   Hemmings N, Birkhead TR. 2017 Differential sperm storage by female zebra

491   finches Taeniopygia guttata. *Proc. R. Soc. B* **284**, 20171032.

492   (doi:10.1098/rspb.2017.1032)

493   29.   Froman DP. 2003 Deduction of a model for sperm storage in the oviduct of the

494   domestic fowl (Gallus domesticus). *Biol. Reprod.* **69**, 248–53.

(doi:10.1095/biolreprod.102.013482)

30.   Froman DP, Pizzari T, Feltmann AJ, Castillo-Juarez H, Birkhead TR. 2002 Sperm

mobility: mechanisms of fertilizing efficiency, genetic variation and phenotypic

relationship with male status in the domestic fowl, Gallus gallus domesticus. *Proc.*

R. Soc. B **269**, 607–12. (doi:10.1098/rspb.2001.1925)

31.   Birkhead TR, Pellatt JE, Hunter FM. 1988 Extra-pair copulation and sperm

competition in the zebra finch. *Nature* **334**, 60–2. (doi:10.1038/334060a0)

32.   Birkhead TR, Biggins JD. 1998 Sperm competition mechanisms in birds: models

and data. *Behav. Ecol.* **9**, 253–260. (doi:10.1093/beheco/9.3.253)

33.   Matsuzaki M *et al.* 2015 Lactic acid is a sperm motility inactivation factor in the

sperm storage tubules. *Sci. Rep.* **5**, 17643. (doi:10.1038/srep17643)

34.   Holm L, Ekwall H, Wishart GJ, Ridderstråle Y, Ridderstrale Y. 2000 Localization of

calcium and zinc in the sperm storage tubules of chicken, quail and turkey using

X-ray microanalysis. *J. Reprod. Fertil.* **118**, 331–336.

(doi:10.1530/reprod/118.2.331)

35.   Bakst MR, Richards MP. 1985 Concentrations of selected cations in turkey serum

and oviductal mucosae. *Poult. Sci.* **64**, 555–563.

36.   Holm L, Ridderstråle Y. 1998 Localization of carbonic anhydrase in the sperm-

storing regions of the turkey and quail oviduct. *Histochem. J.* **30**, 481–488.

(doi:10.1023/A:1003247504288)

37.   Holm L, Wishart GJ. 1998 The effect of pH on the motility of spermatozoa from

chicken, turkey and quail. *Anim. Reprod. Sci.* **54**, 45–54. (doi:10.1016/S0378-

4320(98)00142-0)

38.   Holm L, Ruziwa SD, Dantzer V, Ridderstråle Y. 2000 Carbonic anhydrase in the

utero-vaginal junction of immature and mature ostriches. *Br. Poult. Sci.* **41**, 244–9.

520  (doi:10.1080/713654906)

521 39. Freedman SL, Akuffo VG, Bakst MR. 2001 Evidence for the innervation of sperm

522  storage tubules in the oviduct of the turkey (Meleagris gallopavo). *Reproduction*

523  **121**, 809–814. (doi:10.1530/reprod/121.5.809)

524 40. Atherton RW, Cisson CM, Wilson BW, Golder TK. 1980 Quantitation of avian

525  spermatozoan motility: Neurochemical regulation. *Gamete Res.* **3**, 17–24.

526  (doi:10.1002/mrd.1120030104)

527 41. Hiyama G, Matsuzaki M, Mizushima S, Dohra H, Ikegami K, Yoshimura T, Shiba K,

528  Inaba K, Sasanami T. 2014 Sperm activation by heat shock protein 70 supports

529  the migration of sperm released from sperm storage tubules in Japanese quail

530  (Coturnix japonica). *Reproduction* **147**, 167–178. (doi:10.1530/rep-13-0439)

531 42. Lindquist S, Craig EA. 1988 the Heat -Shock Proteins. *Annu. Rev. Genet.* **22**, 631–

532  677. (doi:https://doi.org/10.1146/annurev.ge.22.120188.003215)

533 43. Van Krey HP, Ogasawara FX, Pangborn J. 1967 Light and Electron Microscopic

534  Studies of Possible Sperm Gland Emptying Mechanisms. *Poult. Sci.* **46**, 69–78.

535  (doi:10.3382/ps.0460069)

536 44. Gilbert AB, Reynolds ME, Lorenz FW. 1968 Distribution of spermatozoa in the

537  oviduct and fertility in domestic birds. V. Histochemistry of the uterovaginal

538  sperm-host glands of the domestic hen. *J. Reprod. Fertil.* **16**, 433–44.

539  (doi:10.1530/jrf.0.0160433)

540 45. Hirokawa N, Keller TCS, Chasan R, Mooseker MS. 1983 Mechanism of brush

541  border contractility studied by the quick-freeze, deep-etch method. *J. Cell Biol.* **96**,

542  1325–1336. (doi:10.1083/jcb.96.5.1325)

543 46. Keller TCS, Conzelman KA, Chasan R, Mooseker MS. 1985 Role of myosin in

544  terminal web contraction in isolated intestinal epithelial brush borders. *J. Cell*

545   *Biol.* **100**, 1647–1655. (doi:10.1083/jcb.100.5.1647)

546 47. Owaribe K, Kodama R, Eguchi G. 1981 Demonstration of contractility of

547   circumferential actin bundles and its morphogenetic significance in pigmented

548   epithelium in vitro and in vivo. *J. Cell Biol.* **90**, 507–514.

549   (doi:10.1083/jcb.90.2.507)

550 48. Ito T, Yoshizaki N, Tokumoto T, Ono H, Yoshimura T, Tsukada A, Kansaku N,

551   Sasanami T. 2011 Progesterone is a sperm-releasing factor from the sperm-

552   storage tubules in birds. *Endocrinology* **152**, 3952–3962. (doi:10.1210/en.2011-

553   0237)

554 49. Hemmings N, Birkhead TR, Brillard JP, Froment P, Briere S. 2015 Timing

555   associated with oviductal sperm storage and release after artificial insemination

556   in domestic hens. *Theriogenology* **83**, 1174–1178.

557   (doi:10.1016/j.theriogenology.2014.12.022)

558 50. Bakst MR. 1998 Structure of the avian oviduct with emphasis on sperm storage in

559   poultry. *J. Exp. Zool.* **282**, 618–626. (doi:10.1002/(SICI)1097-

560   010X(199811/12)282:4/5<618::AID-JEZ11>3.0.CO;2-M)

561 51. Van Drimmelen GC. 1946 'Spermnests' in the oviduct of the domestic hen. *J. South*

562   *African Vet. Med. Assoc* **17**, 42–52.

563 52. Bakst MR, Bauchan G. 2015 Apical blebs on sperm storage tubule epithelial cell

564   microvilli: Their release and interaction with resident sperm in the turkey hen

565   oviduct. *Theriogenology* **83**, 1438–44.

566   (doi:10.1016/j.theriogenology.2015.01.016)

567 53. Bakst MR. 1992 Observations on the turkey oviductal sperm-storage tubule using

568   differential interference contrast microscopy. *J. Reprod. Fertil.* **95**, 877–883.

569 54. Huisken J, Swoger J, Del Bene F, Wittbrodt J, Stelzer EHK. 2004 Optical Sectioning

Deep Inside Live Embryos by Selective Plane Illumination Microscopy. *Science*

(80-.). **305**, 1007–1009. (doi:10.1126/science.1100035)

55. Reynaud EG, Krzic U, Greger K, Stelzer EHK. 2008 Light sheet-based fluorescence

microscopy: more dimensions, more photons, and less photodamage. *HFSP J.* **2**,

266–75. (doi:10.2976/1.2974980)

56. Birkhead TR, Pellatt JE, Brekke P, Yeates R, Castillo-Juarez H. 2005 Genetic effects

on sperm design in the zebra finch. *Nature* **434**, 383–7.

(doi:10.1038/nature03374)

57. Bennison C, Hemmings N, Slate J, Birkhead T. 2015 Long sperm fertilize more

eggs in a bird. *Proc. R. Soc. B* **282**, 20141897. (doi:10.1098/rspb.2014.1897)

58. Pitrone PG, Schindelin J, Stuyvenberg L, Preibisch S, Weber M, Eliceiri KW,

Huisken J, Tomancak P. 2013 OpenSPIM: an open-access light-sheet microscopy

platform. *Nat. Methods* **10**, 598–599. (doi:10.1038/nmeth.2507)

59. Mendonca T, Birkhead TR, Cadby AJ, Forstmeier W, Hemmings N. 2018 A trade-

off between thickness and length in the zebra finch sperm mid-piece. *Proc. R. Soc.*

B Biol. Sci. **285**, 20180865. (doi:10.1098/rspb.2018.0865)

60. Schindelin J *et al.* 2012 Fiji: an open-source platform for biological-image analysis.

Nat. Methods **9**, 676–82. (doi:10.1038/nmeth.2019)

61. Legland D, Arganda-Carreras I, Andrey P. 2016 MorphoLibJ: integrated library

and plugins for mathematical morphology with ImageJ. *Bioinformatics* **32**, 3532–

3534. (doi:10.1093/bioinformatics/btw413)

62. Longair MH, Baker DA, Armstrong JD. 2011 Simple neurite tracer: Open source

software for reconstruction, visualization and analysis of neuronal processes.

Bioinformatics **27**, 2453–2454. (doi:10.1093/bioinformatics/btr390)

63. Teng P. 2011 Extract Slice from Volume.

64.   D'Errico J. 2012 arclength.

65.   R Development Core Team. 2015 *R: A Language and Environment for Statistical*

Computing. R Foundation for Statistical Computing, Vienna, Austria. See

https://www.r-project.org/.

66.   Bates DM, Machler M, Bolker BM, Walker SC. 2014 Fitting Linear Mixed-Effects

Models using lme4. *J. Stat. Softw.* **67**, 1--48. (doi:10.18637/jss.v067.i01)

67.   Kuznetsova A, Brockhoff PB, Christensen RHB. 2017 lmerTest Package: Tests in

Linear Mixed Effects Models. *J. Stat. Softw.* **82**, 1–26. (doi:10.18637/jss.v082.i13)

68.   Birkhead TR, Pellatt JE, Hunter FM. 1990 Numbers and Distribution of Sperm in

the Uterovaginal Sperm Storage Tubules of the Zebra Finch. *Condor* **92**, 508–516.

(doi:10.2307/1368249)

69.   Mero KN, Ogasawara FX. 1970 Dimensions of uterovaginal sperm-storage tubules

of the chicken and their possible significance in sperm release. *Poult. Sci.* **49**,

1304–1308. (doi:10.3382/ps.0491304)

70.   Burke WH. 1968 Release of spermatozoa from storage sites in the hen's oviduct.

Univ. of California, Davis.

71.   Gilbert AB, Lake PE. 1963 Terminal Innervation of the Uterus and Vagina of the

Domestic Hen. *Reproduction* **5**, 41–NP. (doi:10.1530/jrf.0.0050041)

72.   Birkhead TR, Hunter FM. 1990 Numbers of sperm-storage tubules in the Zebra

Finch (Poephila guttata) and Bengalese Finch (Lonchura striata). *Auk* **107**, 193–

197.

73.   Druart X *et al.* 2009 In vivo imaging of in situ motility of fresh and liquid stored

ram spermatozoa in the ewe genital tract. *Reproduction* **138**, 45–53. (doi:Doi

10.1530/Rfp-09-0108)

74.   DaCosta RS. 2005 Autofluorescence characterisation of isolated whole crypts and

primary cultured human epithelial cells from normal, hyperplastic, and

adenomatous colonic mucosa. *J. Clin. Pathol.* **58**, 766–774.

(doi:10.1136/jcp.2004.023804)

75.　Brookner CK, Follen M, Boiko I, Galvan J, Thomsen S, Malpica A, Suzuki S, Lotan R,

Richards-Kortum R. 2000 Autofluorescence Patterns in Short-term Cultures of

Normal Cervical Tissue. *Photochem. Photobiol.* **71**, 730–736. (doi:10.1562/0031-

8655(2000)071<0730:apistc>2.0.co;2)

76.　Schuppin GT, Van Krey HP, Denbow DM, Bakst MR, Meyer GB. 1984

Ultrastructural analyses of uterovaginal sperm storage glands in fertile and

infertile turkey breeder hens. *Poult Sci* **63**, 1872–1882.

(doi:10.3382/ps.0631872)

77.　Burke WH, Ogasawara FX, Fuqua CL. 1972 A study of the ultrastructure of the

uterovaginal sperm-storage glands of the hen, Gallus domesticus, in relation to a

mechanism for the release of spermatozoa. *J. Reprod. Fertil.* **29**, 29–36.

(doi:10.1530/jrf.0.0290029)

78.　Chiba A, Nakamura M. 2001 Microscopic structure of the sperm storage tubules in

the polygynandrous alpine accentor, Prunella collaris (Aves). *Acta Zool.* **82**, 299–

306. (doi:10.1046/j.1463-6395.2001.00091.x)

79.　Renden JA, May EB, Benoff FH. 1981 Histochemistry of uterovaginal sperm-host

glands in Japanese Quail (Coturnix coturnix japonica) with reference to the period

of oviposition. *Poult. Sci.* **60**, 2529–2535. (doi:10.3382/ps.0602529)

80.　Gilbert AB, Reynolds ME, Lorenz FW. 1968 Distribution of spermatozoa in the

oviduct and fertility in domestic birds. 8. The effect of a foreign object in the

uterus on secretions of the sperm-host glands and the survival of spermatozoa in

the oviduct of the domestic hen. *J. Reprod. Fertil.* **17**, 311–4.

(doi:10.1530/jrf.0.0160433)

81.   Pal D. 1977 Histochemistry of the utero-vaginal junction with special reference to

the sperm-host glands in the oviduct of the domestic duck. *Folia Histochem.*

Cytochem. (Krakow). **15**, 235–42.

82.   Heryanto B, Yoshimura Y, Tamura T, Okamoto T. 1997 Involvement of Apoptosis

and Lysosomal Hydrolase Activity in the Oviducal Regression During Induced

Molting in Chickens : A Cytochemical Study for End Labeling of Fragmented DNA

and Acid Phosphatase Treatment of Birds and Collection of Samples. *Poult. Sci.* **76**,

67–72. (doi:10.1093/ps/76.1.67)