Prediction of environmental response in field-grown rice using

expression-dynamics-QTL

Makoto Kashima[1], Ryota L. Sakamoto[2], Hiroki Saito[3,4], Satoshi Ohkubo[3,5], Ayumi

Tezuka[1], Ayumi Deguchi[1], Yoichi Hashida[1], Yuko Kurita[1] & Atsushi J. Nagano[6]*

[1] Research Institute for Food and Agriculture, Ryukoku University, Shiga Japan

[2] Seibi Senior High School, Gifu, Japan

[3] Graduate school of Agriculture, Kyoto University, Kyoto, Japan

[4] Tropical Agriculture Research Front, Japan International Research Center for

Agricultural Sciences, Okinawa, Japan

[5] Institute of Global Innovation Research, Tokyo University of Agriculture and

Technology, Fuchu, Tokyo, Japan

[6] Faculty of Agriculture, Ryukoku University, Shiga, Japan

*Corresponding author: Atsushi J. Nagano (email: anagano@agr.ryukoku.ac.jp)

Organisms respond to the fluctuations in field environments differently throughout their lives depending on their genetic variations. However, how such genetic variations affect environmental responses of organisms under field conditions, that is, the genotype × environment interaction (G×E), remains unclear. In the present study, we developed a method to identify the loci influencing gene expression dynamics (GED) in field-grown rice. We cultivated chromosome segment substitution lines (CSSLs) from Koshihikari and Takanari rice varieties[1] in the field, and conducted approximately 900 time-series RNA sequencing (RNA-Seq) (bihourly-sampling for 16 d during 4 months). This novel analytical approach identified more than one thousand expression dynamics quantitative trait loci (edQTLs), which cause differences in GED between Koshihikari and Takanari. Based on the edQTLs, we successfully predicted the GED under environments different from training environments and in rice cultivars with more complex genotypes than the CSSLs. Results demonstrated that integrating genotype and environment information via the transcriptome can help predict the environmental response of GED under field conditions. We anticipate that edQTLs identification will help understand the G×E of organisms under natural conditions, which is difficult to assess with laboratory equipments[2]. As genetic variation and

environmental conditions converge on transcriptome and transcriptome-based trait prediction functions effectively[3,4], the prediction of GED based on genetic variations and environmental information will contribute to crop breeding by increasing the accuracy of trait prediction under diverse conditions.

To understand the responses of different plant lines under various conditions, it is essential to elucidate G×E. Although the expression quantitative trait locus (eQTL) approach is useful to assess the linkage between genetic variation and transcriptome, it requires transcriptome data for the complete set of a QTL mapping population under given conditions[5–9]. Due to high cost and labor, this approach can cover G×E under limited conditions. Contrarily, mathematical models have succeeded in predicting GED under fluctuating field conditions based on meteorological information[10–12], although only predicting transcriptome dynamics in genotypes used as training data[10–12]. Thus, no study has clarified the relationship between GED and G×E.

Rice (*Oryza sativa* L.) is a major food crop[13–15]. The CSSLs and backcross inbred lines (BILs) between Takanari (a high-yield indica variety)[1,16] and Koshihikari (a leading variety in Japan) are available[1,17], and useful for examining G×E. To understand the broad range of G×E in field-grown rice (Fig. 1a), we conducted mathematical analysis of time-series RNA-Seq of Koshihikari, Takanari, and their reciprocal CSSLs

under field conditions to identify edQTLs and predict transcriptome dynamics under

field conditions. Our approach consisted of four steps (See Methods and Fig. 1b). We

first cultivated 78 reciprocal CSSLs between Koshihikari and Takanari[1], in which a part

of a chromosome was partially substituted with the other parental chromosome, and the

two parental lines (Fig. 1b). Each substituted fragment was categorized into Koshihikari

and Takanari haplotypes using 141 DNA markers (Fig. 1b). To consider the effects of

developmental stage, we transplanted seedlings on four different days (Fig. 1c), and 16

sets of bihourly sampling for 24 h were conducted from May to September 2015

(cropping season in Japan). Two genotypes were collected from each transplant set per

sampling time point along corresponding local meteorological data (air temperature and

global solar radiation) (Fig. 1c). The youngest fully expanded leaf from individual

plants was used for RNA-Seq, and transcriptome data was obtained for 926 samples.

After filtering samples and genes (Extended Data Fig. 1a, b) and confirming genotypes

(Extended Data Fig. 1c), 23,924 expressed genes (EGs) from 854 samples were used for

edQTL detection. Correlation analysis of the transcriptome data showed an obvious

transcriptome-wide diurnal variation in EGs, albeit genotype variation (Fig. 1d).

Second, because lines were not sampled simultaneously, it was difficult to directly

compare expression between genotypes at each time point, and we therefore constructed

predictive models for GED in Koshihikari and Takanari. As most of the genome in the

CSSLs is not substituted, GED of most genes is expected to be the same as in a

background genotype. Thus, the RNA-Seq data of Koshihikari- and

Takanari-background (BG) samples were used to construct BG models in which GED of

Koshihikari and Takanari could be predicted based on meteorological and circadian

clock data (Extended Data Fig. 2a). In addition, "scaled age" was applied to adjust

differences in heading date among rice genotypes (Extended Data Fig. 2). A comparison

of the prediction by the two BG models showed expression dynamics polymorphism

(EDP) in 3696 genes (15.4% of the EGs; Extended Data Fig. 1e). For example, the

predicted GED of *Os09g0343200* differed between the two BG models. Notably, in

some samples, *Os09g0343200* expression appeared to be similar to the prediction of the

other BG model, rather than their own BG model (Fig. 2a), suggesting that chromosome

substitution affects *Os09g0343200* expression in such samples.

Third, to detect edQTLs, we then evaluated the improvement in the residual error of

each gene by assuming the existence of edQTLs around each marker (Extended Data

Fig. 3a). In *Os09g0343200*, the sum of residual errors (SRE) was only significantly

improved assuming edQTLs on chromosome 9 (Fig. 2b and Extended Data Fig. 3b, c),

indicating that *Os09g0343200* edQTLs are cis-edQTLs. As expected, edQTL-based

prediction of *Os09g0343200* was better than BG-based prediction (Fig. 2c). Overall,

edQTLs were identified for 1675 genes, including 222 genes affected by *trans*-edQTLs

(Fig. 2d) and 43 genes affected by multiple edQTLs. For 33 genes, the SRE of GED

based on all edQTLs was smaller than that based on the most significant edQTL only

(Extended Data Fig. 4), suggesting accuracy is higher when considering all edQTLs.

To assess if our approach could detect the edQTLs affecting GED at a specific time

of day and developmental stage, we calculated the variation in median expression level

of each gene with time and scaled age (Extended Data Fig. 5a). The edQTLs affected

393 fluctuating genes depending on the time of day (standard deviation of median

expression at each hour in either BG model was > 1) and 275 genes depending on

scaled age (standard deviation of median expression in every 10 scaled age in either BG

model was > 1) (Extended Data Fig. 5). For example, the GED of *Os01g0537250*

depended on time in both BG models but on scaled age only in the Koshihikari-BG

model, which was caused by the *cis*-edQTL (Fig. 2e, f and Extended Data Fig. 5, 6).

The expression level of *Os01g0537250* in the Koshihikari-BG model differed from that

in the Takanari-BG model during the early developmental stages, but not during the late

stage (Fig. 2e). Thus, the edQTL model can only explain the difference in GED under

limited conditions.

Fourth, we then evaluated the predictive performance of the edQTL model in

different environments from where training data were obtained for model construction.

Koshihikari, Takanari, and the CSSLs of two different transplant sets were cultivated in

a field different from that used in 2015 and sampled in August 2016; 139 RNA-Seq

datasets were obtained. Environmental factors differed between the 2015 and 2016

fields, especially on the third sampling date (Fig. 3a). The prediction of 1537 genes

among the 1675 genes influenced by edQTL was improved by the edQTL model

compared with BG models (Fig. 3b, c and Extended Data Fig. 7). Furthermore, the

prediction of EGs for the validation dataset showed comparable accuracy to that for the

training data set (Extended Data Fig. 8). Thus, our edQTL-based approach can predict

GED in different years and locations.

Finally, GED was predicted for more complex genotypes than CSSLs. For this

purpose, two BILs between Koshihikari and Takanari (HP-a and HP-b)[18] were

cultivated and sampled from June to September 2015, and five and six RNA-Seq

datasets were obtained for HP-a and HP-b, respectively. Although HP-a and HP-b are

Takanari-BG lines, 16.3% and 19.9% of their markers were Koshihikari alleles (Fig. 4a),

respectively. As the genotype of edQTL for *Os12g0406000* presented the Koshihikari

allele in the genome of HP-a and the Takanari allele in the genome of HP-b (Fig. 4a),

Koshihikari- and Takanari-BG models were used to predict *Os12g0406000* expression

in HP-a and HP-b edQTL models, respectively. The GED of *Os12g0406000* fluctuated

with time and it was higher in Koshihikari than in Takanari (Extended Data Fig. 9a).

The edQTL-based prediction of *Os12g0406000* expression was better than the

BG-based prediction (Fig. 4b). Regarding *OsKS3* (*Os04g0611700*)[19] in HP-b, the

edQTL-based prediction was worse than the BG-based prediction (Extended Data Fig.

9b, c). Probably, true edQTLs do not exist around the substituted DNA markers in HP-b

(Extended Data Fig. 9b). Because, in CSSLs, several flanking markers are linked to the

substituted regions of genome, with respect to some genes, the resolution of edQTL

detection in CSSLs was not sufficient for BILs. To evaluate the predictive performance

of the edQTL model, we calculated the sum of prediction errors of all

edQTL-influenced genes using edQTL, Koshihikari-BG, and Takanari-BG models.

Overall, edQTL-based prediction was best (Fig. 4c, Extended Data Fig. 10).

Permutation analysis of markers in HP-a and HP-b genomes showed the significant

advantage of edQTL-based prediction ($p < 0.001$) (Fig. 4c, Extended Data Fig. 10).

Thus, we succeeded in predicting environmental response by integrating genotype and

environment information.

There is a complementary relationship between edQTL and eQTL approaches

regarding sensitivity and environmental conditions (Fig. 1a). Although eQTLs in all

samples showed the same influence on gene expression at a sampling time point, some

edQTLs showed different influences depending on their environments. Generally,

trans-QTLs are involved in regulating sets of genes under a specific environment and/or

at a specific developmental stage. Therefore, the eQTL approach might be more

sensitive for *trans*-QTL than the edQTL approach. This is reflected by the observed

differences in the ratios of *trans*-QTLs to all QTLs (13.3% for edQTL approach; 62–

71% for eQTL approach[6,8,9]).

Phenotypic plasticity via G×E has key roles in the adaptation of plants to field

environment[20,21]. Nevertheless, due to fluctuating environmental factors, the genetic

architecture underlying G×E and phenotypic plasticity under field conditions remains

largely unexplored. By identifying thousands of edQTLs, our novel method provides a

comprehensive picture of the genetic architecture underlying G×E in Koshihikari and

Takanari in field. In addition, our method demonstrated that rice GED can be predicted

based on genotypic and meteorological data. In crop breeding, genetic variations in

response to environment result in unexpected performance of a bred crop under

environments differing from experimental fields. Considering that transcriptome-based

trait prediction functioned effectively in recent studies[3,4], our approach will contribute

for crop breeding by actualizing transcriptome prediction under various conditions.

METHODS

Overview of our approach to identify edQTLs and predict GED. Our approach

consisted of four steps (Fig. 1b). First, rice lines were cultivated in a paddy field and

sparsely sampled at several time points (environments) for RNA-Seq. Second,

predictive models were constructed for GED of the parental lines based on

environmental information (BG-based prediction). Third, the dependency of EDP on

genetic variation was statistically evaluated based on the comparison between RNA-Seq

data and prediction results (edQTL detection). Fourth, by integrating BG-based

prediction and edQTL information, GED was predicted based on environmental

information and genome-wide genotype data (edQTL-based prediction).

Biological samples. The japonica variety Koshihikari, the indica variety Takanari, and

78 CSSLs (40 Koshihikari BG lines, except for SL1213 plus 38 Takanari BG lines,

except for SL1306) were used[1]. Each variety and line was sown in nursery trays. After

approximately one month from sowing, seedlings were transplanted to a paddy field at

Takatsuki, Osaka, Japan (34°51′19″N, 135°37′51″E) in 2015 and at Kizugawa, Kyoto,

Japan (34°4′05″N, 135°50′33″E) in 2016. To integrate the effects of developmental

stages (age) into the prediction of GED with the R package 'FIT'[10], four transplant sets

of the 80 genotypes were prepared (Fig. 1c). Seed sowing date, transplanting date, and

sampling time for each sample are listed in

https://github.com/naganolab/Rice_edQTL-analysis_and_edQTL-based-prediction/Os2

015/Os2015_input/SampleAttribute_2015_Takatsuki.txt and

https://github.com/naganolab/Rice_edQTL-analysis_and_edQTL-based-prediction/Os2

016/Os2016_input/SampleAttribute_2016_Kizu.txt. Briefly, seed sowing and

transplanting were conducted according to the following schedules (Fig. 1c). In 2015,

Transplant set 1: Seed sowing date "2015-04-03", Transplantation date "2015-05-01";

Transplant set 2: Seed sowing date "2015-04-17", Transplantation date "2015-05-08";

Transplant set 3: Seed sowing date "2015-05-01", Transplantation date "2015-05-22";

Transplant set 4: Seed sowing date "2015-05-15", Transplantation date "2015-06-05". In

2016, Transplant set 1: Seed sowing date "2016-04-21", Transplantation date

"2016-05-16"; Transplant set 2: Seed sowing date "2016-05-19", Transplantation date

"2016-06-08".

Sampling and RNA extraction. Sixteen sets (2015) and three sets (2016) of bihourly

sampling for 24 h were conducted on the following dates: 5–6 May, 16–17 June, 22–23

June, 29–30 June, 6–7 July, 13–14 July, 20–21 July, 27–28 July, 3–4 August, 19–20

August, 24–25 August, 31 August to 1 September, 7–8 September, 14–15 September,

21–22 September, and 28–29 September in 2015; and 4–5, 11–12, and 25–26 August in

2016. Two plants from the 80 genotypes of each transplant set were randomly selected

at each sampling point (Fig. 1c). The youngest fully expanded leaf from each plant was

collected, immediately frozen in liquid nitrogen, and stored at -80 °C until RNA

isolation for RNA-Seq. Individual plants were only sampled once. Thus, RNA-Seq data

derived from independent plants. Frozen samples were homogenized with TissueLyser

II (Qiagen, Hilden, Germany), and total RNA was then extracted using the Maxwell 16

LEV Plant RNA Kit (Promega, Tokyo, Japan) and Maxwell 16 Automated Purification

System (Promega, Tokyo, Japan). The concentration of RNA was measured using the

Quant-iT RNA Assay Kit (Thermo Fisher Scientific, Waltham, MA, USA).

RNA-Seq library preparation and sequencing. One microgram of total RNA was

used to prepare each RNA-Seq library. An automated liquid handling system Freedom

EVO 150 (TECAN, Zurich, Switzerland) and a thermal cycler ODTC 384 (INHECO,

Martinsried, Germany) were utilized to prepare the RNA-Seq library of 384 samples

simultaneously. The protocol of RNA-Seq library preparation is described below.

Aliquots of 252 and 192 kinds of 100-μM 60-mer antisense DNA (IDT, San Jose, CA,

USA) covering rRNA and the top eight most frequently-detected transcripts

(*Osp1g00110.1*, *Osp1g00180.1*, *Osp1g00420.1*, *Osp1g00170.1*, *Osp1g00340.1*,

Osp1g00330.1, *Osp1g00600.1*, and *Os11t0707000-02*) in RNA-Seq of *O. sativa* were

mixed in equal amounts (SDRNA oligo pool)[22,23]. One microgram of total RNA, 1.5 μL

of the SDRNA oligo pool, 3 μL of 5× hybridization buffer (0.5 M Tris-HCl (pH 7.4)

and 1 M NaCl), and RNase-free water were mixed to a total volume of 10 μL. The

mixture was incubated to anneal the antisense DNA with the target transcripts using the

thermal cycler ODTC 384 with the following program: 95 °C for 2 min, slow cooled to

45 °C (0.1 °C/s), followed by 45 °C for 5 min. Subsequently, 1 μL of Tli Thermostable

RNase H (TaKaRa, Kusatsu, Japan), 2 μL of 10× RNase H digestion buffer [500 mM

Tris-HCl (pH 7.4), 1 M NaCl, and 200 mM MgCl], and 7 μL of RNase-free water were

added and mixed well. This mixture was incubated at 45 °C for 30 min to selectively

digest the RNA of the DNA/RNA hybrid double strand. Then, 24 μL of AMpure XP

beads (Beckman Coulter, Brea, CA, USA) was added and the mixture was purified in a

384-well magnetic plate following the manufacturer's instructions. The RNA was then

eluted with 14 μL of RNase-free water. Ten microliters of RNA-containing water, 1 μL

of DNaseI (5 unit/μL) (Promega, Tokyo, Japan), 1 μL of 10× DNaseI buffer (Promega,

Tokyo, Japan), 1 μL of 100 mM DTT (Invitrogen, Carlsbad, CA, USA), and 7 μL of

RNase-free water were mixed well. This mixture was incubated at 37 °C for 30 min to

degrade the SDRNA oligo pool. Subsequently, 24 μL of AMpure XP beads (Beckman

Coulter, Brea, CA, USA) was added and purified according to the manufacturer's

instructions. The RNA was eluted with 14 μL of RNase-free water. Five microliters of

the purified mRNA obtained was mixed with 4 μL of 5× SS buffer (Invitrogen,

Carlsbad, CA, USA), and 1 μL of 100 mM DTT (Invitrogen, Carlsbad, CA, USA).

Fragmentation of mRNA was carried out at 94 °C for 4.5 min and immediately cooled

on ice. Subsequently, 0.6 μL of 100 μM random primer $(N)_6$ (Promega, Tokyo, Japan)

and 0.9 μL of RNase-free water were added to the mixture, which was incubated at

50 °C for 5 min and immediately chilled on ice to relax the secondary structure of the

RNA. The fragmented RNA with random hexamer and the reverse transcription master

mix [1 μL of 100 mM DTT, 0.4 μL of dNTPs (25 mM each; Promega, Tokyo, Japan),

0.1 μL of SuperScript (Invitrogen, Carlsbad, CA, USA), 0.2 μL of actinomycin D (1000

ng/μL) (Nacalai Tesque, Kyoto, Japan), and 5.9 μL of RNase-free water] were mixed.

For the reverse transcription step, the mixture was incubated at 25 °C for 10 min,

followed by 50 min at 50 °C. SuperScript was inactivated by heating the mixture at

75 °C for 15 min. Subsequently, 24 μL of AMPure XP (Beckman Coulter, Brea, CA,

USA) and 12 μL of 99.5% ethanol were added, and the mixture was purified according

to the manufacturer's protocol. The reverse transcription product was eluted with 14 μL

of RNase-free water. The purified DNA/RNA hybrid solution without beads and the

second strand synthesis master mix [2 μL of 10× Blue Buffer (Enzymatics, Beverly,

MA, USA), 1 μL of dUTP/NTP mix (Fermentas, Burlington, Canada), 0.5 μL of 100

mM DTT, 0.5 μL of RNase H (Enzymatics, Beverly, MA, USA), 1 μL of DNA

polymerase I (Enzymatics, Beverly, MA, USA), and 5 μL of RNase-free water] were

mixed. This mixture was incubated at 16 °C for 4 h, and then purified with 24 μL of

AMPure XP (Beckman Coulter, Brea, CA, USA) according the manufacturer's protocol.

The purified dsDNA was eluted with 14 μL of RNase-free water. Five microliters of the

dsDNA solution was used in the following step. End-repair, A-tailing, and adapter

ligation were carried out using the KAPA Hyper prep kit (KAPA BIOSYSTEMS,

Wilmington, MA, USA) with 1/10× volume of the solutions according the

manufacturer's protocol. One microliter of 0.1 μM Y-shape adapter[22] was used in the

adapter ligation step for 15 min. Then, size selection of the ligation product was

performed with 5.5 μL of AMPure XP (Beckman Coulter, Brea, CA, USA). The

purified dsDNA was eluted with 10 μL of RNase-free water and a second round of size

selection was performed with 10 μL of AMPure XP (Beckman Coulter, Brea, CA,

USA). The size-selected ligation product was eluted with 15 μL of 10 mM Tris-HCl

(pH 8.0). One microliter of uracil DNA glycosylase (UDG; Enzymatics, Beverly, MA,

USA) was added to the size-selected ligation product. The mixture was incubated at

37 °C for 30 min to exclude the second-strand DNA. For library amplification, 2 μL of

the UDG-digested DNA, 1 μL of 2.5 μM index primer[22], 1 μL of 10 μM universal

primer[22], 0.5 μL of RNase-free water, and 5 μL of KAPA HiFi HotStart ReadyMix (2×)

(KAPA BIOSYSTEMS, Wilmington, MA, USA) were mixed. The DNA fragments

with the adapters and an index sequence were amplified using a thermal cycler with the

following program: denaturation at 94 °C for 2 min, 16 cycles at 98 °C for 10 s, 65 °C

for 30 s, and 72 °C for 30 s for replication, and 72 °C for 5 min for the final extension.

Two rounds of size selection were then performed to remove adapter dimers with equal

volume of AMPure XP (Beckman Coulter, Brea, CA, USA) to the library solution. The

purified library was eluted with 10 μL of RNase-free water. The concentration of each

library was measured by qPCR with the KAPA SYBR FAST qPCR Master Mix (2×)

Kit (KAPA BIOSYSTEMS, Wilmington, MA, USA) and LightCycler 480 II (Roche

Diagnostics, Basel, Switzerland) to mix equal amounts of libraries. One microliter of

the mixed library was used for electrophoresis using a Bioanalyzer 2100 with Agilent

High Sensitivity DNA kit (Agilent Technologies, Santa Clara, CA, USA) to check for

quality. Sequencing of 50-bp single-ends using HiSeq 2500 (Illumina, San Diego, CA,

USA) was carried out by Macrogen Co (Seoul, South Korea).

Calculation and normalization of RNA-Seq count data. All reads obtained were

trimmed with Trimmomatic (version 0.3.3)[24] using the following parameters:

TOPHRED33 ILLUMINACLIP: TruSeq3-SE.fa:2:30:10 LEADING:19 TRAILING:19

SLIDINGWINDOW:30:20 AVGQUAL:20 MINLEN:40. This procedure allowed

obtaining reads with length > 39 nucleotides and average quality score > 19. The

trimmed reads were then mapped to the reference sequences of

IRGSP-1.0_transcript[13,14], rRNAs, and transcripts coded in the mitochondria and

chloroplast genomes (NC_001320.1, NC_011033.1) with RSEM (version 1.3.0)[25], using

Bowtie (version 1.1.2)[26] with default parameters. Expected counts of each gene in the

RSEM outputs were used with R (version 3.4.2)[27] in the following analysis. According

to a previous study[28], we removed the effects of misidentified reads in multiplex

sequencing experiments with HiSeq 2500 (Illumina, San Diego, CA, USA). Of the 926

samples from 2015 and 143 samples from 2016, we used 887 and 139 RNA-Seq

transcriptome datasets from 2015 and 2016, respectively, with more than 10^5 total read

counts for all genes, except for the targets of selective depletion (Extended Data Fig. 1a).

We then filtered out rarely detected genes (number of samples with read count > 0; $\leq$

20% of all samples in the 2015 dataset) from the following analyses to focus on the

23,924 EGs only (Extended Data Fig. 1b).

Identification of Koshihikari and Takanari specific single nucleotide

polymorphisms. The paired-end reads of Koshihikari (SRR1630927) and Takanari

(DRR018360) genomes were trimmed with Trimmomatic (version 0.3.3)[24] using the

parameters stated in the previous section, except for TruSeq3-PE.fa for adapter

trimming. The reads were then mapped to the reference genome (IRGSP-1.0, plastid:

NC_001320.1, mitochondrion: NC_011033.1)[13] using Bowtie2 with option -N 1, which

allows a mismatch in seed alignment (version 2.2.9)[29]. By using SAMtools (version

1.3.1)[30], the reads mapped at multiple loci and the duplicated reads, except for each

representative,   were   discarded.   Picard   (version   2.1.0)

(https://broadinstitute.github.io/picard/) was used to add read group to each SAM file.

All SAM files of each sample were merged and a VCF file was generated using

SAMtools (version 1.3.1)[30] and BCFtools (version 1.1)[30]. Single nucleotide

polymorphisms (SNPs) were filtered using VCFtools (version 0.1.14)[31] with option

--minDP 5, which indicates that all alleles supported by more than four reads are

reported. The VCF file generated was analysed using R (version 3.4.2)[27] and the R

package "vcfR" (version 1.5.0)[32]. We used 15,117 homozygous SNPs distinguishing

Koshihikari and Takanari for genotype confirmation of each sample based on RNA-Seq

data.

Genotype confirmation. To confirm the genotype of each sample, we typed each DNA

marker based on the SNPs obtained by RNA-Seq. First, to type Koshihikari- or

Takanari-specific SNPs in each RNA-Seq sample, the trimmed RNA-Seq reads of each

sample were mapped to the reference genome as described in the previous section using

Bowtie2. The SNPs were filtered using VCFtools (version 0.1.14)[31] with option

--minDP 3, which indicates that all alleles supported by more than two reads are

reported. To type each DNA marker in samples, Koshihikari- and Takanari-type SNPs

were counted from each DNA marker until one of the counts reached "7" or all the

SNPs in 1.5 Mbp from each DNA marker had been counted. In addition, imputation for

typing alleles was performed for SL1201, SL1229, and SL1321 due to the low density

of SNPs around the DNA markers distinguishing genotypes (See

https://github.com/naganolab/Rice_edQTL-analysis_and_edQTL-based-prediction/anal

ysis.R for details). Each DNA marker was categorized into Koshihikari-type,

Takanari-type, or undetermined based on the majority of SNP counts. Each sample was

labeled with the genotype with the most similar DNA marker composition. Thus, 865

samples were confirmed as Koshihikari, Takanari, or CSSLs and 11 samples as BILs in

2015; and 139 samples as Koshihikari, Takanari, or CSSLs in 2016.

Correlation plot of transcriptomes. Pearson's correlations of transcriptome data for

854 samples in 2015 in the time-series order were calculated with the "cor" function of

R, and the heatmap representing such correlations was generated with the "image.plot"

function in the R package "fields" (version 9.0)[33].

Meteorological data. Data on average global solar radiation per 10 min at the Hirakata

Weather Station in 2015 for Takatsuki and Kyotanabe Weather Station in 2016 for

Kizugawa were obtained from the Japan Meteorological Agency. The average air

temperature per minute was then interpolated with the "approxfun" function in R. Data

on average global solar radiation per minute at the Osaka Weather Station in 2015 for

Takatsuki and Nara Weather Station in 2016 for Kizugawa were obtained from the

Japan Meteorological Agency. The time points with global solar radiation < 0.3 kJ m^{-2}

min^{-1} were defined as 'dark condition'.

Conversion of age to scaled age. To consider the difference in heading date among

genotypes, we defined scaled age (SA) based on sampling dates (d), seeding dates (SD),

and genotypes (G). Under the assumption that rice grows lineally before heading at

various speeds among genotypes, flowers at scaled age 85, and ages at the same speed

after flowering, SA (d, SD, G) was calculated using the formula:

$$SA(d, SD, G) = \begin{cases} 85 \times \dfrac{d - SD}{FD(G, SD)} & ((d - SD) \leq FD(G, SD)) \\ 85 + (d - SD) - FD(G, SD) & ((d - SD) > FD(G, SD)) \end{cases}$$

where *FD(G,SD)* is the flowering date of a genotype *G* seeded at seeding date *SD*.

BG-based prediction of GEDs in Koshihikari and Takanari. A predictive model for

GED in Koshihikari and Takanari was constructed with the R package "FIT" (version

0.0.4)[10], based on the meteorological data of Takatsuki in 2015, log2reads per million

(log2rpm) in 2015, and scaled age of each sample. The GED data of Koshihikari- and

Takanari-BG genotypes were used for the Koshihikari and Takanari models,

respectively. Precision weight was calculated according to the voom method[10,34] and

using data from 887 samples in 2015, and then used for FIT. Details on other parameters

are   listed   in

https://github.com/naganolab/Rice_edQTL-analysis_and_edQTL-based-prediction/anal

ysis.R. Further, log2rpm of each gene in each sample was predicted with the constructed

predictive models corresponding to all possible combinations of the meteorological data

and scaled ages of each genotype for the four transplant sets in Takatsuki in 2015 and

two transplant sets in Kizugawa in 2016.

Detection of edQTL. We focused on 3696 genes with EDP (Euclidean distance > 40

between Koshihikari and Takanari based on expression levels at the observed time

points in 2015). According to the genotype of backgrounds or each marker, the SRE

between the observed log2rpm and predicted log2rpm in the Koshihikari- or

Takanari-BG models were calculated (Extended Data Fig. 3a, b). To evaluate the

statistical significance of improvement in SRE, 1000 permutations of samples with

Koshihikari- and Takanari-BG were conducted, and SRE was calculated for each test.

Lognormal distribution was then fitted to the obtained distributions of SREs using the

fitdist function in the R package "fitdistrplus"[35], with the fitting method of quantile

matching of 70% and 1% (Extended Data Fig. 3c). For almost all genes with EDP, the

top 0.1% values in the fitted lognormal distribution were higher than those in the

distributions generated by permutation (Extended Data Fig. 3d), indicating that the

p-value obtained from lognormal distribution is conservative. The p-values for

improvement in the SRE were calculated with the fitted lognormal distribution for each

gene. The adjustment for multiple comparisons with Benjamini-Hochberg method[36] was

performed with the p.adjust function in R. Finally, we defined the markers with the

smallest adjusted p-value (< 0.05) in each peak for each gene as the edQTL affecting

each GED.

edQTL-based prediction of GED. For edQTL-affected genes, the Koshihikari- or

Takanari-BG model was selected based on the genotype of the edQTL(s). For other

genes, the Koshihikari- or Takanari-BG model was selected based on the genotype of

backgrounds.

Statistical test for the accuracy of GED prediction in BILs. The SPEs of all genes

affected by each edQTL were calculated based on BG-based prediction (the

Koshihikari- and Takanari-BG models) and edQTL-based prediction (edQTL model).

To evaluate the statistical significance of improvement in the SPEs, 10,000

permutations were performed for the markers in each BIL genome, and the value

corresponding to 0.1% percentile in each permutation was calculated.

Code availability. The scripts used in this study are available from PRJDB7234.

Data availability. All datasets generated and/or used in this study are available from

PRJDB7234.

1.   Takai, T. et al. Genetic mechanisms underlying yield potential in the rice

high-yielding cultivar Takanari, based on reciprocal chromosome segment

substitution lines. *BMC Plant Biol.* **295**, 14 (2014).

2.   Poorter, H. et al. Pampered inside, pestered outside? Differences and similarities

between plants growing in controlled conditions and in the field. *New Phytol.* **212**,

838–855 (2016).

3.   Kremling, K. A. G. G. et al. Dysregulation of expression correlates with

rare-allele burden and fitness loss in maize. *Nature* **555**, 520–523 (2018).

4.   Horinouchi, T. et al. Prediction of cross-resistance and collateral sensitivity by

gene expression profiles and genomic mutations. *Sci. Rep.* **14009**, 7 (2017).

5.   Horiuchi, Y. et al. Global expression differences and tissue specific expression

differences in rice evolution result in two contrasting types of differentially

expressed genes. *BMC Genomics* **1099**, 16 (2015).

6.   Wang, J. et al. An expression quantitative trait loci-guided co-expression analysis

for constructing regulatory network using a rice recombinant inbred line

population. *J. Exp. Bot.* **65**, 1069–1079 (2014).

7.   Jansen, R. C. & Nap, J. P. Genetical genomics: The added value from segregation.

Trends Genet. **17**, 388–391 (2001).

8.   Wang, J. et al. A global analysis of QTLs for expression variations in rice shoots

at the early seedling stage. *Plant J.* **63**, 1063–1074 (2010).

9.   Kuroha, T. et al. eQTLs regulating transcript variations associated with rapid

internode elongation in deepwater rice. *Front. Plant Sci.* **8**, 1753 (2017).

10.   Iwayama, K., Aisaka, Y., Kutsuna, N. & Nagano, A. J. FIT: Statistical modeling

tool for transcriptome dynamics under fluctuating field conditions.

Bioinformatics **33**, 1672–1680 (2017).

11.   Nagano, A. J. et al. Deciphering and prediction of transcriptome dynamics under

fluctuating field conditions. *Cell* **151**, 1358–1369 (2012).

12.   Matsuzaki, J., Kawahara, Y. & Izawa, T. Punctual transcriptional regulation by

the rice circadian clock under fluctuating field conditions. *Plant Cell* **27**, 633–648

(2015).

13.   Kawahara, Y. et al. Improvement of the *Oryza sativa* Nipponbare reference

genome using next generation sequence and optical map data. *Rice* **6**, 4 (2013).

14.   Sakai, H. et al. Rice annotation project database (RAP-DB): an integrative and

interactive database for rice genomics. *Plant Cell Physiol.* **54**, 1–11 (2018).

15.   Huang, X. et al. A map of rice genome variation reveals the origin of cultivated

rice. *Nature* **490**, 497–501 (2012).

16.   Ookawa, T. et al. Precise estimation of genomic regions controlling lodging

resistance using a set of reciprocal chromosome segment substitution lines in rice.

Sci. Rep. **30572**, 6 (2016).

17.   Adachi, S. et al. Introgression of two chromosome regions for leaf photosynthesis

from an indica rice into the genetic background of a japonica rice. *J. Exp. Bot.* **65**,

2049–2056 (2014).

18.   Adachi, S. et al. The mesophyll anatomy enhancing CO_2 diffusion is a key trait

for improving rice photosynthesis. *J. Exp. Bot.* **64**, 1061–1072 (2013).

19.   Sakamoto, T. An overview of gibberellin metabolism enzyme genes and their

related mutants in rice. *Plant Physiol.* **134**, 1642–1653 (2004).

20.   Sultan, S. E. Phenotypic plasticity and plant adaptation. *Acta Bot. Neerl.* **44**, 363–

383 (1995).

21.   Fournier-Level, A. et al. Predicting the evolutionary dynamics of seasonal

adaptation to novel climates in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. U. S.*

A. **113**, E2812-21 (2016).

Supplementary Information is available in the online version of the paper.

Acknowledgements. We thank the field work support team for help at field sites, Fumie

Kobayashi for assistance with laboratory experiments, and Yasuhiro Sato for valuable

discussions. We thank Toshio Yamamoto (Institute of Crop Science, NARO) for

providing the CSSLs and BILs between Koshihikari and Takanari. This work was

supported by the JST CREST JPMJCR15O2 and JSPS KAKENHI Grants JP16H06171

and JP16H01473 to A.J.N.

Author contributions. M.K. analysed the data. R.L.S., H.S., and S.O. performed field

work. M.K., A.T., A.D., Y.H., and Y.K. conducted RNA-Seq experiments. A.J.N.

designed the study. M.K. and A.J.N. wrote the manuscript with input from all

co-authors.

Declaration of interests. The authors declare no competing interests.

22.   Nagano, A. J., Honjo, M. N., Mihara, M., Sato, M. & Kudoh, H. Detection of

plant viruses in natural environments by using RNA-Seq. *Plant Virol. Protoc.*

Methods Mol. Biol. **1236**, 89–98 (2015).

23.   Morlan, J. D., Qu, K. & Sinicropi, D. V. Selective depletion of rRNA enables

whole transcriptome profiling of archival fixed tissue. *PLoS One* **7**, e42882

(2012).

24.   Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for

Illumina sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

25.   Li, B. & Dewey, C. N. RSEM: accurate transcript quantification from RNA-Seq

data with or without a reference genome. *BMC Bioinformatics* **12**, 323 (2011).

26.   Langmead, B., Trapnell, C., Pop, M. & Salzberg, S. L. Ultrafast and

memory-efficient alignment of short DNA sequences to the human genome.

Genome Biol. **10**, R25 (2009).

27.   R Core Team. R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. (2017).

28.   Kamitani, M., Nagano, A. J., Honjo, M. N. & Kudoh, H. RNA-Seq reveals

virus-virus and virus-plant interactions in nature. *FEMS Microbiol. Ecol.* **92**, 1–

11 (2016).

29.   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie2. *Nat.*

Methods **9**, 357–359 (2012).

30.   Li, H. et al. The sequence alignment / map (SAM) format and SAMtools.

Bioinformatics **25**, 2078–2079 (2009).

31.   Danecek, P. et al. The variant call format and VCFtools. *Bioinformatics* **27**,

2156–2158 (2011).

32.   Knaus, B. J. & Grünwald, N. J. VCFR: a package to manipulate and visualize

variant call format data in R. *Mol. Ecol. Resour.* **17**, 44–53 (2017).

33.   Nychka, D., Furrer, R., Paige, J. & Sain, S. fields: Tools for spatial data. (2017).

doi:10.5065/D6W957CT

34.   Law, C. W., Chen, Y., Shi, W. & Smyth, G. K. VOOM: precision weights unlock

linear model analysis tools for RNA-seq read counts. *Genome Biol.* **15**, R29

(2014).

35.   Delignette-Muller, M. L. & Dutang, C. fitdistrplus : An R package for fitting

distributions. *J. Stat. Softw.* **64**, 1–34 (2015).

36.   Benjamini, Y. & Hochberg, Y. Controlling the false discovery rate : a practical

and powerful approach to multiple testing. *J. R. Stat. Soc. Ser. B* **57**, 289–300

(1995).

Figures

Figure 1 | Concept and workflow of expression dynamics quantitative trait loci

(edQTLs) detection. a, Conceptual differences between expression quantitative trait

locus (eQTL)- and edQTL-based prediction of genotype × environment interaction

(G×E). **b**, Workflow of edQTL detection using chromosome segment substitution lines

(CSSLs) and prediction of expression dynamics. **c**, Summary of the sampling design for

edQTL detection. The left panels show plots of meteorological data [air temperature

(Temp., °C) and global solar radiation (Rad., kJ m^{-2} min^{-1})] and sampling time points in

Takatsuki from May to September in 2015. Vertical red lines represent the sampling

time points. **d**, Pearson's correlations among the 854 samples used for constructing the

predictive model based on expression data of 23,294 expressed genes. White lines

indicate the border of each bihourly sampling set.

Figure 2 | Expression dynamics quantitative trait loci (edQTLs) detection revealed

***cis*- and *trans*-edQTLs are involved in environmental responses. a, c, e,** Prediction

of *Os09g0343200* (**a,c**) and *Os01g0537250* (**e**) expression levels in Takatsuki in 2015

based on environmental information (time, air temperature, and global solar radiation).

━ ━ ━ and ━ ━ ━ indicate predicted expression levels by 'FIT' in Koshihikari and

Takanari in Transplant sets 1, 2, 3, and 4, respectively. ●●●● and ●●●● indicate the

expression level obtained by RNA sequencing for samples in Transplant sets 1, 2, 3, and

4 of individuals with Koshihikari background (BG) and Takanari BG in (**a**) or

Koshihikari- and Takanari-type edQTLs in (**c,e**), respectively. In (**c,e**), strong colors

emphasized by the arrow heads indicate samples harboring edQTLs different from BG,

and light colors indicate the samples harboring edQTLs identical to BG. The upper ▬

indicates dark conditions (global solar radiation < 0.3 kJ m^{-2} min^{-1}). **b,f**, edQTLs

regulating *Os09g0343200* (**b**) and *Os01g0537250* (**f**). **d**, X-axis and Y-axis represent

statistical significance of each marker (FDR = 0.05) and the genes influenced by the

markers in order of position on each chromosome, respectively.

Figure 3 | Expression dynamics quantitative trait locus (edQTL)-based prediction

of gene expression dynamics in different environments. a, Plots of meteorological

data [air temperature (Temp., °C) and global solar radiation (Rad., kJ m^{-2} min^{-1})] on

sampling dates in August. ▬ indicates data in Takatsuki in 2015 and ▬ indicates data in

Kizugawa in 2016. **b,** Examples of prediction of expression dynamics in Kizugawa in

2016 based on environmental information and edQTL in Transplant set 1. Points in

strong colors emphasized by arrowheads indicate samples harboring edQTL different

545   from background (BG) genotypes and light colors indicate samples harboring edQTL

546   identical to BG genotypes. The upper ▬ indicates dark conditions (global solar radiation

547   < 0.3 kJ m-^{2} min^{-1}). **c,** Effects of edQTL over prediction of gene expression dynamics in

548   Kizugawa in 2016. Genes influenced by edQTL are in order of position on the

549   chromosomes along the horizontal axis.

Figure 4 | Expression dynamics quantitative trait locus (edQTL)-based prediction

of gene expression dynamics with more complex genotypes than the chromosome

segment substitution lines (CSSLs). a, edQTL for *Os12g0406000* and genotypes of

HP-a and HP-b. ● indicate significant Koshihikari-type markers. **b,** Prediction of

Os12g0406000 expression in HP-a and HP-b. Strongly colored lines are applied models

for HP-a and HP-b. **c,** Prediction accuracy of edQTL model for HP-a. The blue, red, and

orange vertical lines indicate the sums of prediction errors based on Koshihikari,

Takanari, and edQTL models. The histogram shows the distribution of sums of

prediction errors based on the edQTL model in 10,000 permutations of markers in HP-a

genome. The dashed vertical line indicates the 0.1% percentile of the distribution.