Determinants of *Staphylococcus aureus* PSMα3 Cross-α Fibrils Toxicity

Einav Tayeb-Fligelman[1,§], Nir Salinas[1], Orly Tabachnikov[1], and Meytal Landau[1,*]

Affiliations:

[1] Department of Biology, Technion-Israel Institute of Technology, Haifa 3200003, Israel.

[§] Present affiliation: Departments of Chemistry and Biochemistry and Biological Chemistry, UCLA-DOE Institute, UCLA, Los Angeles, California 90095, USA.

[*] mlandau@technion.ac.il

Abstract

Members of the phenol-soluble modulin (PSM) peptide family secreted by *Staphylococcus aureus* perform various virulence activities, some of which are mediated by the formation of amyloid fibrils of diverse architectures. Specifically, PSMα1 and PSMα4 structure the *S. aureus* biofilm by assembling into robust cross-β amyloid fibrils composed of tightly mated β-sheets. PSMα3, in contrast, the most cytotoxic member of the family, assembles into cross-α amyloid-like fibrils in which α-helices stack perpendicular to the fibril axis into tightly mated sheets. In this work, we have conducted extensive analyses of the structural and toxic properties of PSMα3 derivatives and mutants, along with crystal structure determination of three single-point alanine mutants showing cross-α structural polymorphism. The findings suggest that interactions with the cell membrane and cytotoxicity are regulated by inter- and intra-helical electrostatic interactions within the cross-α fibril. Using confocal and scanning electron microscopy, we demonstrated co-localization of PSMα3 aggregates and membranes, which was

associated with massive cell deformation. Taken together, the process of cross-α fibrillation, along with positive charges, serve as critical determinants of PSMα3 cytotoxicity against human cells. Overall, our results provide mechanistic insight into PSMα3 cytotoxicity, and offer the basis for design of agents to combat staphylococcal virulence.

Introduction

Phenol-soluble modulins (PSMs) comprise a family of virulent peptides secreted by the pathogenic bacterium *Staphylococcus aureus (1)*. PSMs stimulate inflammatory responses, contribute to biofilm formation and are capable of lysing human cells, including erythrocytes and leukocytes, such as T cells *(2,3)*. Several PSM members were shown to form amyloid fibrils, and this configuration has been shown to be essential to their activity *(4–7)*. Amyloids are structured protein aggregates mostly known for their association with systemic and neurodegenerative diseases *(8–10)*, but are also involved in various physiological functions in humans and in microbes *(6,11–19)*. They are known to form highly stable cross-β fibrils composed of tightly mated β-sheets, with β-strands oriented perpendicularly to the fibril axis *(20,21)*. PSMα1 and PSMα4 comprise the robust cross-β fibrils, which confer high stability to the bacterial biofilm *(5–7)*, rendering it a resilient physical barrier against antibiotics and the immune system. However, PSMα3, the most cytotoxic member of the family *(2,22,23)*, forms unique cross-α fibrils *(4)*. These fibrils are composed of amphipathic α-helices oriented perpendicularly to the fibril axis, which stack to form elongated 'sheets', stabilized by inter-helical polar interactions between side chains. The 'helical sheets' mate via an extensive hydrophobic core,

providing stability to the fibril *(4)*. PSMα3 cross-α fibrils exhibit unbranched fibril morphology, bind the amyloid dye-indicator thioflavin T (ThT), show rapid aggregation kinetics, and are highly toxic to human T2 cells *(4)*. Taken together, the secondary-structure polymorphism in amyloid-like structures formed by the homologous PSM family members (PSMα1, PSMα4 and PSMα3) is exceptional within dozens of amyloids studied thus far and is indicative of the specific function of each member.

The crucial role of PSMα3 in aggressive *S. aureus* infections *(22)*, and its unique cross-α fibril architecture *(4)* encouraged us to further explore its structure-function-fibrillation relationships in order to identify the critical determinants in its mode of action. We previously reported that the fibrillation of PSMα3 enhances its cytotoxicity to human cells *(4)*. Using an array of derivatives and mutations, and assessment of their structural and biophysical features, we herein substantiate the critical role of fibrillation and positively charged residues in cytotoxicity. We suggest that cross-α fibrillation regulates intra- and inter-helical electrostatic interactions to modulate activity, and that Lys17 is especially crucial for cytotoxicity, as it provides the only positive charge along the fibril available to interact with cell membrane lipid components. Moreover, we show that PSMα3 aggregates on the plasma membrane, and co-localizes with membrane particles of affected cells. The presented elucidation of structural and functional properties of the highly virulent PSMα3 is expected to provide a template for the development of novel antivirulence agents to combat *Staphylococcal* infections.

Results

Fibrillation and positive charges are key determinants of PSMα3 cytotoxicity

To identify the sequence and structural determinants crucial for PSMa3 fibrillation and activity, several alanine substitutions were generated and tested for their fibril-forming capacities (F3A, L7A, F8A, K9A, F10A, F11A, K12A, D13A, L15A, G16A, and K17A) (Table 1). Among these mutants, only F3A and F8A failed to form fibrils, as visualized via transmission electron microscopy (TEM) (Figure 1A), demonstrating the robust tendency of PSMα3 to fibrillate. The non-fibrillating F3A and F8A mutants showed significantly reduced toxicity towards T2 cells (Figure 1B), while alanine substitutions in other non-polar residues (L7A, F10A, F11A, L15A, and G16A), along with D13A, maintained fibril-formation (Figure 1A) and toxic properties (Figure 1B). Moreover, introduction of a cysteine residue to either termini of PSMα3 showed that while C-cys-PSMα3 formed fibrils with fast kinetics and retained cytotoxicity, N-cys-PSMα3 lost the ability to fibrillate and correspondingly, was not cytotoxic (Figure S1). Furthermore, alanine substitutions of positively charged lysine residues (K9A, K12A, K17A) significantly reduced toxicity (Figure 1B), in accordance with a previous report *(23)*. These lysine to alanine mutants maintained the ability to form fibrils (Figure 1A), indicating the role of positively charged residues in cytotoxicity but not in fibrillation (Table 1).

WT PSMα3, and the K9A, L15A, G16A, and K17A mutants displayed various fibrillation kinetics, as assessed by monitoring ThT fluorescence (Figure S2). However, other mutants (L7A, F10A, F11A, K12A and D13A) failed to bind ThT (Figure S2), in accordance with a previous report *(24)*, despite clear fibril-forming ability visualized by TEM (Figure 1A). These findings emphasize the importance of integrating different

methods to assess fibrillation propensity, and accordingly, point to fibrillation and positive charges as the dominant factors governing cytotoxicity.

Secondary structure properties of PSMα3 mutants in soluble and fibrillar states

The secondary structure properties of PSMα3 mutants were explored in the soluble state and in fibrils. WT PSMα3, as well as most mutants substituting non-polar residues (F3A, L7A, F10A, L15A and G16A), and the all D-enantiomeric amino acid PSMα3 (D-PSMα3), maintained helicity in solution. In contrast, mutants in which a polar residue was substituted to alanine (K9A, K12A, K17A, and D13A), thereby disturbing the amphipathic nature of the peptide, exhibited reduced helicity in solution compared to the WT PSMα3 (Figure 2 and Figure S3). WT PSMα3 and the K17A mutant secondary structures, analyzed using ATR-FTIR (Figure 2), showed predominant α-helical content when dried as a thin film on the ATR module, both on the day of solubilization, and after seven days of incubation in solution. Helicity was even more dominant in the fibrils obtained by washing the fibril pellets after seven days of incubation (Figure 2). In contrast, the K9A, K12A and D13A mutants showed an increase in their β-structure content upon incubation, especially in the fibril pellet (Figure 2).

The X-ray fiber diffraction of WT PSMα3 and of D-PSMα3, displayed the cross-α diffraction pattern, with orthogonal reflection arcs at ~10.5 Å and ~12 Å (Figure 3), corresponding to the spacing between α-helices along each sheet, and the inter-sheet distance, respectively, in the PSMα3 crystal structure (Figure 4). An additional major reflection was present at ~25 Å, which may be attributable to the length of the α-helix, or to the inter-sheet distance over the dry and wet interfaces (Table S1). This pattern was observed in fibrils prepared from freshly dissolved peptide and in fibrils prepared from a

peptide solution incubated for 10 h at room temperature, indicating that the fibrils remained stable in the cross-α form. The X-ray fiber diffraction profile of the K17A mutant showed a very similar cross-α pattern (Figure 3), in accordance with the helicity observed for WT PSMα3 and K17A fibrils in the FTIR spectra (Figure 2). The diffraction pattern of the L7A, K9A, F11A, and L15A mutants indicated additional structural features characteristic of β-rich structures. This was especially pronounced in the moderately toxic L7A mutant which showed additional strong orthogonal reflections at 4.7 Å and 9-12 Å (Figure 3). The diffraction pattern of these four mutants may be indicative of mixed populations of both cross-α and cross-β (Table 1). Other mutants showed only diffuse reflections that could not be interpreted (Figure S4). Taken together, WT PSMα3 consistently maintained its α-helical conformation in all experiments in solution, in the fibril and in the crystal, whereas its mutants demonstrated different degrees of α-helical stability.

Crystal structures of three PSMα3 mutants reveal packing polymorphism in their cross-α architecture

The crystal structure of the K12A, G16A, and L15A PSMα3 mutants demonstrated cross-α fibrils of tightly mated sheets composed of parallel amphipathic α-helices facing each other via their hydrophobic face (Figure 4, Table S2). The structures demonstrated packing polymorphism, namely translations of the mated sheets from a staggered orientation of slightly shifted sheets in the WT PSMα3 and the L15A mutant, to a directly facing orientation in the K12A and G16A mutants. The structures of WT PSMα3 and L15A also displayed a larger inter-sheet distance and a smaller distance between the α-helices along the sheets, in comparison to G16A and K12A (Figure 4 and Table S1).

Electrostatic interactions within the cross-α fibril regulate PSMα3 toxicity

The crystal structures of PSMα3 and its mutants provided the opportunity to explore properties stabilizing the α-helices and their interactions within the cross-α fibrils. Generally, the formation of salt-bridges between positively and negatively charged residues located 3-4 positions apart on the sequence are associated with increased α-helical stability *(3,25)*. In the twisted helical configuration of monomeric PSMα3 observed by NMR in solution *(26)*, such a salt bridge can form between Asp13 and Lys17. However, in the fibrillary crystal structures of PSMα3 and its mutants, demonstrating canonical α-helices, a salt bridge can form between Asp13 and Lys9. In the L15A mutant, an additional intra-helical electrostatic bond formed between Glu2 and Lys6. In PSMα3 and the L15A mutant fibrils, Lys12, from an adjacent α-helix in the same sheet, constitutes a possible second salt-bridge partner for Asp13, which may stabilize the fibril (Figure 5). In the structure of the WT and the K12A and G16A mutants, an inter-helix salt bridge may form along the fibril between Glu2 and Lys6. In all four crystal structures, Lys17 is the only lysine residue fully deprived of potential salt-bridges (Figure 5). Accordingly, the K17A mutation, the least toxic peptide among all tested alanine mutations (Figure 1B), did not affect the electrostatic interactions within the fibril, leading to similar stability of its α-helical configuration in fibrils to that of WT PSMα3 (Figures 2&3). We therefore hypothesized that in the fibril form, Lys17 contributes an exposed and available positive charge that might mediate cytotoxicity via membrane interactions. The D13A mutant eliminates potential electrostatic interactions with lysine residues and possesses an increased net positive charge, rendering it the most cytotoxic derivative (Figure 1B). Overall, our results provide further support of the proposed role of positively charged residues in cell

membrane interactions, and suggest that fibrillation regulates the availability of these charges to interact with the membrane.

PSMα3 aggregates co-localize with cell membranes and lead to cell death

Incubation of PSMα3 with T2 cells elicited PSMα3 binding and aggregation on the cell membranes, as visualized by confocal microscopy using a mixture of WT PSMα3 and a fibrillating fluorescein isothiocyanate (FITC)-labeled PSMα3 derivative (Figure 6 & S5). Scanning electron micrographs demonstrated the massive deformation and rapid cell death occurring within a few minutes of exposure to PSMα3 (Figure 7). The interactions of PSMα3 with the membrane, and resulting membrane rupture and cell death, were also evident by the appearance of co-localization of membrane particles and PSMα3 aggregates around affected cells, as observed by continuous live cell confocal microscopy (Figure 8).

Discussion

S. aureus-secreted PSMα3 is a cytotoxic peptide acting as a functional cross-α fibril *(4)*. In contrast to most typical amyloids that transition from various disordered or ordered conformations into β-rich species upon fibrillation *(20,27,28)*, PSMα3 lacks secondary structure transition and remains helical upon fibrillation (Figures 2-4). We expect that the initial aggregation of these amphipathic peptides involves interactions via their hydrophobic face to form helical bundles, which will eventually elongate to form the extensive hydrophobic core between the α-helical mated sheets. The distinctive secondary structure observed for the PSMα3 cross-α fibrils, particularly when compared to PSMα1 and PSMα4, which are involved in *S. aureus* biofilm structuring, and transition

to form the canonical cross-β amyloid fibrils *(5)*, is indicative of structurally-encoded functional specificity. The role of fibrillation in PSMα3 activities is yet not fully understood, and its impact on cytotoxicity, in particular, has been under debate *(24,29)*. The results presented here demonstrate a critical role of fibrillation in cytotoxicity, and provide structural and mechanistic insight into the involvement of inter- and intra-helical polar interactions within the fibril in regulating toxicity.

Fibrillation is a robust feature of PSMα3, as shown by preserved fibrillation capacities in nine of the eleven single point alanine mutations tested (Figure 1A). F3A and F8A, the two mutants that failed to form fibrils, were also the least toxic, with the exception of those containing substitutions for lysine residues. In addition, while the C-cys-PSMα3 derivative formed fibrils and was cytotoxic, the N-cys-PSMα3 did not form fibrils and was not cytotoxic (Figure S1). This radical difference between peptides with the same amino-acid composition endorses the central role of fibrillation in PSMα3 toxic activity. Furthermore, the deleterious effect of F3A, F8A and N-cys-PSMα3 on fibrillation and toxicity, and the structural changes observed for fibrils of the moderately toxic L7A mutant (Figure 3), points to the N-terminal part as a critical region in cross-α fibrillation.

The presented observations suggest that membrane disruption and cytotoxicity of PSMα3 are regulated by fibrillation-dependent intra- and inter-helix electrostatic interactions. The complete lack of cytotoxicity of the K17A mutant, in contrast to other mutants in lysine residues that retained some cytotoxic activity (Figure 1B), corresponds with the absence of potential electrostatic interactions involving Lys17 within the fibril. This putatively allows the free positive lysine charge to interact with the cell membrane. Interactions between PSMα3 and membrane lipids leads to toxicity, as shown with the

cross-α forming, all-D-amino-acid PSMα3 (D-PSMα3), which exhibited a similar level of cytotoxicity as WT PSMα3 (Figures 1&3), as was also recently reported by others *(29)*. This lack of enantiomer stereo-specificity in PSMα3 toxicity rules out direct interactions with membrane proteins, which are chiral in contrast to the achiral lipids. Aside from the lipids, additional membrane components, particularly in lipid rafts, such as cholesterol, glycoproteins and glycolipids, may also be involved in such peptide-membrane interactions *(3,28,30–32)*.

The activity of PSMα3 involves reciprocal interactions with cell membranes *(33)*. Specifically, eukaryotic membrane mimetics accelerate PSMα3 fibrillation, and the fibrils, in turn, affect the membranes and penetrate deep into the lipid bilayers *(33)*. This corresponds to other membrane-deforming peptides whose aggregation is often induced, or accelerated, by the membrane itself *(34–36)*. An earlier work on pathological amyloids suggested a process of fibril formation on membrane surfaces *(37,38)*. For example, α-synuclein fibrils were shown to interact with the plasma membranes, and act as seeds, where the addition of monomers allows continuous aggregation within the membrane and increased toxicity *(38)*. Co-aggregation of toxic amyloids with lipids was also supported by in-vivo studies of several human pathological diseases *(39,40)*. Such co-aggregation may constitute a universal mechanism of toxicity, in both pathological *(41)* and physiological states, ensuring formation of functional toxic amyloids. This corresponds with our observation that PSMα3 aggregates co-localized with the membranes of affected cells (Figures 6&8). This putative dynamic co-aggregation process would require a mixed population of soluble and fibrillar PSMα3, whereas stabilized mature fibrils lacking equilibrium with soluble species would be less toxic. This might explain the reduced

toxicity of an incubated racemic PSMα3 mixture, which showed robust fibrillation compared to the highly toxic L- and D-PSMα3 enantiomers *(29)*. Nevertheless, the reduced toxicity of the racemic mixture *(29)* could also result from differences in the packing of the hydrophobic core and in the electrostatic interactions along the fibril.

While the helical conformation of the PSMα3 fibrils is stable over extended incubation times, several mutants exhibited secondary structure transitions (Figures 2-3). This was particularly evident for L7A, which formed a mixed population of both α- and β-structures. This residue is indeed central in forming the inter-sheet hydrophobic core observed in the cross-α structure (Figure 4), and its substitution to alanine likely destabilized the cross-α fibril. Overall, several single-point PSMα3 mutants (Figures 2-3), as well as truncations showing antibacterial activity *(5)*, identified the PSMα3 sequence as that which dictates the ability to form β-rich structures. It is nevertheless important to note that interactions with cell membranes can stabilize the helical configuration of PSMα3 *(33)*, and thus, the structural instability observed for the mutants in-vitro might be less significant upon peptide contact with human cells.

The structural variations between PSMα3 mutants might affect the binding of the amyloid indicator dye ThT, which can explain the lack of ThT fluorescence for many of the mutants (L7A, F10A, F11A, K12A and D13A) (Figure S2), despite clear fibril-forming ability visualized by TEM (Figure 1A). Failure to bind ThT was previously reported for amyloid fibrils *(42,43)*, and was attributed to alteration in cavities along the fibril *(44,45)*. Hence, this assay is not sufficient to rule out fibril formation. The absence of ThT binding to PSMα3 mutants indeed led to the misinterpretation that these mutants do not form fibrils, and that toxicity is independent of fibrillation *(24)*. The results presented here

further stress the importance of integration of methods in the evaluation of amyloid properties.

Toxicity of pathological amyloids is often attributed to the prefibrillar and soluble oligomeric conformations, while the mature β-rich fibrils detoxify the amyloid *(46)*. Our study suggests that cross-α fibrillation of PSMα3 enhances toxicity, which may be related to the retained helical nature of its different species. Correspondingly, several eukaryotic pathological amyloids contain α-helices in their prefibrillar intermediate states (e.g., *(27)*), including in initial stages of interactions with lipid membrane surfaces *(36)*. Overall, our work elucidated the key determinants governing PSMα3 cytotoxicity, namely positive charges and cross-α fibrillation that regulates electrostatic interactions and leads to membrane rupturing. This provides a template for designing novel therapeutics against PSMα3 cytotoxicity to reduce virulence and combat resistant infections.

Figure 1. Fibril formation of PSMα3 derivatives and their toxicity against T2 cells

A. Fibril formation of PSMα3 mutants as visualized by transmission electron microscopy (TEM). WT PSMα3 and derivatives demonstrating the highest cytotoxicity levels, including D-PSMα3, L7A, F10A, F11A, D13A, L15A and G16A, form fibrils with various morphologies (top two rows). The F3A and F8A mutants (third row), showing diminished toxicity, did not form fibrils. The K9A, K12A and K17A mutants (bottom row), which also display diminished toxicity, formed fibrils. A scale bar of 100 nm is presented in all images. **B**. PSMα3-induced cell death was assessed using the lactate dehydrogenase (LDH) colorimetric assay upon 30 min of incubation of T cells with 4 µM of each PSMα3 variant. The experiment was conducted in triplicates and repeated at least three times on different days, with similar results. The error bars represent the standard error of the mean. * $p < 0.05$ and ** $p < 0.001$ compared to WT PSMα3, accounting for the data from all three repeats.

Figure 2. Secondary structure analyses of PSMα3 and mutants with polar residue substitutions

A. The CD spectrum of WT PSMα3 shows helical configuration with minima at 208 and 222 nm. The K9A, K12A, D13A and K17A mutants demonstrate diminished helicity compared to WT PSMα3. **B**. ATR-FTIR spectroscopy analyses of the peptides upon solubilization (day one, dotted pink line), after one week of incubation at room temperature (day seven, dashed green line), and isolation of the pellet from the seven-day incubation (pellet, smooth black line). The experiments were repeated three times and the amide I' region of representative curves is shown. The secondary derivative of each curve was calculated and depicted to resolve and identify the major peaks contributing to the ATR-FTIR overlapping signal in this region. Peaks in the region of 1645–1662 cm^{-1} indicate the existence of α-helices, 1637–1645 cm^{-1} indicate disordered species, 1630–1643 cm^{-1} correlate with small and disordered β-rich amyloid fibrils with absorbance which is typical of bent β-sheets in native proteins, and 1611–1630 cm^{-1} as well as ~1685–1695 cm^{-1} are indicative of rigid cross-β fibrils *(47–50)*.

Figure 3. X-ray fiber diffraction of PSMα3 and derivatives

X-ray fiber diffraction patterns, with major reflections labeled. Peptide powders were dissolved in ultra-pure water to a concentration of 20 mg/ml and a microliter drop was applied between two sealed-end glass capillaries. The fiber diffraction was measured upon drying of the fibrils or after up to 10 hours of incubation on the capillaries.

Figure 4. The crystal structures of PSMα3 and its mutants display packing polymorphism

The crystal structures of PSMα3 (PDB ID 5I55) *(4)*, and the L15A (PDB ID 6GQ5), G16A (PDB ID 6GQC) and K12A (PDB ID 6GQ2) mutants determined at 1.45 Å, 1.5 Å, 1.4 Å and 1.5 Å resolution, respectively. The view is down the fibril axis. Two pairs of sheets are depicted in gray or orchid ribbons, side chains are represented as sticks and labeled by their three-letter amino acid code and position number. Heteroatoms are colored by atom type, and alanine substitution sites are indicated by cyan labels. The mated sheets form a hydrophobic dry interface (lacking water molecules) with inter-sheet distances shown on the right. The distances between α-helices along each sheet are shown on the left.

Figure 5. Potential salt bridges in the crystal structures of PSMα3 and its mutants

Crystal structures of PSMα3 and the K12A, L15A, and G16A mutants emphasizing potential intra- and inter-helical electrostatic interactions, with a view perpendicular to the fibril axis. Side chains are shown as sticks and labeled by their three-letter amino acid code and position number, and heteroatoms are colored according to atom type. Potential salt bridges are depicted by dashed lines.

Figure 6. Confocal microscopy of PSMα3 aggregation on the cell membrane

Representative confocal microscopy images of untreated T2 cells (left panels), and cells

incubated for 10 min (**A**) or 30 min (**B**) with a peptide mixture of 5 μM WT PSMα3 and 5

μM FITC-labeled PSMα3 (right panels) . The FITC channel (top) was merged with the

bright-field channel (bottom) to indicate peptide location in the cell. A 20 µm scale bar is shown for all images.

Figure 7. Scanning electron micrographs showing deformation of PSMα3-treated T2 cells

Scanning electron microscopy (SEM) images of (**A**) untreated T2 cells, and (**B**) T2 cells incubated for 5 min with 10 µM PSMα3. The micrographs were taken at different magnifications, as indicated by the scale bars (white on a black background) (10 µm (zoom-out) or 500 nm (zoom-in)).

Figure 8. PSMα3 co-aggregates with membrane particles around affected cells

T2 cells were treated with a mixture of 5 µM WT and 10 µM FITC-labeled PSMα3 peptides (green) in the presence of propidium iodide (PI, red). The cell membranes were stained with wheat gram agglutinin (WGA, blue). Representative images from the live-cell imaging were taken after 50 min. In the composite image of the stains, orange arrows indicate particles of extracted cell membrane co-localized with the aggregated peptide. Bright field (BF) image show cell damage.

Table 1. The biophysical and toxic properties PSMα3 and its derivatives

Peptide	Fibrils visualized by TEM**	ThT binding ***	Relative cytotoxicity ** (a)	Main secondary structure in solution (CD)***	Secondary structure in the fibril (FTIR)***	Main reflections in X-ray fiber diffraction**	Interpretation of fiber diffraction pattern
WT PSMα3*	Fibrils	+	+++	α-helical	α-helical	10.5/12/25 Å	Cross-α
D-PSMα3	Fibrils	+	+++	α-helical	N.D	10/12/25 Å	Cross-α
F3A	No fibrils	-	+	α-helical	N.D.	Diffuse	Ambiguous
L7A	Fibrils	-	++	α-helical	N.D.	4.7/11.5/9-12/28.5 Å	A mixture of cross-α/β
F8A	No fibrils	-	+	α-helical	N.D.	Diffuse	Ambiguous
K9A	Fibrils and crystals	+	+	α-helical with reduced helicity compared to WT	α-helical with partial transition to β-rich	4.6/10/12/24 Å	A mixture of cross-α/β
F10A	Fibrils	-	++	α-helical	N.D.	Diffuse	Ambiguous
F11A	Fibrils	-	+++	α-helical with reduced helicity compared to WT	N.D.	4.7/10/12.5/23-32 Å	A mixture with predominant cross-α diffraction
K12A*	Fibrils	-	+	α-helical with reduced helicity compared to WT	α-helical with partial transition to β-rich	4.3/9 Å	Could not interpret
D13A	Fibrils	-	++++	α-helical with reduced helicity compared to WT	α-helical with partial transition to β-rich	Diffuse	Ambiguous
L15A*	Fibrils	+	++	α-helical	N.D.	4.6/10.5/12.5/25 Å	A mixture with predominant cross-α diffraction
G16A*	Fibrils	+	+++	α-helical	N.D.	Diffuse	Ambiguous
K17A	Fibrils	+	-	α-helical with reduced helicity compared to WT	α- helical	10/12.5/25 Å	Cross-α
C-cys-PSMα3	Fibrils	+	+++(b)	N.D.	N.D.	N.D	N.D.
N-cys-PSMα3	No fibrils	-	+(b)	N.D.	N.D.	N.D	N.D.

N.D. - Not determined

* Crystal structure solved and showed cross-α fibrils

** No pre-treatment of the peptides

*** Peptides were pretreated to dissolve preformed aggregates, as detailed in the Methods section

(a) Results not influenced by peptide pretreatment.

(b) Toxicity induced by 8μM peptide; all other peptides were tested at 4μM

Experimental Section

Table 2. List of PSMα3 derivatives

Peptide Name	Sequence
PSMα3 WT	MEFVAKLFKFFKDLLGKFLGNN
D-PSMα3	**d**(MEFVAKLFKFFKDLLGKFLGNN)
C-FITC-PSMα3*	MEFVAKLFKFFKDLLGKFLGNN-**K-FITC**
N-cys-PSMα3	**C**MEFVAKLFKFFKDLLGKFLGNN
C-cys-PSMα3	MEFVAKLFKFFKDLLGKFLGNN**C**
F3A-PSMα3	ME**A**VAKLFKFFKDLLGKFLGNN
L7A-PSMα3	MEFVAK**A**FKFFKDLLGKFLGNN
F8A-PSMα3	MEFVAKL**A**KFFKDLLGKFLGNN
K9A-PSMα3	MEFVAKLF**A**FFKDLLGKFLGNN
F10A-PSMα3	MEFVAKLFK**A**FKDLLGKFLGNN
F11A-PSMα3	MEFVAKLFKF**A**KDLLGKFLGNN
K12A-PSMα3	MEFVAKLFKFF**A**DLLGKFLGNN
D13A-PSMα3	MEFVAKLFKFFK**A**LLGKFLGNN
L15A-PSMα3	MEFVAKLFKFFKDL**A**GKFLGNN
G16A-PSMα3	MEFVAKLFKFFKDLL**A**KFLGNN
K17A-PSMα3	MEFVAKLFKFFKDLLG**A**FLGNN

*FITC is linked to the side chain of C-terminally added Lysine, hence net charge is unaltered.

PSMα3 WT and mutant peptides (custom synthesis at >98% purity) were purchased from GL Biochem (Shanghai, China) Ltd. or from GenScript. C-FITC-labeled PSMα3 was purchased from GL Biochem. All chemicals and reagents were of analytical grade. Chemicals were purchased from Sigma-Aldrich Ltd., including: trifluoroacetic acid (TFA), hexafluoroisopropanol (HFIP), thioflavin T (ThT), and deuterium-oxide (D_2O). Ultra-pure water and Dullbeco's phosphate buffered saline (PBS) were purchased from Biological Industries (Kibbutz Beit-Haemek, Israel). Phosphate dibasic anhydrous and di-sodium hydrogen phosphate were purchased from Carlo Erba, distributed by Romical Ltd. (Beer-Sheva, Israel). Sodium chloride was purchased from Merck. Additional reagents and consumable are mentioned below.

Peptide pretreatment for disaggregation

In experiments that included a time-resolved aspect or secondary structure determination in different peptide states, for example measuring aggregation kinetics, and CD and FTIR measurements, the peptides were pretreated to dissolve any pre-formed aggregates, and to ensure reproducibility of the results *(51–53)*. Lyophilized peptide powder was freshly dissolved in TFA-HFIP (1:1), to a concentration of 1-5 mg/ml, sonicated for 5-10 min in a sonication bath, and then allowed to evaporate under vacuum or in a chemical hood, for 2-3 days. Residual solvent was evaporated using a speed vac. For certain experiments, the peptides were incubated for 1h in TFA-HFIP at room temperature (RT) and dried directly in an acid-resistant speed vac (RVC 2-18 CDplus, CHRIST, distributed by Dr Golik, Tel-Aviv, Israel). Unless immediately tested, the treated peptides were stored at -20°C.

Fibrillation assays

Thioflavin T fluorescence kinetic assay

Fresh ThT reaction solution was prepared by diluting (1:10) the filtered 2 mM ThT stock, made in ultra-pure water, into the reaction buffer containing 10 mM sodium phosphate buffer and 150 mM NaCl, pH 8.0. PSMα3 and its derivatives (Table 2), pretreated as described above, were dissolved in ultra-pure water to 1 mM, on ice, and sonicated for 5 min using a sonication bath. The peptides were then diluted, on ice, in a filtered reaction buffer, and centrifuged (5 min, 4 °C, 10,000 rpm). As only non-centrifuged K17A samples, as opposed to centrifuged sample, bound ThT binding, K17A centrifugation was avoided. The supernatant of each mutant (or a sample of K17A solution) was separately mixed with ThT reaction solution in the wells of a Greiner bio-one black, 96-well, flat-bottom plate (distributed by De Groot Medical Ltd. Rosh Ha'ain, Israel), to final concentrations of 50 µM peptide and 200 µM ThT. The plate was immediately covered with a silicone sealing film (ThermalSeal RTS, Sigma-Aldrich), and incubated in a plate reader (CLARIOstar or FLUOstar Omega, BMG LABTECH. Distributed by Zotal Ltd. Tel-Aviv, Israel) at 37 °C, with 500 rpm, with shaking for 15 sec before each cycle, and up to 1000 cycles of 3-5 min each. ThT fluorescence was measured at an excitation wavelength of 438±20 nm and emission wavelength of 490±20 nm. Measurements were made in triplicates and each experiment was repeated at least three times. All triplicate values were averaged, appropriate blanks were subtracted, and the resulting values were plotted against time. Calculated standard errors of the mean are presented as error bars. Of note, the measurement of L15A was separated from the measurement of the other mutants since L15A produced a very high signal which caused signal overflow, which required digital

gain reduction in the plate reader. Cysteine derivatives were also tested in a separate experiment, including PSMα3 for comparison, and according to the exact same protocol.

<u>Transmission electron microscopy (TEM)</u>

Lyophilized (untreated) peptide powders were separately dissolved in ultra-pure water to a concentration of 10 mM, and incubated for 7-14 days at RT, or for 3-7 days at 37 °C. All samples were then centrifuged (20 min, 14,800 rpm). The supernatant was discarded and the pellet was resuspended to the same volume and diluted 10-fold in ultra-pure water. Samples (5 µl) were then directly applied onto 400-mesh copper TEM grids with support films of Formvar/Carbon (Ted Pella, distributed by Getter Group Bio Med, Petah-Tikva, Israel), that were charged by high-voltage, alternating current glow-discharge (PELCO easiGlow, Ted Pella), immediately before use. Grids were allowed to adhere for 2 min. Excess solution was blotted with filter paper, and the grids were negatively stained with 2% uranyl acetate for 30 sec, blotted again, and dried under vacuum. Specimens were examined with a FEI Tecnai T12 G2 or FEI Tecnai G2 T20 S-Twin transmission electron microscope (FEI, Hillsboro, Oregon, United States), at an accelerating voltage of 120 or 200 KeV, respectively.

Circular dichroism

All peptides were pretreated TFA: HFIP 1:1 mixture as described above in a concentration of 1 mg/ml in, in order to dissolve any pre-formed aggregates, and then fully dried. The treated peptides were dissolved in ultra-pure water to 1.5 mM and then sonicated for 10 min, in a sonication bath. For each derivative tested, 250 µl of 10 mM sodium phosphate, 150 mM NaCl pH 7.4 buffer was first measured as a blank, after which, 20 µl peptide was diluted into the cuvette (to a calculated concentration of 111 µM). Samples were

thoroughly mixed to ensure homogeneity immediately prior to measurement. CD spectra were recorded on either the Applied Photophysics Chirascan qCD or Pi-star 180 (Applied Photophysics, Surrey, United Kingdom), using a 1 mm path-length fused quartz cell. The presented measurements are an average of four scans for each peptide or two scans for blank, captured at a scan speed of 1 sec, bandwidth and step size of 1 nm, over a wavelength range of 180-260 nm. High-voltage (HV) and absorbance measurements were taken simultaneously, and CD values with corresponding HV or absorbance values above the recommended limit (evident only in wavelengths shorter than 200 nm), were omitted from the spectra. Due to peptide aggregation upon dilution in the tested buffer, which significantly lowered the soluble peptide concentration for certain PSMα3 derivatives, the concentration of each sample at real time was experimentally determined by UV absorbance, which was measured in parallel to CD spectra measurements, in the same machine. Since PSMα3 sequence lacks tryptophan and tyrosine residues, the soluble peptide concentration (in molar units) was calculated using the Beer-Lambert equation at 205 nm: $A_{205} = \varepsilon_{205} \times l \times C$. The pathlength l = 0.1 cm. A_{205} was determined by averaging the blank corrected absorbance values at 205 nm from all four scans. ε_{205} (M^{-1} x cm^{-1}) is a sequence-specific extinction coefficient calculated using the webserver (http://spin.niddk.nih.gov/clore/) *(54)*. The molar ellipticity per residue (Ө, in mdeg*cm^2 *$dmol^{-1}$*$residu^{-1}$) was derived from the raw data (δ, in mdeg.), using the following formula: Ө = δ / (L*C*N), where L is the path length of the cuvette, C is the real-time calculated molar concentration of PSMα3, and N is the number of residues in PSMα3 (i.e., 22).

Attenuated total-internal reflection Fourier transform infrared spectroscopy

PSMα3 and derivatives, pretreated as described above, were separately dissolved in 5 mM HCl to 1 mg/ml, sonicated for 3 min in a sonication bath, frozen in liquid nitrogen and lyophilized. Upon complete dryness, the peptide was subjected to the same treatment without sonication. The latter step was repeated four times. The dry peptide was then dissolved in D_2O to 1 mg/ml, frozen and lyophilized. This process was repeated at least twice. The dry peptide was further dissolved in D_2O to 25 mg/ml and incubated at RT for 2-6 h. Sample (5 µl) were then separately applied on a surface of the ATR module (MIRacle Diamond w/ZnSe lens 3-Reflection HATR Plate; Pike Technologies, Fitchburg, Wisconsin, USA) and left to dry under nitrogen, to form a thin film. Absorption spectra were recorded with a Tensor 27 FTIR spectrometer, equipped with a high-sensitivity LN-cooled MCT detector (Bruker Optics, Bruker Scientific Israel Ltd., Rehovot, Israel). Measurements were performed as an accumulation of 100 scans with 2 cm^{-1} resolution over a range of 1500-1800 cm^{-1}. The same procedure was repeated after incubation of samples for seven days, at RT, in a sealed tube. The residual samples were then centrifuged for 10 min at 17,000g, at RT, to separate the fibrils from the supernatant. The pellet was resuspended with the same amount of D_2O, centrifuged again under the same conditions, and the remaining pellet was resuspended in 10 µl D_2O and measured under the same conditions. Background (ATR module) and blank (D_2O) subtraction, and baseline correction, was performed using the Bruker OPUS software. ATR-FTIR curves were normalized to unity and their second derivatives were calculated and rendered using the Origin 2018b graphing & analysis software (OriginLab®, distributed by Tashtit

Scientific Consultants Ltd, Rishon LeZion, Israel). The amide I' region of the spectra (1600 -1700 cm^{-1}) is presented in the curves.

Tissue culture and cell toxicity

Our calibration experiments verified that T2 cytotoxicity results were not affected by peptide pretreatment, therefore, in this work, cell toxicity assays were conducted with the non-treated peptide powders. T2 cells (174 x CEM.T2) (ATCC® CRL-1992™, distributed by Biological Industries) were cultured in RPMI 1640 medium containing L-glutamine (Gibco, distributed by Rhenium Ltd. Modi'in, Israel), supplemented with 20% heat-inactivated fetal calf serum, penicillin (100 U/ml) and streptomycin (0.1 mg/ml), at 37ºC in 5% CO_2. Before each experiment, the cells were washed and resuspended in assay medium, which was similar to T2 growth medium, except that it contained 0.5% fetal calf serum. The cells were incubated for two hours at 37 °C, in 5% CO_2. PSMα3 and its derivatives were separately dissolved to 1 mM in ultra-pure water, sonicated for 10 min, divided to aliquots and stored at -20ºC. One aliquot was thawed immediately before each experiment. Peptides were serially diluted in assay medium to double the final concentration used in the assay (as denoted in the figures). Then, 50 µl of each diluted sample was pipetted into a well of a 96-well plate in triplicates, and incubated at 37 °C, in 5% CO_2, without shaking, for 1 h. Cells were then diluted in assay medium to 0.15×10^6 viable cells/ml and 50 µl were transferred to each well using a multichannel pipette. The plate was incubated for 30 min, at 37 °C, in 5% CO_2. Cell death and cell lysis were quantified using the LDH colometric assay, according to the manufacturer's instructions, including all recommended controls (LDH; Cytotoxicity Detection Kit Plus, Roche Applied Sciences, Penzberg, Germany). Cell-free culture medium was measured as background.

Cells subjected to the same experimental conditions apart from peptide addition, were used as the 'low control' to account for spontaneous LDH release. Cells subjected to the same experimental conditions apart from peptide addition, and treated with lysis buffer prior to color development reaction, according to the manufacturer instructions, were used as the high positive control to account for maximum LDH release. The average absorbance values of each triplicate and controls, measured at 490 nm with a reference wavelength at 690 nm using the FLUOstar Omega (BMG LABTECH) plate reader, was calculated, and background was subtracted. The following equation was used to calculate %cytotoxicity according to the kit's protocol: $\%Cytotoxicity = \frac{(exp.value - low\ control) \times 100}{high\ control - low\ control}$. The data were analyzed using Excel. The experiments were repeated three times on different days. The PSMα3 cysteine derivatives were examined in separate experiments, with WT PSMα3 for comparison. Statistical analysis was conducted using the Mann-Whitney U (two tailed) test.

Confocal microscopy

All solutions used for the confocal microscope assays, apart for cell media, were prepared with filtered DDW and/or were filtered twice using a 0.22 μm syringe filter.

<u>PSMα3 aggregation on the membranes of fixed cells</u>

Untreated WT and FITC-labeled PSMα3 peptides were separately dissolved in ultra-pure water to a concentration of 2 mM, mixed to a final concentration of 1 mM, and then incubated together at RT just prior to the experiment. T2 cells (174 x CEM.T2) (ATCC® CRL-1992™) grown as described above, were washed and diluted in unsupplemented RPMI 1640 medium (Gibco) to 5×10^6 viable cells/ml and gently pipetted onto dry, small-diameter coverslips pre-coated with 0.1 % poly-lysine solution, in a 24-well plate. Samples

were incubated on the coverslip for 20 min, at 37 °C, in 5% CO_2. In parallel, the peptide mixture was diluted in the cytotoxicity assay medium, comprised of RPMI 1640 medium supplemented with 0.5 % heat-inactivated fetal calf serum, penicillin (100 U/ml) and streptomycin (0.1 mg/ml), to a final concentration of 5 µM of each (WT and FITC-labeled peptides). The cell-coated coverslips were then washed three times with fresh, pre-warmed (to 37 °C) and unsupplemented RPMI medium, and either cytotoxicity assay medium alone (for control), or the above described peptide mixture were applied on the coverslips. The cells were incubated for 10 min at 37 °C in 5% CO_2, rinsed three times with filtered PBS (Biological Industries), and fixed at RT, for 1 h, in aqueous solution containing 100 mM sucrose, 100 mM sodium phosphate buffer, pH 7.4, 2.5% glutaraldehyde (from a 8% stock solution, Electron Microscopy Sciences, Hatfield, Pennsylvania, USA), and 2% paraformaldehyde (from 16% stock solution, Electron Microscopy Sciences). After fixation, the cells were washed three times with filtered PBS. Of note, all solution exchange steps were carried out rapidly, to avoid any contact of the cells with air. Finally, the cells were immersed in freshly filtered PBS and stored at 4°C until visualized with a Leica SP8 confocal laser scanning microscope, equipped with a HCPL APO CS 2 40x/1.10 water objective. The images were taken at a zoom of either 0.75 or 2.3. Images were processed and rendered with ImageJ.

PSMα3 co-localization and clamping with cell membranes in suspension

On the day of the experiment, untreated WT and FITC-PSMα3 were separately dissolved in ultrapure water to a concentration of 1 mM, and mixed together in PBS to concentrations of 10 µM and 20 µM, respectively. Wheat germ agglutinin (WGA) Alexa Fluor™ 350 Conjugate stock solution (ThermoFisher Scientific) was prepared in PBS to

a concentration of 1 mg/ml and then diluted in Hank's balanced salt solution (HBSS) to a concentration of 5 µg/ml, according to the manufacturer's instructions. T2 cells (174 x CEM.T2) (ATCC® CRL-1992™) were grown as described above in RPMI 1640 medium containing L-glutamine (Gibco), supplemented with 20% heat inactivated fetal calf serum, penicillin (100 U/ml) and streptomycin (0.1 mg/ml), at 37 ºC in 5% CO_2. Before the experiment, 1×10^6 cells were washed and resuspended in assay medium, similar to T2 growth medium, except that it contained 0.5% fetal calf serum. After two hours of incubation at 37 °C, in 5% CO_2, the cells were washed and suspended in the WGA staining solution and incubated for 10 min, under the same conditions described for membrane staining. The cells were then washed twice and diluted in assay medium to a density of $0.15*10^6$ cell/ml. Stained cells were transferred to a chamber of a 6-channel microscope µ-slide (µ-Slide VI 0.4, #cat 80606, ibidi) and imaged. The incubated peptide mixture was supplemented with propidium iodide (PI) from a stock of 1 mg/ml in PBS, and then injected to the µ-slide channel and allowed to diffuse toward the cells. Final concentrations in the chamber were 5 µM and 10 µM WT and FITC-PSMα3 peptides, respectively, and 0.02 mg/ml PI. Cells were continuously imaged with an LSM 710, AnxioObserver laser scanning confocal microscope (SEM; Zeiss), equipped with a C-Apochromat 40x/120 W Korr M27 water objective, for 50 min, at 37 ºC and 5% CO_2. Data analysis was performed using the ZEN 2.3 lite (Zeiss) and a representative image was rendered with ImageJ.

Scanning electron microscopy

All solutions used for this assay, apart from cell medium, were prepared with filtered DDW and/or were filtered twice using a 0.22 µm syringe filter. SEM chips (silicon chips, 5x5

mm) were pre-coated with 0.1% poly-lysine solution, washed three times with DDW and dried. Untreated PSMα3 peptide was freshly dissolved in ultra-pure water to a concentration of 1 mM and sonicated for 5 min in a sonication bath, and then further diluted to a concentration of 10 µM, in cytotoxicity assay medium, comprised of RPMI 1640 medium supplemented with 0.5 % heat inactivated fetal calf serum, penicillin (100 U/ml) and streptomycin (0.1 mg/ml). As control, ultra-pure water was added to the medium in the same volume as the peptide. T2 cells (174 x CEM.T2) (ATCC® CRL-1992™) grown in RPMI 1640 medium containing L-glutamine (Gibco), supplemented with 20% heat inactivated fetal calf serum, penicillin (100 U/ml) and streptomycin (0.1 mg/ml), were washed and diluted in unsupplemented RPMI medium to a concentration of ~10x10^6 viable cells/ml. The cell suspension was applied onto the poly-lysine coated chips, and incubated for 20 min. The chips were then rinsed three times with fresh RPMI medium, and immediately immersed, for 5 min, in either 10 µM PSMα3 or the control solutions. The chips were then rinsed twice in filtered PBS (Biological Industries). Then, the chips were incubated, for 1 h, at RT, in aqueous fixation solution containing 100 mM sucrose, 100 mM sodium phosphate buffer, pH 7.4, 2.5% glutaraldehyde (from a 8% stock solution, Electron Microscopy Science) and 2% paraformaldehyde (from 16% stock solution, Electron Microscopy Science). Chips were then washed in three 2-min cycles, with 0.1 M cacodylate buffer, pH 7.4, and incubated, for 15 min, at RT, in freshly prepared secondary fixative containing 1% OsO4 in 0.1 M cacodylate buffer. The chips were then subjected to three 2-min washing cycles with DDW, followed by dehydration by washing the chips twice in a row in each concentration of a graded ethanol series of 15%, 30%, 50%, 70%, 90%, 95% and 100% with 3 min incubation for each wash, apart from the last step which

lasted 5 min. Finally, the chips were dried using the critical point drying (CPD) method, coated with 5 nm of chromium and kept under vacuum until they were scanned with a Zeiss ULTRA *plus* field emission SEM using either the InLens or SE2 detectors (Carl Zeiss, Oberkochen, Germany).

Fiber X-ray diffraction

Peptides were dissolved in ultra-pure water, to a concentration of 20 mg/ml. Several microliters were applied between two sealed glass capillaries and incubated for up to 10 h, at RT. For some peptides, the X-ray fiber diffraction was collected immediately upon complete drying of the fibrils on the capillaries. For WT PSMα3, the diffraction pattern was tested using a freshly dissolved peptide, as well as a peptide solution incubated in the tube for 10 h, at RT, prior to sample preparation. The fresh (Figure 3) and incubated (data not shown) fibrils yielded with the same diffraction pattern. However, for the peptides presented in Figure S4, all conditions tested yielded only diffuse diffraction patterns. X-ray diffraction was collected at the micro-focused beam P14 at the high brilliance 3rd Generation Synchrotron Radiation Source at DESY: PETRA III, Hamburg, Germany or on beamlines ID23-EH2 and ID29 at the European Synchrotron Radiation Facility (ESRF), Grenoble, France.

Peptide crystallization

Peptides were crystallized with no pretreatment. After dissolving the peptides, the solutions were sonicated in a sonication bath for 5-10 min, following by 5 min centrifugation at 13,700 rpm, at 4 °C. The peptides were kept on ice until crystallization plates were set. All crystals were grown at 20 °C, using the hanging-drop vapor diffusion technique. K12A crystals were grown from a 10 mM (26.2 mg/ml) solution in ultra-pure

water, mixed with a reservoir (Qiagen, Classics Suite I crystallization screen, condition B8) containing 0.1 M HEPES pH 7.5, 0.5 M ammonium sulfate, and 30% v/v 2-methyl-2,4-pentanediol (MPD). Crystals were flash-frozen before X-ray data collection. No cryo-protectant was added. L15A crystals were grown from a 7.5 mM (19.2 mg/ml) solution in ultra-pure water, in a reservoir (Molecular Dimensions, JCSG + HTS crystallization screen, condition E5) containing 0.1 M CAPS, pH 10.5 and 40% v/v MPD. Crystals were flash-frozen before X-ray data collection. No cryo-protectant was added. G16A crystals were grown from a 10 mM (26.2 mg/ml) solution in ultra-pure water, in a reservoir (Qiagen, Classics Suite I crystallization screen, condition A1) containing 0.1 M sodium acetate, pH 4.6, 0.01 M cobalt(II) chloride, and 1 M 1,6-hexanediol. Before data collection, the crystals were immersed in a cryo-protectant solution in which the ingredients of the crystallization reservoir were supplemented with 20% ethylene glycol, and then flash-frozen in liquid nitrogen.

Structure determination and refinement

X-ray diffraction data of PSMα3 mutants were collected at 100 °K at the micro-focal beamlines ID23-2 (G16A) and ID30a-3 (K12A, L15A) at the European Synchrotron Radiation facility (ESRF), Grenoble, France, at wavelengths of 0.8729 and 0.9677Å, respectively. Data indexing, integration and scaling were performed using XDS/XSCALE *(55)*. Phases were obtained using ARCIMBOLDO *(56)* (K12A) or by molecular replacement with Phaser *(57)* (L15A, G16A). For the molecular replacement of G16A and L15A, the structure of WT PSMα3 was used as a search model. Crystallographic refinements were performed with Refmac5 *(58)*. Model building for all structures was performed with Coot *(59)* and illustrated with Chimera *(60)*. There were no residues

detected in the disallowed region of the Ramachandran plot. All PSMα3 mutant structures contained one α-helix in the asymmetric unit. The K12A structure contained three water molecules, and one sulfate anion in the asymmetric unit. The L15A structure contained 11 water molecules and one MRD ((4R)-2-methylpentane-2,4-diol) molecule. The G16A structure contained two water molecules, two sodium cations, and one chloride anion in the asymmetric unit. Two alternative conformations of Leu7 were detected. Crystallographic data and statistics for each structure are listed in Table S2.

Calculations of structural properties

The inter-sheet distance in the structures was calculated using Chimera *(60)*. First, a plane was defined for each sheet of α-helices containing at least four helices, using only the backbone atoms of the helix and excluding flanking residues in each structure. Then, the distance between the planes was measured using the "measure distance" command. This method was used to measure distances of both the dry and the wet interfaces. To calculate the distance between α-helices along the sheet, an axis was defined for two adjacent helixes, again using only backbone atoms that participate in the helical portion of the peptide, and the distance between the two axes was measured in the "define axis/plane/centroid" window. All calculated values, as well as the lengths of the axes calculated for the α-helix of each peptide structure, are summarized in Table S1. Of note, the values calculated here for WT PSMα3 are slightly different than those published earlier *(4)*, due to differences in calculation methods.

Notes

The authors declare no competing financial interests.

Data availability

The data that support the findings of this study are available on request from the corresponding author. Coordinates and structure factors for the X-ray crystal structures have been deposited in the protein data bank (PDB) with accession codes 6GQ2, 6GQ5, and 6GQC, for K12A, L15A and G16A respectively.

Acknowledgements

We acknowledge the guidance and technical support provided by Yael Pazy-Benhar and Dikla Hiya at the Technion Center for Structural Biology (TCSB), Yaron Kauffmann from the MIKA electron microscopy center of the department of material science & engineering at the Technion, Israel, Einat Nativ-Roth and Alexander Upcher from the Ilse Katz Institute for Nano-Science and Technology at Ben-Gurion University of the Negev, Israel, Einat Zelinger from The Interdepartmental Equipment Facility of the faculty of agriculture, food and environment, of the Hebrew university of Jerusalem, Israel, and of Nitsan Dahan and Yael Lupu-Haber from the LS&E infrastructure center of the Lokey center of the Technion, Israel. We are grateful for the service provided by Lihi Shaulov from the Rappaport Faculty of Medicine of the Technion, Israel, and by Na'ama Koifman from the Technion Center for Electron Microscopy of Soft Matter of the Russell Berrie Nanotechnology Institute, Israel. This research was supported by the I-CORE Program of the Planning and Budgeting Committee and The Israel Science Foundation, Center of Excellence in Integrated Structural Cell Biology (grant no. 1775/12), DFG: Deutsch-IsraelischeProjektkooperation (DIP) (grant no. LA 3655/1-1), Israel Science Foundation

(grant no. 560/16), University of Michigan – Israel Collaborative Research Grant, and BioStruct-X, funded by FP7. The synchrotron MX data collection experiments were performed at beamlines ID23-EH2, ID29, and MASSIF-3 (ID30A-3) at the European Synchrotron Radiation Facility (ESRF), Grenoble, France, and at beamline P14, operated by EMBL Hamburg at the PETRA III storage ring (DESY, Hamburg, Germany). We are grateful to the teams at ESRF and EMBL Hamburg.

References

(1)   Mehlin, C.; Headley, C. M.; Klebanoff, S. J. An Inflammatory Polypeptide Complex from Staphylococcus Epidermidis: Isolation and Characterization. *J. Exp. Med.* **1999**, *189* (6), 907–918.

(2)   Cheung, G. Y. C.; Joo, H.-S.; Chatterjee, S. S.; Otto, M. Phenol-Soluble Modulins – Critical Determinants of Staphylococcal Virulence. *FEMS Microbiol. Rev.* **2014**, *38* (4).

(3)   Laabei, M.; Jamieson, W. D.; Yang, Y.; van den Elsen, J.; Jenkins, A. T. A. Investigating the Lytic Activity and Structural Properties of Staphylococcus Aureus Phenol Soluble Modulin (PSM) Peptide Toxins. *Biochim. Biophys. Acta - Biomembr.* **2014**, *1838* (12), 3153–3161.

(4)   Tayeb-Fligelman, E.; Tabachnikov, O.; Moshe, A.; Goldshmidt-Tran, O.; Sawaya, M. R.; Coquelle, N.; Colletier, J. P.; Landau, M. The Cytotoxic Staphylococcus Aureus PSMα3 Reveals a Cross-α Amyloid-like Fibril. *Science.* **2017**, *355* (6327), 831–833.

(5)   Salinas, N.; Colletier, J. P.; Moshe, A.; Landau, M. Extreme Amyloid Polymorphism in Staphylococcus Aureus Virulent PSMα Peptides. *Nat. Commun.* **2018**, *9* (1), 3512.

(6)   Schwartz, K.; Syed, A. K.; Stephenson, R. E.; Rickard, A. H.; Boles, B. R. Functional Amyloids Composed of Phenol Soluble Modulins Stabilize Staphylococcus Aureus Biofilms. *PLoS Pathog.* **2012**, *8* (6), e1002744.

(7)   Marinelli, P.; Pallares, I.; Navarro, S.; Ventura, S. Dissecting the Contribution of Staphylococcus Aureus α-Phenol-Soluble Modulins to Biofilm Amyloid Structure. *Sci. Rep.* **2016**, *6* (1), 34552.

(8)   Eisenberg, D.; Jucker, M. The Amyloid State of Proteins in Human Diseases. *Cell* **2012**, *148* (6), 1188–1203.

(9)   Tycko, R. Amyloid Polymorphism: Structural Basis and Neurobiological Relevance. *Neuron* **2015**, *86* (3), 632–645.

(10)   Knowles, T. P. J.; Vendruscolo, M.; Dobson, C. M. The Amyloid State and Its Association with Protein Misfolding Diseases. *Nat. Rev. Mol. Cell Biol.* **2014**, *15* (6), 384–396.

(11)   Fowler, D. M.; Koulov, A. V; Alory-Jost, C.; Marks, M. S.; Balch, W. E.; Kelly, J. W. Functional Amyloid Formation within Mammalian Tissue. *PLoS Biol.* **2005**, *4* (1), e6.

(12)   Jacob, R. S.; Das, S.; Ghosh, S.; Anoop, A.; Jha, N. N.; Khan, T.; Singru, P.; Kumar, A.; Maji, S. K. Amyloid Formation of Growth Hormone in Presence of Zinc: Relevance to Its Storage in Secretory Granules. *Sci. Rep.* **2016**, *6*, 23370.

(13)   Maji, S. K.; Perrin, M. H.; Sawaya, M. R.; Jessberger, S.; Vadodaria, K.; Rissman, R. A.;

Singru, P. S.; Nilsson, K. P. R.; Simon, R.; Schubert, D.; et al. Functional Amyloids as Natural Storage of Peptide Hormones in Pituitary Secretory Granules. *Science* **2009**, *325* (5938), 328–332.

(14) Pham, C. L. L.; Kwan, A. H.; Sunde, M. Functional Amyloid: Widespread in Nature, Diverse in Purpose. *Essays Biochem.* **2014**, *56* (1), 207–219.

(15) DePas, W. H.; Chapman, M. R. Microbial Manipulation of the Amyloid Fold. *Res. Microbiol.* **2012**, *163* (9–10), 592–606.

(16) Otzen, D.; Nielsen, P. H. We Find Them Here, We Find Them There: Functional Bacterial Amyloid. *Cell. Mol. Life Sci.* **2008**, *65* (6), 910–927.

(17) Chapman, M. R.; Robinson, L. S.; Pinkner, J. S.; Roth, R.; Heuser, J.; Hammar, M.; Normark, S.; Hultgren, S. J. Role of Escherichia Coli Curli Operons in Directing Amyloid Fiber Formation. *Science.* **2002**, *295* (5556), 851–855.

(18) Soragni, A.; Yousefi, S.; Stoeckle, C.; Soriaga, A. B.; Sawaya, M. R.; Kozlowski, E.; Schmid, I.; Radonjic-Hoesli, S.; Boutet, S.; Williams, G. J.; et al. Toxicity of Eosinophil MBP Is Repressed by Intracellular Crystallization and Promoted by Extracellular Aggregation. *Mol. Cell* **2015**, *57* (6), 1011–1021.

(19) Hughes, M. P.; Sawaya, M. R.; Boyer, D. R.; Goldschmidt, L.; Rodriguez, J. A.; Cascio, D.; Chong, L.; Gonen, T.; Eisenberg, D. S. Atomic Structures of Low-Complexity Protein Segments Reveal Kinked β Sheets That Assemble Networks. *Science.* **2018**, *359* (6376), 698–701.

(20) Eisenberg, D. S.; Sawaya, M. R. Structural Studies of Amyloid Proteins at the Molecular Level. *Annu. Rev. Biochem.* **2017**, *86* (1), 69–95.

(21) Knowles, T. P.; Fitzpatrick, A. W.; Meehan, S.; Mott, H. R.; Vendruscolo, M.; Dobson, C. M.; Welland, M. E. Role of Intermolecular Forces in Defining Material Properties of Protein Nanofibrils. *Science.* **2007**, *318* (5858), 1900–1903.

(22) Wang, R.; Braughton, K. R.; Kretschmer, D.; Bach, T.-H. L.; Queck, S. Y.; Li, M.; Kennedy, A. D.; Dorward, D. W.; Klebanoff, S. J.; Peschel, A.; et al. Identification of Novel Cytolytic Peptides as Key Virulence Determinants for Community-Associated MRSA. *Nat. Med.* **2007**, *13* (12), 1510–1514.

(23) Cheung, G. Y. C.; Kretschmer, D.; Queck, S. Y.; Joo, H.-S.; Wang, R.; Duong, A. C.; Nguyen, T. H.; Bach, T.-H. L.; Porter, A. R.; DeLeo, F. R.; et al. Insight into Structure-Function Relationship in Phenol-Soluble Modulins Using an Alanine Screen of the Phenol-Soluble Modulin (PSM) α3 Peptide. *FASEB J.* **2014**, *28* (1), 153–161.

(24) Zheng, Y.; Joo, H. S.; Nair, V.; Le, K. Y.; Otto, M. Do Amyloid Structures Formed by Staphylococcus Aureus Phenol-Soluble Modulins Have a Biological Function? *Int. J. Med. Microbiol.* **2018**, *308* (6), 675–682.

(25) Kumar, S.; Nussinov, R. Close-Range Electrostatic Interactions in Proteins. *ChemBioChem* **2002**, *3* (7), 604.

(26) Towle, K. M.; Lohans, C. T.; Miskolzie, M.; Acedo, J. Z.; van Belkum, M. J.; Vederas, J. C. Solution Structures of Phenol-Soluble Modulins A1, A3, and B2, Virulence Factors from Staphylococcus Aureus. *Biochemistry* **2016**, *55* (34), 4798–4806.

(27) Ghosh, D.; Singh, P. K.; Sahay, S.; Jha, N. N.; Jacob, R. S.; Sen, S.; Kumar, A.; Riek, R.; Maji, S. K. Structure Based Aggregation Studies Reveal the Presence of Helix-Rich Intermediate during α-Synuclein Aggregation. *Sci. Rep.* **2015**, *5* (1), 9228.

(28) Hebda, J. A.; Miranker, A. D. The Interplay of Catalysis and Toxicity by Amyloid Intermediates on Lipid Bilayers: Insights from Type II Diabetes. *Annu. Rev. Biophys.* **2009**, *38* (1), 125–152.

(29) Yao, Z.; Cary, B. P.; Bingman, C. A.; Wang, C.; Kreitler, D. F.; Satyshur, K. A.; Forest, K. T.; Gellman, S. H. Use of a Stereochemical Strategy To Probe the Mechanism of Phenol-Soluble Modulin A3 Toxicity. *J. Am. Chem. Soc.* **2019**, *141* (19), 7660–7664.

(30) Stefani, M. Biochemical and Biophysical Features of Both Oligomer/Fibril and Cell Membrane in Amyloid Cytotoxicity. *FEBS J.* **2010**, *277* (22), 4602–4613.

(31) Walsh, P.; Vanderlee, G.; Yau, J.; Campeau, J.; Sim, V. L.; Yip, C. M.; Sharpe, S. The Mechanism of Membrane Disruption by Cytotoxic Amyloid Oligomers Formed by Prion Protein(106-126) Is Dependent on Bilayer Composition. *J. Biol. Chem.* **2014**, *289* (15), 10419–10430.

(32) Habchi, J.; Chia, S.; Galvagnion, C.; Michaels, T. C. T.; Bellaiche, M. M. J.; Ruggeri, F. S.; Sanguanini, M.; Idini, I.; Kumita, J. R.; Sparr, E.; et al. Cholesterol Catalyses Aβ42 Aggregation through a Heterogeneous Nucleation Pathway in the Presence of Lipid Membranes. *Nat. Chem.* **2018**, *10* (6), 673–683.

(33) Malishev, R.; Tayeb-Fligelman, E.; David, S.; Meijler, M. M.; Landau, M.; Jelinek, R. Reciprocal Interactions between Membrane Bilayers and S. Aureus PSMα3 Cross-α Amyloid Fibrils Account for Species-Specific Cytotoxicity. *J. Mol. Biol.* **2018**, *430* (10), 1431–1441.

(34) Pokorny, A.; Birkbeck, T. H.; Almeida, P. F. F. Mechanism and Kinetics of δ-Lysin Interaction with Phospholipid Vesicles. *Biochemistry* **2002**, *41* (36), 11044–11056.

(35) Zhang, M.; Zhao, J.; Zheng, J. Molecular Understanding of a Potential Functional Link between Antimicrobial and Amyloid Peptides. *Soft Matter* **2014**, *10* (38), 7425–7451.

(36) Kinnunen, P. K. J. Amyloid Formation on Lipid Membrane Surfaces. *Open Biol. J.* **2010**, *2* (2), 163–175.

(37) Engel, M. F. M.; Khemtemourian, L.; Kleijer, C. C.; Meeldijk, H. J. D.; Jacobs, J.; Verkleij, A. J.; de Kruijff, B.; Killian, J. A.; Hoppener, J. W. M. Membrane Damage by Human Islet Amyloid Polypeptide through Fibril Growth at the Membrane. *Proc. Natl. Acad. Sci.* **2008**, *105* (16), 6033–6038.

(38) Mahul-Mellier, A.-L.; Vercruysse, F.; Maco, B.; Ait-Bouziad, N.; De Roo, M.; Muller, D.; Lashuel, H. A. Fibril Growth and Seeding Capacity Play Key Roles in α-Synuclein-Mediated Apoptotic Cell Death. *Cell Death Differ.* **2015**, *22* (12), 2107–2122.

(39) Gai, W. P.; Yuan, H. X.; Li, X. Q.; Power, J. T. H.; Blumbergs, P. C.; Jensen, P. H. In Situ and in Vitro Study of Colocalization and Segregation of α-Synuclein, Ubiquitin, and Lipids in Lewy Bodies. *Exp. Neurol.* **2000**, *166* (2), 324–333.

(40) Sharon, R.; Bar-Joseph, I.; Frosch, M. P.; Walsh, D. M.; Hamilton, J. A.; Selkoe, D. J. The Formation of Highly Soluble Oligomers of Alpha-Synuclein Is Regulated by Fatty Acids and Enhanced in Parkinson's Disease. *Neuron* **2003**, *37* (4), 583–595.

(41) Dharmadana, D.; Reynolds, N. P.; Conn, C. E.; Valéry, C. Molecular Interactions of Amyloid Nanofibrils with Biological Aggregation Modifiers: Implications for Cytotoxicity Mechanisms and Biomaterial Design. *Interface Focus* **2017**, *7* (4), 20160160.

(42) Nilsson, M. R. Techniques to Study Amyloid Fibril Formation in Vitro. *Methods* **2004**, *34* (1), 151–160.

(43) Cloe, A. L.; Orgel, J. P. R. O.; Sachleben, J. R.; Tycko, R.; Meredith, S. C. The Japanese Mutant Aβ (ΔE22-Aβ 1−39) Forms Fibrils Instantaneously, with Low-Thioflavin T Fluorescence: Seeding of Wild-Type Aβ 1−40 into Atypical Fibrils by ΔE22-Aβ 1−39. *Biochemistry* **2011**, *50* (12), 2026–2039.

(44) Biancalana, M.; Koide, S. Molecular Mechanism of Thioflavin-T Binding to Amyloid Fibrils. *Biochim. Biophys. Acta* **2010**, *1804* (7), 1405–1412.

(45) Groenning, M.; Olsen, L.; van de Weert, M.; Flink, J. M.; Frokjaer, S.; Jørgensen, F. S. Study on the Binding of Thioflavin T to β-Sheet-Rich and Non-β-Sheet Cavities. *J. Struct. Biol.* **2007**, *158* (3), 358–369.

(46) Stefani, M. Structural Features and Cytotoxicity of Amyloid Oligomers: Implications in Alzheimer's Disease and Other Diseases with Amyloid Deposits. *Prog. Neurobiol.* **2012**, *99* (3), 226–245.

(47) Sarroukh, R.; Goormaghtigh, E.; Ruysschaert, J.-M.; Raussens, V. ATR-FTIR: A "Rejuvenated" Tool to Investigate Amyloid Proteins. *Biochim. Biophys. Acta - Biomembr.* **2013**, *1828* (10), 2328–2338.

(48) Zandomeneghi, G.; Krebs, M. R. H.; McCammon, M. G.; Fändrich, M. FTIR Reveals Structural Differences between Native β-Sheet Proteins and Amyloid Fibrils. *Protein Sci.* **2009**, *13* (12), 3314–3321.

(49) Moran, S. D.; Zanni, M. T. How to Get Insight into Amyloid Structure and Formation from Infrared Spectroscopy. *J. Phys. Chem. Lett.* **2014**, *5* (11), 1984–1993.

(50) Goormaghtigh, E.; Cabiaux, V.; Ruysschaert, J. M. Secondary Structure and Dosage of Soluble and Membrane Proteins by Attenuated Total Reflection Fourier-Transform Infrared Spectroscopy on Hydrated Films. *Eur. J. Biochem.* **1990**, *193* (2), 409–420.

(51) O'Nuallain, B.; Thakur, A. K.; Williams, A. D.; Bhattacharyya, A. M.; Chen, S.; Thiagarajan, G.; Wetzel, R. Kinetics and Thermodynamics of Amyloid Assembly Using a High-Performance Liquid Chromatography–Based Sedimentation Assay. *Methods Enzymol.* **2006**, *413*, 34–74.

(52) Chen, S.; Wetzel, R. Solubilization and Disaggregation of Polyglutamine Peptides. *Protein Sci.* **2001**, *10* (4), 887–891.

(53) Jao, S.-C.; Ma, K.; Talafous, J.; Orlando, R.; Zagorski, M. G. Trifluoroacetic Acid Pretreatment Reproducibly Disaggregates the Amyloid β-Peptide. *Amyloid* **1997**, *4* (4), 240–252.

(54) Anthis, N. J.; Clore, G. M. Sequence-Specific Determination of Protein and Peptide Concentrations by Absorbance at 205 Nm. *Protein Sci.* **2013**, *22* (6), 851–858.

(55) Kabsch, W. *XDS*. *Acta Crystallogr. Sect. D Biol. Crystallogr.* **2010**, *66* (2), 125–132.

(56) Rodríguez, D. D.; Grosse, C.; Himmel, S.; González, C.; de Ilarduya, I. M.; Becker, S.; Sheldrick, G. M.; Usón, I. Crystallographic Ab Initio Protein Structure Solution below Atomic Resolution. *Nat. Methods* **2009**, *6* (9), 651–653.

(57) McCoy, A. J.; Grosse-Kunstleve, R. W.; Adams, P. D.; Winn, M. D.; Storoni, L. C.; Read, R. J.; IUCr. *Phaser* Crystallographic Software. *J. Appl. Crystallogr.* **2007**, *40* (4), 658–674.

(58) Murshudov, G. N.; Vagin, A. A.; Dodson, E. J. Refinement of Macromolecular Structures by the Maximum-Likelihood Method. *Acta Crystallogr. Sect. D Biol. Crystallogr.* **1997**, *53* (3), 240–255.

(59) Emsley, P.; Cowtan, K. *Coot*: Model-Building Tools for Molecular Graphics. *Acta Crystallogr. Sect. D Biol. Crystallogr.* **2004**, *60* (12), 2126–2132.

(60) Pettersen, E. F.; Goddard, T. D.; Huang, C. C.; Couch, G. S.; Greenblatt, D. M.; Meng, E. C.; Ferrin, T. E. UCSF Chimera-a Visualization System for Exploratory Research and Analysis. *J. Comput. Chem.* **2004**, *25* (13), 1605–1612.

Supporting Figures

Figure S1. Fibrillation and cytotoxicity of cysteine derivatives of PSMα3

Fibrillation and cytotoxicity of PSMα3 derivatives with cysteine residues added to either their N-terminus (N-cys-PSMα3) or C-terminus (C-cys-PSMα3) as compared to WT PSMα3; sequences are depicted. (**A**) ThT fibrillation kinetic curve showing the fast fibrillation rate of C-cys-PSMα3 compared to WT PSMα3, and almost no ThT binding to N-cys-PSMα3. (**B**) TEM micrographs showing fibrils of C-cys-PSMα3 and no fibril formation for N-cys-PSMα3. (**C**) C-cys-PSMα3 is toxic to T cells while N-cys-PSMα3 showed significantly reduced cytotoxicity. The experiment was performed in triplicates and repeated three times, on different days. A representative graph is presented. Error bars represent standard error of the mean of each triplicate. * $p < 0.05$ and ** $p < 0.001$ were

calculated using the Mann-Whitney U (two tailed) test in comparison to the values of WT

PSMα3, accounting for the data from all three repeats.

Figure S2. Fibrillation kinetics of PSMα3 mutants

ThT binding to PSMα3 derivatives shows the typical fibrillation kinetics, which includes a

lag time followed by aggregation. The peptides (50 µM) were mixed with ThT (200 µM) in

a buffered solution (pH 8). WT PSMα3, as well as D-PSMα3, G16A, and K17A mutants

bound ThT with different kinetics (top left). The inset on the right presents a weaker

binding of ThT to the K9A mutant, but nevertheless clear fibrillation kinetics were

measured. L15A was associated with an especially high ThT signal and its fibrillation curve is therefore depicted separately from the others (bottom). Overall, PSMα3 and five derivatives were shown to bind ThT, while L7A, F10A, F11A, K12A, and D13A did not bind ThT, but did form fibrils visualized by TEM (Figure 1).

Figure S3. Secondary structure analysis of WT PSMα3 mutants in solution

The circular dichroism (CD) spectra of PSMα3 mutants showed that they maintained α-helicity, as depicted by the double minima near 208 nm and 222 nm typical of a α-helical conformation. F8A and F11A exhibited somewhat reduced helicity but were still mainly helical in solution. The concentrations of the peptides were calculated during the measurement to account for the effect of aggregation on the actual concentration.

Figure S4. X-ray fiber diffraction of PSMα3 mutants showed diffuse patterns

X-ray fiber diffraction of WT PSMα3 exhibited a cross-α pattern. The fiber diffraction of the mutants F3A, F8A, D13A, and G16A all showed diffuse fiber diffraction, hindering definitive interpretation of the data, despite repeated attempts. Main reflections, although diffuse, are depicted. Of note, in contrast to WT PSMα3 that formed clear fibrous species on the edge of the capillary when prepared for fibril diffraction, the non-fibrillating mutants, F3A and F8A, did not form any clear fibrous structures; thus, X-ray data were collected from a dried sample that remained on the capillary.

Figure S5. FITC-labeled PSMα3 forms fibrils

The micrograph shows fibrils formed by FITC-labeled PSMα3. FITC-PSMα3 was dissolved in ultra-pure water to a concentration of 10 mM and incubated at RT for 7 days. Fibrils were then separated from the supernatant, diluted to 1mM and deposited on a carbon-coated grid. A 200 nm scale bar is depicted.

Table S1. Calculated distances in the structures of WT PSMα3 and mutants

Peptide		WT-PSMα3	L15A	G16A	K12A
Inter-sheet distances (Å)	**Dry interface**	12.6	13.0	10.2	10.2
	Wet interface	15.6	13.6	13.9	13.9
Inter-helix distance along each sheet (Å)		10.5	10.6	13.1	13.0
Length of α-helical axis		25.5	26.9	25.1	24.9

The dry interface corresponds to the hydrophobic packing between mated sheets. The amphipathic nature of the peptides results in alternating rows of dry and wet interfaces in the crystal packing. In the wet interface, interactions between sheets are mediated by water molecules (not shown). The distances were calculated in Chimera *(1)* using a different method than in our previous report *(2)*, therefore resulting in a slightly different inter-sheet value for the WT PSMα3.

Table S2. Data collection and refinement statistics

	PSMα3 L15A	PSMα3 K12A	PSMα3 G16A
Beamline	ESRF ID30a-3	ESRF ID30a-3	ESRF ID23_2
Date	November 10th 2016	November 10th 2016	September 17th 2016
PDB ID	6GQ5	6GQ2	6GQC
Data collection			
Space group	P1 21 1	C 1 2 1	C 1 2 1
Cell dimensions			
a, b, c (Å)	28.41 10.67 29.60	52.94 13.15 25.97	53.27 13.30 25.80
α, β, γ (°)	90.0 105.9 90.0	90.0 112.3 90.0	90.0 111.9 90.0
Wavelength (Å)	0.96770	0.96770	0.8729
Resolution (Å)	28.469-1.500 (1.54-1.50)	24.496-1.540 (1.58-1.54)	24.711-1.400 (1.44-1.40)
R-factor observed (%)	13.7 (55.9)	11.4 (83.6)	12.4 (75.4)
$^aR_{meas}$ (%)	17.1 (69.2)	14.2 (103.2)	13.5 (82.2)
$I/\sigma I$	4.09 (1.56)	4.83 (1.20)	8.13 (2.28)
Total reflections	7650 (549)	6862 (468)	21920 (1637)
Unique reflections	2854 (207)	2477 (172)	3478 (267)
Completeness (%)	96.3 (95.0)	94.1 (91.5)	99.0 (100.8)
Redundancy	2.68 (2.65)	2.77 (2.7)	6.30 (6.13)
$^bCC1/2$ (%)	98.7 (85.1)	98.3 (77.2)	99.4 (84.9)
Refinement			
Resolution (Å)	17.47-1.50 (1.68-1.50)	14.61-1.54 (1.72-1.54)	14.67-1.40 (1.44-1.40)
Completeness (%)	96.45 (97.42)	94.58 (96.23)	99.29 (100.00)
^{C} No. reflections	2568 (713)	2229 (620)	3129 (238)
$^dR_{work}$ (%)	19.4 (33.4)	20.3 (37.9)	15.1 (23.9)
R_{free} (%)	24.50 (36.1)	23.6 (38.0)	19.3 (26.4)
No. atoms	201	189	199
Protein	182	181	194
Ligand/ion	8	5	3
Water	11	3	2
B-factors			
Protein	17.428	25.961	17.827
Ligand/ion	41.255 (MRD)	85.652 (SO_4)	43.355 (Na^+) 53.659(Cl^-)
Water	23.327	33.584	24.645
R.m.s. deviations			
Bond lengths (Å)	0.019	0.013	0.011
Bond angles (°)	1.692	1.419	1.385
Clash score *(3)*	2.55	0.00	0.00
Molprobity score *(3)*	1.04	0.5	1.03
Molprobity percentile *(3)*	99th percentile	100th percentile	99th percentile
Number of xtals used for scaling	One crystal	One crystal	One crystal

Values in parentheses are for highest-resolution shell.

[a] R-meas is a redundancy-independent R-factor defined in Reference *(4)*.

[b] $CC_{1/2}$ is the percentage of correlation between intensities from random half datasets

(5).

[c] Number of reflections corresponds to the working set.

[d] Rwork corresponds to the working set

References

(1)   Pettersen, E. F.; Goddard, T. D.; Huang, C. C.; Couch, G. S.; Greenblatt, D. M.; Meng, E. C.; Ferrin, T. E. UCSF Chimera-a Visualization System for Exploratory Research and Analysis. *J. Comput. Chem.* **2004**, *25* (13), 1605–1612.
(2)   Tayeb-Fligelman, E.; Tabachnikov, O.; Moshe, A.; Goldshmidt-Tran, O.; Sawaya, M. R.; Coquelle, N.; Colletier, J. P.; Landau, M. The Cytotoxic Staphylococcus Aureus PSMα3 Reveals a Cross-α Amyloid-like Fibril. *Science.* **2017**, *355* (6327), 831–833.
(3)   Chen, V. B.; Arendall, W. B.; Headd, J. J.; Keedy, D. A.; Immormino, R. M.; Kapral, G. J.; Murray, L. W.; Richardson, J. S.; Richardson, D. C. *MolProbity* : All-Atom Structure Validation for Macromolecular Crystallography. *Acta Crystallogr. Sect. D Biol. Crystallogr.* **2010**, *66* (1), 12–21.
(4)   Weiss, M. S.; IUCr. Global Indicators of X-Ray Data Quality. *J. Appl. Crystallogr.* **2001**, *34* (2), 130–135.
(5)   Karplus, P. A.; Diederichs, K. Linking Crystallographic Model and Data Quality. *Science.* **2012**, *336* (6084), 1030–1033.