The Heterotaxy Gene, *CCDC11*, is Essential for Cytokinesis and Cell-Cell Adhesion Via RhoA Regulation

Saurabh Kulkarni[1], Rachel Stephenson[2], Sarah Amalraj[1], Ewelina Betleja[3], James J. Moresco[4], John R. Yates III[4], Moe R. Mahjoub[3,5], Ann Miller[2], Mustafa K. Khokha[1*]

[1]Pediatric Genomics Discovery Program
Department of Pediatrics and Genetics, Yale University School of Medicine, New Haven, CT

[2]Department of Molecular, Cellular and Developmental Biology, University of Michigan, Ann Arbor, MI

[3]Department of Medicine (Nephrology), Washington University in St. Louis, MO

[4]Department of Molecular Medicine, Scripps Research Institute, La Jolla, CA

[5]Department of Cell Biology and Physiology, Washington University in St. Louis, MO

Keywords: congenital heart disease, *Xenopus*, cilia, RhoA, cytokinesis, left-right patterning, mitosis

SUMMARY

RhoA is critical for cytokinesis and cell-cell adhesion. While we have made significant progress in identifying the regulators of RhoA, our understanding remains incomplete. In multiple patients with heterotaxy (Htx), a disorder of left-right (LR) patterning of the internal organs, mutations in CCDC11 were identified. In *Xenopus*, depletion of *ccdc11* causes defects in LR patterning, recapitulating the patients' phenotype. Interestingly, with *ccdc11* depletion, normally monociliated cells of the Left-Right Organizer (LRO) exhibited multiple cilia per cell. While investigating this multiciliated phenotype, we unexpectedly discovered that *ccdc11* is critical for cytokinesis. In addition, CCDC11 depletion alters cell-cell adhesion with loss of junctional adhesion molecules. Based on these phenotypes, reminiscent of RhoA loss of function, we demonstrate that CCDC11 colocalizes with RhoA during cytokinesis and regulates total RhoA protein levels. In conclusion, CCDC11 surprisingly connects cytokinesis and LR patterning via RhoA regulation, providing a potential mechanism for disease pathogenesis.

RESULTS AND DISCUSSION

Ccdc11 is important in establishing left-right asymmetry

The embryo breaks bilateral symmetry beginning at the Left-Right Organizer (LRO) [1-3]. Here, a subset of cells with motile monocilia drive extracellular fluid towards the left, which is sensed by immotile monocilia[4-6]. As a consequence, *dand5* (Cerl2 in mouse, *coco* in frog), which is initially expressed bilaterally at the margin of the LRO, is repressed on the left[2, 7, 8]. Repression of *dand5* leads to expression of *pitx2* in the left lateral plate mesoderm[1, 9, 10]. Pitx2 drives asymmetric organogenesis such that the heart, which initiates as a midline tube, loops to the right while other organs like the gut also undergo asymmetric morphogenesis[11]. Therefore, a signaling cascade including cilia driven extracellular fluid flow, *dand5*, and *pitx2* establish laterality of internal organs such as the heart.

CCDC11 plays a key role in establishing LR asymmetry[12-15]. A number of Htx patients have CCDC11 alleles, and depletion of *ccdc11* causes LR patterning defects in zebrafish and *Xenopus* [12-15]. Consistent with these studies, we also show that *ccdc11* is important for LR patterning in *Xenopus*. We depleted Ccdc11 using F0 CRISPR and examined cardiac looping (Figure 1A). We found that embryos depleted of *ccdc11* showed significant cardiac looping defects (~22%) compared to uninjected controls. In addition, overexpression of human CCDC11 also showed significant cardiac looping defects (~15%) compared to uninjected controls suggesting that precise regulation of CCDC11 levels is necessary for its proper function (Figure 1A). To place CCDC11 in the LR patterning cascade, we next examined *pitx2* and *coco*. Using a *ccdc11* morpholino oligo (MO), we found that *ccdc11* knockdown led to abnormal patterns of both *pitx2* (~25%) and *coco* (~80%) (Figures 1B-C). Therefore, using either depletion strategy, F0 CRISPR or MO, *ccdc11* depletion is critical for global LR patterning consistent with previous studies [12-14].

Previous studies have shown that CCDC11 is important for cilia although the identified mechanisms differ [13, 14]. In the first study, CCDC11 appeared critical for cilia motility, specifically, docking of the outer dynein arm but not ciliogenesis more generally [14]. On the other hand, a second study demonstrated that CCDC11 is critical for ciliogenesis[13]. In that study, CCDC11 interacts with core components of the centriolar satellite proteins that are essential for ciliogenesis. Loss of CCDC11 also causes defective assembly of cilia in multiciliated cells [13]. Based on these studies, we sought to examine cilia morphology in the LRO of *ccdc11* morphants. The LRO is composed of monociliated cells where the cilium is normally ~5-6 microns in length. Using immunofluorescence labeling, we found that cilia in morphants were abnormal compared to WT embryos (Fig. 1D). Specifically, in *ccdc11* morphants, the cilia were shorter (~4 microns) (Figures 1D-E). Thus, consistent with the second study [13], our data suggested that Ccdc11 is important for general cilia structure. Interestingly, when we marked cell boundaries with phalloidin (that labels filamentous actin), we noted two additional striking findings. First, some of the LRO cells were dramatically larger (Fig 1D,F, S1) in Ccdc11 depleted cells compared to wildtype controls. In addition, many of these larger cells had multiple cilia per cell in some cases as many as 10-12 cilia per cell (Figures 1D,G, S1).

CCDC11 regulates cytokinesis

We considered two hypotheses as to why Ccdc11 depletion causes multiciliation in LRO cells: 1) Ccdc11 is important for regulation of centriole numbers or 2) Ccdc11 is important for cell division. Since alterations in cell division would also lead to changes in centriole numbers, we first wanted to eliminate cell division as a possibility. When we labeled nuclei of LRO cells, we surprisingly found that cells with multiple cilia also had multiple nuclei suggesting Ccdc11 was important for cell division, likely cytokinesis (Figure S1). We quantified the multinucleation in Ccdc11 depleted embryos. We labeled nuclei with H2B-RFP and marked the cell membrane with phalloidin. Using a MO to deplete Ccdc11, approximately 30% of cells at stage 10 (early gastrulation) embryos had evidence of multiple nuclei (Figures 2A,B). Using F0 CRISPR, nearly 20% of cells in the epidermis of stage 28 (post neurulation) embryos had evidence of multinucleation (Figures 2C,D).

To visualize the temporal dynamics of cell division, we performed live imaging in *Xenopus* embryos. We labeled F-actin (and the cell membranes) with utrophin-GFP. In wildtype embryos, F-actin accumulated at the contractile ring during cytokinesis. In contrasts, cells depleted of Ccdc11 failed to accumulate F-actin in the contractile ring. Cytokinesis appeared to be initiated properly but stalled and failed to complete (Figure 2E).

To test if CCDC11 is also important for cytokinesis in humans, we depleted CCDC11 using siRNA in human U2OS cells [13]. *CCDC11* siRNA transfected cells showed significantly more multinucleation compared to cells transfected with control siRNA (Figures 2F-G). We confirmed CCDC11 depletion in U2OS cells using western blot (Figure 2H). In order to test the specificity of the antibody, we overexpressed human CCDC11 RNA in

Xenopus embryos. Our antibody does not detect native *Xenopus* Ccdc11 protein but does detect the human CCDC11 protein in a dose dependent manner (Figure 2I).

CCDC11 is important for cell-cell adhesion

While examining multinucleation in *CCDC11* siRNA transfected cells, we noticed that cells did not appear to form proper cell-cell junctions; the cells seemed to "pile-up" on one another. Therefore, we examined adherence junction molecules by immunofluorescence (IF) using antibodies against β-catenin and pan-cadherin (Figure S2)[16]. We quantified the signal intensity by drawing a line from the center of one cell to the next (line scan) (Figure S2A). In control cells, the signal peaked at the cell junction with lower signal intensity in the cytoplasm (Figure S2B). However, in CCDC11 siRNA transfected cells, both β-catenin and pan-cadherin signals were significantly reduced at the cell-cell junction (Figure S2A-E). Our results suggest that CCDC11 is not only important for cytokinesis but also for cell-cell adhesion

CCDC11 localizes to the contractile ring during cytokinesis

CCDC11, which is also known as cilia and flagella associated protein 53 (CFAP53), localizes to the ciliary base[13, 14]. However, given its role in cytokinesis and cell-cell adhesion, we decided to re-examine its localization. We began by examining ciliated cells such as human retinal-pigmented epithelium (RPE) and pig kidney tubular epithelial cells (LLCPK). As expected, we detected CCDC11 at the base of cilia (Figures S3A-D), and consistent with a role in cytokinesis, CCDC11 was also localized to the midbody of recently divided cells (Figures S3A-B). To test antibody specificity, we depleted CCDC11 protein using siRNA in RPE cells and examined its fluorescence signal. siRNA depletion significantly reduced the CCDC11 signal both at the base of the cilium and the midbody demonstrating the specificity of our localization (Figures S4A-B).

While CCDC11 localizes to the midbody, our live imaging of contractile ring ingression suggests that Ccdc11 acts at a much earlier stage in cytokinesis (Figure 2E). Therefore we decided to examine CCDC11 localization using live imaging in the dividing cells of *Xenopus* embryos. To that end, we tagged mNEON to the N-terminal of human CCDC11 and injected this mRNA into *Xenopus* embryos (Figure 3). We found that CCDC11 was not only localized to the midbody but also to the contractile ring of dividing cells. These observations suggest that CCDC11 is expressed at the earliest steps of contractile ring ingression (Figure 3).

The c-terminal of CCDC11 is necessary and sufficient to localize to the contractile ring

CCDC11 appears to be a multifunctional protein that localizes to different structures such as the ciliary base, midbody and contractile ring (Figures S3 and 3)[13, 14]. However, whether any specific domains regulate differential localization is unknown. Because we were interested in understanding the role of CCDC11 in cytokinesis, we sought to identify domains critical for its localization to the midbody and contractile ring. To that end, we made five different deletion constructs fused to mNEON, separating each coiled coil (CC) domain

(total 3 CC domains), Domain 1 (D1: 1-172aa), D2 (173-345 aa), D3 (346-514 aa), D1/D2 (1-345 aa) and D2/D3 (173-514 aa) (Figure 3). We injected the mRNA of each deletion construct and examined mNEON signal in dividing cells. D1, which contains the first CC domain, was weakly localized to the contractile ring towards the end of cytokinesis but was strongly localized to the midbody. D2, on the other hand, failed to localize to either the contractile ring or the midbody. The third CC region strongly localized to the contractile ring but weakly localized to the midbody. The D1/D2 construct localized similarly to D1 and the D2/D3 construct localized similarly to D3 suggesting that D2 is not a repressive nor instructive towards localization of CCDC11. Taken together our results demonstrate that while the first CC domain is necessary and sufficient for localization to the midbody, the third CC domain is necessary and sufficient for localization to the contractile ring during cytokinesis.

CCDC11 co-localizes with RhoA during cytokinesis

To understand how CCDC11 regulates cytokinesis and cell-cell adhesion, we immunoprecipitated CCDC11 (using two different tags, GFP and Myc) and identified proteins using liquid chromatography and mass spectrometry (LC-MS/MS). Interestingly, a significant number of cytoskeleton proteins were highly enriched in our IP samples (Table S1). The gene ontology (GO) analysis identified significant enrichment for processes related to the actin-cytoskeleton organization (6 out of top 10) (Table S2). Further, pathway maps identified processes related to Protein Kinase cAMP-dependent (PKA) and Rho GTPAse mediated cytoskeleton organization and integrin mediated cell adhesion.

Cells depleted of CCDC11 had two main phenotypes in addition to the cilia defects: a cytokinesis failure and loss of cell adhesion molecules. Therefore, we sought to further investigate Rho GTPases as they are master regulators of cytokinesis and cell-cell adhesion. Specifically, different Rho GTPases, such as RhoA, Rac1 and Cdc42, are critical for the initiation, maintenance and function of cell-cell adhesion[17, 18]. On the other hand, their activity becomes more compartmentalized during cytokinesis[17]. While Cdc42 is critical for mitosis, RhoA is critical for cytokinesis[17]. Since we identified defects in cell-cell adhesion and cytokinesis, we decided to investigate a role for CCDC11 in regulating RhoA. We began by examining the localization of CCDC11 in relation to RhoA in U2OS cells. We focused our attention on the contractile ring and midbody where RhoA is enriched. We found that CCDC11 localization completely overlapped with RhoA localization both in the contractile ring and midbody supporting our hypothesis that CCDC11 may be affecting cytokinesis via manipulating RhoA activity (Figure 4A).

CCDC11 is essential for maintaining RhoA protein levels independent of transcription

To understand how CCDC11 might affect RhoA, we examined RhoA expression in the cell membrane and contractile ring in CCDC11 depleted U2OS cells. Using a line scan, we found that CCDC11 depletion led to a significant reduction in RhoA signal at the cell-cell junctions (Figures 4B, 4D). For the contractile ring, we

normalized RhoA signal at the contractile ring to membrane RhoA signal and found that CCDC11 depletion also led to significant depletion of RhoA in the contractile ring (Figures 4C, 4E). Finally, in CCDC11 depleted cells, total RhoA was reduced using Western blot (Figure 4F).

Loss of RhoA protein suggested two possibilities: 1) CCDC11 is regulating the transcription of RhoA or 2) CCDC11 affects the stability of RhoA protein. To examine these possibilities, first we compared RhoA transcript levels between control cells and CCDC11 depleted cells using quantitative PCR. In CCDC11 depleted cells, RhoA transcripts are in fact mildly elevated compared to controls suggesting that CCDC11 does not regulate RhoA transcription but protein stability (Figure 4G).

In summary, our results confirm previous findings that CCDC11 plays a key role in establishing organ laterality, a role conserved from fish to humans[12-15]. As evidenced by abnormal *coco* and *pitx2* expression, cilia mediated signaling is aberrant in *ccdc11* depleted embryos. Previous studies suggested that CCDC11 is vital for ciliogenesis, and our results confirm defects in cilia length[13]. However, we discovered a "multiciliated" cell phenotype in the LRO that we traced to a defect in cytokinesis. We propose that depletion of CCDC11 leads to cytokinesis defects in LRO cells, resulting in multiple nuclei and multiple centrioles. Multiple centrioles give rise to multiple cilia, which are structurally abnormal and have polarity defects. In addition to this, the defect in cilia signaling is amplified by the loss of CCDC11 from the bases of cilia, where it is required for the composition of core components of satellite proteins[13]. Thus, the resulting LR patterning and cilia defects arise from a combination of CCDC11 function both in cytokinesis and ciliary base structure.

A growing number of studies have shown a strong connection between cytokinesis and ciliogenesis, as both processes are actin and microtubule dependent [19-31]. Recently, a study showed that the midbody hosts multiple proteins important for cilia assembly. These proteins travel to the apical surface of a cell near the centrosome to enable cilia formation providing a direct biological mechanism connecting cytokinesis and ciliogenesis [20]. Interestingly, the first coiled coil domain of CCDC11 is required for localization to the centrioles, basal body and midbody[13]. On the other hand, the third coiled coil domain is necessary for the localization to the contractile ring and centriolar satellites[13]. Both results taken together provide some insights into how CCDC11 localization is regulated during the cell cycle. Others have also identified a large number of proteins that localize both to the contractile ring and the base of cilia. In fact, the core cilia proteome and cytokinesis proteins overlap by more than 17 proteins [24]. For example, central spindle proteins essential for cytokinesis like PRC1, MKLP1 and INCENP are directly localized to the bases of cilia [24]. The master regulator of cytokinesis, RhoA, also localizes to the base of the cilium[28-30]. Recently, we and others discovered that the chromatin modifier WDR5 is also localized to the bases of cilia, the spindle microtubules and the midbody [19, 21, 31]. Further, intraflagellar transport proteins necessary for ciliogenesis and cilia maintenance are also critical for cleavage furrow ingression [22, 23, 25-27]. Thus, localization and functional

analysis of CCDC11 at the bases of cilia and at the cleavage furrow further highlights the importance of these multifunctional proteins that are essential in two critical biological processes: cytokinesis and ciliogenesis.

Cytokinesis failure can lead to tetraploidy or aneuploidy and potentially tumorigenesis[32]. In fact defects in centrosome numbers is often a characteristic of cancer cells. For example, cancers of the breast, prostate, bladder, and pancreas are a subset that is often associated with centrosomal abnormalities[33]. Interestingly, CCDC11 is implicated to play a role in cancer, specifically breast and prostate cancer [34, 35]. Further, in a few cases, an association between heterotaxy and cancer has been identified [36-38], however, no clear mechanism has been identified. Our study suggests a mechanism where cytokinesis defects may explain the connection between heterotaxy and cancer.

Finally, our results emphasize the importance of patient driven gene discovery coupled with basic cell and developmental biology to discover new biology. Identification of multiple CCDC11 variants in heterotaxy as well as cancer patients suggests a common mechanism. Indeed, we discovered that the role of CCDC11 in RhoA regulation and cytokinesis could potentially link both heterotaxy and cancer. Importantly, the cilia and cytokinesis phenotypes can have important implications for the patient management.

ACKNOWLEDGMENTS

This project was funded in part by grants from the National Heart, Lung and Blood Institute K99/R00 (5K99HL133606) award to SSK, National Heart, Lung and Blood Institute (R01-HL128370) and National Institutes of Diabetes and Digestive and Kidney Diseases (R01-DK108005) to MRM, National Institute of Child Health and Human Development (R01HD081379) to MKK, National Institute of General Medical Sciences (P41 GM103533 to JRY), NIH grant R01GM112794 to ALM, and NSF Graduate Research Fellowship (DGE #1256260) to RES. MKK is a Mallinckrodt Scholar.

METHODS AND MATERIALS

Animal husbandry:

Xenopus tropicalis were housed and cared for in our aquatics facility according to established protocols that were approved by the Yale Institutional Animal Care and Use Committee (IACUC). *Xenopus laevis* were housed and cared for by the lab of Dr. Ann Miller according to established protocols that were approved by the University of Michigan Animal Care and Use Committee (IACUC).

Cell culture:

Human retinal-pigmented epithelial (RPE) and pig kidney tubular epithelial cells (LLCPK) cells were obtained from Brueckner lab from Yale School of Medicine. U2OS cells were obtained from Carroll lab at Yale School of Medicine. Cells were cultured in DMEM/Ham's F12 media supplemented with 10% fetal bovine serum (FBS).

Microinjection of morpholino oligonucleotides and mRNA in *Xenopus*:

Embryos were produced by *in vitro* fertilization and raised to appropriate stages in 1/9xMR + gentamycin according to established protocols [39, 40]. Staging of *Xenopus* tadpoles was as previously described [41]. Morpholino oligonucleotides or mRNA were injected into one-cell or two-cell embryos as described previously [39]. The morpholino oligonucleotides for *ccdc11* translation blocking was injected at the concentration 7.5-10ng (5'-CATGCTTTCTCCCCAGCCGTGCTGT -3'). The CRISPR for *ccdc11* was injected at the concentration 800pg (5'-AAAGGAGCCGCCCCCGCCTT -3'). Alexa488 (Invitrogen) or green fluorescent protein (100pg) were injected as tracers when required. We generated *in vitro* capped mRNA using the mMessage machine kit (Ambion) and followed the manufacturer's instructions. Full length *Xenopus ccdc11* image clone (IMAGE: 7730599) and human *CCDC11* image clone (IMAGE: 4831081) was subcloned into the pCS2+ vector using PCR amplification and cloning. Full length and mutant constructs for domain analysis were generated using PCR amplification and cloning into the PCS2+ vector with mNEON fluorescent protein. *Xenopus* WT CCDC11 was injected (100pg) at one-cell embryos for overexpression. *Xenopus laevis* embryos were injected with mRNA encoding mNeon-CCDC11 at following concentrations: full length (FL) 412.5 pg, the single domains (D1, D2, D3) 137.5 pg, and the doubles (D1/D2, D2/D3) 275 pg at the one cell stage. H2B-RFP and Utrophin-GFP were injected at 150pg at one cell stage of *X. tropicalis* embryos.

SiRNA in cell culture:

For siRNA experiments, cells were transfected with *CCDC11* siRNA described before in [13] using Lipofectamine RNAiMAX transfection Reagent (Life Technologies) following the manufacturer's instructions. U2OS cells with ~30% confluency were incubated with siRNA mix for 6 hours in the media without FBS. Cells were then transferred to the media supplemented with 10% FBS for 72 h before fixation. RPE cells with ~90% confluency were incubated with siRNA mix overnight in the media without FBS. The next day, cells were transferred to the new media without FBS for 72 hours to grow cilia.

Cardiac looping in *Xenopus*:

Frog embryos at stage 45 were treated with benzocaine and ventrally scored for cardiac looping using a light dissection microscope as previously described [42]. Loop direction is defined by the position of the outflow tract relative to the inflow of the heart: outflow to the right – D loop; outflow to the left – L loop; outflow midline, fails to loop – A loop.

Image analysis:

Images were captured using a Zeiss 710 Live confocal microscope except domain analysis imaging was done using Olympus Fluoview F1000 inverted confocal microscope. Images were processed in Fiji, Image J, or Adobe Photoshop. Quantification of GRP cilia length, cell number and LRO area was done using Velocity software on 3D image stacks. Number of cilia per cell in the *Xenopus* LRO was counted manually. Percent cells with multinucleation in *Xenopus* and U2OS cells were counted manually. Final figures were made in Adobe illustrator.

Quantitative PCR:

RNA expression levels were quantified using real-time quantitative PCR. Total RNA was isolated from 80-90%

confluent U2OS cells treated either with control siRNA or *CCDC11* siRNA using RNaesy Plus Mini kKit (Qiagen). Three replicate RNA isolations were prepared independently for each treatment. Complementary DNA was generated from 3 µg of total RNA by reverse transcription using iScript cDNA synthesis Kit (BIORAD) and IQ Supermix (BIORAD). Three technical replicates were performed. Gene specific primers were designed to amplify 100 – 150 bp fragments. To exclude genomic contamination, primers were designed spanning an intron. GAPDH was used for normalization of all data. Transcript abundance was quantified using the Applied Biosystems 7900HT. Differences in mRNA expression levels were determined using the standard curve method.

Immunofluorescence:

Xenopus tropicalis embryos at stage 16 were fixed in 4% paraformaldehyde/PBS overnight at 4^0C to harvest LROs. Dissected LROs were then assayed for acetylated α-tubulin and phalloidin. U2OS cells were fixed in 4% paraformaldehyde/PBS for 2 hours at room temperature to assay, β-catenin, pan-cadherin with phalloidin and Hoechst. U2OS cells were fixed in 2% TCA (Trichloroacetic acid)/ PBS for 10 minutes and then washed with PBS to assay RhoA. RPE and LLCPK cells were fixed in 100% chilled methanol overnight at -20^0C for γ-tubulin and CCDC11.

Immunoprecipitation and Mass Spectrometry:

For immnoprecipitation of GFP- or Myc-Ccdc11, RPE-1 cells stably expressing GFP-Ccdc11 or Myc-Ccdc11 at near endogenous levels[13] were grown on twenty 150-mm cell culture dishes to ~ 90% confluence, washed in PBS and trypsinized for 5 min in a 37°C incubator. After centrifugation, the cellular pellet (~ 1000-1500 µL packed cell volume) was lysed in ice-cold Lysis Buffer (50 mM HEPES, pH 7.4, 1 mM $MgCl_2$, 1 mM EGTA, 150mM NaCl and 10% glycerol) containing protease inhibitors (Mammalian Protease Arrest, G Biosciences), phosphatase inhibitors (PhosSTOP, Roche), and 1% NP-40 (Sigma) for 30 min on ice. Lysates were cleared by centrifugation at 3,000 g for 15 min, then at 50,000 x *g* for 30 min in an OPTIMA LE-80K ultracentrifuge using SW 41-Ti rotor (Beckman Coulter). Protein concentration of the clarified cellular extract was determined using a Pierce 660 nm Protein Assay kit. Equal amounts of the clarified cellular extract [~ 1 mg/mL] were mixed with 150 µL of affinity resin composed of the following antibodies: 55 µg goat anti-GFP (Rockland), 55 µg control goat IgG (Santa Cruz Biotechnology), 20 µg of rabbit anti-Myc (Sigma), or 20 µg control rabbit IgG (Santa Cruz Biotechnology). All antibodies were covalently coupled to Protein-G or Protein-A Magnetic Beads via imidoester crosslinker (DMP; Pierce) per manufacturers protocol. Samples were incubated for 2 h at 4°C with rotation. The beads were magnetically pelleted and washed five times with ice-cold Lysis Buffer for 5 min each. Protein complexes were eluted in 240 µL of Lysis Buffer.

MudPIT analysis of immunoprecipitated complexes were performed from protein pellets dissolved in buffer (8 M Urea 100 mM Tris pH 8.5) reduced with TCEP (Tris[2-Carboxyethyl]-Phosphine Hydrochloride), and alklyated with chloroacetamide. After dilution of urea to 2 M, proteins were digested with trypsin. Digested peptides were analyzed by LC/LC/MS/MS using an LTQ-Orbitap mass spectrometer. Multidimensional chromatography was performed online [43]. Tandem mass spectra were collected in a data-dependent manner

with up to 10 ms2 scans performed for each initial scan (m/z range 300-2000). The search program Prolucid was used to match data to a human protein database [44]. Peptide identifications were filtered using the DTASelect program requiring two peptides per protein and a protein false discovery rate of 1% [45]. Interactors were selected as proteins that had more than five cumulative spectral counts in the experimental samples and were not detected in the controls.

Statistical analysis:

Sample size (n) is defined in every figure legend. Statistical analysis was done using PRISM and Vassarstats software. Heart looping, *pitx2c* and *dand5* were compared using Chi-square analysis. "Multiciliation" in the LRO cells, multinucleation in *Xenopus* and U2OS cells and CCDC11 localization signal intensity in response to siRNA in RPE cells were also compared using Chi-square analysis. All other comparisons were made using t-tests after confirming the normal distribution of the data. We randomly picked one cell *X. tropicalis* embryos from fertilization as uninjected controls or for morpholino oligo, CRISPR or RNA injections.

Antibodies:

Mouse monoclonal Anti-acetylated α-tubulin (Sigma, T-6793; 1:1000) and Alexa 488 or 647 phalloidin (1:40) were used to mark the ciliary axoneme and F-actin respectively. Mouse monoclonal Anti-γ tubulin (Sigma, T6557; 1:100) was used for immunofluorescence. Rabbit polyclonal Anti-CCDC11 (Sigma, HPA041069) was used 1:100 for immunofluorescence and 1:500 for western blot. Rabbit polyclonal Anti-CCDC11 (Sigma, HPA040595, 1:100) was used for immunofluorescence. Mouse monoclonal Anti-RhoA (Neweast biosciences, 26007) was used 1:50 was used for immunofluorescence and 1:1000 was used for western blot. Rabbit polyclonal Anti β-catenin (Santa Cruz, sc-7199, 1:100) and Rabbit polyclonal Anti pan-cadherin (Abcam, ab-6529, 1:100) was used for immunofluorescence.

REFERENCES

1.	Burdine, R.D., and Schier, A.F. (2000). Conserved and divergent mechanisms in left-right axis formation. Genes & development *14*, 763-776.
2.	Hamada, H., Meno, C., Watanabe, D., and Saijoh, Y. (2002). Establishment of vertebrate left-right asymmetry. Nature reviews. Genetics *3*, 103-113.
3.	Blum, M., Schweickert, A., Vick, P., Wright, C.V., and Danilchik, M.V. (2014). Symmetry breakage in the vertebrate embryo: when does it happen and how does it work? Developmental biology *393*, 109-123.
4.	McGrath, J., Somlo, S., Makova, S., Tian, X., and Brueckner, M. (2003). Two populations of node monocilia initiate left-right asymmetry in the mouse. Cell *114*, 61-73.
5.	Schweickert, A., Weber, T., Beyer, T., Vick, P., Bogusch, S., Feistel, K., and Blum, M. (2007). Cilia-driven leftward flow determines laterality in Xenopus. Current biology : CB *17*, 60-66.
6.	Doerks, T., Copley, R.R., Schultz, J., Ponting, C.P., and Bork, P. (2002). A two-cilia model for vertebrate left-right axis specification. Genome research *12*, 47-56.
7.	Schweickert, A., Vick, P., Getwan, M., Weber, T., Schneider, I., Eberhardt, M., Beyer, T., Pachur, A., and Blum, M. (2010). The nodal inhibitor Coco is a critical target of leftward flow in Xenopus. Current biology : CB *20*, 738-743.
8.	Vonica, A., and Brivanlou, A.H. (2007). The left-right axis is regulated by the interplay of Coco, Xnr1 and derriere in Xenopus embryos. Developmental biology *303*, 281-294.

9.	Ryan, A.K., Blumberg, B., Rodriguez-Esteban, C., Yonei-Tamura, S., Tamura, K., Tsukui, T., de la Pena, J., Sabbagh, W., Greenwald, J., Choe, S., et al. (1998). Pitx2 determines left-right asymmetry of internal organs in vertebrates. Nature *394*, 545-551.

10.	Yoshioka, H., Meno, C., Koshiba, K., Sugihara, M., Itoh, H., Ishimaru, Y., Inoue, T., Ohuchi, H., Semina, E.V., Murray, J.C., et al. (1998). Pitx2, a bicoid-type homeobox gene, is involved in a lefty-signaling pathway in determination of left-right asymmetry. Cell *94*, 299-305.

11.	Lin, C.R., Kioussi, C., O'Connell, S., Briata, P., Szeto, D., Liu, F., Izpisua-Belmonte, J.C., and Rosenfeld, M.G. (1999). Pitx2 regulates lung asymmetry, cardiac positioning and pituitary and tooth morphogenesis. Nature *401*, 279-282.

12.	Gur, M., Cohen, E.B., Genin, O., Fainsod, A., Perles, Z., and Cinnamon, Y. (2017). Roles of the cilium-associated gene CCDC11 in left-right patterning and in laterality disorders in humans. Int J Dev Biol *61*, 267-276.

13.	Silva, E., Betleja, E., John, E., Spear, P., Moresco, J.J., Zhang, S., Yates, J.R., 3rd, Mitchell, B.J., and Mahjoub, M.R. (2016). Ccdc11 is a novel centriolar satellite protein essential for ciliogenesis and establishment of left-right asymmetry. Molecular biology of the cell *27*, 48-63.

14.	Narasimhan, V., Hjeij, R., Vij, S., Loges, N.T., Wallmeier, J., Koerner-Rettberg, C., Werner, C., Thamilselvam, S.K., Boey, A., Choksi, S.P., et al. (2015). Mutations in CCDC11, which encodes a coiled-coil containing ciliary protein, causes situs inversus due to dysmotility of monocilia in the left-right organizer. Human mutation *36*, 307-318.

15.	Perles, Z., Cinnamon, Y., Ta-Shma, A., Shaag, A., Einbinder, T., Rein, A.J., and Elpeleg, O. (2012). A human laterality disorder associated with recessive CCDC11 mutation. Journal of medical genetics *49*, 386-390.

16.	Nelson, W.J. (2008). Regulation of cell-cell adhesion by the cadherin-catenin complex. Biochem Soc Trans *36*, 149-155.

17.	Derksen, P.W.B., and van de Ven, R.A.H. (2017). Shared mechanisms regulate spatiotemporal RhoA-dependent actomyosin contractility during adhesion and cell division. Small GTPases, 1-9.

18.	Evers, E.E., Zondag, G.C., Malliri, A., Price, L.S., ten Klooster, J.P., van der Kammen, R.A., and Collard, J.G. (2000). Rho family proteins in cell adhesion and cell migration. Eur J Cancer *36*, 1269-1274.

19.	Ali, A., Veeranki, S.N., Chinchole, A., and Tyagi, S. (2017). MLL/WDR5 Complex Regulates Kif2A Localization to Ensure Chromosome Congression and Proper Spindle Assembly during Mitosis. Developmental cell *41*, 605-622 e607.

20.	Bernabe-Rubio, M., Andres, G., Casares-Arias, J., Fernandez-Barrera, J., Rangel, L., Reglero-Real, N., Gershlick, D.C., Fernandez, J.J., Millan, J., Correas, I., et al. (2016). Novel role for the midbody in primary ciliogenesis by polarized epithelial cells. The Journal of cell biology *214*, 259-273.

21.	Bailey, J.K., Fields, A.T., Cheng, K., Lee, A., Wagenaar, E., Lagrois, R., Schmidt, B., Xia, B., and Ma, D. (2015). WD Repeat-containing Protein 5 (WDR5) Localizes to the Midbody and Regulates Abscission. The Journal of biological chemistry *290*, 8987-9001.

22.	Borovina, A., and Ciruna, B. (2013). IFT88 plays a cilia- and PCP-independent role in controlling oriented cell divisions during vertebrate embryonic development. Cell reports *5*, 37-43.

23.	Wood, C.R., Wang, Z., Diener, D., Zones, J.M., Rosenbaum, J., and Umen, J.G. (2012). IFT proteins accumulate during cell division and localize to the cleavage furrow in Chlamydomonas. PloS one *7*, e30729.

24.	Smith, K.R., Kieserman, E.K., Wang, P.I., Basten, S.G., Giles, R.H., Marcotte, E.M., and Wallingford, J.B. (2011). A role for central spindle proteins in cilia structure and function. Cytoskeleton (Hoboken) *68*, 112-124.

25.	Delaval, B., Bright, A., Lawson, N.D., and Doxsey, S. (2011). The cilia protein IFT88 is required for spindle orientation in mitosis. Nature cell biology *13*, 461-468.

26.	Pedersen, L.B., and Rosenbaum, J.L. (2008). Intraflagellar transport (IFT) role in ciliary assembly, resorption and signalling. Curr Top Dev Biol *85*, 23-61.

27.	Pedersen, L.B., and Rosenbaum, J.L. (2008). Chapter Two Intraflagellar Transport (IFT). Current Topics in Developmental Biology *85*, 23-61.

28. Park, T.J., Mitchell, B.J., Abitua, P.B., Kintner, C., and Wallingford, J.B. (2008). Dishevelled controls apical docking and planar polarization of basal bodies in ciliated epithelial cells. Nature genetics *40*, 871-879.
29. Pan, J., You, Y., Huang, T., and Brody, S.L. (2007). RhoA-mediated apical actin enrichment is required for ciliogenesis and promoted by Foxj1. Journal of cell science *120*, 1868-1876.
30. Ioannou, A., Santama, N., and Skourides, P.A. (2013). Xenopus laevis nucleotide binding protein 1 (xNubp1) is important for convergent extension movements and controls ciliogenesis via regulation of the actin cytoskeleton. Developmental biology *380*, 243-258.
31. Kulkarni, S.S., Griffin, J.N., Date, P.P., Liem, K.F., Jr., and Khokha, M.K. (2018). WDR5 Stabilizes Actin Architecture to Promote Multiciliated Cell Formation. Developmental cell *46*, 595-610 e593.
32. Sagona, A.P., and Stenmark, H. (2010). Cytokinesis and cancer. FEBS letters *584*, 2652-2661.
33. Godinho, S.A., and Pellman, D. (2014). Causes and consequences of centrosome abnormalities in cancer. Philos Trans R Soc Lond B Biol Sci *369*.
34. Jaiswal, R., Jauhari, S., and Rizvi, S.A.M. (2017). WDR88, CCDC11, and ARPP21 genes indulge profoundly in the desmoplastic retort to prostate and breast cancer metastasis. bioRxiv.
35. Yamamoto, F., and Yamamoto, M. (2007). Scanning copy number and gene expression on the 18q21-qter chromosomal region by the systematic multiplex PCR and reverse transcription-PCR methods. Electrophoresis *28*, 1882-1895.
36. Pagkratis, S., Kryeziu, S., Lin, M., Hoque, S., Bucobo, J.C., Buscaglia, J.M., Georgakis, G.V., Sasson, A.R., and Kim, J. (2017). Case report of intestinal non-rotation, heterotaxy, and polysplenia in a patient with pancreatic cancer. Medicine (Baltimore) *96*, e8599.
37. Park, S.S., Min, B.W., Kim, W.B., Han, H.J., Yong, H.S., Kim, S.J., Kim, C.S., and Mok, Y.J. (2005). Double cancer of the stomach and oesophagus with situs ambiguus with polysplenia: the importance of preoperative evaluation. Dig Liver Dis *37*, 799-802.
38. Iwamura, T., Shibata, N., Haraguchi, Y., Hisashi, Y., Nishikawa, T., Yamada, H., Hayashi, T., and Toyoda, K. (2001). Synchronous double cancer of the stomach and rectum with situs inversus totalis and polysplenia syndrome. J Clin Gastroenterol *33*, 148-153.
39. Khokha, M.K., Chung, C., Bustamante, E.L., Gaw, L.W., Trott, K.A., Yeh, J., Lim, N., Lin, J.C., Taverner, N., Amaya, E., et al. (2002). Techniques and probes for the study of Xenopus tropicalis development. Developmental dynamics : an official publication of the American Association of Anatomists *225*, 499-510.
40. del Viso, F., and Khokha, M. (2012). Generating diploid embryos from Xenopus tropicalis. Methods Mol Biol *917*, 33-41.
41. Nieuwkoop, P.D.F., J. (1994). Normal Table of Xenopus Laevis (Daudin), 1st Edition.
42. Boskovski, M.T., Yuan, S., Pedersen, N.B., Goth, C.K., Makova, S., Clausen, H., Brueckner, M., and Khokha, M.K. (2013). The heterotaxy gene GALNT11 glycosylates Notch to orchestrate cilia type and laterality. Nature *504*, 456-459.
43. MacCoss, M.J., McDonald, W.H., Saraf, A., Sadygov, R., Clark, J.M., Tasto, J.J., Gould, K.L., Wolters, D., Washburn, M., Weiss, A., et al. (2002). Shotgun identification of protein modifications from protein complexes and lens tissue. Proceedings of the National Academy of Sciences of the United States of America *99*, 7900-7905.
44. Xu, T.P., S. K.; Venable, J. D.; Wohlschlegel, J. A.; Diedrich, J. K.; Cociorva, D.; Lu, B.; Liao, L.; Hewel, J.; Han, X.; Wong, C. C. L.; Fonslow, B.; Delahunty, C.; Gao, Y.; Shah, H.; Yates, J. R., III (2015). ProLuCID: An improved SEQUEST-like algorithm with enhanced sensitivity and specificity. Journal of Proteomics *129*, 16-24.
45. Tabb, D.L., McDonald, W.H., and Yates, J.R., 3rd (2002). DTASelect and Contrast: tools for assembling and comparing protein identifications from shotgun proteomics. J Proteome Res *1*, 21-26.

FIGURE LEGEND

Figure 1: Ccdc11 depletion alters left-right patterning and cilia morphology in *Xenopus tropicalis*

(A, B, C) Percentage of embryos that have abnormal (A) looping of the outflow tube, (B) *pitx2,* or (C) *coco* expression. Embryos were injected at the one-cell stage with *ccdc11* CRISPR, *Xenopus ccdc11* RNA, or *ccdc11* MO. L, outflow tube loops towards left; A, outflow tube fails to loop; Bilateral: signal present on both sides; Absent, signal absent on both sides; Reduced: signal reduced on both sides; Right only: signal is present on only right side, Left>Right: Signal is more on the left side. Number of embryos used is mentioned on the graph.

(D) Cilia in control and *ccdc11* morphant *Xenopus* LROs marked by anti-acetylated α-tubulin (red) and F-actin marked by phalloidin (green).

(E) Length of cilia between uninjected control and *ccdc11* morphant *Xenopus* LROs. Data is presented as box plot with 95% confidence interval. Number of embryos used is mentioned on the graph.

(F) Average cell area between uninjected control and *ccdc11* morphant *Xenopus* LROs. Data is presented as box plot with 95% confidence interval. Number of embryos used is mentioned on the graph.

(G) Percent cells with multiple cilia between uninjected controls and *ccdc11* morphant *Xenopus* LROs. Each experiment was repeated three times. Number of embryos used is mentioned on the graph. ★ = P < 0.001.

Figure 2: CCDC11 depletion affects cytokinesis in *Xenopus* and human cells

(A, B, C, D) Percentage of cells with multiple nuclei in response to depletion of Ccdc11 by (A, B) *ccdc11* morpholino, or (C, D) *ccdc11* CRISPR. *Ccdc11* morpholino or CRIPSR was injected in embryos at one cell stage. Number of cells assayed is mentioned on the graph.

(E) Live imaging of *Xenopus* embryo at Nieuwkoop and Faber stage 10 (pre-gastrulation) to show that cytokinesis is affected during contractile ring ingression in *ccdc11* morphant.

(F, G) Percentage of cells with multiple nuclei in response to CCDC11 depletion by siRNA. Number of cells assayed is mentioned on the graph.

(H) Western blot showing loss of CCDC11 protein in human U2OS cells transfected with *CCDC11* siRNA.

(I) Western blot showing the antibody does not detect native *Xenopus* Ccdc11 protein but can specifically detect human CCDC11 microinjected in *Xenopus* embryos in a dose dependent manner. Each experiment was repeated three times. ★ = P < 0.001.

Figure 3: Domains important for Ccdc11 localization to the contractile ring and midbody during cytokinesis.

Live imaging of mNeon-CCDC11 and fragments in blastomeres of *Xenopus laevis* embryos (Nieuwkoop and Faber stage 5-7). Time 0 is the time membrane ingression is first detectable. Arrowheads indicate the position of the contractile ring. Green arrowheads indicate strong accumulation of CCDC11 at the contractile ring, while white arrowheads indicate weak CCDC11 accumulation at the contractile ring. Yellow arrowheads indicate

accumulation of CCDC11 at the midbody. Scale bars are 100 µm. Full length CCDC11 localizes to the cleavage furrow prior to ingression and the midbody following cytokinesis. D1 of CCDC11 accumulates weakly at the contractile ring and strongly at the midbody. D2 accumulates weakly at the contractile ring. D3 accumulates strongly at the contractile ring. D1/D2 accumulates weakly at the contractile ring, indicating that D3 is necessary and sufficient for strong CR localization. D2/D3 accumulates strongly at the contractile ring and at the midbody, indicating that while D1 is sufficient for contractile ring localization, it is not required.

Figure 4: Ccdc11 co-localizes with RhoA to the contractile ring and is important for RhoA protein stability

(A) CCDC11 co-localizes with RhoA in the contractile ring and midbody during cytokinesis in U2OS cells.

(B, C, D, E) CCDC11 depletion affects RhoA accumulation at (B, D) cell-cell junction, and (C, E) the contractile ring in the U2OS cells. Number of cells assayed is mentioned on the graph.

(F) Western blot showing loss of RhoA protein in human U2OS cells transfected with *CCDC11* siRNA.

(G) Quantitative PCR data showing fold change mRNA levels of RhoA in *CCDC11* siRNA compared to controls. Each experiment was repeated three times. ★ = $P < 0.001$.

Figure 1

Ccdc11 depletion affects left-right patterning

A — Heart looping

% embryos (y-axis: 0, 4, 8, 12, 16, 20, 24)

Legend: A loop (red), L loop (blue)

- Control: 133
- *ccdc11* CRISPR: 73 ★
- *ccdc11* RNA: 183 ★

B — *pitx2* expression

% embryos (y-axis: 0, 5, 10, 15, 20, 25)

Legend: Right only (blue), Absent (red), Bilateral (green)

- Control: 218
- *ccdc11* MO: 169 ★

C — *coco* expression

(y-axis: 0, 10, 20, 30, 40, 50, 60, 70, 80, 90)

Legend: Left > Right (green), Bilateral (blue), Absent (red)

- Control: 106
- *ccdc11* MO: 46 ★

Ccdc11 depletion affects cilia in the left-right organizer

D — Ac. tubulin (red), Phalloidin (green)

Control | *ccdc11* MO

50 μm scale bars

E

Cilia length (uM) (y-axis: 2, 4, 6)

- Control: 23
- *ccdc11* MO: 24 ★

F

Area/cell (a.u.) (y-axis: 100, 200, 300, 400, 500)

- Control: 23
- *ccdc11* MO: 22 ★

G — Cilia/cell

% cells (y-axis: 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10)

Legend: 6-12 cilia (magenta), 5 cilia (green), 4 cilia (red), 3 cilia (blue)

- Control: 4628
- *ccdc11* MO: 3605 ★

Figure 2

Figure 3

Figure 4