Affiliations:

CR: School of Biological Sciences, The University of Queensland, St Lucia QLD, Australia;
c.riginos@uq.edu.au
KH: School of Biological Sciences, The University of Queensland, St Lucia QLD, Australia;
k.hock1@uq.edu.au
AMM: School of Biological Sciences, The University of Queensland, St Lucia QLD, Australia;
Institute of Biology, University of the Philippines Diliman 1101, Philippines;
ambrocio.matias@uqconnect.edu.au
PJM: School of Biological Sciences, The University of Queensland, St Lucia QLD, Australia;
p.j.mumby@uq.edu.au
MJHvO: School of BioSciences, The University of Melbourne, VIC 3010, Australia; Australian
Institute for Marine Sciences, Cape Cleveland QLD 4810, Australia;
madeleine.van@unimelb.edu.au
VL: ARC Centre of Excellence for Coral Reef Studies, James Cook University, Townsville
QLD 4811, Australia; vimoksalehi.lukoschek@jcu.edu.au

Abstract

Aim: Widespread coral bleaching, crown-of-thorns seastar outbreaks, and tropical storms all threaten foundational coral species of the Great Barrier Reef, with impacts differing over time and space. Yet, dispersal via larval propagules could aid reef recovery by supplying new settlers and enabling the spread of adaptive variation among regions. Documenting and predicting spatial connections arising from planktonic larval dispersal in marine species, however, remains a formidable challenge.

Location: The Great Barrier Reef, Australia

Methods: Contemporary biophysical larval dispersal models were used to predict long-distance multigenerational connections for two common and foundational coral species (*Acropora tenuis* and *Acropora millepora*). Spatially extensive genetic surveys allowed us to infer signatures of asymmetric dispersal for these species and evaluate concordance against expectations from biophysical models using coalescent genetic simulations, directions of inferred gene flow, and spatial eigenvector modelling.

Results: At long distances, biophysical models predicted a preponderance of north to south connections and genetic results matched these expectations: coalescent genetic simulations rejected an alternative scenario of historical isolation; the strongest signals of inferred gene flow were from north to south; and asymmetric eigenvectors derived from north to south connections in the biophysical models were significantly better predictors of spatial genetic patterns than eigenvectors derived from symmetric null spatial models.

Main conclusions: Results are consistent with biophysical dispersal models yielding approximate summaries of past multigenerational gene flow conditioned upon directionality of connections. For *A. tenuis* and *A. millepora,* northern and central reefs have been important sources to downstream southern reefs over the recent evolutionary past and should continue to provide southward gene flow. Endemic genetic diversity of southern reefs suggests substantial local recruitment and lack of long distance gene flow from south to north.

Key words: Approximate Bayesian Computation, asymmetric eigenvector modelling, connectivity, gene flow, marine landscape, Moran eigenvector modelling, oceanography, planktonic dispersal, seascape genetics

Introduction

Recurrent mass bleaching events on the Great Barrier Reef (GBR) have been increasing in severity and extent (Hughes et al., 2017) against a backdrop of multidecadal coral decline arising from tropical storms, crown-of-thorns seastar (COTS) predation, terrestrial runoff, and fishing pressure (De'ath, Fabricius, Sweatman, & Puotinen, 2012). Locations of severe impacts, however, differ over time. For example, northern reefs have been less affected by water quality problems and COTS outbreaks (De'ath et al., 2012), but yet the highest rates of bleaching in 2016 (the most recent and extensive mass bleaching event) were reported from these reefs (with declines of up to 60% of coral cover: Hughes et al., 2018). Within the GBR, central reefs have been most affected by terrestrial runoff, episodic COTS outbreaks (Pratchett, Caballes, Rivera-Posada, & Sweatman, 2014), tropical storms, and bleaching (De'ath et al., 2012). Southern reefs also have been subject to COTS outbreaks and tropical storms (De'ath et al., 2012), yet largely escaped bleaching in 2016 (Hughes et al., 2018).

Corals, like most benthic marine animals, have planktonic larvae potentially capable of extensive dispersal. External supplies of settlers can replenish populations; for example, following local extirpation of mature *Acropora* colonies by Cyclone Yasi, recruitment of juvenile *Acropora* was high (Lukoschek, Cross, Torda, Zimmerman, & Willis, 2013). Dispersal connections arising from planktonic larval movements can also enable the spread of adaptive variation. With temperatures and extreme heating events projected to increase in frequency (Wolff, Mumby, Devlin, & Anthony, 2018), resolving the GBR-wide capacity for adaptive gene flow (especially involving loci contributing to heat tolerance) will contribute to emerging debates regarding assisted migration and genetic rescue (Anthony et al., 2017). Of particular interest is to uncover routes of natural connections as well as pathways resistant to gene exchange.

Unfortunately, dynamics of planktonic larval dispersal over space and time remain poorly resolved (Carr et al., 2003; Kritzer & Sale, 2004) especially considering an extensive and geographically complex seascape such as the GBR. Currents and other oceanographic phenomena are inherently dynamic such that dispersal-mediated connections among populations are difficult to predict and vary by place and time (Cowen & Sponaugle, 2009; Gaggiotti, 2017; Liggins, Treml, & Riginos, 2013; Watson, Kendall, Seigel, & Mitarai, 2012). Yet, knowledge regarding the sources and destinations of dispersive larvae underpin fundamental

ecological and evolutionary dynamics and inform optimal management of marine resources

for fishing and the protection of key habitats (Beger et al., 2010; Gaines, White, Carr, &

Palumbi, 2010; Hock et al., 2017; Krueck et al., 2017).

Coupled biological-physical models (Cowen & Sponaugle, 2009; Werner & Cowen, 2007)

relying on simulations based on species attributes and physical oceanography are

increasingly being used to predict spatial and temporal aspects of planktonic larval dispersal.

The flexibility of both spatio-temporal scale and outputs makes biophysical models extremely

well-suited for alignment against various sources of empirical dispersal data (Cowen &

Sponaugle, 2009; Jones, 2015; Kool, Moilanen, & Treml, 2013; Liggins et al., 2013) and for

generating detailed spatial predictions useful for management assessments (Hock et al., 2017;

as in 2016; Krueck et al., 2017).

The extent to which biophysical models accurately approximate real dispersal phenomena,

however, remains an open question. As models of open natural systems, strict validation or

rejection of biophysical dispersal models is not feasible (Oreskes, Shrader-Frechette, & Belitz,

1994). Rather, such models can be evaluated against empirical data whereby alignment of the

model and empirical data confers increased confidence that both the model and empirically-

derived statistics describe the phenomenon of interest, in this case dispersal of planktonic

larvae. Geographic surveys of intraspecific genetic variation can provide important insights

regarding dispersal (Hellberg, 2009; Riginos & Liggins, 2013; Selkoe et al., 2016a) and indeed

several studies have considered predictions arising from biophysical dispersal models

alongside observed spatial genetic patterns, typically for coastal marine taxa. Although some

individual studies report correlations (Benestan et al., 2016; Dalongeville et al., 2018; Foster

et al., 2012; Galindo, Olson, & Palumbi, 2006; Schunter et al., 2011; Thomas et al., 2015;

Truelove et al., 2016; White et al., 2010; Xuereb et al., 2018), a recent review of the field found

only moderate to low concordance between biophysical predictions and empirical genetic

patterns (Selkoe, Scribner, & Galindo, 2016b).

There are many reasons why biophysical dispersal models and genetic data may not align.

First, biophysical models typically focus solely on dispersal and do not consider differential

recruitment and/or reproductive success of settlers (Pineda, Hare, & Sponaugle, 2007).

Second, a biophysical model might not sufficiently account for important biological attributes

of larvae or complex near shore oceanography, which is notoriously difficult to parameterise

(Metaxas & Saunders, 2009; Pineda et al., 2007; Werner & Cowen, 2007). Third, there is a

mismatch in time scales, where biophysical models draw upon oceanographic information

collected within recent years or decades whereas genetic inferences, especially those based

on population allele frequency differences, arise from long term processes (thousands of

years and longer). Fourth, population allele frequencies do not solely reflect gene flow

resulting from dispersal but are also shaped by changes in population sizes, range expansions,

colonization and so forth (Whitlock & McCauley, 1999). Lastly, although biophysical models

typically predict asymmetric directionality of dispersal, typical descriptors of genetic diversity

are symmetric so that statistical comparisons between biophysical models and empirical

genetic data nearly exclusively rely on transforming modelled dispersal predictions into

symmetric measures, greatly reducing the information content (Kool et al., 2013; Riginos,

Crandall, Liggins, Bongaerts, & Treml, 2016).

Only a few studies have attempted to ascertain whether empirical genetic patterns are

specifically consistent with the asymmetric dispersal processes inherent to biophysically

derived dispersal predictions. For example, migration matrices derived from biophysical

modelling were used to inform gene flow in forward population genetic simulations (Galindo

et al., 2006) and to analytically predict genetic differentiation based on forward matrix

projections (Kool, Paris, Andréfouët, & Cowen, 2010; Kool, Paris, Barber, & Cowen, 2011),

yielding outcomes that qualitatively matched empirical genetic patterns. For the Caribbean

coral *Orbicellea annularis*, genetic distances derived from a forward matrix project were

shown to be well correlated (ρ = 0.49) to empirical genetic distance estimates (Foster et al.,

2012). Another strategy draws upon historical demographic simulations to quantify

migration values consistent with empirically observed allele frequency spectra or DNA

sequences. For example, Matz *et al.* (2018) documented correlation between population

pairwise estimates of migration derived from allele frequency spectra against biophysically

derived migration probabilities for five populations for the coral *Acropora millepora* from the

Great Barrier Reef (GBR) (Mantel R = 0.58, P = 0.05), with a preponderance of southward

migration consistent between both measures. Comparing all sampled populations holistically,

Crandall *et al.* (2012) constructed a series of coalescent models of gene flow and contrasted

the likelihood of observed DNA haplotype distributions from the biophysically-informed gene

flow models against various other null geographic models; they demonstrated greater

likelihood of biophysically informed gene flow models across three species of nerite snails in

the South Pacific. Recently, three studies turned to asymmetric eigenvector mapping (AEM: Blanchet, Legendre, Maranger, Monti, & Pepin, 2011) where asymmetric processes (such as biophysical migration probabilities) are statistically modelled as spatial autocorrelation structures; inclusion of AEMs substantially improved predictions of spatial genetic structure for American lobster (Benestan et al., 2016), California sea cucumbers (Xuereb et al., 2018), and Mediterranean striped red mullet (Dalongeville et al., 2018). Thus, these first few studies that quantitatively incorporate asymmetric biophysical predictions suggest that directions of larval dispersal are important elements of marine population connectivity.

Here, we comprehensively assess the alignment between biophysical models of dispersal and observed spatial genetic patterns for two common broadcast spawning coral species on the GBR, *Acropora tenuis* and *Acropora millepora*, drawing upon methods based on historical demographic simulations, patterns of shared alleles, and AEM spatial autocorrelation structures. We capitalise upon spatially rich genetic data sets for the two species (Lukoschek, Riginos, & van Oppen, 2016; van Oppen, Peplow, Kinnimonth, & Berkelmans, 2011) with sampling encompassing most of the 2300 km extent of the GBR. Oceanographic patterns suggest that directional gene flow is likely for GBR species such as *A. tenuis* and *A. millepora* (Fig. 1a).

Although present-day oceanography implies that larval dispersal can create connections among distant reefs, several species show differentiation between central/northern vs. southern GBR locations (reviewed by van Oppen et al., 2011), indicative of possible historical vicariance associated with past sea level changes. This divergence is most notable for the spiny chromis damselfish for which genetically distinct colour morphs abut at the north end of Hydrographer's Channel (Planes & Doherty, 1997). Recent geological investigations, however, have identified reef structures along the Australian shelf edge that may have provided coral reef habitat during intermediate and low sea level stands (Abbey, Webster, & Beaman, 2011; Hinestrosa, Webster, & Beaman, 2016), implying that reef species may have shifted their range margins through multiple glacial cycles following the depth contours of available coastline but maintaining their approximate latitudinal positions.

No single approach can simultaneously infer modern and historical gene flow regimes across numerous sites spanning large spatial scales. Rather than discounting historical influences (as

is implicit in many spatial genetic methods), we use coalescent demographic simulations to

evaluate competing scenarios of long-standing gene flow vs. late Quaternary divergence. The

resultant affirmation of gene flow-dominated demography along the GBR justifies subsequent

frequentist analyses focusing on directional gene flow inferred from population allele

frequencies and using all sampled reefs. Treating the biophysical models as hypotheses of

spatial connections among GBR reefs for *Acropora* corals, we test whether projected

asymmetric directions and dispersal strengths are superior predictors of spatial genetic

patterns as compared to simple symmetric null predictors. In addition, we examine the spatial

scales and regions for which biophysical predictors best align to observed genetic patterns.

This study provides a framework for aligning spatially rich population genetic data against *a*

priori predictions of asymmetric dispersal and represents the most comprehensive analysis of

asymmetric gene flow along the full length of the GBR to date.

METHODS

Multi-step connectivity pathways based on larval dispersal models

Connectivity modelling follows protocols described in detail by Hock et al. (in prep) with an

overview in the supplementary materials. The resultant species-specific connectivity

networks (based on all 3806 GBR reefs) were used to predict multistep paths between the

reefs from which the genetic samples were obtained, focusing on 1) the minimum number of

stepping-stones between two reefs (Dijkstra's 1959), 2) the *maximum flow* capacities of links

necessary to connect the two nodes in a network (Ford & Fulkerson, 1956; Boykov &

Kolmogorov, 2004), and 3) the *reliable path* represented by the maximum product of link

weights representing the greatest chance of (direct or multi-step) larval exchange (Hock &

Mumby, 2015). See supplemental methods and Sup. Fig. 1 for details and further explanation.

Genetic data

We capitalise on the extensive microsatellite datasets for *Acropora tenuis* (Lukoschek et al.,

2016) and *Acropora millepora* (van Oppen et al., 2011). Some populations were merged or

omitted to match the larval dispersal model, some individuals with high missing data were

excluded, and conformity to Hardy-Weinberg expectations and linkage equilibria were

verified. See supplemental methods.

Resolving historical influences using coalescent-based ABC

If allele frequencies among *Acropora* populations have been strongly influenced by past

interruptions to gene flow, then we would need to evaluate observed geographic

differentiation in light of such a divergence history. In contrast, if gene flow has dominated the

modern distributions of allelic diversity then dispersal interpretations based on observed

allele frequencies are reasonable. We used an approximate Bayesian computation (ABC)

approach to test the competing hypotheses of historical divergence vs. stepping-stone gene

flow, with the location of historical divergence set at the northern opening of Hydrographer's

Passage (Fig. 1). Owing to the high computational requirement in simulating genetic data to

match our full datasets, we used representative populations for our ABC analyses (Fig. 2). A

custom R script drew parameter values from the prior distributions, which were used with

fastSimcoal2 v2.6.0.3 (Excoffier, Dupanloup, Huerta-Sánchez, Sousa, & Foll, 2013) to generate

a total of 500,000 simulations, which were summarised using statistics calculated by Arlequin

(arlsumstat v3.5.2.2: Excoffier & Lischer, 2010). Details of simulation conditions, model

support, and cross-validation procedures are in the supplementary methods.

Asymmetric gene flow and larval dispersal connectivity

To estimate directionality of gene flow we use information from semi private alleles between

pairs of populations (Sundqvist, Keenan, Zackrisson, Prodöhl, & Kleinhans, 2016) as

implemented by divMigrateOnline (https://popgen.shinyapps.io/divMigrate-online/) using

the G_{ST} option (with no threshold for filtering out non-significant values). We assessed the

correlation between these asymmetric (directional) estimates of gene flow with asymmetric

measures of connectivity derived from the larval dispersal model (described above).

Significances of the correlations were evaluated using 10,000 permutations. Maximum flow

and reliable path metrics were (natural) log transformed for these and subsequent

evaluations.

Spatial eigenvector mapping

Although isolation-by-distance approaches have long been applied in spatial genetic contexts,

they fail to account for spatial autocorrelation structures. Moran Eigenvector Maps (MEMs)

model orthogonal spatial structures and, when combined with multivariate analyses, can test

for spatial regression (Borcard & Legendre, 2002; Dray, Legendre, & Peres-Neto, 2006) and

are well-suited for genetic investigations (Diniz-Filho, Nabout, Telles, Soares, & Rangel, 2009).

Asymmetric eigenvector maps (AEMs) extend spatial modelling to describe spatial structures

arising from directional processes such as ocean currents (Blanchet et al., 2011; Blanchet, Legendre, & Borcard, 2008). Here we use AEMs derived from the biophysical model to test whether asymmetric (directional) connectivity distances predict genetic structure, with AEM model fits contrasted to null symmetric expectations described by MEMs. Creation of link structures, weighting schemes, and connexion diagrams (Blanchet et al., 2008; 2011) are detailed in the supplemental methods and Supp. Fig. 1. Observed population allele counts were chi-square transformed (somewhat upweighting rare alleles: Legendre & Gallagher, 2001) to form the response variables for MEM and AEM analyses. Selection of MEM and AEM model was undertaken with forward model selection based on adjusted R^2 values following Dray *et al.* 2006 and Blanchet *et al.* 2011. Significance values of individual AEMs for the final model were determined using redundancy analysis.

RESULTS

Larval dispersal model

Larval dispersal models for *Acropora tenuis* and *Acropora millepora* suggest that GBR populations are well connected when multi-step (i.e. multigenerational) connections are considered. Overall southward connections were more prevalent, especially long-distance connections: this can be visualised by vectored arcs curving the right in Fig. 1B and C along with greater dispersal probabilities for southwards movements (Fig. 1 D and E). However, some strong northward connections were also present (vectored arcs curving the left) especially among central reefs and some central to northern reefs. All three multistep connectivity metrics considered were highly correlated ($\rho \geq 0.86$ for *A. tenuis*; $\rho \geq 0.80$ for *A. millepora*) with the greatest correlations between stepping stone distance and reliable paths ($\rho \geq 0.98$ for both species). Given these high correlations, we present results primarily for the reliable path metric, which arguably best aligns with biological intuitions regarding population connectivity across time (Hock & Mumby, 2015).

Resolving historical influences using coalescent-based ABC

In the present study, it was necessary to verify that gene flow has shaped *Acropora* spatial genetic patterns and exclude vicariance as an alternative scenario before proceeding with analyses that implicitly ignore divergence. Indeed, ABC analyses yielded higher support for the gene flow only stepping-stone model (*A. tenuis*: posterior probability, PP= 0.999; Bayes Factor, BF = 1210; *A. millepora*: PP= 0.647; BF = 3) compared to a model of historical

divergence vicariance. Applying model selection with the pseudo-observed data (POD)

yielded high accuracy (> 99 ; proportion of POD that were correctly supported) and high

robustness (> 99 with thresholds above 0.6 PP). Overall, these results indicate strong support

(sensu Roux et al., 2016) for the gene flow only stepping-stone model for both species.

Asymmetric gene flow and larval dispersal connectivity

For *A. tenuis*, all directional predictors of connectivity yielded significant correlations with

relative gene flow estimates from divMigrate (ρ values for stepping stone distance: -0.39;

maximum flow: -0.31; reliable path: -0.37; in all cases P << 0.001). For *A. millepora*, reliable

path (ρ = -0.16; P < 0.005) and stepping stone distances (ρ = - 0.13; P < 0.03) were

significantly correlated with relative gene flow but maximum flow was not (ρ = -0.06; NS).

For both species, inspecting the relationship between reliable path probabilities and gene

flow (Fig. 3) shows that for populations predicted to be well connected by the larval dispersal

model (i.e. P ≥ 10^{-20}) estimates of relative gene flow were highly variable (~0-1), whereas

sites predicted to require many stepping stone connections (P ≤ 10^{-40}) had consistently lower

gene flow estimates. This pattern was stronger in *A. tenuis* than *A. millepora*.

Spatial eigenvector mapping

Directional spatial autocorrelation as assessed by AEMs explained a greater proportion of

variance in allele frequencies for both species (Table 1), where the best models were based on

alongshore north to south movements with inshore connections from the Swains to Keppel

Island and to the Capricorn-Bunker group (for *A. tenuis*, R^2 = 0.37 with 9 AEMs retained, and

for *A. millepora*, R^2 = 0.27 with 5 AEMs retained). There was no consistency in terms of best-

performing link weighting (i.e. stepping stone connections, maximum flow, and reliable path)

and notably the binary weighting (presence vs. absence, no weighting by distance) for north

to south connections in *A. millepora* was the best performing, indicating that simply

recognising strong connections (i.e. those with reliable path probabilities in the top 50th

percentile) yields good approximations of spatial genetic patterns. Fig. 4 depicts the highest

scoring AEMs for *A. tenuis* where AEM 1 (3.5% of allele frequency variance; P = 0.001) as the

largest scale of positive autocorrelation showed a gradient across the entire sampled region of

the GBR and AEM 5 (2.3% of variance; P = 0.001) described more local scale autocorrelation

patterns. For *A. tenuis*, AEMs 1, 5, 17, 3, 2, and 12 were individually significant at a P = 0.05

threshold based on an RDA of the final model where higher numbers (i.e. AEM17) indicate

finer spatial grain sizes out of a total of 19 possible AEMs of positive spatial autocorrelation.

For distance-based *A. tenuis* MEM models, the best model (PCNM) retained MEMs 4, 15, 35, 22

indicating a mix of large- and fine-scale autocorrelations. Similar results were found for *A.*

millepora (Supp Fig. 3) with AEM 1 describing the greatest amount of spatial variance in allele

frequencies (5.9%, P = 0.003) followed by AEM 8 (5.7%, P = 0.007), and AEMs 1, 8, 5, and 3

individually significant below a P = 0.05 threshold out of 9 AEMs evaluated. For distance-

based *A. millepora* MEM models, the best model (custom saturated) retained only MEM 12.

For *A. tenuis*, sampling within regions was sufficiently dense to evaluate fits for adjacent

regions of competing AEM models. In all instances links based on north to south alongshore

flow returned the best scoring models, with fit to allelic frequency patterns notably lower in

the far northern and northern reefs (Table 2).

DISCUSSION

Asymmetric dispersal is undoubtedly a common attribute of benthic marine species. For the

corals *Acropora tenuis* and *Acropora millepora*, a preponderance of north to south movement

along the GBR was predicted by biophysical models and this directional signal was matched in

patterns of microsatellite allele frequencies. Although previous studies have noted distinct

characteristics of northern + central vs. southern + Swains + Capricorn-Bunker regions

(Lukoschek et al., 2016; van Oppen et al., 2011), here we can confirm asymmetric gene flow as

a contributor to these regional differences. Alternative scenarios involving historical

divergence and symmetric gene flow were poor matches to observed spatial genetic patterns

relative to scenarios based on directions and extent of larval exchange predicted by

biophysical models. Thus, at least at the regional scale within the GBR, oceanography is likely

to strongly influence connections arising from planktonic larval dispersal.

Regional connections among GBR populations

Whereas geographically restricted alleles are a hallmark of long-standing isolation, highly

directional dispersal such as among coastal populations can result in regionally private alleles

in downstream populations if dispersal is sufficiently rare and when accompanied by high

self-recruitment of settlers to their parental populations (Pringle & Wares, 2007). This

expectation qualitatively matched diversity gradients for both *Acropora* species (as noted by

Lukoschek et al., 2016). Here, coalescent demographic simulations allowed us to formally

evaluate a hypothesis of Quaternary divergence against a scenario of long-standing gene flow

without divergence. For *A. tenuis,* we found very high support for gene flow without

divergence (posterior probability: 0.999; BF = 1210) and moderate support for this scenario

in *A. millepora* (PP= 0.647; BF = 3). Thus, if there were past divergence events, the imprint of

such events is no longer discernible from extant allele frequencies and present-day

population genetic structure should be strongly influenced by recent gene flow.

Indeed, contemporary biophysical expectations for gene flow were strong predictors of

genetic patterns especially at large spatial scales. Using shared alleles, the weakest genetically

connected population pairs (negligible gene flow from reef i to reef j) match pairs having the

weakest predicted multigenerational connection links (Fig. 3). Where the biophysical

dispersal models predicted stronger connections, in contrast, inferred gene flow was highly

variable indicating that at shorter distances the biophysical models and gene flow were not

consistently aligned. Formal statistical testing for spatial structure of allelic distributions with

MEMs and AEMs reinforced these findings, where eigenvectors derived from predicted

asymmetric connections explained considerably more variance than eigenvectors derived

from null geographic models of symmetric connections (Table 1). Additionally, for both *A.*

tenuis and *A. millepora* AEM 1, which describes the largest scale pattern of spatial

relationships (Fig. 4 and Sup. Fig. 3), explained the greatest proportion of variance in allelic

distributions (3.5 and 5.9% , respectively). AEM analyses, however, were also consistent with

spatial patterning at smaller spatial scales arising from asymmetric connections (Table 1).

That biophysical models and spatial genetic structure were misaligned at small scales does

not seem overly surprising. Small-scale oceanography is likely to be highly variable and may

lead to higher levels of mixing among local clusters of reefs than a limited number of dispersal

years would suggest. Moreover, the time periods are incongruent as the dispersal models

spanned four years (2008-2012), whereas genetic data were derived from mature colonies

(collected in 2009-2013 for *A. tenuis* and 2002-2009 for *A. millepora*). Additionally, chaotic

genetic patterns at small scales such as those observed here (Fig. 3) are well-known for

marine taxa, perhaps reflecting stochasticity associated with planktonic dispersal and/or

post-settlement selection (e.g. Johnson & Black, 1984). Finally, biophysical larval dispersal

models implicitly assume that post-settlement mortality rates would not be affected by

origins of settlers, but larval origins could influence juvenile fitness (Marshall et al. 2010).

Although spatial distributions of microsatellite alleles for both *A. tenuis* and *A. millepora* were
broadly consistent with asymmetric dispersal predicted by biophysical models, the best
alignments were obtained only considering the approximately north to south connections
along the length of the GBR (Table 1). Notably biophysical models for both species suggested
substantial counterflows (south to north) especially among inshore reefs (Fig. 1), yet adding
these connections to AEM models did not increase model fit (Table 1). Thus, either the
stronger north to south connections are sufficient for capturing most of the microsatellite
allelic variance or the predicted south to north movements are not realised: for example,
larvae may conceivably disperse northwards but could have low fitness and therefore not
make substantive contributions to gene flow. Matz *et al.* (2018) also detected greater
southwards migration among five populations of GBR *A. millepora* based on ~11,500
nucleotide polymorphisms, albeit with some northwards migration and only considering an
inshore southern reef location (Keppel Island). Southward spread also characterises Crown-
of-Thorns starfish outbreaks that appear to originate in northern reefs (Pratchett et al., 2014).
Therefore, our finding of a strong southward dispersal signal is compatible with these other
results, but the sensitivity of AEM based analyses to bidirectional flows is unclear.
400
For *A. tenuis,* we were able to further assess correlations between biophysical predictions and
allelic spatial distributions between adjacent regions (Table 2). For southern and central
reefs, north to south connections models yielded the best fits to the data ($R^2 \geq 0.32$). These
observations spatially align with major jets from the Coral Sea encountering the GBR and
driving southward advective flows along the GBR (i.e. surface waters of the South Caledonia
Jet at the top of the central region and Eastern Australian Current at the Swains: Choukroun,
Ridd, Brinkman, & McKinna, 2010; Mao & Luick, 2014). In contrast, far northern and northern
reefs showed reduced concordance between predictions and genetic observations ($R^2 = 0.17$)
with a model based on bidirectional connections yielding the best fit to this regional data;
surface waters of the North Caledonia Jet encounter the GBR here and creates both northward
and southward flows and thus might explain a possible signal of bidirectional connections
among *A. tenuis* populations (Choukroun et al., 2010; Mao & Luick, 2014). For both species,
biophysical models predict rare dispersal connections especially among far northern reefs
(Fig. 1) and yet negligible population structure was observed for either *A. tenuis* or *A.*
millepora among reefs in the far north (Lukoschek et al., 2016) suggesting a weaker fit
between dispersal models and observed genetic patterns in this region.

In summary, for the two acroporid corals considered here, we find strong support for recent gene flow in a predominantly north to south direction. This result implies that northern and central reefs are important sources for downstream southern reefs, at least over evolutionary time scales of many generations. Southern outer shelf reef populations, however, harbour greater genetic diversity than northern reefs (Lukoschek et al., 2016) and are therefore both valuable in terms of distinctive diversity and, as the present results show, likely to be strongly self-recruiting so that they may be more self-sustaining than central and northern reefs. Dynamics of inshore reefs may be different from outer shelf reefs, for example, Lukoschek et al. (2016) highlighted the genetic distinctiveness of the inshore Keppel Island population for both *A. tenuis* and *A. millepora* and Matz et. al. (2018) found low genetic diversity in the Keppels but with no sampling of southern outer reef populations as a basis for comparison. Matz et. al. (2018) also reported low-levels of northward gene flow among their inner reef populations (e.g. Keppel to Magnetic and Orpheus Islands). Further genomic studies with more comprehensive spatial sampling may yield greater detailed resolution regarding rates and directions of gene flow, as the microsatellite data we examined here lack the power to confidently estimate specific gene flow rates.

With temperatures and extreme heating events projected to increase in frequency (Wolff et al., 2018), resolving the GBR-wide capacity for gene flow including the directionality of naturally occurring gene flow provides information relevant to discussions regarding assisted migration and genetic rescue (Anthony et al., 2017). For example, although the southern outer reefs, especially the Swains, have been largely spared from bleaching events, our results indicate that naturally occurring dispersal from south to north is most likely ecologically insignificant (Fig. 3), and thus would indicate that prospective migrants from the southern regions would be unlikely to recolonise central and northern reefs in sufficient quantities over decadal time scales. Because naturally occurring genetic rescue relies on evolutionarily significant (and not ecologically significant) gene flow, the prospects for this phenomenon are better, although predicated on receiving locations hosting healthy populations. Should adaptive variants (such as conferring heat tolerance or bleaching resistance) exist in some reefs then they might be able to spread naturally, with southward spread more probable. In any event, increased knowledge regarding the directionality and magnitudes of genetic exchange among GBR reefs will help forecast possible future evolutionary dynamics and can

help identify dispersal barriers that could be circumvented via assisted migration (Hoffmann

et al. 2015). In the case of GBR corals, assisted movements in a northward direction would

bypass natural dispersal limitations.

Detecting contemporary asymmetric planktonic larval dispersal

Relatively few studies have evaluated the effect of asymmetric larval dispersal on spatial

genetic patterns in marine species (Riginos et al., 2016). Perhaps not surprisingly, these

studies, like our results here with *A. tenuis* and *A. millepora,* consistently uncover substantial

asymmetries in inferred dispersal (e.g., Benestan et al., 2016; Dalongeville et al., 2018; Xuereb

et al., 2018). Clearly, conventional analyses, especially those based on summary statistics like

F_{ST} where all directional information is obscured, will fail to uncover important elements

regarding relationships among populations (Kool et al., 2013; Riginos et al., 2016).

An additional challenge is to clarify the time frame for processes yielding asymmetric spatial

genetic patterns. Empirical genomic data paired with specific historical demographic

simulations can sometimes put bounds on the timing and directionality of gene flow (as in

Duranton et al., 2018; Matz et al., 2018), but such approaches are largely constrained to

examining a few populations at a time due to exponential increase in number of potential free

parameters (for example, if migration is free to vary among n populations then there will be

n!/(n-2)! parameters for directional migration). Moreover, there is greater uncertainty

surrounding parameter estimation for recent events (Robinson, Coffman, Hickerson, &

Gutenkunst, 2014). Simple frequentist approaches based on linear models such as AEMs

represent an attractive alternative especially as they have been conceived for uncovering

contemporary structuring processes in species composition of ecological communities (Dray

et al., 2006), but if applied to intraspecific genetic data these approaches are likely to be

misleading when historical events have strongly shaped spatial genetic patterns.

In the present study, we attempted to bridge this historical-contemporary dilemma by using

constrained historical demographic simulations to reject the historical scenario most likely to

influence observable spatial genetic patterns for GBR species, giving us greater confidence

that the spatial genetic patterns we observe for *A. tenuis* and (to a lesser extent) *A. millepora*

reflect evolutionary processes from the recent past. Using prior information based on geology,

historical habitat shifts, or histories of co-distributed species to identify relevant alternative

hypotheses and then evaluating those hypotheses against hypotheses based on contemporary

gene flow (such as simple stepping-stone gene flow as used here) is a reasonable check of

data before proceeding to interpret results from linear models (including AEMs) at face value.

Biophysical models as summaries of larval dispersal

Biophysical models are increasingly being used to make detailed spatial predictions of

planktonic larval dispersal often with the aim of guiding management actions (as in Hock et

al., 2016; 2017; 2019; Krueck et al., 2017). Although such complex models of natural systems

cannot be truly tested (Oreskes et al., 1994), alignment of model outputs against independent

biological data can provide greater confidence that a biophysical larval dispersal model

captures elements of biological reality. In the present study, we find that directions and

relative magnitudes of dispersal connections derived from biophysical models are better

predictors of spatial genetic structure in *A. tenuis* and *A. millepora* than null models of

symmetric relationships, providing some confirmation that real biological processes mirror

expectations arising from biophysical simulations. The superior predictive power of

asymmetric predictors provides further evidence of the need to consider directional flow

when analysing connectivity in marine systems.

For our *Acropora* species on the GBR, the strongest alignments between the models and

empirical genetic data are at large spatial scales (among regions) and involve north to south

connections. Because small numbers of migrants can homogenize allele frequencies between

populations, inferences based on allele frequencies are poorly suited to distinguishing

strengths of demographically significant connections (Waples, 1998). Allele frequency based

methods, however, can theoretically discriminate between rare and very rare connections

such as those that are likely to link geographically distant populations (including via

multigenerational connections). Thus, the appropriate spatial scale for evaluating biophysical

larval dispersal models with population genetic data is at large spatial distances such a

regional connections among GBR coral populations where we find the greatest concordance

between predicted and observed patterns. Comprehensive evaluation of biophysical models

will best be undertaken with a variety of complementary empirical data types (Gaggiotti,

2017; Jones, 2015), where geographic patterns of allelic distributions contribute valuable

information about rare long-distance connections. Here we have shown that the directionality

of such rare long-distance connections discerned from spatial distributions of alleles provides

additional useful information for gauging concordance of larval dispersal models against past
gene flow.
518

519   **Acknowledgements:**
520   We thank FG Blanchet for guidance on creating connexion diagrams and C Doropoulos for
521   help with parameterisation of the dispersal models, the QRIS computing cluster for enabling
522   simulation modelling, and funding support from the Great Barrier Reef Foundation.

523
524
525   **Data accessibility statement:** Raw data available at datadryad.org/resource/doi:10.5061/
526   dryad.h8gh3 for *Acropora tenuis* and datadryad.org/resource/doi:10.5061/dryad.q0834p71
527   for *Acropora millepora.* Scripts, datafiles, and example results are available on GitHub:
528   github.com/khock1/CoralPaths; github.com/dinmatias; github.com/CRiginos1.
529

REFERENCES

Abbey, E., Webster, J. M., & Beaman, R. J. (2011). Geomorphology of submerged reefs on the shelf edge of the Great Barrier Reef: The influence of oscillating Pleistocene sea-levels. *Marine Geology*, *288*(1-4), 61–78. http://doi.org/10.1016/j.margeo.2011.08.006

Anthony, K. R. N., Bay, L., Costanza, R., Firn, J., Gunn, J., Harrison, P., et al. (2017). New interventions are needed to save coral reefs. *Nature Ecology & Evolution*, *1*(10), 1–3. http://doi.org/10.1038/s41559-017-0313-5

Beger, M., Linke, S., Watts, M., Game, E., Treml, E. A., Ball, I., & Possingham, H. P. (2010). Incorporating asymmetric connectivity into spatial decision making for conservation. *Conservation Letters*, *3*(5), 359–368. http://doi.org/10.1111/j.1755-263X.2010.00123.x

Benestan, L., Quinn, B. K., Maaroufi, H., Laporte, M., Clark, F. K., Greenwood, S. J., et al. (2016). Seascape genomics provides evidence for thermal adaptation and current-mediated population structure in American lobster (*Homarus americanus*). *Molecular Ecology*, *25*(20), 5073–5092. http://doi.org/10.1111/mec.13811

Blanchet, F. G., Legendre, P., & Borcard, D. (2008). Modelling directional spatial processes in ecological data. *Ecological Modelling*, *215*(4), 325–336. http://doi.org/10.1016/j.ecolmodel.2008.04.001

Blanchet, F. G., Legendre, P., Maranger, R., Monti, D., & Pepin, P. (2011). Modelling the effect of directional spatial ecological processes at different scales. *Oecologia*, *166*(2), 357–368. http://doi.org/10.1007/s00442-010-1867-y

Borcard, D., & Legendre, P. (2002). All-scale spatial analysis of ecological data by means of principal coordinates of neighbour matrices. *Ecological Modelling*, *153*(1-2), 51–68. http://doi.org/10.1016/S0304-3800(01)00501-4

Boykov, Y., & Kolmogorov, V. (2004). An experimental comparison of min-cut/max- flow algorithms for energy minimization in vision. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, *26*(9), 1124–1137. http://doi.org/10.1109/TPAMI.2004.60

Carr, M. H., Neigel, J., Estes, J., Andelman, S., Warner, R., & Largier, J. (2003). Comparing marine and terrestrial ecosystems: Implications for the design of coastal marine reserves. *Ecological Applications*, *13*(1), S90–S107.

Choukroun, S., Ridd, P. V., Brinkman, R., & McKinna, L. I. W. (2010). On the surface circulation in the western Coral Sea and residence times in the Great Barrier Reef. *Journal of Geophysical Research*, *115*(C6), 239. http://doi.org/10.1029/2009JC005761

Cowen, R. K., & Sponaugle, S. (2009). Larval dispersal and marine population connectivity. *Annual Review of Marine Science*, *1*(1), 443–466. http://doi.org/10.1146/annurev.marine.010908.163757

Crandall, E. D., Treml, E. A., & Barber, P. H. (2012). Coalescent and biophysical models of stepping-stone gene flow in neritid snails. *Molecular Ecology*, *21*(22), 5579–5598. http://doi.org/10.1111/mec.12031

Dalongeville, A., Andrello, M., Mouillot, D., Lobreaux, S., Fortin, M.-J., Lasram, F., et al. (2018). Geographic isolation and larval dispersal shape seascape genetic patterns differently according to spatial scale. *Evolutionary Applications*, *11*(8), 1437–1447. http://doi.org/10.1111/eva.12638

De'ath, G., Fabricius, K. E., Sweatman, H. P. A., & Puotinen, M. (2012). The 27-year decline of coral cover on the Great Barrier Reef and its causes. *Proceedings of the National Academy of Sciences of the United States of America*, *109*(44), 17995–17999. http://doi.org/10.1073/pnas.1208909109

Dijkstra, E. W. (1959). A note on two problems in connexion with graphs. *Numerische Mathematik*, *1*(1), 269–271. http://doi.org/10.1007/BF01386390

Diniz-Filho, J. A. F., Nabout, J. C., Telles, M. P. de C., Soares, T. N., & Rangel, T. F. L. V. B. (2009).

A review of techniques for spatial modeling in geographical, conservation and landscape
genetics. *Genetics and Molecular Biology*, *32*(2), 203–211. http://doi.org/10.1590/S1415-
47572009000200001
Dray, S., Legendre, P., & Peres-Neto, P. R. (2006). Spatial modelling: a comprehensive
framework for principal coordinate analysis of neighbour matrices (PCNM). *Ecological*
Modelling, *196*(3-4), 483–493. http://doi.org/10.1016/j.ecolmodel.2006.02.015
Duranton, M., Allal, F., Fraïsse, C., Bierne, N., Bonhomme, F., & Gagnaire, P.-A. (2018). The
origin and remolding of genomic islands of differentiation in the European sea bass.
Nature Communications, *9*(1), 1–11. http://doi.org/10.1038/s41467-018-04963-6
Excoffier, L., & Lischer, H. E. L. (2010). Arlequin suite ver 3.5: a new series of programs to
perform population genetics analyses under Linux and Windows. *Molecular Ecology*
Resources, *10*(3), 564–567. http://doi.org/10.1111/j.1755-0998.2010.02847.x
Excoffier, L., Dupanloup, I., Huerta-Sánchez, E., Sousa, V. C., & Foll, M. (2013). Robust
demographic inference from genomic and SNP data. *PLoS Genetics*, *9*(10), e1003905.
http://doi.org/10.1371/journal.pgen.1003905
Ford, L. R., & Fulkerson, D. R. (1956). Maximal flow through a network. *Canadian Journal of*
Mathematics, *8*(0), 399–404. http://doi.org/10.4153/CJM-1956-045-5
Foster, N. L., Paris, C. B., Kool, J. T., Baums, I. B., Stevens, J. R., Sanchez, J. A., et al. (2012).
Connectivity of Caribbean coral populations: complementary insights from empirical and
modelled gene flow. *Molecular Ecology*, *21*(5), 1143–1157.
http://doi.org/10.1111/j.1365-294X.2012.05455.x
Gaggiotti, O. E. (2017). Metapopulations of marine species with larval dispersal: a
counterpoint to Ilkka's Glanville Fritillary metapopulations. *Annales Zoologici Fennici*,
54(1-4), 97–112. http://doi.org/10.5735/086.054.0110
Gaines, S. D., White, C., Carr, M. H., & Palumbi, S. R. (2010). Designing marine reserve networks
for both conservation and fisheries management. *Proceedings of the National Academy of*
Sciences of the United States of America, *107*(43), 18286–18293.
http://doi.org/10.1073/pnas.0906473107
Galindo, H. M., Olson, M. V., & Palumbi, S. R. (2006). Seascape genetics: A coupled
oceanographic-genetic model predicts population structure of Caribbean corals. *Current*
Biology, *16*(16), 1622–1626. http://doi.org/10.1016/j.cub.2006.06.052
GBRMPA. Great Barrier Reef Marine Park Zoning Plan 2003. Report No. ISBN9781876945237,
1-211 (Great Barrier Reef Marine Park Authority, Townsville, QLD, Australia, 2004).
Hellberg, M. E. (2009). Gene flow and isolation among populations of marine animals. *Annual*
Review of Ecology, Evolution, and Systematics, *40*(1), 291–310.
http://doi.org/10.1146/annurev.ecolsys.110308.120223
Hinestrosa, G., Webster, J. M., & Beaman, R. J. (2016). Postglacial sediment deposition along a
mixed carbonate-siliciclastic margin: New constraints from the drowned shelf-edge reefs
of the Great Barrier Reef, Australia. *Palaeogeography, Palaeoclimatology, Palaeoecology*,
446, 168–185. http://doi.org/10.1016/j.palaeo.2016.01.023
Hock, K., Doropoulos, C., Gorton, R., Condie, S.A., & Mumby, P.J. (2019) Split spawning
increases robustness of coral larval supply and inter-reef connectivity. *Nature*
Communications. In press.
Hock, K., & Mumby, P. J. (2015). Quantifying the reliability of dispersal paths in connectivity
networks. *Journal of the Royal Society Interface*, *12*(105), 20150013–20150013.
http://doi.org/10.1098/rsif.2015.0013
Hock, K., Wolff, N. H., Beeden, R., Hoey, J., Condie, S. A., Anthony, K. R. N., et al. (2016).
Controlling range expansion in habitat networks by adaptively targeting source
populations. *Conservation Biology*, *30*(4), 856–866. http://doi.org/10.1111/cobi.12665

Hock, K., Wolff, N. H., Ortiz, J. C., Condie, S. A., Anthony, K. R. N., Blackwell, P. G., & Mumby, P. J.
(2017). Connectivity and systemic resilience of the Great Barrier Reef. *PLoS Biology*,
15(11), e2003355–23. http://doi.org/10.1371/journal.pbio.2003355
Hoffmann, A. A., Griffin, P. C., Dillon, S., Catullo, R. A., Rane, R., Byrne, M., et al. (2015). A
framework for incorporating evolutionary genomics into biodiversity conservation and
management. *Climate Change Responses*, *2*(1), 1–24. http://doi.org/10.1186/s40665-014-
0009-x
Hughes, T. P., Kerry, J. T., Álvarez-Noriega, M., Álvarez-Romero, J. G., Anderson, K. D., Baird, A.
H., et al. (2017). Global warming and recurrent mass bleaching of corals. *Nature*,
543(7645), 373–377. http://doi.org/10.1038/nature21707
Hughes, T. P., Kerry, J. T., Baird, A. H., Connolly, S. R., Dietzel, A., Eakin, C. M., et al. (2018).
Global warming transforms coral reef assemblages. *Nature*, *556*(7702), 492–496.
http://doi.org/10.1038/s41586-018-0041-2
Johnson, M. S., Black, R. 1984. Pattern beneath the chaos: the effect of recruitment on genetic
patchiness in an intertidal limpet. *Evolution* 38: 1371-1383.
Jones, G. P. (2015). Mission impossible: unlocking the secrets of coral reef fish dispersal. In C.
Mora (Ed.), *Ecology of Fishes on Coral Reefs* (pp. 16–27). Cambridge: Cambridge University
Press. http://doi.org/10.1017/cbo9781316105412.004
Kool, J. T., Moilanen, A., & Treml, E. A. (2013). Population connectivity: recent advances and
new perspectives. *Landscape Ecology*, *28*(2), 165–185. http://doi.org/10.1007/s10980-
012-9819-z
Kool, J. T., Paris, C. B., Andréfouët, S., & Cowen, R. K. (2010). Complex migration and the
development of genetic structure in subdivided populations: an example from Caribbean
coral reef ecosystems. *Ecography*, *105*, 26191–10. http://doi.org/10.1111/j.1600-
0587.2009.06012.x
Kool, J. T., Paris, C. B., Barber, P. H., & Cowen, R. K. (2011). Connectivity and the development
of population genetic structure in Indo-West Pacific coral reef communities. *Global
Ecology and Biogeography*, *20*(5), 695–706. http://doi.org/10.1111/j.1466-
8238.2010.00637.x
Kritzer, J. P., & Sale, P. F. (2004). Metapopulation ecology in the sea: from Levins' model to
marine ecology and fisheries science. *Fish and Fisheries*, *5*(2), 131–140.
http://doi.org/10.1111/j.1467-2979.2004.00131.x
Krueck, N. C., Ahmadia, G. N., Green, A., Jones, G. P., Possingham, H. P., Riginos, C., et al. (2017).
Incorporating larval dispersal into MPA design for both conservation and fisheries.
Ecological Applications 27(3), 925–941. http://doi.org/10.1002/eap.1495
Legendre, P., & Gallagher, E. D. (2001). Ecologically meaningful transformations for ordination
of species data. *Oecologia*, *129*(2), 271–280. http://doi.org/10.1007/s004420100716
Liggins, L., Treml, E. A., & Riginos, C. (2013). Taking the plunge: an introduction to
undertaking seascape genetic studies and using biophysical models. *Geography Compass*,
7(3), 173–196. http://doi.org/10.1111/gec3.12031
Lukoschek, V., Cross, P., Torda, G., Zimmerman, R., & Willis, B. L. (2013). The importance of
coral larval recruitment for the recovery of reefs impacted by Cyclone Yasi in the central
Great Barrier Reef. *PLoS ONE*, *8*(6), e65363–11.
http://doi.org/10.1371/journal.pone.0065363
Lukoschek, V., Riginos, C., & van Oppen, M. J. H. (2016). Congruent patterns of connectivity can
inform management for broadcast spawning corals on the Great Barrier Reef. *Molecular
Ecology*, *25*(13), 3065–3080. http://doi.org/10.1111/mec.13649
Mao, Y., & Luick, J. L. (2014). Circulation in the southern Great Barrier Reef studied through an
integration of multiple remote sensing and in situ measurements. *Journal of Geophysical*

Research: Oceans, *119*(3), 1621–1643. http://doi.org/10.1002/2013JC009397

Marshall, D. J., Monro, K., Bode, M., Keough, M. J., & Swearer, S. (2010). Phenotype-
environment mismatches reduce connectivity in the sea. *Ecology Letters*, *13*(1), 128–140.
http://doi.org/10.1111/j.1461-0248.2009.01408.x

Matz, M. V., Treml, E. A., Aglyamova, G. V., & Bay, L. (2018). Potential and limits for rapid
genetic adaptation to warming in a Great Barrier Reef coral. *PLoS Genetics*, *14*(4),
e1007220–19. http://doi.org/10.1371/journal.pgen.1007220

Metaxas, A., & Saunders, M. (2009). Quantifying the "bio-" components in biophysical models
of larval transport in marine benthic invertebrates: advances and pitfalls. *The Biological*
Bulletin, *216*(3), 257–272. http://doi.org/10.1086/BBLv216n3p257

Oreskes, N., Shrader-Frechette, K., & Belitz, K. (1994). Verification, validation, and
confirmation of numerical models in the Earth sciences. *Science*, *263*(5147), 641–646.
http://doi.org/10.1126/science.263.5147.641

Pineda, J., Hare, J. A., & Sponaugle, S. (2007). Larval transport and dispersal in the coastal
ocean and consequences for population connectivity. *Oceanography*, *20*(3), 22–39.

Planes, S., & Doherty, P. J. (1997). Genetic and color interactions at a contact zone of
Acanthochromis polyacanthus: a marine fish lacking pelagic larvae. *Evolution*, *51*(4), 1232–
1243. http://doi.org/10.1111/j.1558-5646.1997.tb03970.x

Pratchett, M. S., Caballes, C. F., Rivera-Posada, J. A., & Sweatman, H. P. A. (2014). Limits to
understanding and managing outbreaks of crown-of-thorns starfish (*Acanthaster spp.*).
Oceanography and Marine Biology, 1–68.

Pringle, J. M., & Wares, J. P. (2007). Going against the flow: maintenance of alongshore
variation in allele frequency in a coastal ocean. *Marine Ecology Progress Series*, *335*, 69–84.
http://doi.org/10.3354/meps335069

Riginos, C., & Liggins, L. (2013). Seascape genetics: populations, individuals, and genes
marooned and adrift. *Geography Compass 7*(3), 197–216.
http://doi.org/10.1111/gec3.12032

Riginos, C., Crandall, E. D., Liggins, L., Bongaerts, P., & Treml, E. A. (2016). Navigating the
currents of seascape genomics: how spatial analyses can augment population genomic
studies. *Current Zoology*, *62*(6), 581–601. http://doi.org/10.1093/cz/zow067

Robinson, J. D., Coffman, A. J., Hickerson, M. J., & Gutenkunst, R. N. (2014). Sampling strategies
for frequency spectrum-based population genomic inference. *BMC Evolutionary Biology*,
14(1), 254. http://doi.org/10.1186/s12862-014-0254-4

Roux, C., Fraïsse, C., Romiguier, J., Anciaux, Y., Galtier, N., & Bierne, N. (2016). Shedding light on
the grey zone of speciation along a continuum of genomic divergence. *PLoS Biology*,
14(12), e2000234–22. http://doi.org/10.1371/journal.pbio.2000234

Schunter, C., Carreras-Carbonell, J., Macpherson, E., Tintoré, J., Vidal-Vijande, E., Pascual, A., et
al. (2011). Matching genetics with oceanography: directional gene flow in a
Mediterranean fish species. *Molecular Ecology*, *20*(24), 5167–5181.
http://doi.org/10.1111/j.1365-294X.2011.05355.x

Selkoe, K. A., D'Aloia, C. C., Crandall, E. D., Iacchei, M., Liggins, L., Puritz, J. B., et al. (2016a). A
decade of seascape genetics: contributions to basic and applied marine connectivity.
Marine Ecology Progress Series, *554*, 1–19. http://doi.org/10.3354/meps11792

Selkoe, K. A., Scribner, K. T., & Galindo, H. M. (2016b). Waterscape genetics – applications of
landscape genetics to rivers, lakes, and seas. In N. Balkenhol, S. A. Cushman, A. Storfer, & L.
P. Waits (Eds.), *Landscape Genetics: Concepts, Methods, Applications* (pp. 1–27). John Wiley
& Sons, Ltd.

Sundqvist, L., Keenan, K., Zackrisson, M., Prodöhl, P., & Kleinhans, D. (2016). Directional
genetic differentiation and relative migration. *Ecology and Evolution*, *6*(11), 3461–3475.

http://doi.org/10.1002/ece3.2096

Thomas, L., Kennington, W. J., Stat, M., Wilkinson, S. P., Kool, J. T., & Kendrick, G. A. (2015).
Isolation by resistance across a complex coral reef seascape. *Proceedings of the Royal*
Society: Biological Sciences 282(1812), 20151217.
http://doi.org/10.1098/rspb.2015.1217

Truelove, N. K., Kough, A. S., Behringer, D. C., Paris, C. B., Box, S. J., Preziosi, R. F., & Butler, M. J.
(2016). Biophysical connectivity explains population genetic structure in a highly
dispersive marine species. *Coral Reefs*, 1–13. http://doi.org/10.1007/s00338-016-1516-y

van Oppen, M. J. H., Peplow, L. M., Kinnimonth, S., & Berkelmans, R. (2011). Historical and
contemporary factors shape the population genetic structure of the broadcast spawning
coral, *Acropora millepora*, on the Great Barrier Reef. *Molecular Ecology*, *20*(23), 4899–
4914. http://doi.org/10.1111/j.1365-294X.2011.05328.x

Waples, R. S. (1998). Separating the wheat from the chaff: patterns of genetic differentiation
in high gene flow species. *The Journal of Heredity*, *89*(5), 438–450.
http://doi.org/10.1093/jhered/89.5.438

Watson, J. R., Kendall, B. E., Seigel, D. A., & Mitarai, S. (2012). Changing seascapes, stochastic
connectivity, and marine metapopulation dynamics. *American Naturalist*, *180*(1), 99–112.
http://doi.org/10.1086/665992

Werner, F., & Cowen, R. K. (2007). Coupled biological and physical models: present
capabilities and necessary developments for future studies of population connectivity.
Oceanography, *20*, 54–69.

White, C., Selkoe, K. A., Watson, J., Siegel, D. A., Zacherl, D. C., & Toonen, R. J. (2010). Ocean
currents help explain population genetic structure. *Proceedings. Biological Sciences / the*
Royal Society, *277*(1688), 1685–1694. http://doi.org/10.1098/rspb.2009.2214

Whitlock, M. C., & McCauley, D. E. (1999). Indirect measures of gene flow and migration: FST
not equal to 1/(4Nm + 1). *Heredity*, *82*, 117–125. Retrieved from
http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=10098262&r
etmode=ref&cmd=prlinks

Wolff, N. H., Mumby, P. J., Devlin, M., & Anthony, K. R. N. (2018). Vulnerability of the Great
Barrier Reef to climate change and local pressures. *Global Change Biology*, *24*(5), 1978–
1991. http://doi.org/10.1111/gcb.14043

Xuereb, A., Benestan, L., Normandeau, E., Curtis, J. M., Bernatchez, L., & Fortin, M.-J. (2018).
Asymmetric oceanographic processes mediate connectivity and population genetic
structure, as revealed by RADseq, in a highly dispersive marine invertebrate
(*Parastichopus californicus*). *Molecular Ecology*, 1–46.

761
762

Biosketch: The authors are linked by a desire to better understand how planktonic larval
dispersal connects marine populations. Advances in measuring and predicting dispersal
connections are urgently needed in the context of the Great Barrier Reef where there have
been recent widespread outbreaks of coral bleaching. Thus, progress in uncovering the
dynamics of long-distance dispersal connections contributes to an ultimate goal of identifying
locations best-able to supply propagules and seed downstream reefs. Author contributions:
CR and VL conceived the overarching concept and KH, AMM, PJM, & MJHvO contributed ideas
fundamental to the study; VL, MJHvO, and KH collected the data; CR, KH, and AMM analysed
the data and developed the figures; and CR led the writing with contributions from KH, AMM,
PJM, MJHvO & VL.

TABLES

Table 1: Spatial eigenvector map modelling of genetic structure. The models explaining the greatest amount of genetic variance are

those that incorporate directional link structures derived from the larval dispersal model, especially those that are based on north to

south alongshore flow.

		Acropora tenuis			*Acropora millepora*		
Eigenmodel	Link structure	Link weighting of best model [a]	Numb EMs in best model [b]	adjusted R^{2} [b]	Link weighting of best model [a]	Numb EMs in best model [b]	adjusted R^{2} [b]
MEM	Delaunay	Binary	1	0.02	Binary	1	0.02
MEM	Delaunay	log	1	0.03	log	1	0.03
MEM	PCNM	y=1	**4**	**0.24***	log	**1**	**0.09**
MEM	Custom saturated	y=1	3	0.16*	log	1	0.10
AEM	Alongshore N to S only [c]	MF, y=1	**9**	**0.37**	Binary	**5**	**0.27**
AEM	Alongshore both N to S and S to N	SS, y=1	8	0.29	MF, y=1	2	0.15

[a] Binary distances represent a single model. The link structures were evaluated for link distance weightings based on the formula 1-

(D/Dmax)^{y} for y=[1-3] or with a log transformation to approximate a typical isolation-by-distance analysis. For AEMs, links were limited

to those in the top 50 percentile of reliable path weights. Link weightings were evaluated based on the number of stepping stone (SS)

connections, maximum flow (MF), and reliable path (RP) strength.

[b] Following forward model selection; asterisks indicate spatial models with **Δ**AICc < 2 as compared to the null (intercept only) model.

[c] Select AEMs shown graphically in Fig. 4 and Sup. Fig. 3

Table 2: Asymmetric eigenvector map modelling regions within *Acropora tenuis*

		Best model by region		
Region	Link structure	Link weighting [a]	Numb EMs in best model [b]	adjusted R^2 [b]
Far north & north	Alongshore both N to S and S to N	MF, y=1	**3**	0.17
North, central, south	Alongshore N to S	MF, y=1	**7**	0.32
Central & south	Alongshore N to S	RP, y=2	**5**	0.35

788   [a] Link structures were evaluated for distance weightings based on the formula $1-(D/Dmax)^y$ for y=[1-3].

789   [b] Following forward model selection.

 FIGURE LEGENDS

Figure 1: Sampling locations and main attributes of biophysical model. A) Coastal Queensland

and the Great Barrier Reef where bathymetry is shown by grey shading and 120 m depth

reflects the approximate land mass exposure at lowest Pleistocene sea level stands.

Designated regions correspond to management areas. Major offshore currents are shown

(NVJ and SVJ: north and south Vanuatu jets, NQC: north Queensland current, EAC: east

Australia current; modified from Coukroun *et al.* 2010; Mao & Luick 2014). B & C) Summary of

top 50 percentile predicted connections based on relative path probabilities for *Acropora*

tenuis (B) and *Acropora millepora* (C). Sampling locations are color coded by latitude with

northern low latitude sites shown in reds (warm) and southern higher latitude sites shown in

blues (cool). Vectors show predicted dispersal probabilities with thicker lines indicating

higher probabilities and colored by source population. D & E) Dispersal probabilities relative

to coastline position for *Acropora tenuis* (D) and *Acropora millepora* (E). The largest positional

changes northwards have the lowest probabilities whereas small positional changes and

movements southwards have higher probabilities. Movement relative to coastline is based on

the difference in relative position between locations as summarised by the first principal

component axis describing Queensland coastline. Example contrasts in northward vs.

southward dispersal strengths shown for example population pairs (a,b and c,d) traversing

the length of the sampling region where a->b and c->d are high probabilities ($P \approx 10^{-20}$) of

southward movement (-4 relative positional change) and b->a and d->c are low probabilities

($P \approx 10^{-40}$) of northward movement (+4 relative positional change).

Figure 2: Competing models of historical scenarios evaluated using Approximate Bayesian

Computation based on historical divergence vs. gene flow between adjacent stepping stone

populations. A) Representative populations chosen for comparison via ABC against simulated

data. B) Historical divergence models with a barrier to gene flow at Hydrographer's Passage.

C) Stepping-stone gene flow models where migration was permitted only between adjacent

populations but with rates free to vary. The prior distributions of the parameters are

summarized in Table S1. ABC analyses provided higher support for the stepping-stone gene

flow models (*A. tenuis*: posterior probability = 0.999; Bayes Factor = 1210; *A. millepora*: PP =

0.647; BF = 3) compared to the model of historical divergence vicariance.

Figure 3: Concordance in directional movement between the larval dispersal model and

genetic estimates for pairs of populations for *Acropora tenuis* (A), and *Acropora millepora* (B).

Routes of low predicted connectivity (small relative path probabilities: $P < 10^{-40}$) experience

low directional gene flow (using the DivMigrate method). Routes of high predicted

connectivity ($P > 10^{-20}$) have variable rates of gene flow.

Figure 4: Leading asymmetric eigenvector maps describing spatial genetic structure in

Acropora tenuis. Values by sampling location are colored by intensity of green hue such that

locations with similar colors have similar AEM values. The greatest proportion of variance

(AEM 1) describes a GBR-wide cline in genetic diversity followed by local scale spatial

autocorrelation structures (AEM 5).

A

NQC

NVJ

Far Northern

Northern

Coral Sea

SCJ

Depth (m)

Central

Southern

Swains

2500

Hydrographer's Passage

Capricorn-Bunker Group

Keppel Island

EAC

Queensland, Australia

B

a

Percentile dispersal strength:

b

Acropora tenuis

C

c

Percentile dispersal strength:

d

Acropora millepora

D

northwards

b -> a

Movement relative to coastline

a -> b

southwards

Acropora tenuis

E

d -> c

c -> d

Acropora millepora

Log$_{10}$ probability of dispersal (reliable path)

Acropora tenius　　　　　　*Acropora millepora*

a) Focal locations

● included in the ABC analysis
· excluded in the ABC analysis

(*n*) number of individuals analysed

b) Historical divergence model

c) Stepping-stone gene flow model

AEM 1
3.5% variance; P = 0.001

AEM 5
2.3% variance; P = 0.001