Title: Different factors shape pigmentation and other physiological trait variation within and between
species in the *Drosophila americana* group.

Running title: Pigmentation variation in the *D. americana* group

Jeremy S. Davis[1]*, Leonie C. Moyle[1]

[1]Dept. of Biology, Indiana University, Bloomington, IN, USA 47405

*Corresponding author: jsd2@indiana.edu

Acknowledgements:

We would like to thank the Drosophila species stock center for help procuring the population lines used

in this study. We would also like to thank the curators of the Worldclim, Climond, and taxodros

databases for providing valuable resources necessary for this study. Research was supported by Indiana

University Department of Biology funding to LCM and JSD.

Author contributions: JSD and LCM conceived and designed the experiment. JSD performed the

experiment and statistical analyses. JSD and LCM wrote and edited the manuscript.

Abstract

Disentangling the factors shaping adaptive trait variation is an important but challenging task. Natural
pigmentation variation in *Drosophila* is one instance where studies support multiple—sometimes
contradictory—proposed mechanisms for this variation, based on conflicting patterns of association
with other traits and/or environmental factors. Here we systematically examined associations between
pigmentation variation, a suite of environmental variables, and two other physiological traits
(desiccation and UV resistance), to critically evaluate several of these proposed mechanisms. Using 3
populations from each of 3 species—*D. americana americana, D. a. texana* and *D. novamexicana*—we
detected significant population- and species-level variation for all three traits and environmental
preferences. Further, we found that patterns of pigmentation variation differed between evolutionary
scales: within species, darker pigmentation was associated with hotter, drier, more irradiated
environments, but an opposing association (i.e., lighter species in these environments) was evident
between species. Weak (desiccation resistance) or no (UV resistance) associations with other trait
variation suggests little evidence that pleiotropy with these traits directly shapes pigmentation variation.
By simultaneously assessing multiple trait-trait and trait-environment relationships, our approach
indicates pigmentation variation is likely shaped by alternative factors at different evolutionary scales,
supporting different selective hypotheses at each of these scales.

Introduction

Differentiating the historical, genetic, and selective factors that cause some traits to vary locally

according to environment, while other traits become fixed between lineages, is of critical importance to

our understanding of how adaptations arise, spread, and persist. One hallmark of local adaptation is a

consistent association between trait variation and one or more aspects of the natural environment.

Classically, these associations have been assessed via the study of clines; by definition, clines exhibit

spatial variation, and geographic space is frequently environmentally heterogeneous, so traits exhibiting

functionally-relevant clinal variation are clear candidates for targets of local selection. Latitudinal or

altitudinal clines have received particular attention in numerous systems, including *Drosophila*,

Arabidopsis thaliana, and humans, where analyses indicate strong trait-environment associations for

several physiological and other variants (Adrian et al. 2015, Fournier-Level et al. 2011, Hancock et al.

2011). Nonetheless, even among these well-characterized examples, the underlying cause of clinal

variation is still not always clear, particularly when trait variation is surveyed across generalized

geographic space as opposed to a specific environmental gradient. In addition, clinal analyses are often

designed to identify traits that are subject to strong local adaptation—that is, that vary among

populations within species. However, some ecologically important traits might instead show discrete

differences between closely related species, especially when closely-related species are themselves

spatially partitioned. Whether these species-level differences dominate patterns of adaptation can

depend upon when variants arose in the history of lineage differentiation, whether they confer a net

fitness benefit in spite of intraspecific environmental variation, and/or whether they are subject to

constraints that prevent them from responding to local conditions. Studying traits that exhibit a range of

variation, both between populations within species and between populations of closely related species,

could therefore be invaluable in understanding how natural selection can differentially shape trait

variation at different spatial and evolutionary scales.

In *Drosophila,* patterns of intraspecific clinal variation and species differences have unveiled

several traits as potential targets of environmentally-mediated selection, including cuticle pigmentation,

and desiccation and UV resistance. Nonetheless, in many cases the environmental factors responsible

for driving variation in these traits is equivocal. For example, multiple studies have shown an association

between pigmentation variation and latitude, including within *D. melanogaster* in Europe (David et al.

1985), Australia (Telonis-Scott et al. 2011), India (Munjal et al. 1997), and sub-saharan Africa (Pool and

Aquadro 2007, Bastide et al. 2014), as well as within *D. simulans* (Capy et al. 1988) and the *D. cardini*

group (Heed and Krishnamurthy 1959). However, the nature of this association varies between studies;

while most find increased melanization at higher latitudes, the opposite pattern (i.e., darker flies at

lower latitudes) has been observed in *D. melanogaster* from sub-saharan Africa (Pool and Aquadro 2007,

Bastide et al. 2014). Moreover, while a number of hypotheses have been proposed to explain latitudinal

clines based on direct physiological effects of pigmentation variation—including selection for increased

pigmentation at higher, colder latitudes in order to increase absorption of solar radiation (the Thermal

Melanism Hypothesis, or Bogert's Rule; Gaston et al. 2009)—direct empirical evidence in support of this

and similar hypotheses remains limited and inconclusive.

Alternative hypotheses propose that selection for pigmentation variation is driven by a

pleiotropic association with other physiological stress resistance responses, especially UV radiation

and/or desiccation resistance responses. For example, selection for increased UV resistance has been

proposed to drive increased (darker) pigmentation, however empirical data indicates that the

mechanistic relationship between these two traits is more complex. Darker pigmentation is associated

with increased UV exposure and altitude in sub-saharan *D. melanogaster* (Pool and Aquadro 2007,

Bastide et al. 2014), but the opposite pattern has been observed for *D. yakuba* and *D. santomea*, with

lighter flies exhibiting greater UV resistance (Matute and Harris 2013). Experimentally, both light

(*yellow*) and dark (*ebony*) mutants have decreased survivorship after UV exposure (relative to wild type),

although dark mutants retained significantly better UV resistance compared to light mutants (Wang et

al. 2008). Similar to UV resistance, pleiotropic associations with desiccation stress have also been

proposed as a causal factor shaping pigmentation variation, with studies showing that increased

desiccation resistance is correlated clinally with darker pigmentation in *D. polymorpha* (Brisson et al.

2005) and Indian *D. melanogaster* and *D. ananassae* (Parkash et al. 2008, Parkash et al. 2010), although

this relationship was not observed in *D. yakuba/D. santomea* (Matute and Harris 2013). Experimental

evidence for a role of melanization in desiccation resistance is also mixed. In *D. melanogaster, D.

simulans, D. subobscura, and D. pseudoobscura*, *yellow* (light pigment) mutants have lower survival after

desiccation stress (relative to wild type), while *ebony D. melanogaster* mutants survive significantly

longer than wildtype (Kalmus 1941). However, Wittkopp et al. (2010) showed that introgression of dark

alleles into a light background and vice versa produced no effect on desiccation resistance among

species in the *D. americana* group. It therefore remains unclear whether these two traits are directly

causally connected or simply covary due to their shared association with other variables that remain

unaccounted for.

Here, our goal was to simultaneously evaluate desiccation resistance, UV resistance, and

pigmentation, the relationships among these traits, and their associations with climate and

environmental variation, to investigate the factors responsible for shaping trait variation within and

between closely related species. To do so, we experimentally assayed physiological and pigmentation

variation in three population lines from each of three focal taxa within the *D. americana* species group—

D. americana americana, D. americana texana, and *D. novamexicana*, and combined these data with

bioclimate information from each species range and focal population. This approach allowed us to

disentangle the contributions of species-wide trait differences, local adaptation, and pleiotropic trait-

interactions to shaping the overall distribution of variation in these important physiological traits.

Methods

Experimental Fly stocks

Three stocks from each focal species, and one line from an outgroup species (*D. virilis*), were

obtained from the University of California San Diego Drosophila Species Stock Center (DSSC). We used

Drosophila novamexicana stocks from San Antonio, NM, Grand Junction, CO, and Moab, UT (15010-

1031.08, 15010-1031.00, and 15010-1031.04 respectively); *D. americana americana* (hereafter *D. a.*

americana) stocks from Chinook, MT, Chadron, NE, and White River, AR (15010-0951.02, 15010-

0951.06, and 15010-0951.17 respectively); and, *D. americana texana* (hereafter *D. a. texana*) stocks

from New Orleans, LA, Jamestown, SC, and Morrilton, AR (15010-1041.24, 15010-1041.29, and 15010-

1041.23 respectively). The *D. virilis* stock originated from Texmelucan, Puebla, Mexico (15010-1051.48).

D. americana is divided into subspecies according to presence of a chromosomal fusion of the X- and 4-

chromosomes in *D. a. americana* that shows a distinct latitudinal cline (McAllister 2002). For simplicity

we shall refer to them by their subspecies names. All fly stocks were reared on standard cornmeal media

prepared by the Bloomington Drosophila Stock Center (BDSC) at Indiana University, and were kept at

room temperature (~22C).

Environmental data

To quantify environmental variation in the natural range of our three focal species, we extracted

bioclimatic variable data from documented collection locations of each species and used these data to

create principal components that summarize the major axes of climate variation. First, geographical

coordinate data were obtained for all known collections using TaxoDros (www.taxodros.uzh.ch) – a

database that compiles latitude and longitude coordinates from published collection records. After

curating for duplicates and erroneous or unspecified coordinates, we retained passport data for 10 *D.*

novamexicana, 73 *D. a. americana*, and 68 *D. a. texana* population locations. For each of these

geographic locations, we extracted bioclimate variable data from two sources. From the Worldclim 2

database we extracted source location data at 30 arcsecond resolution for 19 bioclimatic variables (Fick

and Hijmans 2017); and, from the CliMond archive (Kriticos et al. 2012) we extracted 16 additional

bioclimatic variables at 10 minute resolution (see supplement). The latter were included despite this

relatively coarse resolution because they contained additional data on ultraviolet (UV) radiation and soil

moisture that were not available in the Worldclim registry. Many of these 35 bioclimatic variables

describe alternative aspects of temperature, precipitation, and seasonality over different time intervals,

so we performed a principal component analysis (PCA) in *R* on all 35 variables across all 151 population

locations to reduce non-independence and redundancy in this dataset. PCA uses orthogonal

transformation to generate a set of linearly uncorrelated principal components that summarize major

axes of variation across a dataset of potentially correlated variables. In our data, the first 3 PCs

explained 95% of the environmental variation across populations; variable loadings for these 3 PCs is

provided in the supplement (Table S1). Because the first 3 PCs explained ~85% of the variation across

populations, we used values for these PCs in our subsequent analyses of the relationship between

environmental variation and variation in physiological traits. Based on the variable loadings (Table S1)

we interpret the environmental significance of each of the first 3 PCs as follows: PC1 is strongly

negatively loaded for the majority of the precipitation and moisture index variables – with the exception

of the seasonality terms; therefore higher PC1 values indicate more arid areas with lower rainfall year-

round, and higher daily and seasonal temperature variability. PC2 is most strongly loaded for radiation

and temperature extremes for the positive values, and negatively loaded for ground moisture variables

and temperature/UV seasonality. We therefore interpret high values of PC2 to indicate year-round high

temperature and radiation, with low moisture, and negative values to indicate areas with year-round

high moisture along with high seasonality in temperature and radiation. Last, negative PC3 values

indicate locations where the hottest and wettest periods of year align, and are accompanied by high

radiation and moisture, while positive PC3 values indicate high temperatures and radiation during the

driest times of year when precipitation is low.

Desiccation resistance assay

To assess population, species, and sex specific differences, desiccation resistance was assayed in

replicate for individual males and females of each of our nine focal populations, using custom

desiccation vials. Virgin flies were isolated within 24 hours of eclosion and aged individually for 3 days

prior to the start of the experiment. Flies were then mouth aspirated individually into a modified

Drosophila culture vial which contained a layer of 20g of Drierite, 10g of desiccated cork, and a piece of

cheesecloth, and was sealed by a layer of parafilm. Each fly was placed above the cheese cloth and cork

(in order to avoid direct contact with Drierite that negatively effects survival) and observed every 15

minutes until death. Desiccation resistance was then quantified as the total time in minutes that each

individual survived in a desiccation vial. A minimum of 5 replicates were performed per sex for each

population. Trials were performed in blocks in which one fly of every identity (population x sex) was

assayed simultaneously, to avoid confounding sex or population effects with trial date. At the end of the

survival assay, each individual was weighed to obtain their dry weight (as a proxy for size) to include as a

covariate in survival analyses. Dry (post-death) weight was determined to be an effective proxy for wet

(pre-desiccation) weight in a pilot experiment in which individuals of each population and sex were

weighed before and after individual desiccation trials (Pearson's correlation, females: $r(8) = 0.899$, $P <$

0.001; males: $r(8) = 0.925$, $P < 0.001$).

UV irradiation resistance assay

We assessed UV-B resistance for each sex within each population (including the *D. virilis* line), at

each of four different exposure intensities: 100, 500, 1000, and 5000 Joules/m^2, plus a control assay at 0

J/m^2. UV resistance trials were performed similarly to Matute and Harris (2013) and Aguilar-Fuentes et

al. (2008). Briefly, virgin males and females of each population were isolated and kept in single-sex

groups of 20 for 24 hours prior to experiment start. Each group of 20 flies was then lightly anesthetized

and weighed as a group before being irradiated with UV-B light at one of the four experimental

intensities using an ultraviolet Stratalinker 2000 (Stratagene, La Jolla, CA). For the 0J exposure - which

essentially measures longevity in the absence of acute UV exposure - flies were simply anesthetized,

weighed, and placed in the Stratalinker without UV exposure. Each group was then transferred to a vial

containing standard cornmeal media and scored once daily for number of flies still alive; groups were

transferred to fresh food vials as often as necessary. The experiment continued until all flies in each vial

were dead. For each assayed energy level, trials for both sexes in all ten lines were initiated

simultaneously, to avoid confounding these factors with date effects.

Pigmentation assay

Dorsal abdominal pigmentation was assessed on individual males and females from each focal

population (and for the *D. virilis* line), in a similar manner to Wittkopp et al. (2010, dataset 'A'). Briefly,

individual 7-day old virgin flies for each sex and population were placed in 10:1 ethanol to glycerol

mixture and stored at room temperature for 1-5 days. The dorsal abdominal cuticle (specifically tergites

A3-A5) was dissected from each fly, isolated from other tissues, and mounted in Hoyer's solution. Each

cuticle was then viewed and digitally imaged on an AMG EVOS FL scope (AMG, Bothell, WA, USA) under

standardized light conditions. Body color was quantified on gray-scale images of each cuticle by

calculating the average median pixel intensity of 20 randomly-selected, non-overlapping regions on a 0-

255 scale (avoiding the dorsal midline which has consistently lighter pigmentation), in Image J (NIH,

Bethesda, MD, USA). Five replicate individuals from each sex within each population were assessed.

Statistical analyses

Trait differences between sex, species, and populations

All statistical analyses were performed in R version 3.4.3, as was figure construction. To first

assess the distribution of variation in our traits, we analyzed each physiological trait for differences

between species, populations, and sex. For each of desiccation resistance and pigmentation, we fit a

multi-way ANOVA with sex, species, and population nested within species, as independent variables,

and each trait as the response variable. For desiccation resistance, dry weight was also included as a

fixed effect to account for individual body size. For both of these traits we also performed post-hoc

contrasts between each pair of species using the Tukey test.

For UV resistance, effects of sex, species, and population were assessed differently because the

data for each group within each treatment are best summarized and compared as Meier-Kaplan survival

curves. A cox proportional hazard model (Therneau 2013) was first used to determine the effect of sex

between all survival curves across treatment levels and species, and sexes were found to differ (Wald

test (1) = 105.2, *P* < 0.0001). Survival also consistently differed between sexes within each treatment

level (see supplement), and all subsequent analyses were performed separately on each sex. To assess

species and population effects on UV resistance within each of the five different treatment levels, two

cox proportional hazard models were constructed: one to test for significant effects of species (where

populations were grouped by species) and the other to test for significant differences between

populations irrespective of species. The Wald test was used to evaluate the overall effects of terms

included in each model. Finally, we used the median of the Meier-Kaplan curve estimate—equivalent to

the day in which 50% of the flies in a given trial are dead—as a summary statistic for UV resistance for

each sex in each population at each given treatment level, for use in subsequent analyses of trait-trait

associations and trait associations with environmental PC axes (see below).

Environmental variation and association with physiological traits

We tested for evidence that species significantly differed in environment from one another by

performing one-way ANOVAs with species as the independent variable and each of the first three PC

axes as the dependent variables. These analyses were performed with data from all collection localities

used to generate the PC axes (N=149), and also on the set of nine focal populations used for our trait

analyses. For each analysis with a significant species effect, we also performed Tukey post-hoc contrasts

to determine which species differ from one another on each PC axis.

To look for evidence of local adaptation (specifically trait-environment associations), for each

sex separately we evaluated the relationship between population physiological trait means (or, for UV

resistance, median survival values) and population values for principal component axes of

environmental variation, while taking into account broader environmental differences between species.

Because our PC axes showed statistical separation among species—that is, *D. novamexicana* had

complete separation from the other two taxa along both PC1 and PC2 axes (Figure 3)—for each PC we

first calculated the residuals from a one-way ANOVA with species as the independent variable, and then

used these residual PC values in our analyses of population-level effects. Then, for each of the first three

PCs separately, we fit an ANOVA with residual PC values and species as independent variables, and the

mean population trait value for either desiccation resistance or pigmentation as the response variable. A

similar set of models were performed with UV resistance data from each UV treatment level, but

median Meier-Kaplan curve estimates (as described above) as the response variable. These analyses

allowed us to simultaneously evaluate the contribution of both species differences and local

environmental variation to variation in each physiological trait, and therefore assess how each PC

contributes to variation in a given trait within each species. Because we performed 6 tests for each trait,

the Bonferroni-corrected significance level is p=0.008 for each trait.

Finally, we examined the strength of pairwise associations between each of our phenotypic

traits of interest (desiccation resistance, pigmentation, and UV resistance at each of five levels), using

Pearson's correlation coefficients. Analyses were performed using population means (because each trait

was measured on different individuals and, for UV resistance, groups of individuals), and on each sex

separately (as there was evidence that each trait is moderately to strongly different between sexes; see

above and results).

Results

Species differed along axes of environmental variation

Across all species collection records (N = 151), we found that species differ along PC1 (F(2, 82.2),

$P < 0.001$) and PC2 (F(2, 59.76), $P < 0.001$), but not PC3 (F(2, 3.4), $P = 0.065$), environmental axes. Post-

hoc Tukey tests indicated that all 3 species differed from one another for both PC1 and PC2 (all P values

< 0.001). Despite modest power, species comparisons using environmental data from only our nine

focal populations were similar: Species differed along PC1 (F(2, 17.4), $P = 0.003$) and PC2 (F(2, 6.6), $P =$

0.030), but not PC3 (F(2, 0.7), $P = 0.550$). For PC1, *D. novamexicana* differed from both *D. a. americana*

($P = 0.033$) and *D. a. texana* ($P = 0.003$), although the two *americana* subspecies did not differ ($P =$

0.100); for PC2, *D. novamexicana* differed from *D. a. americana* ($P = 0.026$), but the other two contrasts

(*D.nov – D. tex: P = 0.147; D.am – D.tex: P = 0.400*) were not significant.

Traits differed among both species and populations and, in some cases, sexes

Abdominal pigmentation differed between species (F(2, 11.9), $P < 0.001$) and between

populations within species (F(6, 3.1), $P = 0.008$), and marginally differed between sexes (F(1, 0.06), $P =$

0.063). *D. novamexicana* were the lightest flies, and post-hoc Tukey tests confirmed *D. novamexicana*

was significantly less pigmented than *D. a. americana* ($P < 0.001$) or *D. a. texana* ($P < 0.001$), while the

two americana subspecies had similar pigmentation (P = 0.96) (Figure 2). With respect to sex-specific

differences, females were marginally less pigmented than males.

Desiccation resistance also differed between species (F(2, 29.8); P < 0.001) as well as between

populations within species (F(6, 4.5); P < 0.001), but not between sexes (F(1, 0.8); P = 0.364); dry weight

(body size) had no effect (F(1, 1.3), P = 0.25). Post-hoc Tukey tests indicated significant pairwise

differences between all three species (*D.nov – D. am: P* < 0.001, *D.nov – D. tex*: *P* < 0.001; *D.am – D. tex:*

P = 0.002); *D. novamexicana* had the greatest desiccation resistance, followed by *D. a. texana* and *D. a.*

americana. Consistent with this, *D. novamexicana* populations generally performed better than the

other populations, with the exception of *D. a. texana* from Morrilton, Arkansas, which had the third

highest survival overall (Figure 3).

Species and population differences for UV resistance were assessed within each treatment and

for each sex separately (see methods and supplement for sex-specific analyses). For females, species

and populations differed in survivorship after 0J, 100J, 500J, and 5000J of UV exposure, but not after

1000J (Table 1). For males we found a significant species effect on survival after 0J and 100J of UV

exposure, but not after 500J, 1000J, or 5000J; results for population-level analyses were the same

except UV exposure to 1000J was also significant (Table 1). Despite these significant differences, the

rank order of UV resistance between populations varied idiosyncratically among different UV exposure

levels, and was generally inconsistent between them (Figure S1).

Pigmentation shows different patterns of environmental association within- and between-species, while

desiccation resistance displays predominantly species-level patterns.

In addition to large significant differences in pigmentation among species for both sexes, we

found that intraspecific pigmentation variation was modestly associated with PC2 for both males and

females, although not with PC1 or PC3 (Table 2) (Figure 3). Based on the factor loadings for PC2 (see

methods and Table S1), this latter finding indicates that higher year-round UV exposure and

temperature coupled with low moisture (i.e. positive values of PC2), are associated with relatively

darker pigmentation within each of these species. This association is particularly interesting as the

broader-scale differences between species show the opposite association, where the species with the

lightest pigmentation—*D. novamexicana*—is clearly associated with positive values for this

environmental axis (Figure 3).

In contrast to pigmentation, for all three PCs we found that species differences were modestly

explanatory of desiccation resistance variation among males (but not females) (Table 2), whereas

intraspecific resistance variation was not associated with residuals of any of the three PCs for either sex

(Table 2) (Figure 4). Based on these results, differential desiccation resistance in males is generally

consistent with large-scale environmental differences between species (in particular, *D. novamexicana*

males are more desiccation resistant), but there is little evidence for additional local adaptive variation

within species for either sex for this trait. For UV resistance, associations with environment were

assessed within each treatment level and sex, but we found only one significant association after

correcting for multiple testing: for the 5000J treatment, females showed a significant association with

PC2 ($F(1) = 12.75$; $P = 0.009$). Full results of these analyses are found in the supplemental material (Table

S3).

For associations between traits, we found that lighter pigmentation is positively correlated with

higher desiccation resistance in males ($r(7) = 0.70$, $P = 0.036$), but only marginally for females ($r(7) =

0.59$, $P = 0.091$)(Figure S2). In contrast, UV resistance was not correlated with pigmentation for either

sex at any of the treatment levels (Table S4). UV resistance was modestly negatively correlated with

desiccation resistance for females at the highest exposure (5000J, $r(7) = -0.720$, $P = 0.029$), but this

association does not survive multiple-testing correction.

Discussion

Here we surveyed natural variation in three physiological traits in the *Drosophila americana*

species group, examined how environmental factors have shaped this variation, and how these traits

might influence each other. We found strong differences between species in their environmental

associations, as well as in pigmentation and desiccation resistance variation, but not in assayed UV

resistance. When we evaluated evidence of adaptation by examining trait-environment associations, we

found evidence for a complex role of natural selection in shaping pigmentation variation; within-species

variation suggests that darker flies occur in consistently hotter, drier, higher radiation environments, but

between-species the opposite pattern is evident. Conversely, desiccation resistance was largely

differentiated by species-level patterns, and UV resistance phenotypes showed no clear association with

measured environmental variation or these other traits. These different patterns of association between

environmental and trait variation, both within and between species, suggests that the temporal and

spatial scale at which selection acts on these traits is different, as are the major selective drivers

responsible for patterns of observed variation.

Pigmentation variation is shaped by alternative factors within and between species

The observation that *Drosophila* species, sexes, and/or populations vary in pigment is common

(see Introduction) and has prompted several, sometimes conflicting, hypotheses about the functional

role of this variation. Many of these hypotheses are based on the general observation that pigmentation

is associated with spatial variation, especially latitudinal gradients: Previous analyses along latitudinal

clines in *Drosophila* have shown evidence for both increasing (David et al. 1985, Munjal et al. 1997,

Telonis-Scott et al. 2011, Capy et al. 1988, Heed and Krishnamurthy 1959) and decreasing (Bastide et al.

2014) melanization with latitude. Here, one of our goals was to more directly evaluate this trait variation

by associating pigmentation with axes of bioclimatic variation (rather than solely geographic patterns),

and with other physiological trait variation, to assess the plausibility of these prior hypotheses in our

focal *Drosophila* species. We found both that our significantly less pigmented species—*D.*

novamexicana—is found in consistently hotter, drier, higher radiation environments, and also that

darker pigmentation is moderately positively associated with these same environmental factors within

each species. Our observations suggest that environment variation imposes different directions of

selection on this trait at the species level versus the local level.

These results also provide new data with which to evaluate three alternative hypotheses

proposed to explain natural variation in pigmentation found in other *Drosophila* species. The first of

these hypotheses—the thermal melanism hypothesis (or Bogert's Rule in ectotherms)—proposes that

more pigment allows better absorption of thermal radiation in colder environments. This proposal has

been commonly invoked to explain a geographical association between darker pigmented ectotherms

and higher latitudes (Gaston et al. 2009). In our data, at the species level there is some evidence that

flies in colder regions are darker, at least when comparing *D. a. americana* to *D. novamexicana*.

However, several other aspects of our data are inconsistent with the association between pigmentation

and colder environments proposed by the thermal melanism hypothesis: *D. a. americana* and *D. a.*

texana do not significantly differ in pigmentation, even though *D. a. texana* is both from lower latitudes

and regions with warmer temperatures. Furthermore, at the population level within all three species we

see the opposite association, where darker flies are associated with more consistently warm and higher

radiation areas.

A second hypothesis proposed to explain pigmentation variation is that it plays a direct role in

UV protection. Bastide et al. (2014) showed in Sub-Saharan African populations of *D. melanogaster* that

darker morphs are found at lower latitudes, and that UV exposure (among other environmental factors)

was the factor most highly correlated with pigmentation variation, from which they inferred that there

is strong evidence that pigmentation plays a role in UV protection (Bastide et al. 2014). Within each

species examined here, our data are consistent with Bastide et al.'s observations, as populations from

areas with higher year-round UV radiation and temperatures (PC2) have darker pigmentation. However,

we found little evidence for strong associations between UV resistance *per se* (as measured by our

assay) and either pigmentation or environment (as discussed below), so the exact causal relationship

between UV resistance and pigmentation in our study remains uncertain. It has been previously shown

in *D. melanogaster* that *ebony* mutants (which are more darkly pigmented) live significantly longer than

lighter pigmented *yellow* mutants when both are exposed to UV irradiation (Wang et al 2008), giving

some credence to the proposal that pigmentation could be important for UV resistance. However, even

if selection imposed by UV stress does indeed shape some of the pigmentation variation within

species—specifically by increasing pigmentation in high UV environments—this cannot explain the

pattern of pigmentation we observe *between* species where the least pigmented species is found in

drier, more consistently hot environments with greater UV exposure (high PC2). This apparent

inconsistency between trait-environment associations at alternative evolutionary scales (within and

between species) itself indicates that UV resistance cannot be a complete explanation of the

pigmentation variation we observe here. Instead, additional patterns of selection and/or demographic

history appear to be responsible for species-level differences—particularly the lighter pigmentation of *D.*

novamexicana—although the nature of these forces and when they acted are harder to determine.

Wittkopp et al. (2009) showed that the yellow body color of *D. novamexicana* is a fixed, derived

phenotype determined by alleles at the *tan* and *ebony* loci, and inferred that these alleles pre-date

speciation, as they can also be found segregating across populations of *D. americana*. The broader

distribution of these shared pigmentation-variation alleles suggests that historical selection could have

played a role in shaping genetic variation prior to speciation, although it doesn't indicate the specific

selective agent(s) responsible. Moreover, despite these shared alleles, contemporary *D. novamexicana*

lineages are clearly significantly less pigmented than even the lightest *D. americana* populations. This

suggests that additional species-specific forces have acted since the split of their common ancestors,

that might include the fixation of light alleles via drift, or the influence of other species-wide selective

agents. For example, although evidence in *Drosophila* is currently lacking (Wittkopp and Beldade 2009),

prior studies indicate that pigmentation can play a role in sexual signaling in other insects (True 2003).

Therefore, exaggerated pigmentation differences between species could result from pleiotropic effects

on mating preferences within and between species, a hypothesis that is testable with future work.

Regardless of the actual cause of species pigmentation differentiation, at present our data suggest that

different selective pressures operating at different times and scales could produce complex, and even

opposing, patterns of environmental association with a single trait, especially when the trait has more

than one biological function.

Aside from the above proposals, a third class of hypotheses about the evolution of pigmentation

focuses on its proposed association with desiccation resistance. One prevailing hypothesis is that

increased melanization confers thicker cuticles that increase tolerance to desiccation (Parkash et al.

2008), and increased melanism is strongly correlated with differences in desiccation resistance in some

Drosophila (Brisson et al. 2005, Rajpurohit et al. 2008, Karan et al. 1998, Parkash et al. 2008). While we

did find a modest correlation between desiccation resistance and pigmentation, the relationship we

detected is in the opposite direction than is proposed under this hypothesis. Furthermore, genetic

analysis in this group indicates little evidence that these traits are mechanistically associated. In

particular, Wittkopp et al. 2010 used *D. novamexicana* and *D. americana* to create backcrossed lines

with *ebony* or *tan* alleles introgressed from the opposite species; they found that introgression of darker

or lighter alleles into the background of the opposing species did not alter desiccation resistance,

providing strong evidence that these traits are not mechanistically associated, at least at these large

effect loci. An alternative to a direct mechanistic connection is that these trait associations could simply

emerge due to their shared covariation with other, causal, axes of environmental variance. This

possibility highlights the need for more studies that simultaneously address these associations between

multiple traits, species, and environmental gradients, in order to better disentangle alternative

explanations of the forces shaping complex patterns of trait variation.

Desiccation resistance is shaped by factors acting among species rather than at local spatial scales

Unlike pigmentation variation, the adaptive connection between species trait differences and

the overall climate of their natural ranges is more readily understood for desiccation resistance: *D.*

novamexicana is from the xeric regions of the US and has significantly elevated desiccation resistance

relative to its mesic sister species, presumably because this confers improved survivorship throughout a

substantially drier species range. It is intriguing, then, that there is no consistent relationship between

desiccation resistance and any PC axis within each of our species. As temperature and UV exposure

become increasingly and consistently high (as shown by increasingly positive values of PC2) desiccation

resistance generally increases across our focal populations (Figure 4); however, species differences

clearly determine this pattern (Table 2), with little evidence for additional within-species local

adaptation in response to this axis. This lack of a stronger association within species could be due to

several non-mutually exclusive factors, although in the absence of more direct data they remain

hypotheses. First, our assay only assesses one element of physiological tolerance to desiccation,

whereas there are multiple ways to avoid desiccation stress in the wild. Populations within our species

might therefore be responding to local variation in desiccation stress via strategies or behaviors that are

not observable in the experimental assay we used here. A similar proposal was suggested by Sillero et al.

(2014) to explain a lack of association between source environment and thermal performance in these

same species. Second, while bioclimatic variables can be very effective in describing macro-level

associations between populations and environment, they do not provide a complete view of microscale

differences in abiotic and biotic environments. Consequently, small scale environmental differences,

such as shade availability or proximity to local water sources, will be unaccounted for in these analyses

even though these might be important elements shaping trait-environment relationships. Finally, there

could be additional within-species constraints on the mechanisms underpinning desiccation resistance,

such that they are either unable to respond more finely to local conditions, or the fitness benefits of

doing so are insufficient to outweigh the benefits of optimizing alternative functions. This is especially

relevant for traits that can serve several different functional roles (i.e. are pleiotropic), and that

experience trade-offs between these functions. One such trait that could be relevant to desiccation is

cuticular hydrocarbon (CHC) composition—the blend of waxy compounds on the surface of insects—

that is known to be important for desiccation resistance in multiple species, but can also play a role in

sexual communication (reviewed Chung and Carroll 2015). For example, if maintaining effective species-

wide CHC-mediated sexual signaling imposes constraints on local physiological responses in CHC-

mediated desiccation resistance, this could lead to a pattern in which species strongly differ in

desiccation resistance (in a way that matches broad environmental variation), but there is no local

pattern of adaptation in the same physiological response. In this way, when responses to a general

physiological stress can be achieved by an amalgam of alternative adaptive 'solutions', some of which

are subject to additional functional constraints that differ between evolutionary and geographical scales,

observed patterns of variation in individual traits need not fully reflect the complexity of these adaptive

trait responses.

Life history differences between sexes, populations, and species might have additional ecological

functions

While our results for pigmentation and desiccation resistance reveal differences at the

population and/or species level that are consistent with selective responses to abiotic environmental

variation, the patterns that emerge from our UV resistance assay are less clear. Although we detected

population and species differences in some individual treatments, these results were generally weak and

idiosyncratic with respect to their direction and magnitude. This might reflect the limitations of

extrapolating natural functional differences in UV tolerance from survivorship after an acute single

exposure to UV (a technique that is commonly used; Matute and Harris 2013, Jacobs 1974, Wang et al.

2008). Exposing flies to moderate to high doses of UV radiation in a single exposure may not be

analogous to UV resistance to low levels of radiation on a daily basis; more realistic assays, with periodic

or sustained exposures of lower average dose, might better simulate differences in natural

environmental conditions. Regardless, this assay did reveal at least one interesting life-history difference

among our groups: a consistent pattern of significantly increased longevity in females relative to males,

regardless of species, as well as longevity differences among some of our populations. Because these

differences were also apparent in the control (0J) exposure, it is likely that they largely reflect baseline

longevity differences between these groups. Our finding is generally consistent with a previous analysis

of longevity in this species group that also found that females tended to live longer than males, at least

in some strains (Durbin and Yoon 1986). Greater female longevity is a pattern seen across many, but not

all, *Drosophila* species (Yoon et al. 1990), and several factors could explain both sex and strain

differences in lifespan, including sex-linkage of the underlying causal factors but also life history

differences associated with ecological variation. The lifespan variation we observe here might therefore

provide another opportunity to assess the ecological significance of additional quantitative trait

variation, in the future.

Conclusions

Our study took a multi-faceted approach to assessing the factors responsible for natural

variation in a set of ecologically-relevant physiological traits. By simultaneously assaying variation both

within and between species, we were better able to discern how geospatial variation can influence

patterns of trait variation in ways that can be complex. Our finding that pigmentation-environment

associations differ between evolutionary scales indicates that pigmentation variation is shaped by

different factors within and between species. The contrasting environment associations found for our

other two traits—few clear associations for assayed UV resistance, but species-level differentiation in

acute desiccation resistance that matched species-wide differences in natural climate conditions--

indicate that while pigmentation, desiccation resistance, and UV resistance frequently appear to covary,

this pattern need not indicate that these traits are directly mechanistically connected. Together, the

specific nature of trait and environment associations observed at each scale provided new data to assess

several conflicting explanations previously proposed for pigmentation variation in *Drosophila*, including

whether pleiotropy plays an important role in determining the distribution of this natural variation. The

multilevel approach that we have taken here can therefore potentially provide more nuanced insights

into complex relationships among traits, and between traits and environmental variation, that both

suggest resolutions to apparently contradictory patterns in the literature and generate additional

hypotheses for future mechanistic and ecological study.

References

Adrion, J. R., Hahn, M. W., & Cooper, B. S. (2015). Revisiting classic clines in Drosophila melanogaster in the age of genomics. Trends in Genetics, 91(2), 165–171. https://doi.org/10.1016/j.chemosphere.2012.12.037.Reactivity

Aguilar-Fuentes, J., Fregoso, M., Herrera, M., Reynaud, E., Braun, C., Egly, J. M., & Zurita, M. (2008). p8/TTDA overexpression enhances UV-irradiation resistance and suppresses TFIIH mutations in a Drosophila trichothiodystrophy model. PLoS Genetics, 4(11), 1–9. https://doi.org/10.1371/journal.pgen.1000253

Bachli, Gerhard (2018). Taxodros: The database on Taxonomy of *Drosophilidae*. www.taxodros.uzh.ch

Bastide, H., Yassin, A., Johanning, E. J., & Pool, J. E. (2014). Pigmentation in Drosophila melanogaster reaches its maximum in Ethiopia and correlates most strongly with ultra-violet radiation in sub-Saharan Africa. BMC Evolutionary Biology, 14(1), 1–14. https://doi.org/10.1186/s12862-014-0179-y

Brisson, J. A., De Toni, D. C., Duncan, I., & Templeton, A. R. (2005). Abdominal pigmentation variation in Drosophila polymorpha: Geographic variation in the trait, and underlying phylogeography. Evolution, 59(5), 1046–1059. https://doi.org/10.1111/j.0014-3820.2005.tb01043.x

Capy, P., David, J. R., & Robertson, A. (1988). Thoracic trident pigmentation in natural populations of Drosophila simulans: a comparison with D. melanogaster. Heredity, 61(2), 263–268. https://doi.org/10.1038/hdy.1988.114

Chung, H., & Carroll, S. B. (2015). Wax, sex and the origin of species: Dual roles of insect cuticular hydrocarbons in adaptation and mating. BioEssays, 822–830. https://doi.org/10.1002/bies.201500014

David, J., Capy, P., Payant, V., & Tsakas, S. (1985). Thoracic trident pigmentation in Drosophila melanogaster: Differentiation of geographical populations. Genetics Selection Evolution, 17(2), 211. https://doi.org/10.1186/1297-9686-17-2-211

Durbin, E. J., & Yoon, J. S. (1986). Longevity in the Drosophila virilis species Group. I. The Drosophila virilis phylad. Ohio Journal of Science, 86(1), 14–17.

Fick, S.E. and R.J. Hijmans, 2017. Worldclim 2: New 1-km spatial resolution climate surfaces for global land areas. International Journal of Climatology.

Fournier-Level, A., Korte, A., Cooper, M. D., Nordborg, M., Schmitt, J., & Wilczek, a. M. (2011). A map of local adaptation in Arabidopsis thaliana. Science, 334(6052), 86–89. https://doi.org/10.1126/science.1209271

Gaston, K. J., Chown, S. L., Calosi, P., Bernardo, J., Bilton, D. T., Clarke, A., … van Kleunen, M. (2009). Macrophysiology: A Conceptual Reunification. The American Naturalist, 174(5), 595–612. https://doi.org/10.1086/605982

Hancock, A. M., Witonsky, D. B., Alkorta-Aranburu, G., Beall, C. M., Gebremedhin, A., Sukernik, R., … Di Rienzo, A. (2014). Adaptations to climate-mediated selective pressures in sheep. Molecular Biology and Evolution, 31(12), 3324–3343. https://doi.org/10.1093/molbev/msu264

Heed WB, Krishnamurthy NB: Genetic studies on the cardini group of Drosophila in the West Indies. Univ
Tex Publ 1959, 5914:155–179.

Jacobs, M. E. (1974). Beta-alanine and adaptation in Drosophila. Journal of Insect Physiology, 20(5), 859–
866. https://doi.org/10.1016/0022-1910(74)90175-9

Kalmus, H. (1941). The Resistance to Desiccation of Drosophila Mutants Affecting Body Colour.
Proceedings of the Royal Society B: Biological Sciences, 130(859), 185–201.
https://doi.org/10.1098/rspb.1941.0011

Karan, D., & Parkash, R. (1998). Desiccation tolerance and starvation resistance exhibit opposite
latitudinal clines in Indian geographical populations of Drosophila kikkawai. Ecological
Entomology, 23(4), 391–396. https://doi.org/10.1046/j.1365-2311.1998.00157.x

Kriticos D.J., Webber B.L., Leriche A., Ota N., Macadam I., Bathols J. & Scott J.K. (2012) CliMond: global
high resolution historical and future scenario climate surfaces for bioclimatic modelling.
Methods in Ecology and Evolution, 3: 53-64.

Matute, D. R., & Harris, A. (2013). The influence of abdominal pigmentation on desiccation and
ultraviolet resistance in two species of drosophila. Evolution, 67(8), 2451–2460.
https://doi.org/10.1111/evo.12122

McAllister, B. F. (2002). Chromosomal and allelic variation in Drosophila americana : selective
maintenance of a chromosomal cline. Genome, 45(1), 13–21. https://doi.org/10.1139/g01-112

Munjal, A., Karan, D., Gibert, P., Moreteau, B., Parkash, R., & David, J. (1997). Thoracic trident
pigmentation in Drosophila melanogaster: latitudinal and altitudinal clines in Indian populations.
Genetics Selection Evolution, 29(5), 601. https://doi.org/10.1186/1297-9686-29-5-601

Parkash, R., Kalra, B., & Sharma, V. (2010). Impact of body melanisation on contrasting levels of
desiccation resistance in a circumtropical and a generalist Drosophila species. Evolutionary
Ecology, 24(1), 207–225. https://doi.org/10.1007/s10682-009-9301-9

Parkash, R., Rajpurohit, S., & Ramniwas, S. (2008). Changes in body melanisation and desiccation
resistance in highland vs. lowland populations of D. melanogaster. Journal of Insect Physiology,
54(6), 1050–1056. https://doi.org/10.1016/j.jinsphys.2008.04.008

Pool, J. E., & Aquadro, C. F. (2007). The genetic basis of adaptive pigmentation variation in Drosophila
melanogaster. Molecular Ecology, 16(14), 2844–2851. https://doi.org/10.1111/j.1365-
294X.2007.03324.x

Rajpurohit, S., Parkash, R., & Ramniwas, S. (2008). Body melanization and its adaptive role in
thermoregulation and tolerance against desiccating conditions in drosophilids. Entomological
Research, 38(1), 49–60. https://doi.org/10.1111/j.1748-5967.2008.00129.x

Sillero, N., Reis, M., Vieira, C. P., Vieira, J., & Morales-Hojas, R. (2014). Niche evolution and thermal
adaptation in the temperate species Drosophila americana. Journal of Evolutionary Biology, 27,
1549–1561. https://doi.org/10.1111/jeb.12400

Telonis-Scott, M., Hoffmann, A. A., & Sgrò, C. M. (2011). The molecular genetics of clinal variation: A
case study of ebony and thoracic trident pigmentation in Drosophila melanogaster from eastern
Australia. Molecular Ecology, 20(10), 2100–2110. https://doi.org/10.1111/j.1365-
294X.2011.05089.x

Therneau, T. M, P. M. Grambsch, 2000 Modeling Survival Data: Extending the Cox Model. Statistics for
594   Biology and Health.

595   True, J. R. (2003). Insect melanism: The molecules matter. Trends in Ecology and Evolution, 18(12), 640–
596   647. https://doi.org/10.1016/j.tree.2003.09.006

597   Wang, Z. P., Liu, R. F., Wang, A. R., Du, L. L., & Deng, X. M. (2008). Phototoxic effect of UVR on wild type,
598   ebony and yellow mutants of Drosophila melanogaster: Life Span, fertility, courtship and
599   biochemical aspects. Science in China, Series C: Life Sciences, 51(10), 885–893.
600   https://doi.org/10.1007/s11427-008-0085-5

601   Wittkopp, P. J., Smith-Winberry, G., Arnold, L. L., Thompson, E. M., Cooley, A. M., Yuan, D. C., …
602   McAllister, B. F. (2011). Local adaptation for body color in Drosophila americana. Heredity,
603   106(4), 592–602. https://doi.org/10.1038/hdy.2010.90

604   Wittkopp, P. J., & Beldade, P. (2009). Development and evolution of insect pigmentation: Genetic
605   mechanisms and the potential consequences of pleiotropy. Seminars in Cell and Developmental
606   Biology, 20(1), 65–71. https://doi.org/10.1016/j.semcdb.2008.10.002

607   Wittkopp, P. J., Stewart, E. E., Arnold, L. L., Neidert, A. H., Haerum, B. K., Thompson, E. M., … Shefner, L.
608   (2009). Intraspecific Polymorphism to Pigmentation in Drosophila. Science, 326(5952), 540–544.
609   https://doi.org/10.1126/science.1176980

610   Yoon, J. S., Gagen, K. P., & Zhu, D. L. (1990). Longevity of 68 species of Drosophila. Ohio Journal of
611   Science, 90(1), 16–32. Retrieved from http://hdl.handle.net/1811/23379
612

613 **Table 1: Population differences in UV survivorship within treatment levels. Survivorship curves were**
614 **constructed for each treatment according to a cox proportional hazard model, and statistical**
615 **differences between populations within treatments was assessed using the Wald test. 9 degrees of**
616 **freedom for each analysis. Bonferroni corrected significance level is *P* < 0.005.**
617

Energy	Male		Female	
	F value	*P* value	*F* value	*P* value
0J	32.42	0.0001	62.88	< 0.0001
100J	31.35	0.0003	72.26	< 0.0001
500J	8.68	0.4669	41.2	< 0.0001
1000J	36.98	< 0.0001	16.9	0.0503
5000J	9.74	0.3719	33.42	0.0001

620 **Table 2: Associations between principal component (PC) axes of environmental variation and**
621 **physiological traits for these species. Reported statistics are from multi-way ANOVA models that**
622 **include a PC (d.f.=1) and species (d.f.=2) as independent variables, and a given physiological trait as**
623 **the dependent variable. Bonferroni corrected significance level is *P* < 0.005.**
624

	Desiccation Resistance				Pigmentation			
	Male		*Female*		*Male*		*Female*	
	F	*P*	*F*	*P*	*F*	*P*	*F*	*P*
PC1	0.89	0.39	0.34	0.58	1.17	0.33	1.12	0.34
Species	7.08	0.035	2.61	0.17	46.43	0.00059	81.05	0.00016
PC2	0.20	0.67	0.33	0.58	7.28	0.043	11.56	0.019
Species	6.25	0.044	2.61	0.17	92.42	0.00011	219.13	<0.0001
PC3	3.64	0.12	3.84	0.11	2.65	0.16	2.95	0.15
Species	10.38	0.017	4.31	0.082	57.60	0.00035	105.25	<0.0001

631
632
Figure 1: Distribution of collection locations and major environmental variables for our three focal
species. Panel A shows collection records for each of *D. novamexicana* (green), *D. a. americana* (blue),
and *D. a. texana* (red) as obtained from the TaxoDros database (taxodros.uzh.ch, see methods).
Closed circles indicate the nine sample locations for populations used in this study. Panel B-D show
heatmaps of spatial variation in annual mean temperature (B), annual mean radiation (C), and annual
mean moisture index (D), as obtained from the Worldclim and Climond databases. Cuticle dissection
images inset into Panel A are representative male cuticles from (left to right) Grand Junction,
Colorado, White River, Arkansas, and New Orleans, Louisiana.

Figure 2: Distribution of individual values for pigmentation (top) and desiccation resistance (bottom, in minutes), according to species (X axis) and population within species (shade variation among points). The y-axis values for individual cuticles on the pigmentation graph correspond to a computed greyscale value represented by the gradient bar. The thick bars indicate species means; thin bars

643
644
645
646

indicate 95% confidence intervals around the mean. Pigmentation differs between both species (F(2,
11.86), P < 0.0001) and populations within species (F(6, 3.13), P = 0.0083). Post-hoc contrasts indicate
D. novamexicana is significantly lighter than *D. a. americana* (P < 0.0001) and *D. a. texana* (P <
0.0001); *D. a. americana* and *D. a. texana* do not differ (P = 0.96). Desiccation resistance differs
between species (F(2, 29.76); P < 0.0001) and populations within species (F(6, 4.48); P = 0.0004). Post-
hoc tests indicate significant differences in all pairwise contrasts (*D.nov – D. am: P* < 0.0001, *D.nov – D.*
tex: P = 0.00039; *D.am – D. tex: P* = 0.0023).
654

Figure 3: Relationship between mean male (triangle) and female (circle) pigmentation in each population and PC1 (top panel) or PC2 (bottom panel). The y-axis values on the pigmentation graph

correspond to a greyscale value represented by the gradient bar. Pigmentation was not associated
with PC1 (annual rainfall and moisture) for either males ($F(1) = 1.17$; $P = 0.33$) or females ($F(1) = 1.12$;
$P = 0.34$) when also accounting for species differences. Pigmentation was modestly associated with
year-round high temperature and radiation (PC2) for both males ($F(1) = 7.28$; $P = 0.043$; solid lines)
and females ($F(1) = 11.56$; $P = 0.019$; dashed lines) when also accounting for species differences (Table
2).
665

Figure 4: Relationship between population mean desiccation resistance and PC1 (top) and PC2
(bottom). Desiccation resistance was not associated with PC1 (annual rainfall and moisture) or PC2,
for either sex, when also accounting for species differences (see Table 2). In comparison there was a
modest species effect when analyzing associations between desiccation resistance and PC1(F(2) =
7.08; P = 0.035) , and with PC2 (F(2) = 6.25; P = 0.044) (Table 2).

674