Title: Cyclic, condition-independent activity in primary motor cortex predicts corrective movement

behavior

Abbreviated Title: Cyclic neural activity in corrective movements

Author: Adam G. Rouse[1]*, Marc H. Schieber[2], Sridevi V. Sarma[3]

Affiliations:

[1]Department of Neurosurgery, Department of Molecular and Integrative Physiology, University of Kansas

Medical Center, Kansas City, KS

[2]Department of Neurology, Department of Neuroscience, Del Monte Institute for Neuroscience,

University of Rochester, Rochester, NY

[3]Department of Biomedical Engineering, Institute for Computational Medicine, Johns Hopkins

University, Baltimore, MD

Corresponding Author and Lead Contact: *Please address correspondence to:

Adam G. Rouse

Department of Neurosurgery

University of Kansas Medical Center

3901 Rainbow Blvd, Mailstop 3021

Rochester, NY 66160

E-mail: arouse@kumc.edu

Pages: 34

Figures: 8

Introduction: 620 words (650 max), Discussion: 1,133 words (1500 max)

Conflict of Interest: The authors declare no competing financial interests.

Abstract

Reaching movements have previously been observed to have large condition-independent neural

activity and cyclic neural dynamics. A new precision center-out task was used to test whether cyclic

neural activity in the primary motor cortex (M1) occurred not only during initial reaching movements but

also during subsequent corrective movements. Corrective movements were observed to be discrete with a

time course and bell-shaped speed profile similar to the initial movements. Cyclic trajectories identified

in the condition-independent neural activity were similar for initial and corrective submovements. The

phase of the cyclic condition-independent neural activity predicted when peak movement speeds

occurred, even when the subject made multiple corrective movements. Rather than being controlled as

continuations of the initial reach, a discrete cycle of motor cortex activity encodes each corrective

submovement.

Significance Statement

During a precision center-out task, initial and subsequent corrective movements occur as discrete

submovements with bell-shaped speed profiles. A cycle of condition-independent activity in primary

motor cortex neuron populations corresponds to each submovement whether initial or corrective, such

that the phase of this cyclic activity predicts the time of peak speed. These submovements accompanied

by cyclic neural activity offer important clues into how the we successfully execute precise, corrective

reaching movements and may have implications for optimizing control of brain-computer interfaces.

Introduction

Corrective movements with sensorimotor feedback are critical for elegant motor control. While a

single, discrete movement like a pointing gesture may be mostly ballistic, more precise aiming

movements typically require an error correction phase (Woodworth, 1899; Craik, 1947; Abrams et al.,

1990; Sainburg et al., 1999; Elliott et al., 2010). Yet the way neurons in motor areas of the brain encode

and generate corrective movements to achieve movement precision is relatively unexplored. Most

experimental tasks to study individual neurons in primary motor cortex have used instructed movements

with a clear onset and offset of movement. These single movement tasks have identified a variety of

movement features being encoded (Evarts, 1968; Thach, 1978; Georgopoulos et al., 1982; Kalaska et al.,

1989; Kakei et al., 1999).

In addition to encoded movement features, however, there is also a large condition-independent

component in the firing rate of neurons in motor cortex (Kaufman et al., 2016; Rouse and Schieber,

2018). Condition-independent neural activity is the change in a neuron's firing rate from baseline that

happens regardless of the instructed movement for any given trial within a given task.   Techniques like

demixed principal component analysis partition a neural population's activity into condition-independent

modulation and the more classical neural tuning to task features (Kobak et al., 2016). This condition-

independent activity is not a simple increase and decrease in firing rate for each neuron but a more

complex neural trajectory (Yu et al., 2007) and has been argued to occur for even discrete movements like

point-to-point reaching (Kaufman et al., 2016), in addition to inherently cyclic movements like

locomotion (Grillner, 1985). The inherent cyclical structure is hypothesized to be related to intrinsic

cortical dynamics (Churchland et al., 2012; Hall et al., 2014). These intrinsic dynamics occur presumably

due to the network of synaptic connections leading to consistent relative lags with the firing of some

neurons reliably occurring earlier relative to other neurons with the generation of movement.

Condition-independent neural activity presents a challenge in describing how corrective

movements might be optimized for precision using the classic models typically ascribed to primary motor

cortex (M1). For experimental tasks that have been designed with reaches in different directions but

similar distances and speeds, it is straightforward to subtract away condition-independent firing rate and

use the remaining firing rate to identify the encoding of movement direction. A linear ("cosine-tuning")

model fits this data well (Georgopoulos et al., 1986; Salinas and Abbott, 1994). Experiments where the

speed of instructed movement has been varied has shown larger changes for faster movements suggesting

some condition-independent neural activity is speed tuning (Moran and Schwartz, 1999). However, these

experiments are all single instructed movements and have not sampled smaller amplitude movements

needed for precise correction. It is unclear if these smaller movements follow the same speed tuning with

very small changes in global firing rate or if some activity is truly condition-independent with any

voluntary movement no matter how small. Additionally, it is unknown how the complex, cyclic neural

trajectories observed during reaching might disrupt or enhance precision when subsequent corrective

movements of various speeds and directions are made.

In the present analysis, we test whether neural dynamics in cortex are repeatable between initial

and corrective movements and examine the condition-independent activity across a wide range of speeds.

A precision center-out task that required moving to small targets was used to elicit visuomotor

corrections. We ask whether corrective movements in our task appear to have discrete submovements

and are similar to initial movements. We then ask whether condition-independent activity is affected by

speed or if the neural modulation appears similar for both small and large movements. Finally, we ask

whether common neural dynamics across movements improve our predictions for when initial and

corrective movements occur.

Materials and Methods

Non-human primates

Two male rhesus monkeys, P and Q (weight 11 and 10 kg, ages 7 and 6 years old, respectively),

were subjects in the present study. All procedures for the care and use of these nonhuman primates

followed the Guide for the Care and Use of Laboratory Animals and were approved by the University

Committee on Animal Resources at the University of Rochester, Rochester, New York.

    Experimental Design

A precision center-out task was performed by the monkey using a joystick to control a cursor on a

24" LCD display. The monkey manipulated a custom, 18 cm long, cylindrical rod attached to the joystick

housing. The joystick moved freely with minimal resistance as the spring mechanism for providing

centering, restorative force was removed. The cursor directly represented measured joystick position and

moved within a 1000 x1000 pixel workspace centered on the display. The cursor workspace limits were

within the physical limits of the joystick. The cursor on the display was a small cross that represented a

single point in the workspace. The custom software for task control sampled the joystick data, updated

the scene, and stored the cursor position (equivalent to joystick position) and trial event times at 100 Hz.

The precision center-out task consisted of three sets of eight peripheral targets located

equidistance and equally spaced in 45˚ intervals around a center, home target (Figure 1). The center

target had a radius of 75 pixels. Each center-out target—defined in polar coordinates—was one of three

different sizes i) large targets spanning 45˚ of the workspace and covering 250-450 pixels from the center,

ii) shallower targets also spanning 45˚ but covering a width only 325-375 pixels from the center, and iii)

narrower targets only spanning 15˚ covering 250-450 pixels from the center. All 24 targets (3 sizes x 8

locations) were presented pseudorandomly in equal amounts throughout a session.

For each trial, following the subject acquiring the home target and performing a required initial

hold ranging from 300-500 ms, the instruction occurred with the given trial's correct target changing from

black to green. Following instruction, the monkey could move the cursor immediately to contact the

correct target. At contact, the outline of all targets changed colors from white to black providing visual

feedback when the cursor was within the target boundaries. After contacting the desired target, the cursor

was required to remain within the target for a variable hold time of 500-600 ms. If the cursor left the

target during a final hold, the monkey was allowed to leave and contact the target again and complete a

final hold. Once a successful final hold of 500-600 ms was completed, the animal received a liquid

reward. Both the required initial and final hold times for each trial were randomly sampled from a

uniform distribution.

A peak finding algorithm to identify local maxima (*findpeaks* function in Matlab (Mathworks,

2020)) was used for analysis of the timing of cursor speed peaks. Cursor speeds were identified as peaks

provided they met the following criteria: i) the peak speed was greater than 250 pixels/s and ii) the peak's

prominence— the height difference between the peak and the larger of the two adjacent troughs

(minimum speed before encountering a larger peak)—was at least 50% of the absolute height of the peak.

Neural Recordings

Floating microelectrode arrays (MicroProbes for Life Science) were implanted in the anterior

bank of the central sulcus to record from primary motor cortex (M1) in each monkey, using methods

described in detail previously (Mollazadeh et al., 2011; Rouse and Schieber, 2016). For monkey P,

recordings were collected from six 16-channel arrays implanted in M1. For monkey Q, two 32-channel

arrays and one 16-channel array in M1 were used. Channels with spiking activity were thresholded

manually on-line and spike-snippets and spike times were collected with the Plexon MAP and Blackrock

Cerebus data acquisition systems. The spike snippets were sorted off-line with a custom, semi-automated

algorithm. Both single- and multi-unit recordings were used in the present analysis.

Statistal Analyses

The firing rates for individual units were estimated by convolving spike times with a Gaussian

filter (σ=30 ms). The firing rates were estimated with 10 ms bin resolution for 100 Hz signals to match

the joystick data and events. The firing rates were then square-root transformed to equalize variance

between high and low firing rates (Kihlberg et al., 1972; Snedecor and Cochran, 1980; Ashe and

Georgopoulos, 1994).

We wished to assess how firing rates changed across a large population of neurons as a function

of time for both initial and corrective movements.   To aid in visualizing this high-dimensional neural

space of all recorded spiking units, we chose to use classical multi-dimensional scaling (also known as

Principal Coordinates Analysis) for dimensionality reduction (*cmdscale* in Matlab). Multidimensional

scaling reduces the high-dimensional dataset to a chosen number of dimensions while preserving as well

as possible the distance (differences in firing rates) between original datapoints.

Dynamical Systems Model

To analyze whether there were repeatable temporal dynamics of the neural population that

correlated with movement, the neural data was modeled as a first-order linear, time-invariant system.

This model was built using only the condition-independent activity by averaging the firing rates for

individual spiking units across all trials regardless of the movement condition (i.e. target location). The

condition-independent activity was submitted to jPCA (Churchland et al., 2012) to identify the two-

dimensional neural plane with the most rotational/cyclic activity. Note, this utilization of the jPCA

algorithm on only the condition-independent activity is different than the typical application of jPCA to

data containing the condition-dependent activity. Once this jPC neural plane was identified, this work

introduces a new analysis variable—condition-independent phase (CIφ)—which estimates at each time

the instantaneous phase by using the Hilbert transfer of the firing rates within the identified two-

dimensional plane. More thorough descriptions of the dynamical systems model, jPCA deminsionality

reduction, and CIφ estimation are given in the Mathematics and Equations section.

Firing Rate vs. Speed Model

As a comparison to our dynamic system model and phase analysis, the average firing rate across

the neural population was also used to identify when peak firing rates occurred and predict future speed

peaks. Since some neurons increase their firing rate and others decrease their firing rate on average with

movement, a linear prediction model was used to create a weighted sum of the neural firing rates to

predict cursor speed peaks. The average firing rate of each unit from 300 ms before to 100 ms after each

peak speed was used in a linear regression to predict the peak cursor speed for each submovement. The

regression model thus generated weighting coefficients for each neuron to movement speed. These

coefficients were then used to weight the instantaneous firing rates at any time point to generate a speed

estimate. While this estimate predicts upcoming movement speed, the estimate from the firing rates is

given with no lag added to report the time when peak firing rate occurred, not the peak of actual

movement.

Circular Statistics

The CIφ values were tested for a significant relationship to the instantaneous cursor speed as well

as the occurrence of speed peaks. All circular statistics—mean, variance, correlation, and Rayleigh test

for non-uniformity—were calculated with CircStat, a Circular Statistics Toolbox for Matlab (Berens,

2009).

Results

Motor Behavior

Movement speed was analyzed throughout the trial from instruction until successful completion

of the final target hold.   The two monkeys successfully completed 10,965 (monkey P) and 8,745

(monkey Q) trials across 12 recording sessions each. All cursor speed peaks that exceeded 250 pixels/s

were identified as submovements within a trial. Initial peaks were identified as the first submovement

that ended at least 150 pixels from the center (approximately halfway to the peripheral target). Any small

movements before the initial speed peak—592 (5.4% of trials) for P and 622 (7.1% of trials) for Q—were

discarded from further analysis. Additional speed peaks following the initial speed peak were defined as

corrective submovements. To focus analysis on submovements made to successfully acquire the target,

corrective submovements were only included if some portion of the acceleration phase—time from

preceding speed trough to speed peak—occurred outside the peripheral target. There were 6480 and 3917

corrective submovements identified for monkeys P and Q, respectively.   Across all trials, 68.3% (P) and

71.1% (Q) were completed in a single initial movement, 17.5% (P) and 20.3% (Q) of trials were

completed with one additional corrective submovement, and 14.2% (P) and 8.6% (Q) of trials required

two or more corrective submovements. The location of the identified speed peaks within example trials

and the speed profiles are shown in Figure 1.   When examining the cursor speed in Figure 1B, the speed

peaks tended to be distinctly separated with nearly zero velocity between most peaks.

Figure 1. A) Cursor paths for a subset of example trials to each of the three target sizes: regular (top), narrow (middle), shallow (bottom). Initial submovements from 200ms before to 200ms after speed peaks are plotted in blue with the point when peak speed occurred shown with a blue dot. Corrective movements are similarly identified in red with a red dot. Grey lines connect the rest of a trial before, between, or after submovements with a speed peak. B) Cursor speed plotted versus time for a subset of trials. Initial (blue) and corrective (red) submovements are identified with squares.

The speed profiles were time aligned to peak speed to better examine the identified submovements (Figure 2A). Almost all submovements show a clear bell-shaped profile for both the initial and corrective movements. The mean peak speeds for initial submovements were 1528 (P) and 1187 (Q) pixels/s

while corrective submovement peaks speeds were 504 (P) and 571 (Q) pixels/s. Thus, the average peaks

for corrective submovements were 33.0% and 48.1% of initial submovements. The time points when

speed was one-half of the maximum speed were identified and the relationship between peak speed and

submovement duration was examined. As shown in Figure 2B, the movement duration to half maximum

speed was similar and close to symmetric for both initial and corrective submovements with initial

submovements being slightly longer on average. Since the target was always the same distance from the

center and the animals' initial reaches tended to get close to the target, there was an inverse relationship

between the time duration of initial reaches and the maximum speed observed (Figure 2C). For corrective

submovements, there was no clear relationship between movement duration and speed. Overall, all

submovement durations as measure by the full width at half maximum time occurred within a similar

range with 96.7%/87.5% (P/Q) of all initial and 96.3%/92.8% (P/Q) corrective submovements between

100-350 ms. The time between speed peaks—either initial to first corrective submovement or between

subsequent corrective submovements—is plotted in Figure 2D. The median time between peaks was 570

ms for monkey P and 700 ms for monkey Q with only 3.2% (P) and 0.1% (Q) of speed peaks having a

time between peaks less than 200ms.

Figure 2. Time course of submovements. A) The cursor speeds are plotted aligned to speed peaks for

initial (blue) and corrective (red) submovements. Squares identify speed peaks while circles are located

at half the maximum of the speed peak. N.B. The cursor speeds shown are before the bandpass filter used

for identifying peaks displayed in Figure 1B. Thus, the maximum of each trace may not align exactly

with the plotted peak speed. B) Histogram of the timing of half-maximum speed of all submovements

before and after peak speed. C) Peak cursor speed plotted versus the duration of the submovement as

measured by the full time width between half-maximum speed peaks (FWHM). D) The time duration

between speed peaks including the times from initial submovement to first corrective submovement as

well as between any consecutive pairs of corrective submovements.

Condition-independent Firing Rates

The neural data was analyzed by also time aligning to the movement speed peaks and examining

the firing rates from 500 ms before until 300 ms after the peak speed. The average number of spikes

(binned in 10 ms time bins) for all analyzed units aligned to the peak speed for initial and corrective

submovements are shown in Figure 3A. A clear peak occurs before the peak speed for both initial and

corrective submovements in both monkeys. Monkey P's peak firing rates occurred 190 ms and 110 ms

before initial and corrective submovements, respectively, while monkey Q's occurred 180 and 170 ms

before. Thus, firing rates increased globally for corrective submovements in addition to the initial reach.

The variance of the condition-independent firing rates from a baseline average for corrective

submovements was 25.5% (P) and 34.4% (Q) of the condition-independent variance for initial

submovements. Notably, the total variance of the neural activity for corrective submovements was 97.3%

(P) and 97.9% (Q) of the total variance for initial submovements. Thus, while the average firing rate

changes were smaller for corrective submovements and consistent with the smaller movement speeds, the

total amount of firing rate modulation that occurred for all the various submovements was nearly as large

for the corrective ones as it was for the initial.

A simple speed tuning model with all neurons having the same time lag would show a

synchronized increase and decrease of all condition-independent firing rates simultaneously. However,

when examining average firing rates from 10 example neurons from one recording session from monkey

P aligned to peak speed, we see heterogenous timing of the rise and fall relative to the peak speed (Figure

3B). This relationship tended to be conserved across initial and corrective movements. For example, the

peak firing rate of the orange neuron (identified by gold arrows) peaked 210 and 170 ms before initial and

corrective peak speeds, respectively, while the gold neuron (gold arrows) peak firing rates were 90 and 40

ms peak movement speed. This suggests that the condition-independent neural activity across the

neurons might form a repeatable temporal structure—a neural trajectory—that is more complex than a

simultaneous rise and fall in firing rate across the population.

271

Figure 3. Mean firing rate relative to peak cursor speed. A) The firing rate for all spiking units was

averaged for all initial (blue) and corrective (red) submovements. Circles indicate the time of peak firing

rate for each condition. B) Average firing rates for 10 example spiking units time aligned for all initial

(left) and corrective (right) submovements. Arrows indicate peaks in firing rate for two example units.

To examine the structure of the condition-independent neural trajectory for initial and corrective

submovements, the distances in the neural space—differences in firing rates—between various time

points was calculated and visualized using multidimensional scaling. Multidimensional scaling identifies

a low-dimensional subspace while preserving the distance between original high-dimensional datapoints

as well as possible. If there was global modulation of firing rate, all the change in condition-independent

firing rates would occur along a single dimension. However, a more complex or cyclic trajectory would

indicate repeatable dynamics within the population of neurons. The time points of -450, -300, -150, 0,

and +150ms relative to peak speed were examined. These times were selected because the highest global

firing rate occurred at approximately -150 ms before peak speed, -450 and +150 ms were times of low

firing rates, and -300 and 0 ms were times of intermediate firing rates as shown in Figure 3. Yet we

observe in Figure 4 that the average distance in the neural space (i.e. difference in all spiking units' firing

rates) between the times of intermediate firing rates (-300 and 0 ms), was only slightly less than the

distance between 300 ms intervals with low versus high (-450 and -150 ms) or high versus low (-150 and

+150 ms) firing rates, for both initial and corrective submovements. This almost circular neural

trajectory—a width nearly equal to height—demonstrates that the condition-independent activity does not

occur as a single rise and fall in global firing rate, but rather as a neural trajectory with certain neuron's

firing rates modulating earlier relative to movement while others modulate later.

Figure 4. Neural distance between condition-independent firing rates. The differences between average firing rates (in arbitrary units) for initial (blue) and corrective (red) for times before (-) and after (+) peak speed (0ms). Distances are shown for three timepoint pairs: -450ms & -150ms, -300 & 0ms, and -150ms & 150ms. The trajectories and distances are shown as a two-dimensional approximation of the high-dimensional neural space using multidimensional scaling. The dimensions for each panel were derived independently with no relationship across panels.

To visualize the high-dimensional firing rates for all recorded spiking units, multidimensional scaling was used to plot the average firing rates of initial and corrective movements in a common 6-dimensional subspace. A complex, multidimensional trajectory was indeed observed for both monkeys. With the initial movements, the average neural activity began (blue triangle) in one region of the neural space before curling into a cyclic trajectory. The corrective neural activity tended to start (red triangle) at

a point near the end of the initial trajectory (blue square) and formed a smaller but similar shaped cyclic

trajectory in the same region of neural activity that occurred during the later portion of the initial reach.

The initial and corrective submovements formed cyclic trajectories in a hyperplane with some but not did

not overlap completely in their features, i.e. the neural trajectories were neither completely parallel nor

completely orthogonal.

Figure 5. Neural trajectories shown in 6-dimensions identified with multidimensional scaling.

Firing rates for each unit were time-aligned relative to peak speed and averaged across all initial and

corrective submovements separately to generate two condition-independent trajectories. The two—initial

and corrective—high-dimensional trajectories were concatenated in time and submitted to classical

multidimensional scaling to create a common 6-dimensional neural space. Each trajectory starts at the

open triangle and ends at the open square. The filled circle signify 150 ms steps of -450, -300, -150, 0,

and 150ms relative to peak speed.

Consistent Neural Dynamics

Despite the smaller magnitude of the condition-independent neural activity during corrective

movements, it appeared likely that a portion of neural activity was repeatable and common for both initial

and corrective submovements. To further test this, we built a simple first-order dynamical system model

using the neural firing rates from both initial and corrective submovements to characterize any common

temporal dynamics that might be present. The neural firing rates were again averaged across all

movement directions to examine only the common condition-independent activity. The plane in the

neural space with the most rotational neural activity was identified using jPCA. We labeled the two

neural dimensions with the most rotational condition-independent activity as CIx and CIy. To compare

consistently across recording sessions as well as monkeys, the pair of dimensions were rotated so CIx

aligned most closely with the time course of average increase in firing rate (shown in Fig 3A).

The average firing rate for all initial and corrective submovements in our identified CI plane are

shown in Figure 6. The CIx and CIy dimensions are plotted as a function of time in Figure 6B while the

neural trajectory in the 2-dimensional CIx/CIy plane are shown in Figure 6A. Considering the initial

versus corrective submovements, the peaks and troughs of both CIx and CIy relative to peak speed, while

differing in magnitude were similar in their timing. The peak in the CIx dimension—defined as the

dimension in the plane that best correlated with the global average firing rate of the population—occurred

approximately 150-200 ms before peak speed for both initial and corrective submovements, whereas the

peak in the CIy dimension occurred near the time of peak speed for both. The initial and corrective neural

trajectories are very similar in their shape and direction of rotation within the plane, with the trajectories

for corrective submovements resembling a smaller, scaled version of the larger trajectories for initial

submovements.

Figure 6. Average firing rate in CIx/CIy plane. A) The average firing rate for initial and corrective submovements are plotted in the CIx/CIy plane identified with jPCA. The trajectories start at the triangles and end at the squares. Each filled circle is a 150 ms time step. B). Average CIx (solid lines) and CIy (dashed lines) plotted as a function of time relative to average cursor speed (dotted lines).

The instantaneous phases of CIx and CIy activity were calculated using the Hilbert transform. The average phase of this neural activity is shown in Figure 7A. The phase of CIx (solid lines) and that of CIy (dashed lines) each were similar for both initial and corrective submovements, with the zero phase of CIx occurring about 150 ms before the peak speed while CIy lagged by approximate $\pi/2$ with the zero crossing around peak speed. Histograms of the phase of CIx and of CIy at peak speed on individual trials is shown in Figure 7B. Mean values of CIx were 0.40π (initial) and 0.43π (corrective) for monkey P and 0.31π (initial) and 0.44π (corrective) for monkey Q. Mean values of CIy were -0.09π (initial) and -0.10π

(corrective) for monkey P and -0.11π (initial) and 0.01π (corrective) for monkey Q. Thus, there was a

clear relationship between peak speed and the phase of condition-independent activity. CIx leads peak

speed by ~ π/2 whereas CIy peaks close to peak speed. Similarly, the time when the phase of CIx = 0 and

CIy = - π/2 was observed to occur 150-200 ms before the peak speed for most initial and corrective

submovements.

Figure 7. Phase of CIx and CIy relative to peak cursor speed. A) Average phase of CIx (solid

lines) and CIy (dashed lines) time-aligned to peak speed (Time = 0) for all initial (blue) and corrective

(red) submovements. B) Histogram of the phase of CIx and CIy at the time of peak speed for initial

(blue) and corrective (red) submovements.   All distibutions of CIx and CIy were non-uniformly

distributed at peak speed (Rayleigh test, all p < 0.001).

Since the phase in the CI plane appeared to predict upcoming speed peaks, we next wanted to

compare this estimate of the timing of speed peaks to using the global sum of the firing rates across the

population. We created a metric we call the condition-independent phase (CIφ) by averaging the phase of

CIx and phase of CIy + π/2 to calculate the current phase in the CI plane. Figure 8A illustrates the

temporal distribution of CIφ relative to the time of peak speed. These distributions consistently peaked

100-150 ms before the speed peak for both initial and corrective submovements. A relatively consistent

relationship between neural activity in this CIx/CIy plane and peak speed thus was present for both initial

and corrective submovements across all trials regardless of target size or reach direction.

For comparison with these results from the dynamical system model, we built a simple linear

regression model to estimate peak speed with a weighted sum of the instantaneous firing rate of all

spiking units. Using this model, we could calculate when the peak firing rate across the neural population

occurred relative to the peak speed. Figure 8B shows the temporal distributions of peak firing rate

relative to peak speed for both initial and corrective submovements. Like the distributions using the

dynamical model above (Figure 8A), those for the firing rate model peaked 150 to 100 ms before peak

speed. The peaks were broader by 15-20 ms, however, as characterized by the corresponding standard

deviations (σ) given for each distribution in the figure. Furthermore, for both the dynamical systems and

firing rate models, a small fraction of submovements did not have a CIφ=0 or peak in the firing rate,

respectively, within the -300 to 100 ms time window illustrated. The percentage of these unaligned trials

(given in the figure) was consistently larger for the firing rate model. Thus, using the temporal dynamics

of the neural population significantly improved our prediction of the timing of the submovements as

compared to using only instantaneous firing rates.

Finally, we examined directly the relationship between cursor speed and the phase of neural

activity in the identified rotational CIx/CIy plane. In figure 8C, we plot the cursor speed as a function of

CIφ. While the CIφ is an angle that ranges between +/-π radians when calculated, it was useful to use a

continuous CIφ to count the number of cycles as it progressed during a trial. Each increase of 2π in CIφ

represents one rotation of neural activity in the CIx/CIy plane. The speed of movement clearly oscillates

with the cyclic neural activity, with the cursor speeds for most trials rising and falling in 2π cycles of CIφ.

The circular correlation between speed and CIφ was 0.43 and 0.42 (p<0.001) with the largest speeds

occurring at CIφ = 0.42π and 0.30π (+2kπ) for monkeys P and Q, respectively. Both the cyclic average

speed (white) and the non-uniform occurrence of peak speeds in individual trials (black) demonstrate that

movement speed was consistently correlated with cyclic, condition-independent brain activity.

Figure 8. Relationship between CIφ and cursor speed. Identifying the times of peak speeds with a

dynamical systems model (A) or with an instantaneous firing rates (B). The time point when CIφ = 0 (A)

or peak firing rate (B) was used as a prediction of the upcoming submovement. Each histogram shows

only those submovements for which the neural data aligned with the movement data, i.e. CIφ = 0 (A) or

maximum firing rate (B) occurred within the time range examined (-300 to 100 ms relative to

submovement peak speed). The percentage of total aligned trials is shown for each distribution as well as

the standard deviation (σ) for the aligned trials. In all cases, the dynamical systems model predictions

were more precise, with a narrower standard deviation (F-test, all p<0.001 for both initial and corrective

submovements for both monkeys) and fewer unaligned trials. C) Cursor speed is plotted as a function of

CIφ for all individual trials with at least one corrective submovement. The average speed as a function of

CIφ is shown in white, illustrating the oscillation in cursor speed depending on the phase of neural

activity. The circular correlations between CIφ and cursor speed for all corrective trials were 0.43 and

0.42 for monkeys P and Q, respectively, p<0.001 in both cases.

Discussion

A consistent relationship with cyclic, condition-independent brain activity preceding both initial

and corrective reaching movements was observed. The precision center-out task utilized small targets to

elicit many trials that were not completed in an initial reach and required corrective movements. The

speed profiles of these corrective movements aligned with cyclic activity in primary motor cortex

suggesting a useful neural marker of the encoding of a series of submovements. Discrete submovements

defined by multiple speed peaks have previously been described in behavioral studies of reaching (Pratt

et al., 1994; Lee et al., 1997; Hatsopoulos et al., 2007; Polyakov et al., 2009), turning a knob (Novak et

al., 2000), isometric contractions (Massey et al., 1992; Hall et al., 2014), and object manipulation for

tactile discrimination (Pruszynski et al., 2018). The experimental results and analysis presented here

provide new evidence of a relationship between neural activity and behaviorally observed submovements.

Distinct movement speed peaks with corrective movements

In the precision center-out task presented here, the monkeys' movements showed consistent bell-

shaped speed profiles. These speed profiles were true for both the larger initial movements from the

center toward the target as well as all subsequent corrective movements.   A large majority of both initial

and corrective submovements had durations of 100-350 ms. For initial movements, the movement

duration was longer for slower reach speed, which was likely due to the reach target being the same

distance for all trials. For corrective movements, there was no relationship between movement duration

and speed with all submovements being approximately the same duration regardless of speed.

Similar but smaller condition-independent activity for corrective movements

436   The magnitude of the condition-independent neural activity that occurred for corrective

submovements was approximately one-third to one-half the magnitude of that for the initial

submovements. This decreased change in average firing rate approximately matched the decrease in

movement speed for the corrective compared to the initial submovements suggesting speed tuning of the

condition-independent activity. This does not imply, however, that each individual trial and each

individual neuron have smaller changes of firing rate during smaller amplitude movements. In fact, the

overall total variance of firing rates—the combined condition-independent activity, condition-dependent

encoding of task variables such as reach direction or target location, and unexplained noise—for

corrective submovements was nearly the same magnitude (>95%) of the initial submovement. Only when

averaging firing rates—time aligned to the peak movement speed or the decoded condition-independent

phase—were the differences in modulation in firing rate between initial and corrective movements

apparent. On average, the encoding of movement speed is clearly present in primary motor cortex. By

identifying the neural dimensions with condition-independent activity, it is possible to identify this speed

dependent encoding. Yet precisely identifying encoded speed on a trial by trial basis within the neural

activity remains challenging.

451   Condition-independent neural activity previously has been observed in motor tasks that are

overtrained to be consistent across trials (Kaufman et al., 2016; Kobak et al., 2016; Rouse and Schieber,

2018). Here, in a precision center-out task with considerable trial-to-trial variability, the condition-

independent neural dynamics were observed to be similar for both initial as well as corrective movements

and were cyclic. The orientation and direction of the cyclic neural modulation was similar with the

corrective submovements having a smaller amplitude on average. Large condition-independent activity

has been suggested to make brain dynamics more robust to noise by increasing the differences in neural

signals even when the muscle activation pattern at certain time points look very similar (Russo et al.,

2018). In the context of corrective movements for a precision task, one might speculate cyclic brain

dynamics organize movements into distinct submovements to create time-varying neural and

musculoskeletal dynamics that are more predictable and robust to noise.

Condition-independent phase predictive of cursor speed

The cyclic CIφ values aligned with cursor speed. This further confirmed the observed cyclic

neural dynamics were behaviorally relevant. The cursor speed peaks and troughs corresponded to a

particular CIφ values further suggesting the encoding of the corrective movements were cyclic with

separate submovements. Rather than continuously moving the hand toward the target at all times, the

subjects' movements appeared as pulsatile bursts of joystick movement and presumably muscle

contraction.

It is difficult to determine how much of the cyclic brain activity and behavior observed is caused

by intrinsic neural dynamics in M1 compared to sensorimotor feedback control and/or cognitive strategy.

Observations of additional submovements defined by second or third speed peaks does not necessarily

require a feedback controller with discrete updates. A single, continuous optimal feedback controller with

appropriate delays and signal dependent noise can also generate additional submovements with multiple

speed peaks (Li et al., 2018). Recent results by Jackson et al. (2018) argue that extrinsic effects of a task

interact with the intrinsic dynamics of the brain in a manner consistent with an optimal feedback

controller and may provide a framework for assessing these effects across a variety of tasks including our

precision center-out task. While difficult to determine in a single task how much the observed behavior is

the result of a cognitive strategy for the given task versus intrinsic sensorimotor cortical dynamics control,

the observed results highlight the importance of cyclic neural dynamics in M1 in at least some corrective

tasks. Further work examining neural activity in various tasks and/or additional sensorimotor brain areas

may lead to clearer descriptions of the relationship between sensorimotor control and motor execution of

precise, corrective movements.

The cyclic dynamics of corrective movements have important implications for brain-computer

interfaces (BCI). To date, most BCI decoders are time-invariant to the timing of any single movement

and when multi-part movements or corrective movements occur. The decoders are typically built initially

with observed or imagined movements that assume single, straight-line movements. When algorithms

for updating BCI decoders consider the change in movement direction for corrective movements, it is

assumed the intended path is updated continuously (Gilja et al., 2012; Shanechi et al., 2016).

Experiments have suggested that BCI control can be improved with two states: active control and rest

(Kim et al., 2011; Williams et al., 2013, 2016; Sachs et al., 2016). The ability to predict the phase of a

movement—like CIφ presented here—might further allow for much better prediction of when the subject

intends to move, contact the target, and make a corrective movement. The cyclic dynamics of the

condition-independent neural activity may also lead to better descriptions of the condition-dependent

activity that encode task features, specifically the observation that direction encoding is often not fixed

but shifts with different phases of movement (Churchland and Shenoy, 2007; Suminski et al., 2015;

Suway et al., 2017), enabling decoders of movement direction that change with time. All leading to a

more robust description of the of neural encoding of precise and corrective movements.

References

Abrams RA, Meyer DE, Kornblum S (1990) Eye-hand coordination: oculomotor control in rapid aimed limb movements. J Exp Psychol Hum Percept Perform 16:248–267.

Ashe J, Georgopoulos AP (1994) Movement parameters and neural activity in motor cortex and area 5. Cereb Cortex 6:590–600.

Berens P (2009) CircStat : A MATLAB Toolbox for Circular Statistics. J Stat Softw 31:1–21.

Churchland MM, Cunningham JP, Kaufman MT, Foster JD, Nuyujukian P, Ryu S, Shenoy K V (2012) Neural population dynamics during reaching. Nature 487:51–56.

Churchland MM, Shenoy K V (2007) Temporal complexity and heterogeneity of single-neuron activity in premotor and motor cortex. J Neurophysiol 97:4235–4257.

Craik KJW (1947) Theory of the human operator in control systems. I. the operator as an engineering system. Br J Psychol 38:56–61.

Elliott D, Hansen S, Grierson LE, Lyons J, Bennett SJ, Hayes SJ (2010) Goal-Directed Aiming: Two Components but Multiple Processes. Psychol Bull 136:1023.

Evarts E V (1968) Relation of pyramidal tract activity to force exerted during voluntary movement. J Neurophysiol 31:14–27.

Georgopoulos AP, Kalaska JF, Caminiti R, Massey JT (1982) On the relations between the direction of two-dimensional arm movements and cell discharge in primate motor cortex. J Neurosci 2:1527–1537.

Georgopoulos AP, Schwartz AB, Kettner RE (1986) Neuronal Population Coding of Movement Direction. Science 233:1416–1419.

Gilja V, Nuyujukian P, Chestek CA, Cunningham JP, Yu BM, Fan JM, Ryu S, Shenoy K V (2012) A

brain machine interface control algorithm designed from a feedback control perspective. Conf Proc

IEEE Eng Med Biol Soc 2012:1318–1322.

Grillner S (1985) Neurobiological bases of rhythmic motor acts in vertebrates. Science 228:143–149.

Hall TM, de Carvalho F, Jackson A (2014) A common structure underlies low-frequency cortical

dynamics in movement, sleep, and sedation. Neuron 83:1185–1199.

Hatsopoulos NG, Xu Q, Amit Y (2007) Encoding of movement fragments in the motor cortex. J Neurosci

27:5105–5114.

Kakei S, Hoffman DS, Strick PL (1999) Muscle and movement representations in the primary motor

cortex. Science 285:2136–2139.

Kalaska JF, Cohen DAD, Hyde ML, Prud'homme M (1989) A comparison of movement direction-related

versus load direction-related activity in primate motor cortex, using a two-dimensional reaching

task. J Neurosci 9:2080–2102.

Kaufman MT, Seely JS, Sussillo D, Ryu SI, Shenoy K V, Churchland MM (2016) The largest response

component in motor cortex reflects movement timing but not movement type. eNeuro

3:ENEURO.0085-16.2016.

Kihlberg JK, Herson JH, Schotz WE (1972) Square Root Transformation Revisited. J R Stat Soc 21:76–

81.

Kim S-P, Simeral JD, Hochberg LR, Donoghue JP, Friehs GM, Black MJ (2011) Point-and-click cursor

control with an intracortical neural interface system by humans with tetraplegia. IEEE Trans neural

Syst Rehabil Eng 19:193–203.

Kobak D, Brendel W, Constantinidis C, Feierstein CE, Kepecs A, Mainen ZF, Qi XL, Romo R, Uchida

N, Machens CK (2016) Demixed principal component analysis of neural population data. Elife 5:1–

36.

Lee D, Port NL, Georgopoulos AP (1997) Manual interception of moving targets. Exp brain Res

116:421–433.

Li Z, Mazzoni P, Song S, Qian N (2018) A Single, Continuously Applied Control Policy for Modeling

Reaching Movements with and without Perturbation. Neural Comput 30:397–427.

Massey JT, Lurito JT, Pellizzer G, Georgopoulos AP (1992) Three-dimensional drawings in isometric

conditions: relation between geometry and kinematics. Exp Brain Res 88:685–690.

Mathworks (2020) Natick, Massachusetts: The MathWorks Inc.

Mollazadeh M, Aggarwal V, Davidson AG, Law AJ, Thakor N V, Schieber MH (2011) Spatiotemporal

variation of multiple neurophysiological signals in the primary motor cortex during dexterous reach-

to-grasp movements. J Neurosci 31:15531–15543.

Moran DW, Schwartz AB (1999) Motor Cortical Representation of Speed and Direction During

Reaching. J Neurophysiol 82:2676–2692.

Novak KE, Miller LE, Houk JC (2000) Kinematic properties of rapid hand movements in a knob turning

task. Exp Brain Res 132:419–433.

Polyakov F, Stark E, Drori R, Abeles M, Flash T (2009) Parabolic movement primitives and cortical

states: Merging optimality with geometric invariance. Biol Cybern 100:159–184.

Pratt J, Chasteen AL, Abrams RA (1994) Rapid aimed limb movements: age differences and practice

effects in component submovements. Psychol Aging 9:325–334.

Pruszynski JA, Flanagan JR, Johansson RS (2018) Fast and accurate edge orientation processing during

object manipulation. Elife 7:e31200.

Rouse AG, Schieber MH (2016) Spatiotemporal Distribution of Location and Object Effects in Primary

Motor Cortex Neurons during Reach-to-Grasp. J Neurosci 36:10640–10653.

Rouse AG, Schieber MH (2018) Condition-dependent neural dimensions progressively shift during reach
    to grasp. Cell Rep 25:3158–3168.

Russo AA, Bittner SR, Perkins SM, Seely JS, London BM, Lara AH, Miri A, Marshall NJ, Kohn A,
    Jessell TM, Abbott LF, Cunningham JP, Churchland MM (2018) Motor Cortex Embeds Muscle-like
    Commands in an Untangled Population Response. Neuron 97:953-966.e8.

Sachs NA, Ruiz-Torres R, Perreault EJ, Miller LE (2016) Brain-state classification and a dual-state
    decoder dramatically improve the control of cursor movement through a brain-machine interface. J
    Neural Eng 13:016009.

Sainburg RL, Ghez C, Kalakanis D (1999) Intersegmental dynamics are controlled by sequential
    anticipatory, error correction, and postural mechanisms. J Neurophysiol 81:1045–1056.

Salinas E, Abbott LF (1994) Vector reconstruction from firing rates. J Comput Neurosci 1:89–107.

Shanechi MM, Orsborn AL, Carmena JM (2016) Robust Brain-Machine Interface Design Using Optimal
    Feedback Control Modeling and Adaptive Point Process Filtering. PLoS Comput Biol 12:1–29.

Snedecor G, Cochran W (1980) Statistical Methods, 7th ed. Ames, IA: Iowa State UP.

Suminski AJ, Mardoum P, Lillicrap TP, Hatsopoulos NG (2015) Temporal evolution of both premotor
    and motor cortical tuning properties reflect changes in limb biomechanics. J Neurophysiol
    113:2812–2823.

Susilaradeya D, Xu W, Hall TM, Galan F, Alter K, Jackson A (2018) Extrinsic and Intrinsic Dynamics in
    Movement Intermittency. bioRxiv.

Suway SB, Orellana J, McMorland AJC, Fraser GW, Liu Z, Velliste M, Chase SM, Kass RE, Schwartz
    AB (2017) Temporally Segmented Directionality in the Motor Cortex. Cereb Cortex 12:1–14.

Thach WT (1978) Correlation of Neural Discharge with Pattern and Force of Muscular Activity, Joint

Position, and Direction of Intended Next Movement in Motor Cortex and Cerebellum. J

Neurophysiol 41:654–676.

Williams JJ, Rouse AG, Thongpang S, Williams JC, Moran DW (2013) Differentiating closed-loop

cortical intention from rest: building an asynchronous electrocorticographic BCI. J Neural Eng

10:046001.

Williams JJ, Tien RN, Inoue Y, Schwartz AB (2016) Idle state classification using spiking activity and

local field potentials in a brain computer interface. 2016 38th Annu Int Conf IEEE Eng Med Biol

Soc:1572–1575.

Woodworth RS (1899) The accuracy of voluntary movement. Psychol Rev 3:1–119.

Yu BM, Kemere C, Santhanam G, Afshar A, Ryu S, Meng TH, Sahani M, Shenoy K V (2007) Mixture of

trajectory models for neural decoding of goal-directed movements. J Neurophysiol 97:3763–3780.

Mathematics and Equations

<u>Calculation of jPCA plane and Condition-Independent Phase</u>

Since the trial data contains corrective movements in addition to the large initial movements that
were not precisely time aligned to trial events for averaging condition-independent neural activity, we
developed a novel algorithm to iteratively average the firing rates, calculate CIφ, then average the firing
rates again based on the CIφ. This iterative process involves three steps: i) Each unit's firing rate is
averaged across all trials to determine its condition-independent firing rate. ii) Dimensionality reduction
is performed using PCA and jPCA on the condition-independent firing rates to identify the neural plane
with the most rotational/cyclic condition-independent activity. iii) The instantaneous phase is calculated
using the Hilbert transform on the first two jPC dimensions for all data points. A schematic of the
iterative algorithm is shown in Figure A1.

i)   Trial averaging to identify condition-independent activity

The condition-independent neural activity is the average firing rate for each recorded spiking unit
for all experimental trials regardless of the movement condition (ie target location). For classic
neurophysiology experiments, this can be calculated by averaging time-aligned data. However, since
trials were of varying durations and many included corrective movements, simple time alignment of trials
was difficult. Data was therefore aligned based on each time-point's calculated CIφ rather than absolute
time.

Since CIφ depends on the averaged condition-independent activity and the condition-independent
activity was averaged by CIφ alignment, an iterative approach was required. An initial estimate using
simple time-alignment averaging of all initial and corrective submovements was used for the first
iteration. For each subsequent iteration, the condition-independent firing rates for each unit were
calculated by averaging all data points when the CIφ values were similar. The average firing rate was

estimated using a sliding window of CIφ values with a step size of π/50 and a window size of π/25 to

generate 100 equally spaced samples ranging from -π to π.

ii)   *PCA and jPCA to identify rotational/cyclic neural activity*

Next, the condition-independent firing rates were submitted to PCA and jPCA (Churchland et al.,

2012) to identify the neural dimensions with the most cyclic activity. PCA was performed on the high-

dimensional neural space to reduce the condition-independent firing rates to the six dimensions with the

most variance. jPCA was then performed on this six-dimensional space. jPCA is a dimensionality

reduction technique to identify the neural planes with the most rotational activity and is more fully

described in Churchland et al (2012). Briefly, jPCA fits a first order dynamical system model to the

neural activity : $\dot{X} = AX$

to predict change in firing rate $(\dot{X})$ based on the current firing rate (X). The transform matrix of this

model (A) can be separated into a symmetric matrix representing pure scaling and a skew-symmetric

matrix representing pure rotational dynamics. By taking the eigendecomposition of the skew-symmetric

matrix, we obtain pairs of purely imaginary eigenvalues and corresponding eigenvectors that define

planes of rotation in the neural space rank-ordered from greatest to least rotation. In the present analysis,

only the first plane with the greatest condition-independent rotation was used and we defined the two

dimensions of the plane as CIx and CIy.   Additionally, to obtain a consistent CIφ across recording

sessions, CIx was defined as the dimension in the CIx/CIy plane with the most variance. The positive

CIx direction was defined as having more positive than negative coefficients in the neural space which

corresponds to the direction where more units have an increased firing rate. Choosing this convention

causes an increase in CIx to generally align with the onset of initial movement, since a majority of units

increase firing rates at the onset of movement.

iii)   *Instantaneous phase estimate*

Finally, the instantaneous phase were estimated by i) bidirectional bandpass filtering of the

activity in both the CIx and CIy dimensions between 0.5-5 Hz with a 1[st] order Butterworth filter, ii)

performing the Hilbert transform of both filtered signal (s_x and s_y) to generate a transformed signal

($\hat{s}_x$ and $\hat{s}_y$) that is a 90° phase shift of every Fourier component in the frequency domain to create an

analytic representation of CIx and CIy, and iii) calculating the angle of the resulting analytic signal

($s_1 + i\hat{s}_1$) to estimate the instantaneous phase in each dimension, and iv) which we then sum (with a $\frac{\pi}{2}$

phase shift added to ϕy) to obtain a single instantaneous phase estimate for the neural activity within the

plane which we call the condition-independent phase ($CI\varphi$).

i) $s_x(t) = filt_{0.5-5Hz}[CIx(t)]$;

ii) $\hat{s}_x(t) = \mathcal{H}[s_x(t)]$

iii) $\phi x(t) = \arg\{s_x(t) + i\hat{s}_x(t)\}$

iv) $CI\phi(t) = \phi x + (\phi y + \frac{\pi}{2})$

$Note -$ i, ii, and iii are performed for both x and y

The bandpass filtering reduces the low-frequency drift and high-frequency variability in CIx to generate a

more consistent subsequent phase estimate. The condition-independent phase (CIφ) thus represents the

instantaneous phase in the dimension of the neural space that has the most cyclic, condition-independent

activity.

Matlab code to calculate the CIφ is available at:

https://github.com/arouseKUMC/CIphase

Figure A1. Schematic of CIφ estimation. The cyclic condition-independent neural activity was estimated

using an iterative technique. After initial time-alignment and averaging of the firing rates, PCA and jPCA

was performed to estimate the dimensions neural activity with the greatest cyclic condition-activity

were identified. Using the phase of this first jPC dimension, the firing rates were realigned for

averaging. The PCA and jPCA were again performed a total of three times. The final CIφ was then used

for the presented analysis.

683