Title: Cyclic, condition-independent activity in primary motor cortex predicts corrective movement behavior

Abbreviated Title: Cyclic neural activity in corrective movements

Author: Adam G. Rouse[1]*, Marc H. Schieber[2], Sridevi V. Sarma[3]

Affiliations:

[1]Department of Neurosurgery, Department of Molecular and Integrative Physiology, University of Kansas Medical Center, Kansas City, KS, 66160

[2]Department of Neurology, Department of Neuroscience, Del Monte Institute for Neuroscience, University of Rochester Medical Center, Rochester, NY, 14642

[3]Department of Biomedical Engineering, Institute for Computational Medicine, Johns Hopkins University, Baltimore, MD, 21218

Author Contributions: AR and MS Designed research; AR Performed Research; AR and SS Analyzed data; AR, MS and SS Wrote the paper

Corresponding Author and Lead Contact: *Please address correspondence to:

Adam G. Rouse

Department of Neurosurgery

University of Kansas Medical Center

3901 Rainbow Blvd, Mailstop 3021

Kansas City, KS 66160

E-mail: arouse@kumc.edu

Figures: 7 figures and 3 tables in manuscript, 1 extended data item

Word Counts: Abstract 135, Significance Statement 80, Introduction: 647 words, Discussion: 1,632 words

Acknowledgements: This work was supported by NIH NINDS K99/R00 NS101127.

Conflict of Interest: The authors declare no competing financial interests.

Abstract

Reaching movements are known to have large condition-independent neural activity and cyclic neural dynamics. A new precision center-out task was performed by rhesus macaques to test the hypothesis that cyclic, condition-independent neural activity in the primary motor cortex (M1) occurs not only during initial reaching movements but also during subsequent corrective movements. Corrective movements were observed to be discrete with time courses and bell-shaped speed profiles similar to the initial movements. Condition-independent cyclic neural trajectories were similar and repeated for initial and each additional corrective submovement. The phase of the cyclic condition-independent neural activity predicted the time of peak movement speed more accurately than regression of instantaneous firing rate, even when the subject made multiple corrective movements. Rather than being controlled as continuations of the initial reach, a discrete cycle of motor cortex activity encodes each corrective submovement.

Significance Statement

During a precision center-out task, initial and subsequent corrective movements occur as discrete submovements with bell-shaped speed profiles. A cycle of condition-independent activity in primary motor cortex neuron populations corresponds to each submovement, such that the phase of this cyclic activity predicts the time of peak speeds—both initial and corrective. These submovements accompanied by cyclic neural activity offer important clues into how we successfully execute precise, corrective reaching movements and may have implications for optimizing control of brain-computer interfaces.

Introduction

Corrective movements based on sensorimotor feedback are critical for elegant motor control. While a single, discrete movement like a pointing gesture may be mostly ballistic, more precise aiming movements typically require an error correction phase (Woodworth, 1899; Craik, 1947; Abrams et al., 1990; Sainburg et al., 1999; Elliott et al., 2010). In making an online correction, the brain must respond

to updated sensory information about the current position relative to the desired target. Yet the way

neurons in motor areas of the brain encode and generate corrective movements to achieve movement

precision is relatively unexplored. When examining populations of neurons in primary motor cortex

during instructed movements, predictable dynamics of neural spiking occur with a progression from

initiation to completion of a movement (Maynard et al., 1999; Jackson et al., 2003; Truccolo et al., 2005;

Sarma et al., 2010). Yet behaving animals also respond to updated sensorimotor information, as happens

in tasks that require precision. For corrective movements with new sensory information, does the neural

activity update as a continuation of the initial reach or does it return to an initial state and cycle again for

additional submovements?

We investigated the neural dynamics underlying corrective movements, focusing on two key

features of neural activity in primary motor cortex that have been previously described during reaching: i)

condition-independent neural activity and ii) rotations in neural dynamics. Although individual neurons

in primary motor cortex encode a variety of condition-dependent movement features (Evarts, 1968;

Thach, 1978; Georgopoulos et al., 1982; Kalaska et al., 1989; Kakei et al., 1999), there is also a large

condition-independent component in the firing rate of neurons in motor cortex (Kaufman et al., 2016;

Rouse and Schieber, 2018). Condition-independent neural activity is the change in a neuron's firing rate

from baseline over time that happens regardless of the instructed movement for any given trial within a

given task. Condition-independent activity presumably carries information on the timing of movement as

opposed to specific, condition-dependent features.   Techniques like demixed principal component

analysis can partition a neural population's activity into condition-independent modulation and the more

classically described condition-dependent tuning to task conditions (Kobak et al., 2016). In addition to

being condition-independent or -dependent, changes in firing rate in theory might be temporally

synchronous across a population. But in practice, primary motor cortex neurons have an asynchronous

range of onset latencies before movement, with latencies for most corticomotoneuronal (CM) cells

ranging from 120ms to 0ms (Cheney and Fetz, 1980) while other motor cortex neurons can lead

movement by up to 200ms (Moran and Schwartz, 1999). Because the increases and decreases in firing

rates are not synchronous, the population activity forms a more complex trajectory in neural state space

(Yu et al., 2007; Cunningham and Yu, 2014). While the precise meaning of these features of neural

encoding remains debated (Churchland et al., 2012; Hall et al., 2014; Lebedev et al., 2019), these shifts

between different combinations of active neurons leads to shifts through different dimensions of the

neural space, often forming a cyclic neural trajectory.

We hypothesized that if the primary motor cortex handles online corrections as ongoing

adjustments to a single reach, then one cycle of the neural trajectory would include both the initial and the

corrective submovements. In contrast, if the primary motor cortex handles each correction as a distinct

(albeit smaller) movement, then each corrective submovement would correspond to its own cycle. We

used a precision center-out task that required moving to small targets (either narrow or shallow) to elicit

visuomotor corrections. We examined whether corrective movements in this task were simple

adjustments in the ongoing reach or discrete submovements, behaviorally similar to initial movements.

We then ask whether condition-independent activity—representing the time course of movement

irrespective of its direction or amplitude—is similar for both initial and corrective submovements.

Finally, we ask whether cyclic neural dynamics improve our predictions of when initial and corrective

movements occur.

Materials and Methods

Non-human primates

Two male rhesus macaque monkeys, P and Q (weight 11 and 10 kg, ages 7 and 6 years old,

respectively), were subjects in the present study. All procedures for the care and use of these nonhuman

primates followed the Guide for the Care and Use of Laboratory Animals and were approved by the

University Committee on Animal Resources at the University of Rochester Medical Center.

A precision center-out task was performed by the monkey, using an 18 cm handle attached to a

commercial joystick (M212 series joystick, PQ Controls Inc.) to control a cursor on a 24" LCD display.

The joystick handle moved freely with minimal resistance as the spring mechanism for providing

centering, restorative force was removed. The end of the joystick could move approximately 9.3 cm in

both the forward/backward and left/right directions. Motion of the joystick was transduced linearly by

two Hall effect sensors sliding in both the backward/forward and left/right directions. The cursor viewed

by the monkey directly represented the planar position of these two sensors scaled to fit within a 1000

horizontal x 1000 vertical pixel workspace in the center of the LCD display. The limits of the cursor

workspace were slightly within the physical limits of the joystick, with 110 pixels corresponded to

approximately 1 cm of movement at the end of the joystick. The cursor appeared on the display as a

small cross centered on a single pixel in the workspace. Custom software for task control sampled the

joystick data, updated the scene, and stored the cursor position (equivalent to joystick position) and trial

event times at 100 Hz.

The precision center-out task consisted of three sets of eight peripheral targets located

equidistance and equally spaced in 45˚ intervals around a center, home target (see Figure 2). The center

target had a radius of 75 pixels. Each center-out target—defined in polar coordinates—was one of three

different sizes i) large targets spanning 45˚ of the workspace and covering 250-450 pixels from the center,

ii) shallow targets spanning 45˚ but covering a width of only 325-375 pixels from the center, and iii)

narrow targets spanning 15˚ covering 250-450 pixels from the center. All 24 targets (3 sizes x 8

locations) were presented pseudo-randomly in equal amounts throughout a session.

For each trial, following the subject acquiring the home target and performing a required initial

hold ranging from 300-500 ms, the instruction occurred with the given trial's correct target changing from

black to green. Following this instruction, the monkey could move the cursor immediately to contact the

correct target. At contact, the outline of all targets changed colors from white to black providing visual

feedback that the cursor was within the target boundaries. After contacting the desired target, the cursor

was required to remain within the target for a variable hold time of 500-600 ms. If the cursor left the

target during this hold, the monkey was allowed to enter the target again and complete a final hold. Once

a successful final hold of 500-600 ms was completed, the animal received a liquid reward. Both the

required initial and final hold times for each trial were randomly sampled from a uniform distribution.

Neural Recordings

Floating microelectrode arrays (MicroProbes for Life Science) were implanted in the anterior lip

and bank of the central sulcus to record from primary motor cortex (M1) in each monkey, using methods

described in detail previously (Mollazadeh et al., 2011; Rouse and Schieber, 2016). For monkey P,

recordings were collected from six 16-channel arrays implanted in M1. For monkey Q, two 32-channel

arrays and one 16-channel array in M1 were used. The location of the implanted arrays, spanning the

forelimb representation in M1, have been previously reported (Fig. 2 of Liu and Schieber, 2020) and

spanned the forelimb area of M1. Intracortical microstimulation on single electrodes with a current up to

a maximum of 100 µA (12 biphasic pulses, 0.2ms pulse width per phase, 3ms interpulse interval) with the

animal lightly anesthetized with ketamine evoked a variety of forelimb movements. Of the 96 electrodes

for monkey P, stimulation of 11 sites elicited proximal arm movements, 6 sites elicited wrist movements,

and 21 sites elicited movement of the digits. Of the 80 electrodes for monkey Q, 34 sites were proximal,

9 sites were wrist, and 25 were digits. During recording sessions, channels with spiking activity were

thresholded manually online, and spike-waveform snippets and spike times were collected with Plexon

MAP (Plexon, Inc.) and Cerebus (Blackrock Microsystems, LLC.) data acquisition systems. The spike

snippets were sorted off-line with a custom, semi-automated algorithm. Chronic multielectrode arrays do

not always yield well-isolated single-unit recordings. To define likely single units, we utilized the signal

to noise ratio of the sorted spike waveforms and the percent of true single unit spikes estimated from a

formula using the number of interspike interval (ISI) violations less than 1ms (Hill et al., 2011; Rouse and

Schieber, 2016). Using a signal to noise ratio of SNR > 3 and 100% true single unit spikes (no ISI

violations) to define definite single units and SNR > 2.5 and >90% true single unit spikes to define

probable single units, 543 (monkey P) and 304 (monkey Q) of sorted spike waveforms were classified as

definite single units while 268 (P) and 208 (Q) additional units were probable single units. Thus,

811/1293=63% (monkey P) and 512/1185 = 43% (monkey Q) of all spiking units were classified as likely

single units. Because including multi-units would be unlikely to provide results more significant than

similar numbers of single-units, we included both single- and multi-unit recordings in our analyses.

Behavior Analyses

A peak finding algorithm to identify local maxima was used for analysis of the timing of cursor

speed peaks. Off-line, cursor speed was calculated by filtering the cursor position with a 10-Hz low-pass

1st-order Butterworth filter (bidirectionally for zero phase lag) and then calculating the first derivative

using the 5-point central difference. Local maxima of cursor speeds (identified with *findpeaks* function in

Matlab (Mathworks, 2020)) were identified as peaks if they met the following criteria: i) the peak speed

was greater than 250 pixels/s and ii) the peak's prominence— the height difference between the peak and

the larger of the two adjacent troughs (minimum speed before encountering a larger peak)—was at least

50% of the absolute height of the peak. All such cursor speed peaks with their surrounding ±200 ms time

windows were considered submovements within a trial. Initial peaks were identified as the first

submovement that ended at least 150 pixels from the center (approximately halfway to the peripheral

target). Any small movements before the initial speed peak—506 (4.6% of trials) for P and 616 (7.0% of

trials) for Q—were discarded from further analysis. Speed peaks following the initial speed peak were

defined as corrective submovements. To focus analysis on submovements made to successfully acquire

the target, corrective submovements were only included if some portion of the acceleration phase—time

from preceding speed trough to speed peak—occurred outside the peripheral target.

The speed profiles for individual submovements were analyzed between -200 and 200 ms relative

to peak speed. As a measure of similarity between speed profiles, the Pearson's correlation between these

speed profiles for pairs of submovements was calculated, yielding a similarity score between -1 and 1. To

measure how similar corrective submovements were to initial submovements, the correlation of each

initial submovement to a randomly selected corrective submovement was calculated. As a ceiling

comparison, each initial submovement was also compared to another randomly selected initial

submovement. Thus, the distribution of correlations for initial-corrective submovement pairs was

compared to the distribution of initial-initial pairs.

Identifying condition-independent, rotational neural activity

We focused our neural population analysis on the neural dimensions that contained the most

condition-independent, rotational activity. A schematic illustration of these two features—i) condition-

independent vs. -dependent, and ii) synchronous vs. rotational/asynchronous is shown in Figure 1. The

condition-independent activity is the time-varying average of firing rate across all trials regardless of

condition while the condition-dependent is the specific tuning to task condition like target direction.

Synchronous, time-locked activity represents changes in firing rate that happen simultaneously across the

neural population, while asynchronous activity of varying time course in different neurons can lead to

patterns of traveling waves or oscillations in the population with a predictable progression in time.

Firing rates of the neural population can be visualized as either: i) a function of time (Fig. 1B) or

ii) neural trajectories in a Cartesian neural space where each neuron's firing rate is plotted on an

orthogonal dimension (Fig. 1C). For a complex task with variable corrective movements such as our

precision center-out task, the condition-independent activity provides a useful analysis to identify the

neural activity underlying a submovement. Although synchronous rise and fall of firing rate across the

neural population may provide some information, the rotational activity from sequential firing rate

changes across different neurons has the potential to improve identification of corrective submovements.

171
Figure 1. Idealized representation of both the synchronous and rotational components of condition-

independent and -dependent changes in neuronal firing rate. A) The firing rates for four neurons (blue,

orange, yellow, and purple) are shown for reaches to four target directions (light to dark grayscale). The

overall firing rates differ for both the four neurons and the four target condition. By time averaging

across the four conditions, the condition-independent firing rates and the residual condition-dependent

firing rates are both identified. B) Next, averaging across the population reveals that firing rates are i)

synchronous activity across all neurons at each time point and ii) the remaining, asynchronous/rotational

firing rate changes specific for each neuron. C) The neural space visualizes the population activity by

showing each neuron's firing rate as a point along an orthogonal dimension with time represented as a

trajectory through this space. In this representation, the difference between synchronous and rotational

activity is better appreciated. Synchronous activity is movement along a single neural dimension while

rotational activity is movement between dimensions. Note, the dimensions defined by individual neurons

are shown projected in a 2D plane. Only the given component (synchronous/rotational and condition-

independent/-dependent) are shown for these four example neurons for visualization purposes. In a much

higher dimensional space when recording from a large number of neurons, the possibility of finding

dimensions with little overlap between components is much greater.

Dynamical Systems Model

We employed dynamical system modeling to characterize repeated changes in firing rates across

our recorded neural population. To identify and analyze potential repeatable temporal dynamics of the

neural population that correlated with movement, our neural data was modeled as a linear, time-invariant

system using a system of coupled first-order ordinary differential equation defined by a transform matrix.

This model was built using only the condition-independent activity by averaging the firing rates for

individual spiking units across all trials regardless of the movement condition (i.e. target location). The

condition-independent activity was then submitted to jPCA (Churchland et al., 2012) to identify the two-

dimensional neural plane with the most rotational/cyclic activity. In this model, the changes in firing rate

can grow/shrink along a single dimension (synchronous) as well as rotate across dimensions

(asynchronous). The eigen decomposition of the transform matrix yields eigenvalues with the real part

representing growing or shrinking away from the origin while the imaginary part represents rotations.

Note, this utilization of the jPCA algorithm on only the condition-independent activity is different than

the typical application of jPCA to data containing the condition-dependent activity. Additionally, we find

the results of the dynamical system are more stable when the firing rates are square-root transformed to

equalize variance between high and low firing rates (Kihlberg et al., 1972; Snedecor and Cochran, 1980;

Ashe and Georgopoulos, 1994) and thus performed this transform before submitting firing rates to jPCA.

We call the plane with the most rotation the condition-independent (CI) plane and define the two

neural dimensions that define this plane as CIx and CIy. To consistently define CIx and CIy across

recording sessions and monkeys, we defined the +CIx direction as the neural dimension that had the

maximum average firing rate. This was performed by calculating the population averaged firing rate at

all angles in the plane and rotating the CIx and CIy axes so that +CIx aligned with the largest firing rate.

Having identified this jPC neural plane, our work introduces a new analytic variable—condition-

independent phase (CIφ)—which estimates the instantaneous phase angle within this two-dimensional

plane of the projected population firing rates. We calculate CIφ using the Hilbert transform applied to the

two signals, CIx and CIy, generating a complex, analytical representation of the population signal. The

angle of this complex signal is then used to calculate the instantaneous phase. The Matlab code and

additional documentation about the calculation of CIφ as described in the paper is freely available online

at https://github.com/arouseKUMC/CIphase. The code is also available as Extended Data 1.

The calculation of the jPC plane and the CIφ was performed using 5-fold cross-validation. Each

recording session was divided into 5 testing sets of trials each containing 20% of the data. The jPC plane

was calculated by training on the other 80% of the data and then tested on each test set. All presented

results for CIφ are using the test data projected into the jPC dimensions identified by the separate training

set.

Firing Rate vs. Speed Model

For comparison with our dynamical system model and phase analysis, we also used a standard

linear regression model to predict the timing of speed peaks from firing rates across the neural population,

using the synchronous, condition-independent neural activity. The average firing rate of each unit from

300 ms before to 100 ms after each peak speed was submitted to a linear regression to predict the peak

cursor speed for each submovement. The regression model thus generated weighting coefficients for each

neuron's prediction of movement speed. These coefficients were then used to weight the instantaneous

firing rates at any time point to generate a speed estimate.

Statistics

Several statistical analyses (Table 1) were used to assess how similar corrective submovements

were to initial submovements and whether there were repeated cycles of neural activity and if these cycles

corresponded to behavior. For correlations between submovement speed profiles, movement times, and

average spike times, non-parametric tests were used. Since CIφ values represent an angle ranging from -π

to π, circular distribution statistics—mean, variance, correlation, and Rayleigh test for non-uniformity—

were used. All circular statistics were calculated with CircStat, a Circular Statistics Toolbox for Matlab

(Berens, 2009).

Table 1. Statistical tests and confidence intervals reported throughout this study referenced with letter

superscripts.

	Data Structure	Type of test	Confidence intervals
a	Correlation between speed profiles from --200 to 200 ms relative to peak speed, nonparametric	Wilcoxon rank sum test	[25th, 75th] percentiles
b	Initial vs. Corrective movement times, nonparametric	Two-sided Wilcoxon rank sum test	Percentage of submovements within 100-350 ms
c	Initial vs. Corrective Average spike times between -200 to 100ms, nonparametric	Spearman's rank correlation	95% confidence interval by bootstrapping (1000 repetitions)
d	Circular distribution of phase	Rayleigh test	Circular standard deviation
e	CIφ-angle Speed – linear random variable	Circular correlation between angle and linear variable	Minimum and maximum across 12 recording sessions
f	Ratio of Standard deviations of times estimated with CIφ and Firing rate model	F-test	95% confidence interval

Results

Motor behavior – initial and corrective submovements

Movement speed was analyzed throughout the center-out task from instruction until successful

completion of the final target hold.   The two monkeys successfully completed 10,963 (monkey P) and

8,737 (monkey Q) trials across 12 recording sessions each. In addition to the peaks in speed with the

initial reach after instruction, additional peaks in speed were observed and labeled as corrective

submovements. There were 6478 and 3912 corrective submovements identified for monkeys P and Q,

respectively.   Across all trials, 68.3% (P) and 71.1% (Q) were completed in a single initial movement,

17.5% (P) and 20.3% (Q) of trials were completed with one additional corrective submovement, and

14.2% (P) and 8.6% (Q) of trials required two or more corrective submovements. The location of the

identified speed peaks within example trials and the speed profiles for monkey P are shown in Figure 2A

and 2B, respectively.   The speed peaks tended to be distinct with nearly zero velocity between most

peaks. As shown in Figure 2C, 99.0% (P) and 97.7% (Q) of the minimum speed trough following the

initial speed peak were less than 20% of the peak. Similarly, 82.4% (P) and 85.8% (Q) of the troughs

were less than 20% of the preceding peak between sequential corrective speed peaks. The mean peak

speeds for initial submovements were 1533 (P) and 1182 (Q) pixels/s while corrective submovement peak

speeds were 460 (P) and 400 (Q) pixels/s. Thus, the average peaks for corrective submovements were

30.0% and 33.8% of initial submovements, and a low-speed trough almost always occurred between two

speed peaks making it reasonable to analyze submovements defined by their peak speeds.

Figure 2. The precision center-out task. A) Cursor paths for four example trials to each target for the three target sizes: regular (top), narrow (middle), shallow (bottom). Initial submovements from 200ms before to 200ms after speed peaks are plotted in blue with the point when peak speed occurred shown with a blue dot. Corrective movements are similarly identified in red with a red dot. Grey lines connect the rest of a trial before, between, or after submovements with a speed peak. B) Cursor speed plotted versus time for a subset of trials. Initial (blue) and corrective (red) submovement speed peaks are identified with squares. Gray squares identify speed peaks that were thrown out because they i) were small initial movements that did not move outside the center or ii) occurred entirely within the peripheral target. C) Top) Distribution of peak speeds for initial (blue) and corrective (red) submovements. Bottom)

Distribution of the trough-to-peak ratio for the troughs following an initial submovement before a

corrective submovements and following a corrective submovement before another corrective

submovement. Data is shown for monkey P. Data for monkey Q, which had similar results, is not shown.

The speed profiles were time aligned to peak speed to better examine the identified

submovements (Figure 3A). Almost all submovements show a clear bell-shaped profile for both the

initial and corrective movements. The similarity between initial and corrective speed profiles was

assessed by using the correlation between randomly selected pairs of movements. For random pairs of

one initial and one corrective submovement, the median correlation was 0.78 [0.58 0.89] (monkey P) and

0.83 [0.70, 0.90] (monkey Q). As a ceiling comparison, the correlation between randomly selected pairs

of initial submovements was observed to be 0.93 [0.86 0.96] (P) and 0.91 [0.80, 0.96] (Q). Thus, initial-

corrective pairs were significantly correlated (p<0.001[a]) and although this correlation was significantly

less than the initial-initial pairs, corrective submovements still had a high similarity of 0.78/0.93=84%

and 0.82/0.91=91% to the speed profile shape observed in initial submovements.

The time duration and timing of submovements was also examined. The onset and offset of

submovements were defined as the time points when speed was one-half of the maximum speed both

before and after the speed peak. As shown in Figure 3B, the movement duration at half maximum speed

was similar and close to symmetric for both initial and corrective submovements. The initial

submovements were slightly longer having a median time of 220ms (P) and 270ms (Q) compared to

corrective submovements with medians of 180ms (P) and 220ms (Q). This difference in median

movement times was statistically significant (p<0.001[b]) but the difference of 40 and 50ms was small,

especially given the peak speed was only one-third the magnitude for the smaller corrective movements.

Overall, all submovement durations, as measured by the full width at half maximum, occurred within a

similar range with 96.7%/88.0% (P/Q) of all initial and 96.3%/93.0% (P/Q) corrective submovements

between 100-350 ms. The time between speed peaks—either initial to first corrective submovements or

between subsequent corrective submovements—is plotted in Figure 3C. The median time between peaks

were 570 ms for monkey P and 700 ms for monkey Q with the mode time between peaks being 450ms (P)

and 550ms (Q). Only 3.2% (P) and 0.1% (Q) of speed peaks had a time between peaks less than 200ms

and 6.1% (P) and 10.0% (Q) of speed peak pairs had times greater than 1200 ms. These observations

suggest the movement behavior could be divided into submovements with similar bell-shaped velocity

profiles and similar time durations.

Figure 3. Time course of submovements. A) The cursor speeds are plotted aligned to speed peaks for

initial (blue) and corrective (red) submovements. N.B. The cursor speeds shown are before the bandpass

filter used for identifying peaks displayed in Figure 1B. Thus, the maximum of each trace may not align

exactly with the plotted peak speed. B) Histogram of the time at half-maximum speed before and after

peak speed for all initial (blue) and corrective (red) submovements. C) The time duration between speed

peaks including the times from initial submovement to first corrective submovement as well as between

any consecutive pairs of corrective submovements.

Consistent Timing of Neural Firing Rates for Initial and Corrective Submovements

315   Single target acquisition movements thus often consisted of initial and corrective submovements

with similar temporal characteristics. Did neural activity in the primary motor cortex control such target-

acquisition movements as a single movement, or as a series of discrete submovements? The neural firing

rates across the recorded population were time aligned to the submovement speed peaks to examine the

firing rates from 500 ms before until 300 ms after the peak speed. The average firing rate (smoothed with

a Gaussian window, σ=30ms) for all analyzed units aligned to the peak speed for initial and corrective

submovements are shown in Figure 4A. A clear peak occurs before the peak speed for both initial and

corrective submovements in both monkeys. Monkey P's peak firing rates occurred 170 ms and 120 ms

before initial and corrective submovements, respectively, while monkey Q's occurred at 160 and 160 ms

before for both initial and corrective submovements. Thus, firing rates increased and peaked globally for

corrective submovements in addition to the initial reach.

326   If all neurons had the same time lag preceding the upcoming peak in movement speed, there

would be a synchronized increase and decrease of all condition-independent firing rates simultaneously.

However, when examining average firing rates from 10 example neurons from one recording session

from monkey P, all aligned to peak speed, we see heterogenous timing of firing rates relative to the peak

speed (Figure 4B). This relationship tended to be conserved across initial and corrective movements,

with the purple spiking units tending to fire earlier and the orange units later for both initial and corrective

submovements. This suggests that the condition-independent neural activity across the neurons might

form a repeatable temporal structure—a neural trajectory—that is more than a simple simultaneous rise

and fall in firing rate across the population

To quantify the early versus late consistency of spiking units, we calculated the average time of

all spikes that occurred within a window from -200ms before to 100ms after peak speed to determine

whether a unit tended to increase its firing rate earlier (negative time) or later (positive time) relative to

peak speed. We then compared these average spike times for initial versus corrective submovements for

each spiking unit. As shown in Figure 4C, earlier firing units (more negative) for initial submovements

tended to fire earlier for corrective submovements, while units later (more positive) for initial

submovements also tended to fire later for corrective submovements. This correlation was significant for

all spiking units with Spearman correlations of $\rho = 0.40$ [0.35, 0.45] (P) and $\rho = 0.58$ [0.53, 0.62] (Q),

p<0.001[c]. Using only single units, the Spearman correlations were $\rho = 0.37$ [0.31, 0.44] (P) and $\rho = 0.61$

[0.54, 0.68] (Q), p<0.001[c]. Thus, a significant portion of the ordered timing of units was conserved

relative to peaks in movement speed for both initial and corrective submovements.

Figure 4. Neural firing relative to initial and corrective submovements. A) The firing rate for all spiking

units was averaged for all initial (blue) and corrective (red) submovements. The shaded region interval

shows the 95% confidence interval of the calculated mean for all spiking units. Circles indicate the time

of peak firing rate for each condition. B) Average condition-independent firing rates for 10 example

spiking units recorded simultaneously from monkey P time-aligned relative to peak speed for all initial

(left) and corrective (right) submovements. Firing rates are shown relative to the average firing rate

within the given time window (initial or corrective) for each spiking unit. The weighted timing of spikes

(in ms) within the -200ms to 100ms window is given for each unit. Units are colored based on the initial

movement by whether their firing rates were greater early (purple) or late (orange). C) Weighted timing

of spiking relative to peak speed for each unit for initial (abscissa) and corrective (ordinate)

submovements. More negative times represent spiking earlier relative to the peak speed of each

submovement. Single units are shown with filled circles while all other spiking multi-units are shown

with open circles.

Consistent Neural Dynamics for Initial and Corrective Submovements

We next wanted to examine whether these repeatable neural patterns that occurred on average

across all movements could be used to identify submovements on individual trials. Despite the smaller

magnitude of the condition-independent neural activity during corrective movements, the repeated

oscillations in speed and repeated neural dynamics suggested a portion of neural activity was repeatable

and common to initial and corrective submovements. To examine this, we built a simple linear dynamical

system model using the neural firing rates from the entire trial—including both initial and corrective

submovements—to characterize common temporal dynamics that might be present. The neural firing

rates were again averaged across all conditions, i.e. movement directions, and both initial and corrective

portions of the trials so the dynamical system model would identify common condition-independent

activity. Using the jPCA algorithm described in Churchland et al. (2012), i) the first six principal

components of the neural space and ii) the two dimension plane within the space of those six principal

components that captured the most rotational neural activity were identified. We labeled the two neural

dimensions of the plane with the most rotational condition-independent activity as CIx and CIy. To

consistently define CIx and CIy across recording sessions and monkeys, we aligned the +CIx direction

with the neural dimension that had the maximum average firing rate in the plane. This was performed by

calculating the average firing rate across all spiking units for neural activity based on each timepoint's

angle in the CIx/CIy plane (binned in 100 angle intervals) and rotating the CIx and CIy axes so that +CIx

aligned with the angle with largest firing rate. This alignment results in the +CIx dimension closely

aligning with the time course of the global average firing rate across the population (shown in Fig 4A)

while CIy is an orthogonal neural dimension that oscillates with a phase lag of $\pi/2$ compared to CIx.

The average firing rates projected in our identified CI plane for all initial and corrective

submovements are shown in Figure 5, where the neural data was again aligned relative to peak speed for

initial and corrective submovements separately. The neural trajectory in the 2-dimensional CIx/CIy plane

are shown in Figure 5A, while the same CIx and CIy dimensions are plotted as a function of time in

Figure 5B. The initial and corrective neural trajectories (Fig. 5A) are very similar in their shape and

direction of rotation within the plane, with the trajectories for corrective submovements appearing as an

additional cycle resembling a smaller, scaled version of the larger trajectories for initial submovements.

The time courses of CIx (solid) and CIy (dashed) (Fig. 5B) were similar for initial (blue) and corrective

(red) submovements, though they differed in magnitude. The peak in the CIx dimension (denoted with an

X)—defined as the dimension in the plane that best correlated with the global average firing rate of the

population—occurred approximately 150 ms before peak speed for initial and corrective submovements,

whereas the peak in the CIy dimension (also denoted with an X) occurred near the time of peak speed for

both submovement types.

Figure 5. Cyclic neural dynamics related to initial and corrective submovements. A) The average

population firing rates for initial (blue) and corrective (red) submovements are projected in the CIx/CIy

plane identified with jPCA. The trajectories start at the triangles and end at the squares. Each filled circle

is a 150 ms time step and the open corresponds to peak speed. C). Average CIx (solid lines) and CIy

(dashed lines) plotted as a function of time relative to average cursor speed (dotted lines).

Neural cycles improve predictions of behavioral timing

Since the population firing rates in the CI plane appeared to cycle with similar timing for initial

and corrective submovements, despite different magnitudes, we next chose to examine the instantaneous

phases of CIx and CIy activity to see if it was a statistically significant marker of the neural state of motor

cortex and its relationship with upcoming movement. We used a Hilbert transform to create an analytic

representation of the CIx and CIy signals and then calculated the instantaneous phase by taking the angle

between the real component and the Hilbert transformed imaginary component. The average phase of CIx

and CIy for both initial and corrective submovements—time aligned to peak speed—is shown in Figure

6A. The phase of CIx (solid lines) and that of CIy (dashed lines) each were similar for initial and

corrective submovements, with the zero phase of CIx occurring about 150 ms before the peak speed while

CIy lagged CIx with an approximately $\pi/2$ phase lag, with the zero crossing occurring around peak speed.

The slope of the phase for corrective movements was slightly steeper indicating that neural activity cycled

slightly faster for corrective movements than initial. Histograms of the phase of CIx and of CIy at peak

speed on individual trials are shown in Figure 6B. The distributions of phases of CIx and CIy were

significantly non-uniform for both monkeys and the means and standard deviations are given in Table 2.

Thus, there was a clear relationship between peak speed and the phase of condition-independent activity

that occurred with almost all submovements, both initial and corrective, and had similar timing.

Figure 6. Phase of CIx and CIy relative to peak cursor speed. A) Phase of CIx (solid lines) and

CIy (dashed lines) time-aligned to peak speed (Time = 0) and averaged for all initial (blue) and corrective

(red) submovements. B) Histograms of the phase of CIx and Ciy at the time of peak speed for initial

(blue) and corrective (red) submovements.   Means and standard deviations are given in Table 2.

Table 2. Means and standard deviations of the phase of CIx and CIy. All circular distributions of the

phase of CIx and CIy were non-uniform (all $p < 0.001$[d]).

Monkey P	CIx Mean	CIx Std. dev	CIy Mean	CIy Std. dev
Initial	0.35 π	0.26 π	-0.14 π	0.23 π
Corrective	0.43 π	0.33 π	-0.15 π	0.36 π
Monkey Q				
Initial	0.31 π	0.22 π	-0.15 π	0.30 π
Corrective	0.45 π	0.31 π	-0.05 π	0.37 π

Because the phase in the CI plane appeared to define the neural dynamics and predict upcoming

speed peaks, we created a metric we call the condition-independent phase (CIφ) by averaging the phase of

CIx and phase of CIy + π/2 to calculate the current phase in the CI plane. We then examined the

continuous relationship between cursor speed and neural CIφ. In figure 7A, we have plotted the cursor

speed as a function of CIφ. While the CIφ is an angle that ranges between +/-π radians when calculated,

for purposes of display here we have incremented CIφ in steps of 2π to show how successive cycles of

neural activity (abscissa) were related to movement speed (ordinate) as individual trials progressed

through both initial and subsequent corrective submovements. The individual trials for monkey P in

Figure 7A are the same as the trials shown in Figure 2B. However, the speed traces have now been

stretched or condensed in time based on the current brain state measured with the CIφ. This plot now

shows that the speed of movement varied with the cyclic neural activity with the cursor speeds for most

trials rising and falling in 2π cycles of CIφ. Both the speed averaged across all trials (white) and the non-

uniform occurrence of peak speeds in individual trials (black) demonstrate that movement speed was

consistently correlated with the cycles of condition-independent neural activity. The statistically

significant circular correlation between speed and CIφ was 0.44 [0.39,0.53] and 0.42 [0.35,0.50]

(p<0.001^e for both animals) with the largest speeds occurring at CIφ = 0.32π and 0.31π (+2kπ) for

monkeys P and Q, respectively.

Finally, we examined the predictive power of CIφ for estimating when the peak speed occurred.

Figure 7B illustrates the distribution of the time at which CIφ =0 relative to the time of peak speed for

initial submovements (top) and corrective submovements (bottom). These distributions consistently

peaked 100-150 ms before the speed peak for both initial and corrective submovements. Corrective

movements had CIφ =0 at times slightly closer to peak speed indicating that the time delay to peak speed

was slightly less for corrective movements. A relatively consistent relationship between neural activity in

the CIx/CIy plane and peak speed was present for both initial and corrective submovements across all

trials regardless of target size or reach direction.

To examine if incorporating neural dynamics significantly improved prediction, we compared our

CIφ predictions with these population dynamics to predictions using a standard approach of using the

instantaneous firing rate of all units to predict peak speeds. For predictions with the instantaneous firing

rates, we built a linear regression model to estimate peak speed with a weighted sum of the instantaneous

firing rate of all spiking units (see Methods). Using this model, we obtained when the peak in firing rate

occurred to predict the upcoming peak speed.   Figure 7C shows the temporal distributions of these peak

firing rates relative to peak speed for both initial and corrective submovements. Like the distributions

using the dynamical model above (Figure 7B), the firing rate model peaked 150 to 100 ms before peak

speed. The peaks were broader by 10-20 ms, however, as characterized by the greater standard deviations

(σ) given for each distribution. The standard deviations were significantly different in all cases—initial

and corrective for both monkeys (Table 2). Furthermore, although ≥ 84% of submovements were

included in each of these distributions (percentages given in Fig. 7), a small fraction of submovements

could not be aligned, lacking a CIφ=0 in the dynamical systems model and/or a peak in the firing rate

model within the -300 to 100 ms time window examined. The percentage of these unaligned trials was

consistently smaller for the dynamical systems model. Compared to using only the

instantaneous/synchronous firing rates, using the cyclic/asynchronous dynamics of the neural population

thus significantly improved the accuracy and consistency with which the time of peak speed could be

predicted.

Figure 7. Relationship between CIφ and cursor speed. A) Cursor speed is plotted as a function of CIφ for

200 trials with at least one corrective submovement. The average speed of all trials as a function of CIφ

is shown in white, illustrating the oscillation in cursor speed depending on the phase of neural activity.

The circular correlations between CIφ and cursor speed for all corrective trials were 0.44 [0.39, 0.53] and

0.43 [0.36, 0.50] for monkeys P and Q, respectively, p<0.001[e] in both cases. Note, the unwrapped CIφ is

not always a monotonically increasing value as occasionally the neural activity could reverse and move

clockwise rather than counter-clockwise in the neural plane shown in Figure 6B.

B & C) Identifying the times of peak speeds with a dynamical systems model (B) or with an

instantaneous firing rates (C). The time point when CIφ = 0 (B) or peak firing rate (C) was used as a

prediction of the upcoming submovement. Each histogram shows only those submovements for which

the neural data aligned with the movement data, i.e. CIφ = 0 (B) or maximum firing rate (C) occurred

within the time range examined (-300 to 100 ms relative to submovement peak speed). The percentage of

total aligned trials is shown for each distribution as well as the standard deviation (σ) for the aligned

trials. In all cases, the dynamical systems model predictions were more precise, with a narrower standard

deviation (statistics in Table 3) and fewer unaligned trials.

Table 3. Comparison of predication accuracy as measured with standard deviation in predictions using
the dynamical system CIφ model vs. an instantaneous firing rate model.

	σ_1, CIφ, (ms)	σ_2, Firing Rate (ms)	F-stat, $\dfrac{\sigma_1^2}{\sigma_2^2}$	95% Confid. interval	Data Comparison	Statistical Test
Initial					All submovements with a prediction between -300:100 ms, assuming normal distribution	F-test[f], all p<0.001
Monkey P	66.1	84.4	0.61	[0.59, 0.64]		
Monkey Q	75.1	91.1	0.69	[0.65, 0.71]		
Corrective						
Monkey P	88.0	104.2	0.71	[0.68, 0.75]		
Monkey Q	87.0	98.7	2.06	[0.73, 0.83]		

Discussion

500   Our precision center-out task utilized small targets to elicit one or more corrective submovements

in many trials. We found a temporal relationship for both initial and corrective reaching movements with

cyclic, condition-independent neural activity. Rather than a single cycle of neural activity in the primary

motor cortex occurring during each trial, the speed profiles of initial and corrective submovements each

aligned with a cycle of neural activity, providing a useful neural marker encoding the series of

submovements.

In our precision center-out task, the monkeys' movements showed consistent bell-shaped speed

profiles. These speed profiles were evident for both the larger initial movement from the center toward

the peripheral target as well as for each subsequent corrective movement.   A large majority of both initial

and corrective submovements had durations of 100-350 ms, with a low-speed trough separating almost all

submovements. Discrete submovements defined by multiple speed peaks have previously been described

in behavioral studies of reaching (Pratt et al., 1994; Lee et al., 1997; Hatsopoulos et al., 2007; Polyakov et

al., 2009), turning a knob (Novak et al., 2000), isometric contractions (Massey et al., 1992; Hall et al.,

2014), and object manipulation for tactile discrimination (Pruszynski et al., 2018). The experimental

results and analysis presented here provide new evidence of a relationship between previously reported

cyclic neural dynamics (Churchland et al., 2012; Hall et al., 2014) and such behaviorally observed

submovements.

Condition-independent phase predictive of cursor speed

Churchland et al. (2012) originally described a single cycle of rotational dynamics in the activity

of neurons in the primary motor and premotor cortex during both straight reaches and curved reaches

around obstacles. While originally presented as similar rotations but with different starting points and

radial magnitudes for varying conditions, others have pointed out how neurons with various but consistent

temporal lags (Lebedev et al., 2019) may explain many of the results. More recently, Zimnik and

Churchland (2021) demonstrated two repeated cycles of neural activity, each shortened in time, when a

pair of movements were simultaneously instructed to be performed in rapid succession. Here, by

focusing on condition-independent neural activity, we found that similar cycles of neural activity appear

not only for initially planned reaches but also for the highly variable, corrective submovements that are

made online with visual feedback. Our results highlight that the various time lags between individual

cortical neurons' firing and the upcoming reaching movements are conserved, whether large and

instructed or small and made online with feedback.

Similar but smaller cyclic, condition-independent activity for corrective movements

Although the orientation and direction of rotation for the cyclic, condition-independent neural

modulation was similar for initial and corrective submovements, the magnitude of the condition-

independent neural activity that occurred for corrective submovements was approximately one-third to

one-half the magnitude of that for the initial submovements (both in average firing rate, Fig. 4A, and

within our identified rotational CI plane, Fig. 5). On average, the encoding of movement speed is clearly

present in primary motor cortex (Moran and Schwartz, 1999; Paninski et al., 2004), and the smaller

change in average firing rate observed here during corrective movements reflected the lower movement

speed for the corrective compared to the initial submovements, suggesting speed tuning in the magnitude

of the condition-independent activity. This does not imply, however, that each individual trial and each

individual neuron have proportionally smaller changes of firing rate during smaller amplitude

movements. Examination of small, instructed movements has shown that a fraction of primary motor

cortex neurons have similar firing rates for small, precise and for larger wrist movements while others are

selective for only larger movements (Fromm and Evarts, 1981). We too observed similar large changes in

firing rate on individual corrective submovements for certain neurons (data not shown). Only when

averaging firing rates—time aligned to the peak movement speed or the decoded condition-independent

phase—were the population differences in firing rate modulation between initial and corrective

movements readily apparent.   Precisely identifying encoded speed on a trial-by-trial basis with the neural

activity remains challenging as there are often large changes of firing rates for individual neurons that are

variable and idiosyncratic during any particular corrective submovement.

Adding condition-independent activity to condition-dependent activity has been suggested to

make brain dynamics more robust to noise by increasing the differences in neural signals even when the

muscle activation pattern at certain time points are very similar (Russo et al., 2018). In the context of

precise, corrective movements, we speculate cyclic brain dynamics can be used to organize neural activity

that creates distinct submovements with time-varying neural and musculoskeletal dynamics that are more

reliable for motor control. Previous reports of neural activity defining submovements linked together

have used the term movement fragments (Hatsopoulos et al., 2007). In the context of precise movements,

we hypothesize that organizing movement into submovements or movement fragments might allow the

control of particular submovements to have different encoding features, neural processing, or control

policies, for instance, allowing the large initial movements to be larger amplitude and less precise while

the corrective submovements are smaller and more precise. Further studies will be needed to understand

the condition-dependent differences that accompany the condition-independent neural features presented

here.

Though various time lags in different neurons seem likely to be present across many tasks, cyclic,

condition-independent neural dynamics may not be similar for all upper extremity movements. For

instance, whereas during combined reach-and-grasp movements cyclic condition-independent activity

occurs along with more complex condition-dependent dynamics (Rouse and Schieber, 2018), during

separate reaching movements and grasping movements condition-dependent activity was cyclic during

reaching, but was more complex during grasping (Suresh et al., 2020). The neural signals in a given

hemisphere for cyclic movements of the contra- and ipsilateral arms have also have been reported to be in

orthogonal subspaces (Ames and Churchland, 2019). Cyclic neural activity may not be due only to

intrinsic neural dynamics in M1, but also the result of sensorimotor feedback control and/or a cognitive

strategy. With sufficient time delay between each submovement, the neural activity could fit both

descriptions. Observations of additional submovements defined by second or third speed peaks do not

necessarily require a feedback controller with discrete updates. A single, continuous optimal feedback

controller with appropriate delays and signal dependent noise can generate additional submovements with

multiple, sequential speed peaks (Li et al., 2018). Results by Susilaradeya et al. (2019) argue that

extrinsic effects of a task interact with the intrinsic dynamics of the brain in a manner consistent with an

optimal feedback controller, possibly providing a framework for assessing these effects across a variety of

tasks including our precision center-out task. Further work examining neural activity in various tasks

and/or additional sensorimotor brain areas will be needed to advance our understanding of the neural

dynamics of the sensory processing, cognitive planning, and motor execution for precise, corrective

movements.

The cyclic dynamics of corrective movements have important implications for brain-computer

interfaces (BCIs). To date, most BCI decoders are time-invariant, not recognizing when submovements

occur. Decoders are typically first constructed from observed or imagined movements that assume single,

straight-line movements.   When algorithms for updating BCI decoders consider the change in movement

direction for corrective movements, it typically has been assumed the intended path is updated

continuously (Gilja et al., 2012; Shanechi et al., 2016). Experiments have suggested that BCI control can

be improved with two states: active control and rest (Kim et al., 2011; Williams et al., 2013, 2016; Sachs

et al., 2016). Our results suggest that computing the phase of cyclic, condition-independent neural

activity with CIφ (Fig. 7B) can provide better prediction of the timing of corrective submovements than

using the instantaneous firing rates alone (Fig. 7C). This may lead to BCIs that allow the subject to better

signal when they intend to make a corrective movement. With additional information about the typical

neural dynamics and kinematics of submovements, BCI decoders may better estimate natural kinematics

from noisy neural signals. Taking into account the cyclic dynamics of the condition-independent neural

activity may also lead to better descriptions of the condition-dependent activity that encodes task features.

For example, direction encoding has been shown to shift progressively during a single movement (Sergio

and Kalaska, 1998; Churchland and Shenoy, 2007; Suminski et al., 2015; Suway et al., 2017). Accounting

for the phase of a movement with its cyclic, condition-independent activity (i.e. CIφ) could enable

decoders of movement direction that shift progressive during a single movement. Such improvements

could lead to a more robust description of the neural encoding of precise and corrective movements.

Extended Data 1. Matlab code to calculate the CIφ is available on Github.

Since the trial data contains corrective movements in addition to the large initial movements that

were not precisely time aligned to trial events for averaging condition-independent neural activity, we

developed a novel algorithm to iteratively average the firing rates, calculate CIφ, then average the firing

rates again based on the CIφ. This iterative process involves three steps: i) Each unit's firing rate is

averaged across all trials to determine its condition-independent firing rate. ii) Dimensionality reduction

is performed using PCA and jPCA on the condition-independent firing rates to identify the neural plane

with the most rotational/cyclic condition-independent activity. iii) The instantaneous phase is calculated

using the Hilbert transform on the first two jPC dimensions for all data points. Matlab code is available at

https://github.com/arouseKUMC/CIphase. Further details are available in the Readme document attached

to the code.

References

Abrams RA, Meyer DE, Kornblum S (1990) Eye-hand coordination: oculomotor control in rapid aimed limb movements. J Exp Psychol Hum Percept Perform 16:248–267.

Ames KC, Churchland MM (2019) Motor cortex signals for each arm are mixed across hemispheres and neurons yet partitioned within the population response. Elife 8:1–36.

Ashe J, Georgopoulos AP (1994) Movement parameters and neural activity in motor cortex and area 5. Cereb Cortex 6:590–600.

Berens P (2009) CircStat : A MATLAB Toolbox for Circular Statistics. J Stat Softw 31:1–21.

Cheney PD, Fetz EE (1980) Functional classes of primate corticomotoneuronal cells and their relation to active force. J Neurophysiol 44:773–791.

Churchland MM, Cunningham JP, Kaufman MT, Foster JD, Nuyujukian P, Ryu S, Shenoy K V (2012) Neural population dynamics during reaching. Nature 487:51–56.

Churchland MM, Shenoy K V (2007) Temporal complexity and heterogeneity of single-neuron activity in premotor and motor cortex. J Neurophysiol 97:4235–4257.

Craik KJW (1947) Theory of the human operator in control systems. I. the operator as an engineering system. Br J Psychol 38:56–61.

Cunningham JP, Yu BM (2014) Dimensionality reduction for large-scale neural recordings. Nat Neurosci 17:1500–1509.

Elliott D, Hansen S, Grierson LE, Lyons J, Bennett SJ, Hayes SJ (2010) Goal-Directed Aiming: Two Components but Multiple Processes. Psychol Bull 136:1023.

Evarts E V (1968) Relation of pyramidal tract activity to force exerted during voluntary movement. J Neurophysiol 31:14–27.

Fromm C, Evarts E V (1981) Relation of size and activity of motor cortex pyramidal tract neurons during

skilled movements in the monkey. J Neurosci 1:453–460.

Georgopoulos AP, Kalaska JF, Caminiti R, Massey JT (1982) On the relations between the direction of

two-dimensional arm movements and cell discharge in primate motor cortex. J Neurosci 2:1527–

1537.

Gilja V, Nuyujukian P, Chestek CA, Cunningham JP, Yu BM, Fan JM, Ryu S, Shenoy K V (2012) A

brain machine interface control algorithm designed from a feedback control perspective. Conf Proc

IEEE Eng Med Biol Soc 2012:1318–1322.

Hall TM, de Carvalho F, Jackson A (2014) A common structure underlies low-frequency cortical

dynamics in movement, sleep, and sedation. Neuron 83:1185–1199.

Hatsopoulos NG, Xu Q, Amit Y (2007) Encoding of movement fragments in the motor cortex. J Neurosci

27:5105–5114.

Hill DN, Mehta SB, Kleinfeld D (2011) Quality metrics to accompany spike sorting of extracellular

signals. J Neurosci 31:8699–8705.

Jackson A, Gee VJ, Baker SN, Lemon RN (2003) Synchrony between Neurons with Similar Muscle

Fields in Monkey Motor Cortex. Neuron 38:115–125.

Kakei S, Hoffman DS, Strick PL (1999) Muscle and movement representations in the primary motor

cortex. Science 285:2136–2139.

Kalaska JF, Cohen DAD, Hyde ML, Prud'homme M (1989) A comparison of movement direction-related

versus load direction-related activity in primate motor cortex, using a two-dimensional reaching

task. J Neurosci 9:2080–2102.

Kaufman MT, Seely JS, Sussillo D, Ryu SI, Shenoy K V, Churchland MM (2016) The largest response

component in motor cortex reflects movement timing but not movement type. eNeuro

661 3:ENEURO.0085-16.2016.

662 Kihlberg JK, Herson JH, Schotz WE (1972) Square Root Transformation Revisited. J R Stat Soc 21:76–

663 81.

664 Kim S-P, Simeral JD, Hochberg LR, Donoghue JP, Friehs GM, Black MJ (2011) Point-and-click cursor

665 control with an intracortical neural interface system by humans with tetraplegia. IEEE Trans neural

666 Syst Rehabil Eng 19:193–203.

667 Kobak D, Brendel W, Constantinidis C, Feierstein CE, Kepecs A, Mainen ZF, Qi XL, Romo R, Uchida

668 N, Machens CK (2016) Demixed principal component analysis of neural population data. Elife 5:1–

669 36.

670 Lebedev MA, Ossadtchi A, Mill NA, Urpí NA, Cervera MR, Nicolelis MAL (2019) Analysis of neuronal

671 ensemble activity reveals the pitfalls and shortcomings of rotation dynamics. Sci Rep 9:1–14.

672 Lee D, Port NL, Georgopoulos AP (1997) Manual interception of moving targets. Exp brain Res

673 116:421–433.

674 Li Z, Mazzoni P, Song S, Qian N (2018) A Single, Continuously Applied Control Policy for Modeling

675 Reaching Movements with and without Perturbation. Neural Comput 30:397–427.

676 Liu Z, Schieber MH (2020) Neuronal activity distributed in multiple cortical areas during voluntary

677 control of the native arm or a brain-computer interface. eNeuro 7:1–16.

678 Massey JT, Lurito JT, Pellizzer G, Georgopoulos AP (1992) Three-dimensional drawings in isometric

679 conditions: relation between geometry and kinematics. Exp Brain Res 88:685–690.

680 Mathworks (2020) Natick, Massachusetts: The MathWorks Inc.

681 Maynard EM, Hatsopoulos NG, Ojakangas CL, Acuna BD, Sanes JN, Normann RA, Donoghue JP (1999)

682 Neuronal interactions improve cortical population coding of movement direction. J Neurosci

683  19:8083–8093.

684 Mollazadeh M, Aggarwal V, Davidson AG, Law AJ, Thakor N V, Schieber MH (2011) Spatiotemporal

685  variation of multiple neurophysiological signals in the primary motor cortex during dexterous reach-

686  to-grasp movements. J Neurosci 31:15531–15543.

687 Moran DW, Schwartz AB (1999) Motor Cortical Representation of Speed and Direction During

688  Reaching. J Neurophysiol 82:2676–2692.

689 Novak KE, Miller LE, Houk JC (2000) Kinematic properties of rapid hand movements in a knob turning

690  task. Exp Brain Res 132:419–433.

691 Paninski L, Fellows MR, Hatsopoulos NG, Donoghue JP (2004) Spatiotemporal tuning of motor cortical

692  neurons for hand position and velocity. J Neurophysiol 91:515–532.

693 Polyakov F, Stark E, Drori R, Abeles M, Flash T (2009) Parabolic movement primitives and cortical

694  states: Merging optimality with geometric invariance. Biol Cybern 100:159–184.

695 Pratt J, Chasteen AL, Abrams RA (1994) Rapid aimed limb movements: age differences and practice

696  effects in component submovements. Psychol Aging 9:325–334.

697 Pruszynski JA, Flanagan JR, Johansson RS (2018) Fast and accurate edge orientation processing during

698  object manipulation. Elife 7:e31200.

699 Rouse AG, Schieber MH (2016) Spatiotemporal Distribution of Location and Object Effects in Primary

700  Motor Cortex Neurons during Reach-to-Grasp. J Neurosci 36:10640–10653.

701 Rouse AG, Schieber MH (2018) Condition-dependent neural dimensions progressively shift during reach

702  to grasp. Cell Rep 25:3158–3168.

703 Russo AA, Bittner SR, Perkins SM, Seely JS, London BM, Lara AH, Miri A, Marshall NJ, Kohn A,

704  Jessell TM, Abbott LF, Cunningham JP, Churchland MM (2018) Motor Cortex Embeds Muscle-like

Commands in an Untangled Population Response. Neuron 97:953-966.e8.

Sachs NA, Ruiz-Torres R, Perreault EJ, Miller LE (2016) Brain-state classification and a dual-state

decoder dramatically improve the control of cursor movement through a brain-machine interface. J

Neural Eng 13:016009.

Sainburg RL, Ghez C, Kalakanis D (1999) Intersegmental dynamics are controlled by sequential

anticipatory, error correction, and postural mechanisms. J Neurophysiol 81:1045–1056.

Sarma S V, Eden UT, Cheng ML, Williams ZM, Hu R, Eskandar E, Brown EN (2010) Using point

process models to compare neural spiking activity in the subthalamic nucleus of parkinsons patients

and a healthy primate. IEEE Trans Biomed Eng 57:1297–1305.

Sergio LE, Kalaska JF (1998) Changes in the Temporal Pattern of Primary Motor Cortex Activity in a

Directional Isometric Force Versus Limb Movement Task. J Neurophysiol 80:1577–1583.

Shanechi MM, Orsborn AL, Carmena JM (2016) Robust Brain-Machine Interface Design Using Optimal

Feedback Control Modeling and Adaptive Point Process Filtering. PLoS Comput Biol 12:1–29.

Snedecor G, Cochran W (1980) Statistical Methods, 7th ed. Ames, IA: Iowa State UP.

Suminski AJ, Mardoum P, Lillicrap TP, Hatsopoulos NG (2015) Temporal evolution of both premotor

and motor cortical tuning properties reflect changes in limb biomechanics. J Neurophysiol

113:2812–2823.

Suresh AK, Goodman JM, Okorokova E V., Kaufman MT, Hatsopoulos NG, Bensmaia SJ (2020) Neural

population dynamics in motor cortex are different for reach and grasp. Elife 9:1–16.

Susilaradeya D, Xu W, Hall TM, Galán F, Alter K, Jackson A (2019) Extrinsic and intrinsic dynamics in

movement intermittency. Elife 8:1–27.

Suway SB, Orellana J, McMorland AJC, Fraser GW, Liu Z, Velliste M, Chase SM, Kass RE, Schwartz

AB (2017) Temporally Segmented Directionality in the Motor Cortex. Cereb Cortex 12:1–14.

Thach WT (1978) Correlation of Neural Discharge with Pattern and Force of Muscular Activity, Joint

Position, and Direction of Intended Next Movement in Motor Cortex and Cerebellum. J

Neurophysiol 41:654–676.

Truccolo W, Eden UT, Fellows MR, Donoghue JP, Brown EN (2005) A Point Process Framework for

Relating Neural Spiking Activity to Spiking History, Neural Ensemble, and Extrinsic Covariate

Effects. J Neurophysiol 96:1074–1089.

Williams JJ, Rouse AG, Thongpang S, Williams JC, Moran DW (2013) Differentiating closed-loop

cortical intention from rest: building an asynchronous electrocorticographic BCI. J Neural Eng

10:046001.

Williams JJ, Tien RN, Inoue Y, Schwartz AB (2016) Idle state classification using spiking activity and

local field potentials in a brain computer interface. 2016 38th Annu Int Conf IEEE Eng Med Biol

Soc:1572–1575.

Woodworth RS (1899) The accuracy of voluntary movement. Psychol Rev 3:1–119.

Yu BM, Kemere C, Santhanam G, Afshar A, Ryu S, Meng TH, Sahani M, Shenoy K V (2007) Mixture of

trajectory models for neural decoding of goal-directed movements. J Neurophysiol 97:3763–3780.

Zimnik AJ, Churchland MM (2021) Independent generation of sequence elements by motor cortex. Nat

Neurosci 24:412–424.

745