Genetic association of photoplethysmography-derived arterial stiffness index with blood pressure and coronary artery disease

Seyedeh M. Zekavat BS (1-4), Mary Haas PhD (1), Krishna Aragam MD MS (1,4,5,6), Connor Emdin DPhil (1,5), Amit V. Khera MD MSc (1,4,5,6), Derek Klarin MD (1,4), Hongyu Zhao PhD (3,7), Pradeep Natarajan MD MMSc (1,4,5,6)

1. Program in Medical and Population Genetics, Broad Institute of MIT and Harvard, Cambridge, MA;
2. Yale School of Medicine, New Haven, CT;
3. Computational Biology & Bioinformatics Program, Yale University, New Haven, CT;
4. Center for Genomic Medicine, Massachusetts General Hospital, Boston, MA;
5. Harvard Medical School, Boston, MA;
6. Cardiovascular Research Center, Massachusetts General Hospital, Boston, MA;
7. Department of Biostatistics, Yale School of Public Health, New Haven, CT

Correspondence to:
Pradeep Natarajan, MD MMSc
pnatarajan@mgh.harvard.edu
185 Cambridge Street, CPZN 3.184, Boston, MA 02114
Tel: 617-724-3526
Fax: 617-726-2203

Abstract (max 350): 322

Total Word Count (minus Abstract): 4,064

Main text word count: 3488

Abstract

Background: Arterial stiffness index (ASI) has been proposed as an independent

mediator of blood pressure phenotypes and coronary artery disease (CAD) risk given

epidemiologic associations. However, it is unknown whether these associations

represent causal relationships. Here, we leverage human genetics to query whether

genetic predisposition to increased ASI (m/s), acquired from finger

photoplethysmography, is associated with elevated blood pressure and CAD risk.

Methods: Genome-wide association analysis (GWAS) of ASI was performed in 131,686

participants from the UK Biobank. Across 223,061 separate UK Biobank participants not

in the ASI GWAS, an ASI polygenic risk score consisting of the top 6 independent

variants ($P<5\times10^{-7}$, LD $r^2<0.25$) was applied. Mendelian randomization analyses of ASI

were performed with systolic blood pressure (SBP, N=208,897), diastolic blood pressure

(DBP, N=208,894), and incident CAD (7,534 cases) over 10 years follow-up in UK

Biobank. The lack of association observed with respect to CAD was replicated among

184,305 participants (60,810 cases) from the Coronary Artery Disease Genetics

Consortium (CARDIOGRAMplusC4D).

Results: Among the 131,686 individuals with ASI phenotype, median age was 59 (IQR

51-63) years, 70,847 (54%) were female. We replicated previous reports of the

epidemiologic association of ASI with SBP (Beta 0.55mmHg, [95% CI, 0.45-0.65],

$P=5.77\times10^{-24}$), DBP (Beta 1.05mmHg, [95% CI, 0.99-1.11], $P=7.27\times10^{-272}$), and incident

CAD (HR 1.09 [95% CI, 1.05-1.12], $P=1.5\times10^{-6}$) in multivariable models. Genome-wide

association analysis of ASI yielded two significant loci ($P<5\times10^{-8}$) at *TEX41-ZEB2* and

FOXO1, and three suggestive loci ($P<5\times10^{-7}$) at *COL4A2-COL4A1*, *RNF126*, and

TCF20. While each SD increase in genetic predisposition to elevated ASI was highly

associated with SBP (Beta **4.63 mmHg** [95% CI, 2.1-7.2]; *P*=3.37x10^{-4}), and DBP (Beta

2.61 mmHg [95% CI, 1.2-4.0]; *P*=2.85x10^{-4}), no association was observed with incident

CAD in UK Biobank (HR 1.12 [95% CI, 0.55-2.3]; *P*=0.75), or with prevalent CAD in

CARDIOGRAMplusC4D (OR 0.56 [95% CI, 0.26-1.24]; *P*=0.15).

Conclusions: A genetic predisposition to higher ASI was associated with elevated

systolic and diastolic blood pressures, but not associated with increased risk of

developing CAD.

Introduction:

Arterial stiffness index (ASI), as measured non-invasively via pulse wave analysis, has been independently associated with cardiovascular disease risk in multiple epidemiological studies[1-9]. Increased vascular resistance and diminished viscoelasticity are key features of vascular aging which have been previously associated with systolic hypertension[5], coronary artery disease (CAD)[2, 4, 7], and all-cause mortality.[10] Arterial stiffness may be influenced by variations in collagen, elastin, smooth muscle tone, and endothelial dysfunction, in addition to other factors[11-17]. Carotid-femoral (aortic) pulse wave velocity has been the 'gold-standard' approach for assessing arterial stiffness. ASI measurement using finger infrared analysis is a scalable, non-invasive approach to assess ASI and is correlated with aortic pulse wave velocity[18-20].

While arterial stiffness measures are associated with cardiovascular diseases[1-8], whether the associations are causal is not clear. For example, non-causal risk factors, such as high-density lipoprotein cholesterol for CAD, are good risk predictors but have been disappointing therapeutic targets[21-26]. Lifestyle factors are separately linked to arterial stiffness and cardiovascular diseases, potentially confounding the observed relationships[27]. Furthermore, reverse causality could lead to a statistically robust but non-causal relationship. For example, individuals with increased arterial stiffness might develop cardiovascular disease because of reduced exercise[28].

ASI has been proposed as a non-invasive surrogate end point for cardiovascular events largely based on robust epidemiological associations [29-31,2, 32-38]. Understanding whether

ASI causally mediates CAD, independent of blood pressure, may help determine

whether ASI is a suitable surrogate end point for CAD separate from its utility as a risk

predictor. Mendelian randomization uses human genetics for causal inference by

leveraging the random assortment of genetic variants during meiosis at conception[39].

This approach is therefore less susceptible to confounding or reverse causality. Here,

we used Mendelian randomization to determine whether a genetic predisposition to

increased arterial stiffness is associated with elevated blood pressure and increased

risk for incident CAD.

Methods:

UK Biobank study participants and phenotypes

Individual-level genomic data and longitudinal phenotypic data from the UK Biobank, a

large-scale population-based dataset consisting of genotype and phenotype data in

approximately 500,000 volunteer participants collected from 2007-2017, was used.

Consenting individuals where ASI was measured who did not have prevalent peripheral

arterial disease, aortic valve disease, or CAD were included in analysis.

Clinical disease definitions are detailed in **Supplementary Table 1**. In summary, the

main outcome, CAD, was defined by billing codes for heart attack, angina pectoris,

unstable angina, myocardial infarction, coronary atherosclerosis, coronary artery

revascularization, and other acute, subacute, and chronic forms of ischemic heart

disease, or with self-reported angina, heart attack/myocardial infarction, coronary

angioplasty +/- stent, or coronary artery bypass graft (CABG) surgery. We also

assessed systolic and diastolic blood pressures (adjusted for blood pressure

medications by adding 15 and 10 mmHg to systolic and diastolic blood pressures,

respectively[40]).

Arterial stiffness index measurement

ASI was previously measured in the UK Biobank using the PulseTrace PCA2

(CareFusion, San Diego, CA), which uses finger photoplethysmography over a 10- to

15-second timeframe to obtain the pulse waveform from an infrared sensor clipped to

the end of the index finger. ASI (in m/s) was calculated by dividing standing height by

the time between the systolic and diastolic peaks of the pulse waveform. ASI by this

approach was previously correlated with aortic pulse wave velocity, which is regarded

as the gold standard[18]. ASI was inverse rank normalized for analysis (with mean = 0,

SD = 1).

Genotyping and imputation

Genome-wide genotyping was previously performed in the UK Biobank using two

genotyping arrays sharing 95% of marker content: Applied Biosystems UK BiLEVE

Axiom Array (807,411 markers in 49,950 participants) and Applied Biosystems UK

Biobank Axiom Array (825,927 markers in 438,427 participants) both by Affymetrix

(Santa Clara, CA)[41]. Variants used in the present analysis include those also imputed

using the Haplotype Reference Consortium reference panel of up to 39M SNPs[41, 42].

Quality control and variant annotation

Poor quality variants and genotypes were filtered as previously described[41]. We further

filtered out individuals from both genetic and epidemiological analyses using the

following genetic criteria: non-white or not of British ancestry, gender mismatch between

reported and genotypic genders, sex chromosome aneuploidy, or one from each pair of

1st or 2nd degree relatives (**Supplementary Table 2**). Individuals with prevalent

peripheral arterial disease, aortic valve disease, or CAD were excluded, as were

extreme outliers for any of the arterial pulse wave phenotypes listed in **Supplementary**

Table 3. Extreme outliers were determined by adjusting the traditional box and whisker

upper and lower bounds and accounting for skewness in the phenotypic data identified

using the Robustbase package in R (setting range=3) (https://cran.r-

[project.org/web/packages/robustbase/robustbase.pdf](https://cran.r-project.org/web/packages/robustbase/robustbase.pdf)).

After filtering samples, variants were further filtered by the following criteria: not in

Hardy-Weinberg Equilibrium (P<1x10^{-10}), low imputation quality (INFO score < 0.3), call

rate < 95%, and minor allele frequency < 0.05% (minor allele count < 66).

Variant consequences were annotated using with Ensembl's Variant Effect Predictor

(VEP), ascribing the most severe consequence and associated gene among the

canonical transcripts present for each variant[43]. The Hail v0.1 software (https://hail.is)

was used to perform quality control and variant annotation[44].

Epidemiological association analyses with arterial stiffness index

Epidemiological association of ASI with blood pressure phenotypes and incident CAD

was performed using linear regression and Cox proportional hazards model,

respectively, in R (version 3.3, R Foundation, Vienna, Austria). For CAD, adjustment

was performed for age, sex, ever smoking status, heart rate at pulse wave analysis,

prevalent hypertension, prevalent hypercholesterolemia, prevalent diabetes, alcohol

intake (self-reported alcohol intake of at least once per month), exercise (self-reported

exercise of at least 3x per week), and vegetable intake (self-reported intake of at least 6

tablespoons of vegetable intake per day). The same adjustment variables were used for

SBP and DBP, except prevalent hypertension was not included as a covariate.

Blood pressure phenotypes included diastolic and systolic blood pressures (adjusted for

blood pressure medications by adding 10 and 15 mmHg to diastolic and systolic blood

pressures, respectively[40]). Analyses were performed using a Cox proportional hazards

model for incident CAD, and linear regression for the blood pressure traits. The

threshold for significance for the three primary phenotypes was defined as alpha =

0.05/3 tests = 0.017.

Genome-wide association analysis of arterial stiffness

A genome-wide association of ASI was performed using individual-level data from

131,686 individuals of European descent from the UK Biobank, collected from 2007 to

2017. Each variant was individually associated with ASI in an additive linear regression

model and adjusted for sex, age, smoking status, genotyping array type, and the first

ten principal components of ancestry[45]. Only variants with minor allele frequency >

0.05% (minor allele count > 66) were considered. $P < 5\times10^{-8}$ was considered to be

significant. The Hail software version 0.1 (https://hail.is) was used for genome-wide

association analysis[44].

Mendelian randomization

Overall, Mendelian randomization was performed by first creating a genetic instrument

for ASI then evaluating its association with blood pressure traits and incident CAD

among individuals not included in the ASI genome-wide association analysis.

An additive genetic risk score (GRS) was calculated as $\sum_{i=1}^{m} ln(OR_i) \times SNP_{ij}$ were m is

the number of SNPs, $ln(OR_i)$ is the weight for SNP_i from the discovery sample, SNP_{ij} is

the number of alleles (i.e., 0, 1, or 2) for SNP_i in person j in the validation sample. Six

independent variants (linkage disequilibrium $r^2 < 0.25$ within 500kb windows)

demonstrating at least suggestive association with ASI ($P<5x10^{-7}$) were included in the

GRS. The raw GRS was calculated for each individual using PLINK[46], inverse rank

normalized, then re-scaled such that one unit increase in the GRS was equivalent to a

one standard deviation (SD) increase in ASI.

To confirm that the GRS for ASI was a strong instrument for ASI, an F-statistic for the

instrument was calculated in the UK Biobank. An F-statistic is a measure of the

significance of an instrument (the GRS) for prediction of the exposure (ASI), controlling

for additional covariates (age, sex, ever smoked, 10 principal components of ancestry,

and genotyping array type). An F-statistic greater than 10 is evidence of a strong

instrument. Furthermore, sensitivity analyses were performed to evaluate for

associations between the ASI GRS and potential environmental confounders including

sex, ever smoking status, diet (alcohol intake, vegetable intake), and exercise frequency

among individuals not in the ASI genome-wide association analyses.

A linear regression model was used to associate the ASI GRS with systolic and diastolic

blood pressures. A Cox proportional hazards model was used to associate ASI GRS

with incident CAD. For CAD, adjustment was performed for age, sex, ever smoking

status, heart rate at blood pressure measurement, prevalent hypertension, prevalent

hypercholesterolemia, prevalent diabetes, alcohol intake (self-reported alcohol intake of

at least once per month), exercise (self-reported exercise of at least 3x per week), and

vegetable intake (self-reported intake of at least 6 tablespoons of vegetable intake per

day), where indicated. The same adjustment variables were used for SBP and DBP,

except for prevalent hypertension.

2-Sample Mendelian randomization with coronary artery disease

To address potential power limitations from the lack of association between ASI and

CAD, we also pursued 2-sample Mendelian randomization using variant-level summary

statistics from prior genome-wide association analyses of CAD from several

independent case-control studies, specifically 184,305 individuals from the Coronary

Artery Disease Genetics Consortium (CARDIOGRAMplusC4D)[47]. The effect estimates

and standard errors for the six GRS variants for ASI (from UK Biobank) and for CAD

(from CARDIOGramplusC4D) were used to perform robust, penalized inverse variance

weighted (IVW) 2-sample Mendelian randomization using the MendelianRandomization

package in R[48,49]. IVW 2-sample Mendelian randomization uses a weighted linear

regression of the ratio of the SNP effects on the outcomes to the SNP effects on the risk

factor, without using an intercept term. The threshold for significance was defined as

alpha = 0.05.

Additionally, analyses were performed to evaluate the reverse association, of CAD

causally impacting ASI. 77 known, independent, genome-wide significant CAD locus

variants were identified across several published sources[47, 50-52] (**Supplementary Table**

9). These 77 CAD locus variants were used as an instrument in 2-sample Mendelian

randomization to evaluate whether CAD causally affects ASI.

Results:

Baseline characteristics

A total of 131,686 individuals in the UK Biobank had ASI measured, genotype data

available, and passed quality control (**Supplementary Table 2**). Among these

individuals, median age was 59 (IQR 51-63) years, 53.8% were female, 4.6% had

diabetes, 27.1% had hypertension, and 12.9% had hypercholesterolemia. Median

systolic blood pressure was 139 (IQR 127-153) mmHg, median diastolic blood pressure

was 82 (IQR 75-89) mmHg. 44.1% of individuals were prior or current smokers, and

10.1% of individuals were on antihypertensive medications (**Table 1**). The median ASI

was 9 (IQR 7-11) m/s (**Supplementary Table 3**).

Epidemiological associations of ASI

Univariate association of cardiovascular risk factors with ASI showed the following

associations with at least nominal significance ($P<0.05$): for age (0.024 SD/year,

$P<1\times10^{-300}$), sex (0.40 SD higher in males, $P<1\times10^{-300}$), blood pressure medication (0.34

SD, $P=1.4\times10^{-317}$), hypertension (0.21 SD, $P=1.4\times10^{-269}$), hypercholesterolemia (0.20

SD, $P=4.1\times10^{-137}$), diabetes (0.20 SD, $P=9.1\times10^{-54}$), ever smoking (0.18 SD, $P=3.0\times10^{-}$

[250]), exercise $\geq$3x/wk (-0.16 SD, P=2.9x10[-66]), alcohol intake $\geq$1x/mo (0.05 SD,

P=3.3x10[-20]), and $\geq$6 tablespoons vegetable intake per day (-0.063 SD, *P*=3.1x10[-4])

(**Supplementary Table 4**).

For the associations of ASI with SBP and DBP, both univariable and multivariable,

adjusting for age, sex, smoking status, prevalent hypercholesterolemia, prevalent

diabetes, vegetable intake, alcohol intake, and exercise, analyses showed strong

associations (**Figure 1A**). Each SD of ASI was associated with elevated SBP by 0.55

mmHg ([95% CI, 0.45-0.65], *P*=5.77x10[-24]) and DBP by 1.05 mmHg ([95% CI, 0.99-

1.11], *P*=7.27x10[-272]).

ASI was also significantly independently associated with incident CAD, adjusting for

age, sex, ever smoking status, heart rate, prevalent hypertension, prevalent

hypercholesterolemia, prevalent diabetes, vegetable intake, alcohol intake, and exercise

(HR 1.08 per SD ASI [95% CI, 1.04-1.11], *P*=7.67x10[-6]) (**Figure 2A**).

Genome-wide association analysis of ASI

A genome-wide association analysis of ASI was performed among 131,686 individuals

and 13,995,214 variants in the UK Biobank. A quantile-quantile plot of the genome-wide

association statistics did not show substantial genomic inflation ($\lambda = 1.05$)

(**Supplementary Figure 1**). Two genome-wide significant loci were identified (*P*<5x10[-

8]), the top variants of which were in second intron of *TEX41*: rs1006923 (-0.025 SD,

P=3.7x10[-10], minor allele frequency (MAF)=0.32), and first intron of *FOXO1*: rs7331212

(-0.024 SD, P=9.3x10^{-9}, MAF=0.26). Three additional suggestive loci (P<5x10^{-7}) were

also identified, of which the top variants are intronic variants in *COL4A2:* rs872588 (-

0.020 SD, P=2.3x10^{-7}, MAF=0.42), *RNF126:* rs1009628 (-0.027 SD, P=1.2x10^{-7},

MAF=0.15), and *TCF20:* rs55906806 (-0.024 SD, P=2.4x10^{-7}, MAF=0.20). Through

chromatin conformational changes, intronic variants at *TEX41* and *COL4A2* may

influence gene expression at nearby enhancers **Supplementary Results,**

Supplementary Figure 2). Interrogation of disruptive protein-coding variants yielded

moderate association for *HFE* p.Cys282Tyr (MAF 0.076), the most common variant

implicated in hereditary hemochromatosis (**Supplementary Results**, **Supplementary**

Table 4).

Mendelian randomization in the UK Biobank

Six independent and at least suggestive (P<5x10^{-7}) variants were used towards an ASI

genetic risk score (GRS) (**Supplementary Table 5**). The raw ASI GRS was associated

with a 0.85 SD increase in ASI (SE: 0.072; P=8.0x10^{-32}). The F-statistic of the GRS was

123 (recommended F-statistic > 10), suggesting high instrument strength. The inverse-

rank normalized GRS was re-scaled such that each unit reflected one SD in ASI for

comparison with the phenotypic associations (**Supplementary Figure 3**). Sensitivity

analysis was performed to evaluate for associations between the ASI GRS and potential

environmental confounders including sex, ever smoking status, diet (alcohol intake,

vegetable intake), and exercise frequency. No significant associations between the ASI

GRS and environmental confounders were observed (**Supplementary Table 6**).

A 1-SD increase in genetically-mediated ASI was significantly associated with elevated

systolic blood pressure (Beta 4.63 mmHg [95% CI, 2.1-7.2]; P=3.37x10^{-4}), and diastolic

blood pressure (Beta 2.61 mmHg [95% CI, 1.2-4.0]; P=2.85x10^{-4}), independent of

cardiometabolic risk factors (age, sex, and smoking status, prevalent

hypercholesterolemia, prevalent diabetes, heart rate, vegetable intake, alcohol intake,

and exercise frequency) (**Figure 1B**).

The ASI GRS, however, was not associated with incident CAD in UK Biobank in an

unadjusted model (HR 1.3 [95% CI, 0.64-2.6]; P=0.47) or an adjusted model including

age, sex, smoking status, prevalent hypertension, prevalent hypercholesterolemia,

prevalent diabetes, heart rate, vegetable intake, alcohol intake, and exercise frequency

as covariates(HR 1.12 [95% CI, 0.55-2.3]; P=0.75) (**Figure 2B**).

2-Sample Mendelian randomization with coronary artery disease

To address potential power limitations impeding association of ASI GRS with incident

CAD in the UK Biobank, we also pursued 2-sample Mendelian randomization between

ASI and prevalent CAD using variant-level summary statistics from 184,305 separate

individuals in the Coronary Artery Disease Genetics Consortium

(CARDIOGRAMplusC4D)[47]. Robust, penalized inverse-variance weighted 2-sample

Mendelian randomization similarly did not demonstrate an association between

genetically-elevated ASI and CAD (OR 0.56 [95% CI, 0.26-1.24], P=0.15) (**Figure 3**).

Furthermore, the six individual variants showing at least suggestive association with ASI

did not demonstrate a significant positive association with CAD across several different

2-sample Mendelian randomization methods (**Supplementary Table 8**).

77 genome-wide significant CAD loci from past literature were assessed with respect to

both their ASI and CAD risk estimated effects (**Figure 4**). While 3 of 77 previously-

associated CAD loci had variants $P<0.05/77=6.5 \times 10^{-4}$ with respect to ASI, these

showed inconsistent effects between ASI and CAD. For example, the variant

rs9349379-A, an intronic variant in *PHACTR1*, was associated with increased ASI

(0.015 SD, $P=4.5 \times 10^{-5}$) but decreased risk for CAD (OR= 0.87, $P=1.8 \times 10^{-42}$). Similarly,

ASI-raising alleles at the *ZEB2-TEX41* and *ABO* loci decrease CAD risk, while ASI-

raising alleles at *CYP17A1-CNNM2-NT4C2* and *SH2B3* increase CAD risk. Detailed

variant-level summary statistics for these 77 CAD locus variants are provided in

Supplementary Tables 9-10. These 77 CAD locus variants were also used as an

instrument in 2-sample Mendelian randomization to evaluate the reverse association, of

CAD causally affecting ASI. No significant associations were observed across various

2-sample Mendelian randomization methods for the reverse association

(**Supplementary Table 11**).

<u>Discussion:</u>

We performed the largest genome-wide association analysis of a measure of vascular

aging, ASI, in 131,686 individuals, and leveraged these observations to perform causal

inference analyses with blood pressure and risk of CAD in 223,061 separate individuals.

In our genome-wide association analyses, we discover the first genome-wide variants

associated with ASI. We replicate the epidemiologic associations of ASI with blood

pressure and CAD, and find that genetic analyses do indeed support a causal

relationship between arterial stiffness and blood pressure. However, our genetic

analyses do not support a causal relationship between ASI and CAD.

These results permit several conclusions. First, we observe strong epidemiologic and

genetic association between ASI and blood pressure. These data indicate that non-

invasive photoplethysmography, employed by a finger probe or potentially

commercially-available wearable monitors that measure heart rate[53], may be used to

impute continuous blood pressure, and that changes will track with blood pressure

changes. However, given associated risk differences and independent effects, ASI

measurement may complement blood pressure assessments. Second, there is a long-

standing debate whether ASI precedes elevated blood pressure or vice versa[54].

Compared to its phenotypic effect, the effect conferred by genetically-elevated ASI is

8.4-fold higher for SBP (4.63 mmHg for ASI GRS versus 0.55 mmHg for ASI phenotype)

and 2.5-fold higher for DBP (2.61 mmHg for ASI GRS versus 1.05 mmHg for ASI

phenotype), potentially representing the effects of life-long exposure to elevated arterial

stiffness on blood pressure. This supports the notion that arterial stiffness may predate

the onset of elevated blood pressure indicating that ASI may independently identify

individuals at heightened risk for future blood pressure elevations.

Third, our epidemiological and genetic analyses indicate that ASI is an independent,

non-causal risk factor for CAD. Arterial stiffness may be a parallel disrupted pathway in

the setting of CAD, as opposed to an upstream causal mediating factor. Thus, while ASI

monitoring may still serve as a good proxy for blood pressure, therapeutic modulation of

ASI in isolation may not have a meaningful impact on CAD outcomes. Similarly, a

recent study of twins showed that while carotid-femoral pulse wave velocity was

heritable, it did not associate with 5-year progression of carotid intima media

thickness[55]. The lack of significance between genetically-elevated ASI and CAD is also

consistent with prior mixed results in experimental models. Fragmentation of elastin

fibers and deposition of collagen fibers are features of vascular aging implicated in

arterial stiffness[56]. However, murine models lacking elastin do not have endothelial

damage, thrombosis, or inflammation which typically occur with atherosclerosis[57].

Furthermore, we found that while few variants associated with CAD show apparent

association with ASI, our data indicate that ASI may not be mediating the apparent CAD

risk. We observed inconsistent genetic effects between ASI and coronary artery disease

risk, whereby many genetic variants had different directions of effect on arterial stiffness

index versus coronary artery disease. In particular, an intronic variant within *PHACTR1*

(rs9349379-A), which was recently shown to influence endothelin-1 expression in the

vasculature, is associated with decreased risk for CAD[61], increased blood pressure[62],

and increased ASI. Additionally, genetic variants disrupting nitric oxide signaling at the

NOS3 and *GUCY13* loci influence both blood pressure and risk of CAD[63-65]. Notably, in

our study, risk variants at these loci were not strongly associated with ASI. Extensive

prior experimental work has linked nitric oxide signaling and endothelin-1 with

endothelial function and vascular tone[66-70]. Our data suggests that increased risk of

CAD through these pathways is unlikely to be through changes in ASI but potentially

through alternative vascular mechanisms.

While our study has several strengths, some limitations should be considered. First,

lack of ASI genetic risk score association with CAD may be due to limited statistical

power. Our replication of the lack of association using 2-sample Mendelian

randomization, combined with our analysis showing inconsistent effects of individual

variants between CAD and ASI suggests that this is less likely. Second, our imputation

of untreated blood pressure among those with prescribed hypertensives assumes a

homogenous blood pressure effect across the population. Without prescription data in

the UK Biobank, we are unable to account for different medication regimens and

adherence.

Conclusion:

A genetic predisposition to higher ASI was associated with increased blood pressure,

but not increased risk of CAD. Our data support the conclusion that finger

photoplethysmography-derived ASI is an independent, causal risk factor for blood

pressure and an independent, non-causal risk factor for CAD.

Sources of Funding

S.M.Z. and P.N. had full access to all the data in the study and take responsibility for the integrity of the data and accuracy of the data analysis. S.M.Z. and P.N. conceived of and designed the study. P.N. acquired the data. S.M.Z., M.H., and K.A. analyzed the data. S.M.Z., C.E., A.V.K., D.K., H.Z., and P.N. interpreted the data. S.M.Z. and P.N. drafted the manuscript. S.M.Z. is supported by the National Institutes of Health's Medical Scientist Training Program at the Yale School of Medicine and the Paul and Daisy Soros Fellowship. P.N. is supported by a Hassenfeld Scholar Award from the Massachusetts General Hospital, and K08 HL140203 from the National Heart, Lung, and Blood Institute. The authors would like to acknowledge the participants, study staff, and investigators of the UK Biobank and CARDIOGRAMplusC4D Consortium.

Acknowledgements

The authors would like to acknowledge and thank the participants and staff of the UK Biobank and of the CARDIOGramplusC4D consortium cohorts.

Conflicts of Interest

P.N. reports investigator-initiated grants from Amgen and Boston Scientific, and consulting income from Apple.

References:

1.	Ikonomidis I, Ntai K, Kadoglou NP, Papadakis I, Kornelakis M, Tritakis V, Varoudi M, Papadima T, Triantafyllidi H, Parissis J, Paraskevaidis I and Lekakis J. The evaluation of pulse wave velocity using Arteriograph and Complior apparatus across multiple cohorts of cardiovascular-related diseases. *Int J Cardiol*. 2013;168:4890-2.

2.	Mattace-Raso FU, van der Cammen TJ, Hofman A, van Popele NM, Bos ML, Schalekamp MA, Asmar R, Reneman RS, Hoeks AP, Breteler MM and Witteman JC. Arterial stiffness and risk of coronary heart disease and stroke: the Rotterdam Study. *Circulation*. 2006;113:657-63.

3.	Ben-Shlomo Y, Spears M, Boustred C, May M, Anderson SG, Benjamin EJ, Boutouyrie P, Cameron J, Chen CH, Cruickshank JK, Hwang SJ, Lakatta EG, Laurent S, Maldonado J, Mitchell GF, Najjar SS, Newman AB, Ohishi M, Pannier B, Pereira T, Vasan RS, Shokawa T, Sutton-Tyrell K, Verbeke F, Wang KL, Webb DJ, Willum Hansen T, Zoungas S, McEniery CM, Cockcroft JR and Wilkinson IB. Aortic pulse wave velocity improves cardiovascular event prediction: an individual participant meta-analysis of prospective observational data from 17,635 subjects. *J Am Coll Cardiol*. 2014;63:636-46.

4.	Ikonomidis I, Makavos G and Lekakis J. Arterial stiffness and coronary artery disease. *Curr Opin Cardiol*. 2015;30:422-31.

5.	Franklin SS, Gustin Wt, Wong ND, Larson MG, Weber MA, Kannel WB and Levy D. Hemodynamic patterns of age-related changes in blood pressure. The Framingham Heart Study. *Circulation*. 1997;96:308-15.

6.	Chirinos JA, Kips JG, Jacobs DR, Jr., Brumback L, Duprez DA, Kronmal R, Bluemke DA, Townsend RR, Vermeersch S and Segers P. Arterial wave reflections and incident cardiovascular events and heart failure: MESA (Multiethnic Study of Atherosclerosis). *J Am Coll Cardiol*. 2012;60:2170-7.

7.	Sutton-Tyrell K, Najjar SS, Boudreau RM, Venkitachalam L, Kupelian V, Simonsick EM, Havlik R, Lakatta EG, Spurgeon H, Kritchevsky S, Pahor M, Bauer D, Newman A and Health ABCS. Elevated aortic pulse wave velocity, a marker of arterial stiffness, predicts cardiovascular events in well-functioning older adults. *Circulation*. 2005;111:3384-90.

8.	Tsao CW, Lyass A, Larson MG, Levy D, Hamburg NM, Vita JA, Benjamin EJ, Mitchell GF and Vasan RS. Relation of Central Arterial Stiffness to Incident Heart Failure in the Community. *J Am Heart Assoc*. 2015;4.

9.	Said MA, Eppinga RN, Lipsic E, Verweij N and van der Harst P. Relationship of Arterial Stiffness Index and Pulse Pressure With Cardiovascular Disease and Mortality. *J Am Heart Assoc*. 2018;7.

10.	Vlachopoulos C, Aznaouridis K and Stefanadis C. Prediction of cardiovascular events and all-cause mortality with arterial stiffness: a systematic review and meta-analysis. *J Am Coll Cardiol*. 2010;55:1318-27.

11.	Bank AJ and Kaiser DR. Smooth muscle relaxation: effects on arterial compliance, distensibility, elastic modulus, and pulse wave velocity. *Hypertension*. 1998;32:356-9.

12.	Anderson TJ. Arterial stiffness or endothelial dysfunction as a surrogate marker of vascular risk. *Can J Cardiol*. 2006;22 Suppl B:72B-80B.

13.	Ikonomidis I, Lekakis J, Papadopoulos C, Triantafyllidi H, Paraskevaidis I, Georgoula G, Tzortzis S, Revela I and Kremastinos DT. Incremental value of pulse wave velocity in the

determination of coronary microcirculatory dysfunction in never-treated patients with essential hypertension. *Am J Hypertens*. 2008;21:806-13.

14.   McNulty M, Mahmud A, Spiers P and Feely J. Collagen type-I degradation is related to arterial stiffness in hypertensive and normotensive subjects. *J Hum Hypertens*. 2006;20:867-73.

15.   Nigam A, Mitchell GF, Lambert J and Tardif JC. Relation between conduit vessel stiffness (assessed by tonometry) and endothelial function (assessed by flow-mediated dilatation) in patients with and without coronary heart disease. *Am J Cardiol*. 2003;92:395-9.

16.   Wagenseil JE and Mecham RP. Elastin in large artery stiffness and hypertension. *J Cardiovasc Transl Res*. 2012;5:264-73.

17.   Zieman SJ, Melenovsky V and Kass DA. Mechanisms, pathophysiology, and therapy of arterial stiffness. *Arterioscler Thromb Vasc Biol*. 2005;25:932-43.

18.   Woodman RJ, Kingwell BA, Beilin LJ, Hamilton SE, Dart AM and Watts GF. Assessment of central and peripheral arterial stiffness: studies indicating the need to use a combination of techniques. *Am J Hypertens*. 2005;18:249-60.

19.   Millasseau SC, Kelly RP, Ritter JM and Chowienczyk PJ. Determination of age-related increases in large artery stiffness by digital pulse contour analysis. *Clin Sci (Lond)*. 2002;103:371-7.

20.   Alty SR, Angarita-Jaimes N, Millasseau SC and Chowienczyk PJ. Predicting arterial stiffness from the digital volume pulse waveform. *IEEE Trans Biomed Eng*. 2007;54:2268-75.

21.   Brousseau ME, Schaefer EJ, Wolfe ML, Bloedon LT, Digenio AG, Clark RW, Mancuso JP and Rader DJ. Effects of an inhibitor of cholesteryl ester transfer protein on HDL cholesterol. *N Engl J Med*. 2004;350:1505-15.

22.   Emerging Risk Factors C, Di Angelantonio E, Sarwar N, Perry P, Kaptoge S, Ray KK, Thompson A, Wood AM, Lewington S, Sattar N, Packard CJ, Collins R, Thompson SG and Danesh J. Major lipids, apolipoproteins, and risk of vascular disease. *JAMA*. 2009;302:1993-2000.

23.   Group HTC, Landray MJ, Haynes R, Hopewell JC, Parish S, Aung T, Tomson J, Wallendszus K, Craig M, Jiang L, Collins R and Armitage J. Effects of extended-release niacin with laropiprant in high-risk patients. *N Engl J Med*. 2014;371:203-12.

24.   Investigators A-H, Boden WE, Probstfield JL, Anderson T, Chaitman BR, Desvignes-Nickens P, Koprowicz K, McBride R, Teo K and Weintraub W. Niacin in patients with low HDL cholesterol levels receiving intensive statin therapy. *N Engl J Med*. 2011;365:2255-67.

25.   Schwartz GG, Olsson AG, Abt M, Ballantyne CM, Barter PJ, Brumm J, Chaitman BR, Holme IM, Kallend D, Leiter LA, Leitersdorf E, McMurray JJ, Mundl H, Nicholls SJ, Shah PK, Tardif JC, Wright RS and dal OI. Effects of dalcetrapib in patients with a recent acute coronary syndrome. *N Engl J Med*. 2012;367:2089-99.

26.   Voight BF, Peloso GM, Orho-Melander M, Frikke-Schmidt R, Barbalic M, Jensen MK, Hindy G, Holm H, Ding EL, Johnson T, Schunkert H, Samani NJ, Clarke R, Hopewell JC, Thompson JF, Li M, Thorleifsson G, Newton-Cheh C, Musunuru K, Pirruccello JP, Saleheen D, Chen L, Stewart A, Schillert A, Thorsteinsdottir U, Thorgeirsson G, Anand S, Engert JC, Morgan T, Spertus J, Stoll M, Berger K, Martinelli N, Girelli D, McKeown PP, Patterson CC, Epstein SE, Devaney J, Burnett MS, Mooser V, Ripatti S, Surakka I, Nieminen MS, Sinisalo J, Lokki ML, Perola M, Havulinna A, de Faire U, Gigante B, Ingelsson E, Zeller T, Wild P, de Bakker PI, Klungel OH, Maitland-van der Zee AH, Peters BJ, de Boer A, Grobbee DE, Kamphuisen PW, Deneer VH, Elbers CC, Onland-Moret NC, Hofker MH, Wijmenga C, Verschuren WM, Boer JM, van der

Schouw YT, Rasheed A, Frossard P, Demissie S, Willer C, Do R, Ordovas JM, Abecasis GR, Boehnke M, Mohlke KL, Daly MJ, Guiducci C, Burtt NP, Surti A, Gonzalez E, Purcell S, Gabriel S, Marrugat J, Peden J, Erdmann J, Diemert P, Willenborg C, Konig IR, Fischer M, Hengstenberg C, Ziegler A, Buysschaert I, Lambrechts D, Van de Werf F, Fox KA, El Mokhtari NE, Rubin D, Schrezenmeir J, Schreiber S, Schafer A, Danesh J, Blankenberg S, Roberts R, McPherson R, Watkins H, Hall AS, Overvad K, Rimm E, Boerwinkle E, Tybjaerg-Hansen A, Cupples LA, Reilly MP, Melander O, Mannucci PM, Ardissino D, Siscovick D, Elosua R, Stefansson K, O'Donnell CJ, Salomaa V, Rader DJ, Peltonen L, Schwartz SM, Altshuler D and Kathiresan S. Plasma HDL cholesterol and risk of myocardial infarction: a mendelian randomisation study. *Lancet*. 2012;380:572-80.

27.   Tanaka H and Safar ME. Influence of lifestyle modification on arterial stiffness and wave reflections. *Am J Hypertens*. 2005;18:137-44.

28.   Sacre JW, Jennings GL and Kingwell BA. Exercise and dietary influences on arterial stiffness in cardiometabolic disease. *Hypertension*. 2014;63:888-93.

29.   Laurent S, Briet M and Boutouyrie P. Arterial stiffness as surrogate end point: needed clinical trials. *Hypertension*. 2012;60:518-22.

30.   Laurent S. Arterial stiffness: intermediate or surrogate endpoint for cardiovascular events? *Eur Heart J*. 2005;26:1152-4.

31.   Cohn JN, Quyyumi AA, Hollenberg NK and Jamerson KA. Surrogate markers for cardiovascular disease: functional markers. *Circulation*. 2004;109:IV31-46.

32.   Blacher J, Guerin AP, Pannier B, Marchais SJ, Safar ME and London GM. Impact of aortic stiffness on survival in end-stage renal disease. *Circulation*. 1999;99:2434-9.

33.   Laurent S, Boutouyrie P, Asmar R, Gautier I, Laloux B, Guize L, Ducimetiere P and Benetos A. Aortic stiffness is an independent predictor of all-cause and cardiovascular mortality in hypertensive patients. *Hypertension*. 2001;37:1236-41.

34.   London GM, Blacher J, Pannier B, Guerin AP, Marchais SJ and Safar ME. Arterial wave reflections and survival in end-stage renal failure. *Hypertension*. 2001;38:434-8.

35.   Mitchell GF, Moye LA, Braunwald E, Rouleau JL, Bernstein V, Geltman EM, Flaker GC and Pfeffer MA. Sphygmomanometrically determined pulse pressure is a powerful independent predictor of recurrent events after myocardial infarction in patients with impaired left ventricular function. SAVE investigators. Survival and Ventricular Enlargement. *Circulation*. 1997;96:4254-60.

36.   Safar ME. Systolic blood pressure, pulse pressure and arterial stiffness as cardiovascular risk factors. *Curr Opin Nephrol Hypertens*. 2001;10:257-61.

37.   van Popele NM, Grobbee DE, Bots ML, Asmar R, Topouchian J, Reneman RS, Hoeks AP, van der Kuip DA, Hofman A and Witteman JC. Association between arterial stiffness and atherosclerosis: the Rotterdam Study. *Stroke*. 2001;32:454-60.

38.   Weber T, Auer J, O'Rourke MF, Kvas E, Lassnig E, Berent R and Eber B. Arterial stiffness, wave reflections, and the risk of coronary artery disease. *Circulation*. 2004;109:184-9.

39.   Smith GD and Ebrahim S. 'Mendelian randomization': can genetic epidemiology contribute to understanding environmental determinants of disease? *Int J Epidemiol*. 2003;32:1-22.

40.   Warren HR, Evangelou E, Cabrera CP, Gao H, Ren M, Mifsud B, Ntalla I, Surendran P, Liu C, Cook JP, Kraja AT, Drenos F, Loh M, Verweij N, Marten J, Karaman I, Lepe MP, O'Reilly PF,

Knight J, Snieder H, Kato N, He J, Tai ES, Said MA, Porteous D, Alver M, Poulter N, Farrall M, Gansevoort RT, Padmanabhan S, Magi R, Stanton A, Connell J, Bakker SJ, Metspalu A, Shields DC, Thom S, Brown M, Sever P, Esko T, Hayward C, van der Harst P, Saleheen D, Chowdhury R, Chambers JC, Chasman DI, Chakravarti A, Newton-Cheh C, Lindgren CM, Levy D, Kooner JS, Keavney B, Tomaszewski M, Samani NJ, Howson JM, Tobin MD, Munroe PB, Ehret GB, Wain LV, International Consortium of Blood Pressure GA, Consortium B, Lifelines Cohort S, Understanding Society Scientific g, Consortium CHDE, Exome BPC, Consortium TDG, Go TDC, Cohorts for H, Ageing Research in Genome Epidemiology BPEC, International Genomics of Blood Pressure C and group UKBCCBw. Genome-wide association analysis identifies novel blood pressure loci and offers biological insights into cardiovascular risk. *Nat Genet*. 2017;49:403-415.

41.   Bycroft C, Freeman C, Petkova D, Band G, Elliott LT, Sharp K, Motyer A, Vukcevic D, Delaneau O, O'Connell J, Cortes A, Welsh S, McVean G, Leslie S, Donnelly P and Marchini J. Genome-wide genetic data on ~500,000 UK Biobank participants. *bioRxiv*. 2017.

42.   McCarthy S, Das S, Kretzschmar W, Delaneau O, Wood AR, Teumer A, Kang HM, Fuchsberger C, Danecek P, Sharp K, Luo Y, Sidore C, Kwong A, Timpson N, Koskinen S, Vrieze S, Scott LJ, Zhang H, Mahajan A, Veldink J, Peters U, Pato C, van Duijn CM, Gillies CE, Gandin I, Mezzavilla M, Gilly A, Cocca M, Traglia M, Angius A, Barrett JC, Boomsma D, Branham K, Breen G, Brummett CM, Busonero F, Campbell H, Chan A, Chen S, Chew E, Collins FS, Corbin LJ, Smith GD, Dedoussis G, Dorr M, Farmaki AE, Ferrucci L, Forer L, Fraser RM, Gabriel S, Levy S, Groop L, Harrison T, Hattersley A, Holmen OL, Hveem K, Kretzler M, Lee JC, McGue M, Meitinger T, Melzer D, Min JL, Mohlke KL, Vincent JB, Nauck M, Nickerson D, Palotie A, Pato M, Pirastu N, McInnis M, Richards JB, Sala C, Salomaa V, Schlessinger D, Schoenherr S, Slagboom PE, Small K, Spector T, Stambolian D, Tuke M, Tuomilehto J, Van den Berg LH, Van Rheenen W, Volker U, Wijmenga C, Toniolo D, Zeggini E, Gasparini P, Sampson MG, Wilson JF, Frayling T, de Bakker PI, Swertz MA, McCarroll S, Kooperberg C, Dekker A, Altshuler D, Willer C, Iacono W, Ripatti S, Soranzo N, Walter K, Swaroop A, Cucca F, Anderson CA, Myers RM, Boehnke M, McCarthy MI, Durbin R and Haplotype Reference C. A reference panel of 64,976 haplotypes for genotype imputation. *Nat Genet*. 2016;48:1279-83.

43.   McLaren W, Pritchard B, Rios D, Chen Y, Flicek P and Cunningham F. Deriving the consequences of genomic variants with the Ensembl API and SNP Effect Predictor. *Bioinformatics*. 2010;26:2069-70.

44.   Ganna A, Genovese G, Howrigan DP, Byrnes A, Kurki M, Zekavat SM, Whelan CW, Kals M, Nivard MG, Bloemendal A, Bloom JM, Goldstein JI, Poterba T, Seed C, Handsaker RE, Natarajan P, Magi R, Gage D, Robinson EB, Metspalu A, Salomaa V, Suvisaari J, Purcell SM, Sklar P, Kathiresan S, Daly MJ, McCarroll SA, Sullivan PF, Palotie A, Esko T, Hultman C and Neale BM. Ultra-rare disruptive and damaging mutations influence educational attainment in the general population. *Nat Neurosci*. 2016;19:1563-1565.

45.   Hoggart CJ, Parra EJ, Shriver MD, Bonilla C, Kittles RA, Clayton DG and McKeigue PM. Control of confounding of genetic associations in stratified populations. *Am J Hum Genet*. 2003;72:1492-1504.

46.   Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D, Maller J, Sklar P, de Bakker PI, Daly MJ and Sham PC. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet*. 2007;81:559-75.

47.   Nikpay M, Goel A, Won HH, Hall LM, Willenborg C, Kanoni S, Saleheen D, Kyriakou T, Nelson CP, Hopewell JC, Webb TR, Zeng L, Dehghan A, Alver M, Armasu SM, Auro K, Bjonnes A, Chasman DI, Chen S, Ford I, Franceschini N, Gieger C, Grace C, Gustafsson S, Huang J, Hwang SJ, Kim YK, Kleber ME, Lau KW, Lu X, Lu Y, Lyytikainen LP, Mihailov E, Morrison AC, Pervjakova N, Qu L, Rose LM, Salfati E, Saxena R, Scholz M, Smith AV, Tikkanen E, Uitterlinden A, Yang X, Zhang W, Zhao W, de Andrade M, de Vries PS, van Zuydam NR, Anand SS, Bertram L, Beutner F, Dedoussis G, Frossard P, Gauguier D, Goodall AH, Gottesman O, Haber M, Han BG, Huang J, Jalilzadeh S, Kessler T, Konig IR, Lannfelt L, Lieb W, Lind L, Lindgren CM, Lokki ML, Magnusson PK, Mallick NH, Mehra N, Meitinger T, Memon FU, Morris AP, Nieminen MS, Pedersen NL, Peters A, Rallidis LS, Rasheed A, Samuel M, Shah SH, Sinisalo J, Stirrups KE, Trompet S, Wang L, Zaman KS, Ardissino D, Boerwinkle E, Borecki IB, Bottinger EP, Buring JE, Chambers JC, Collins R, Cupples LA, Danesh J, Demuth I, Elosua R, Epstein SE, Esko T, Feitosa MF, Franco OH, Franzosi MG, Granger CB, Gu D, Gudnason V, Hall AS, Hamsten A, Harris TB, Hazen SL, Hengstenberg C, Hofman A, Ingelsson E, Iribarren C, Jukema JW, Karhunen PJ, Kim BJ, Kooner JS, Kullo IJ, Lehtimaki T, Loos RJF, Melander O, Metspalu A, Marz W, Palmer CN, Perola M, Quertermous T, Rader DJ, Ridker PM, Ripatti S, Roberts R, Salomaa V, Sanghera DK, Schwartz SM, Seedorf U, Stewart AF, Stott DJ, Thiery J, Zalloua PA, O'Donnell CJ, Reilly MP, Assimes TL, Thompson JR, Erdmann J, Clarke R, Watkins H, Kathiresan S, McPherson R, Deloukas P, Schunkert H, Samani NJ and Farrall M. A comprehensive 1,000 Genomes-based genome-wide association meta-analysis of coronary artery disease. *Nat Genet*. 2015;47:1121-1130.

48.   Yavorska OO and Burgess S. MendelianRandomization: an R package for performing Mendelian randomization analyses using summarized data. *Int J Epidemiol*. 2017;46:1734-1739.

49.   Burgess S, Bowden J, Dudbridge F and Thompson SG. Robust instrumental variable methods using multiple candidate instruments with application to Mendelian randomization. *ArXiv e-prints*. 2016.

50.   Webb TR, Erdmann J, Stirrups KE, Stitziel NO, Masca NG, Jansen H, Kanoni S, Nelson CP, Ferrario PG, Konig IR, Eicher JD, Johnson AD, Hamby SE, Betsholtz C, Ruusalepp A, Franzen O, Schadt EE, Bjorkegren JL, Weeke PE, Auer PL, Schick UM, Lu Y, Zhang H, Dube MP, Goel A, Farrall M, Peloso GM, Won HH, Do R, van Iperen E, Kruppa J, Mahajan A, Scott RA, Willenborg C, Braund PS, van Capelleveen JC, Doney AS, Donnelly LA, Asselta R, Merlini PA, Duga S, Marziliano N, Denny JC, Shaffer C, El-Mokhtari NE, Franke A, Heilmann S, Hengstenberg C, Hoffmann P, Holmen OL, Hveem K, Jansson JH, Jockel KH, Kessler T, Kriebel J, Laugwitz KL, Marouli E, Martinelli N, McCarthy MI, Van Zuydam NR, Meisinger C, Esko T, Mihailov E, Escher SA, Alver M, Moebus S, Morris AD, Virtamo J, Nikpay M, Olivieri O, Provost S, AlQarawi A, Robertson NR, Akinsansya KO, Reilly DF, Vogt TF, Yin W, Asselbergs FW, Kooperberg C, Jackson RD, Stahl E, Muller-Nurasyid M, Strauch K, Varga TV, Waldenberger M, Wellcome Trust Case Control C, Zeng L, Chowdhury R, Salomaa V, Ford I, Jukema JW, Amouyel P, Kontto J, Investigators M, Nordestgaard BG, Ferrieres J, Saleheen D, Sattar N, Surendran P, Wagner A, Young R, Howson JM, Butterworth AS, Danesh J, Ardissino D, Bottinger EP, Erbel R, Franks PW, Girelli D, Hall AS, Hovingh GK, Kastrati A, Lieb W, Meitinger T, Kraus WE, Shah SH, McPherson R, Orho-Melander M, Melander O, Metspalu A, Palmer CN, Peters A, Rader DJ, Reilly MP, Loos RJ, Reiner AP, Roden DM, Tardif JC, Thompson JR, Wareham NJ, Watkins H, Willer CJ, Samani NJ, Schunkert H, Deloukas P, Kathiresan S, Myocardial Infarction G and Investigators CAEC. Systematic Evaluation

of Pleiotropy Identifies 6 Further Loci Associated With Coronary Artery Disease. *J Am Coll Cardiol*. 2017;69:823-836.

51.   Klarin D, Zhu QM, Emdin CA, Chaffin M, Horner S, McMillan BJ, Leed A, Weale ME, Spencer CCA, Aguet F, Segre AV, Ardlie KG, Khera AV, Kaushik VK, Natarajan P, Consortium CAD and Kathiresan S. Genetic analysis in UK Biobank links insulin resistance and transendothelial migration pathways to coronary artery disease. *Nat Genet*. 2017;49:1392-1397.

52.   Consortium CAD, Deloukas P, Kanoni S, Willenborg C, Farrall M, Assimes TL, Thompson JR, Ingelsson E, Saleheen D, Erdmann J, Goldstein BA, Stirrups K, Konig IR, Cazier JB, Johansson A, Hall AS, Lee JY, Willer CJ, Chambers JC, Esko T, Folkersen L, Goel A, Grundberg E, Havulinna AS, Ho WK, Hopewell JC, Eriksson N, Kleber ME, Kristiansson K, Lundmark P, Lyytikainen LP, Rafelt S, Shungin D, Strawbridge RJ, Thorleifsson G, Tikkanen E, Van Zuydam N, Voight BF, Waite LL, Zhang W, Ziegler A, Absher D, Altshuler D, Balmforth AJ, Barroso I, Braund PS, Burgdorf C, Claudi-Boehm S, Cox D, Dimitriou M, Do R, Consortium D, Consortium C, Doney AS, El Mokhtari N, Eriksson P, Fischer K, Fontanillas P, Franco-Cereceda A, Gigante B, Groop L, Gustafsson S, Hager J, Hallmans G, Han BG, Hunt SE, Kang HM, Illig T, Kessler T, Knowles JW, Kolovou G, Kuusisto J, Langenberg C, Langford C, Leander K, Lokki ML, Lundmark A, McCarthy MI, Meisinger C, Melander O, Mihailov E, Maouche S, Morris AD, Muller-Nurasyid M, Mu TC, Nikus K, Peden JF, Rayner NW, Rasheed A, Rosinger S, Rubin D, Rumpf MP, Schafer A, Sivananthan M, Song C, Stewart AF, Tan ST, Thorgeirsson G, van der Schoot CE, Wagner PJ, Wellcome Trust Case Control C, Wells GA, Wild PS, Yang TP, Amouyel P, Arveiler D, Basart H, Boehnke M, Boerwinkle E, Brambilla P, Cambien F, Cupples AL, de Faire U, Dehghan A, Diemert P, Epstein SE, Evans A, Ferrario MM, Ferrieres J, Gauguier D, Go AS, Goodall AH, Gudnason V, Hazen SL, Holm H, Iribarren C, Jang Y, Kahonen M, Kee F, Kim HS, Klopp N, Koenig W, Kratzer W, Kuulasmaa K, Laakso M, Laaksonen R, Lee JY, Lind L, Ouwehand WH, Parish S, Park JE, Pedersen NL, Peters A, Quertermous T, Rader DJ, Salomaa V, Schadt E, Shah SH, Sinisalo J, Stark K, Stefansson K, Tregouet DA, Virtamo J, Wallentin L, Wareham N, Zimmermann ME, Nieminen MS, Hengstenberg C, Sandhu MS, Pastinen T, Syvanen AC, Hovingh GK, Dedoussis G, Franks PW, Lehtimaki T, Metspalu A, Zalloua PA, Siegbahn A, Schreiber S, Ripatti S, Blankenberg SS, Perola M, Clarke R, Boehm BO, O'Donnell C, Reilly MP, Marz W, Collins R, Kathiresan S, Hamsten A, Kooner JS, Thorsteinsdottir U, Danesh J, Palmer CN, Roberts R, Watkins H, Schunkert H and Samani NJ. Large-scale association analysis identifies new risk loci for coronary artery disease. *Nat Genet*. 2013;45:25-33.

53.   Georgiou K, Larentzakis AV, Khamis NN, Alsuhaibani GI, Alaska YA and Giallafos EJ. Can Wearable Devices Accurately Measure Heart Rate Variability? A Systematic Review. *Folia Med (Plovdiv)*. 2018;60:7-20.

54.   Mitchell GF. Arterial stiffness and hypertension: chicken or egg? *Hypertension*. 2014;64:210-4.

55.   Cecelja M, Jiang B, Keehn L, Hussain T, Silva Vieira M, Phinikaridou A, Greil G, Spector TD and Chowienczyk P. Arterial stiffening is a heritable trait associated with arterial dilation but not wall thickening: a longitudinal study in the twins UK cohort. *Eur Heart J*. 2018;39:2282-2288.

56.   Xu J and Shi GP. Vascular wall extracellular matrix proteins and vascular diseases. *Biochim Biophys Acta*. 2014;1842:2106-2119.

57.   Li DY, Brooke B, Davis EC, Mecham RP, Sorensen LK, Boak BB, Eichwald E and Keating MT. Elastin is an essential determinant of arterial morphogenesis. *Nature*. 1998;393:276-80.

58.	Jimi S, Saku K, Uesugi N, Sakata N and Takebayashi S. Oxidized low density lipoprotein stimulates collagen production in cultured arterial smooth muscle cells. *Atherosclerosis*. 1995;116:15-26.

59.	Jimi S, Sakata N and Takebayashi S. Oxidized LDL induces an increase in the relative collagen synthesis of rabbit aortic smooth muscle cells. *J Atheroscler Thromb*. 1994;1:53-9.

60.	Orekhov AN, Tertov VV, Kudryashov SA and Smirnov VN. Triggerlike stimulation of cholesterol accumulation and DNA and extracellular matrix synthesis induced by atherogenic serum or low density lipoprotein in cultured cells. *Circ Res*. 1990;66:311-20.

61.	Gupta RM, Hadaya J, Trehan A, Zekavat SM, Roselli C, Klarin D, Emdin CA, Hilvering CRE, Bianchi V, Mueller C, Khera AV, Ryan RJH, Engreitz JM, Issner R, Shoresh N, Epstein CB, de Laat W, Brown JD, Schnabel RB, Bernstein BE and Kathiresan S. A Genetic Variant Associated with Five Vascular Diseases Is a Distal Regulator of Endothelin-1 Gene Expression. *Cell*. 2017;170:522-533 e15.

62.	Surendran P, Drenos F, Young R, Warren H, Cook JP, Manning AK, Grarup N, Sim X, Barnes DR, Witkowska K, Staley JR, Tragante V, Tukiainen T, Yaghootkar H, Masca N, Freitag DF, Ferreira T, Giannakopoulou O, Tinker A, Harakalova M, Mihailov E, Liu C, Kraja AT, Fallgaard Nielsen S, Rasheed A, Samuel M, Zhao W, Bonnycastle LL, Jackson AU, Narisu N, Swift AJ, Southam L, Marten J, Huyghe JR, Stancakova A, Fava C, Ohlsson T, Matchan A, Stirrups KE, Bork-Jensen J, Gjesing AP, Kontto J, Perola M, Shaw-Hawkins S, Havulinna AS, Zhang H, Donnelly LA, Groves CJ, Rayner NW, Neville MJ, Robertson NR, Yiorkas AM, Herzig KH, Kajantie E, Zhang W, Willems SM, Lannfelt L, Malerba G, Soranzo N, Trabetti E, Verweij N, Evangelou E, Moayyeri A, Vergnaud AC, Nelson CP, Poveda A, Varga TV, Caslake M, de Craen AJ, Trompet S, Luan J, Scott RA, Harris SE, Liewald DC, Marioni R, Menni C, Farmaki AE, Hallmans G, Renstrom F, Huffman JE, Hassinen M, Burgess S, Vasan RS, Felix JF, Consortium CH-HF, Uria-Nickelsen M, Malarstig A, Reily DF, Hoek M, Vogt T, Lin H, Lieb W, EchoGen C, Traylor M, Markus HF, Consortium M, Highland HM, Justice AE, Marouli E, Consortium G, Lindstrom J, Uusitupa M, Komulainen P, Lakka TA, Rauramaa R, Polasek O, Rudan I, Rolandsson O, Franks PW, Dedoussis G, Spector TD, Consortium EP-I, Jousilahti P, Mannisto S, Deary IJ, Starr JM, Langenberg C, Wareham NJ, Brown MJ, Dominiczak AF, Connell JM, Jukema JW, Sattar N, Ford I, Packard CJ, Esko T, Magi R, Metspalu A, de Boer RA, van der Meer P, van der Harst P, Lifelines Cohort S, Gambaro G, Ingelsson E, Lind L, de Bakker PI, Numans ME, Brandslund I, Christensen C, Petersen ER, Korpi-Hyovalti E, Oksa H, Chambers JC, Kooner JS, Blakemore AI, Franks S, Jarvelin MR, Husemoen LL, Linneberg A, Skaaby T, Thuesen B, Karpe F, Tuomilehto J, Doney AS, Morris AD, Palmer CN, Holmen OL, Hveem K, Willer CJ, Tuomi T, Groop L, Karajamaki A, Palotie A, Ripatti S, Salomaa V, Alam DS, Shafi Majumder AA, Di Angelantonio E, Chowdhury R, McCarthy MI, Poulter N, Stanton AV, Sever P, Amouyel P, Arveiler D, Blankenberg S, Ferrieres J, Kee F, Kuulasmaa K, Muller-Nurasyid M, Veronesi G, Virtamo J, Deloukas P, Wellcome Trust Case Control C, Elliott P, Understanding Society Scientific G, Zeggini E, Kathiresan S, Melander O, Kuusisto J, Laakso M, Padmanabhan S, Porteous D, Hayward C, Scotland G, Collins FS, Mohlke KL, Hansen T, Pedersen O, Boehnke M, Stringham HM, Consortium E-C, Frossard P, Newton-Cheh C, Consortium CECBP, Tobin MD, Nordestgaard BG, Consortium TDG, Go TDC, Exome BPC, Consortium CHDE, Caulfield MJ, Mahajan A, Morris AP, Tomaszewski M, Samani NJ, Saleheen D, Asselbergs FW, Lindgren CM, Danesh J, Wain LV, Butterworth AS, Howson JM and Munroe PB. Trans-ancestry meta-

analyses identify rare and common variants associated with blood pressure and hypertension. *Nat Genet*. 2016;48:1151-1161.

63.   Emdin CA, Khera AV, Klarin D, Natarajan P, Zekavat SM, Nomura A, Haas M, Aragam K, Ardissino D, Wilson JG, Schunkert H, McPherson R, Watkins H, Elosua R, Bown MJ, Samani NJ, Baber U, Erdmann J, Gormley P, Palotie A, Stitziel NO, Gupta N, Danesh J, Saleheen D, Gabriel S and Kathiresan S. Phenotypic Consequences of a Genetic Predisposition to Enhanced Nitric Oxide Signaling. *Circulation*. 2018;137:222-232.

64.   Kessler T, Wobst J, Wolf B, Eckhold J, Vilne B, Hollstein R, von Ameln S, Dang TA, Sager HB, Moritz Rumpf P, Aherrahrou R, Kastrati A, Bjorkegren JLM, Erdmann J, Lusis AJ, Civelek M, Kaiser FJ and Schunkert H. Functional Characterization of the GUCY1A3 Coronary Artery Disease Risk Locus. *Circulation*. 2017;136:476-489.

65.   Erdmann J, Stark K, Esslinger UB, Rumpf PM, Koesling D, de Wit C, Kaiser FJ, Braunholz D, Medack A, Fischer M, Zimmermann ME, Tennstedt S, Graf E, Eck S, Aherrahrou Z, Nahrstaedt J, Willenborg C, Bruse P, Braenne I, Nothen MM, Hofmann P, Braund PS, Mergia E, Reinhard W, Burgdorf C, Schreiber S, Balmforth AJ, Hall AS, Bertram L, Steinhagen-Thiessen E, Li SC, Marz W, Reilly M, Kathiresan S, McPherson R, Walter U, CardioGram, Ott J, Samani NJ, Strom TM, Meitinger T, Hengstenberg C and Schunkert H. Dysfunctional nitric oxide signalling increases risk of myocardial infarction. *Nature*. 2013;504:432-6.

66.   Wilkinson IB, Qasem A, McEniery CM, Webb DJ, Avolio AP and Cockcroft JR. Nitric oxide regulates local arterial distensibility in vivo. *Circulation*. 2002;105:213-7.

67.   Stewart AD, Millasseau SC, Kearney MT, Ritter JM and Chowienczyk PJ. Effects of inhibition of basal nitric oxide synthesis on carotid-femoral pulse wave velocity and augmentation index in humans. *Hypertension*. 2003;42:915-8.

68.   McEniery CM, Qasem A, Schmitt M, Avolio AP, Cockcroft JR and Wilkinson IB. Endothelin-1 regulates arterial pulse wave velocity in vivo. *J Am Coll Cardiol*. 2003;42:1975-81.

69.   Li JS, Lariviere R and Schiffrin EL. Effect of a nonselective endothelin antagonist on vascular remodeling in deoxycorticosterone acetate-salt hypertensive rats. Evidence for a role of endothelin in vascular hypertrophy. *Hypertension*. 1994;24:183-8.

70.   Amiri F, Virdis A, Neves MF, Iglarz M, Seidah NG, Touyz RM, Reudelhuber TL and Schiffrin EL. Endothelium-restricted overexpression of human endothelin-1 causes vascular remodeling and endothelial dysfunction. *Circulation*. 2004;110:2233-40.

Table 1: Baseline characteristics of analyzed participants with arterial stiffness index and genotypes

Category	Phenotype†	
Demographic phenotypes	Age (Median; Q1-Q3 (N))	59; 51-63 (131,686)
	Sex (% Female)	70,847 (53.8%)
Prevalent Disease (Cases/Controls)	Prevalent Diabetes	6019/125667 (4.6%)
	Prevalent Hypertension	35639/96047 (27.1%)
	Prevalent Hypercholesterolemia	17056/114630 (12.9%)
	Prevalent Atrial Fibrillation or Atrial Flutter	1830/129856 (1.4%)
	Prevalent Heart Failure	305/131381 (0.23%)
Blood Pressure (Median; Q1-Q3 (N))	SBP	139; 127-153 (131,084)
	DBP	82; 75-89 (131,086)
Lifestyle factors & Medications N (%)	Previous or Current Smoker	57,974 (44.1%)
	Antihypertensive Medication	13,296 (10.1%)

†these values reflect the 131,686 samples with all pulse wave analysis phenotypes and genotype data present used in the genome-wide association analysis; sample outliers for quantitative phenotypes were removed as described in the methods. SBP=systolic blood pressure, DBP=diastolic blood pressure.

Figure 1: Epidemiologic and genetic associations of arterial stiffness index with blood pressure.

Association between (A) phenotypic ASI, and, (B) genotypic ASI (ie: the ASI GRS), with systolic and diastolic blood pressures in the UK Biobank. Results are presented as both unadjusted and, separately, adjusted by age, sex, smoking status, prevalent hypercholesterolemia, prevalent diabetes, heart rate, vegetable intake, alcohol intake, and exercise frequency. For the ASI GRS instrument, analysis was performed excluding individuals used in the ASI genome-wide association study. Effect estimates represent mmHg increase in blood pressure resulting from (A) 1 SD increase in ASI phenotype, and (B) 1SD increase in genetically-mediated ASI from the ASI GRS. ASI = Arterial stiffness index, DBP = diastolic blood pressure, GRS = genetic risk score, SBP = systolic blood pressure, SD = standard deviation

1 **Figure 2: Epidemiologic and genetic associations of arterial stiffness index with**
2 **incident coronary artery disease.**
3

4
Association between (A) phenotypic ASI, and, (B) the ASI GRS, with incident coronary
artery disease in the UK Biobank. Results are presented as both unadjusted (cyan) and
adjusted (purple) by age, sex, smoking status, prevalent hypertension, prevalent
hypercholesterolemia, prevalent diabetes, heart rate, vegetable intake, alcohol intake,
and exercise frequency. For the ASI GRS instrument, analysis was performed excluding
individuals used in the ASI genome-wide association study. Hazard ratios represent
increased risk of incident CAD resulting from (A) 1 SD increase in ASI phenotype, and
(B) 1SD increase in genetically-mediated ASI from the ASI GRS. Sample sizes for (A)
the phenotypic association are 3,692 cases, 126,615 controls, and for (B) the genotypic
association are 7,534 cases, 215,527 controls.
ASI = Arterial stiffness index, CAD = coronary artery disease, GRS = genetic risk score,
HR = hazard ratio, SD = standard deviation.

Figure 3: One- and two-sample Mendelian randomization analyses of arterial

stiffness index with coronary artery disease.

	CAD Effect per SD ASI GRS	Effect	95% CI	P	Cases (N)	Controls (N)
Incident CAD						
UK Biobank		HR=1.12	[0.55; 2.3]	0.75	7,534	215,527
Prevalent CAD						
CARDIOGRAMplusC4D		OR=0.56	[0.25; 1.2]	0.15	60,810	123,495

0.3   0.5   1   2   3

Association between the ASI GRS and incident CAD in the UK Biobank, as well as
prevalent CAD in the CARDIOGRAMplusC4D consortium. Incident CAD results were
derived using individual-level data from the UK Biobank and adjusting by
cardiometabolic risk factors (age, sex, smoking status, prevalent hypertension,
prevalent hypercholesterolemia, prevalent diabetes, heart rate, vegetable intake,
alcohol intake, and exercise frequency). Individuals used in the ASI genome-wide
association study were excluded in the analyses. Prevalent CAD results were derived
from summary-level genome-wide association data from the CARDIOGRAMplusC4D
consortium using robust, penalized inverse-variance weighted 2-sample Mendelian
randomization. For the ASI GRS instrument, analysis was performed excluding
individuals used in the ASI genome-wide association study.
ASI = Arterial stiffness index, CAD = coronary artery disease, GRS = genetic risk score,
HR = hazard ratio, OR = odds ratio

Figure 4: Comparison of variant level-effects with arterial stiffness index and with coronary artery disease shows inconsistency.

Variant-level effect estimates (from CARDIOGRAMplusC4D) from variants at 77
independent known CAD loci, were compared to their ASI associations. Highlighted are
5 out of the 77 variants with at least suggestive significance with ASI (*P*<0.005),
showing that ASI-raising alleles have inconsistent effects on CAD risk. The variant-level
summary statistics for these 77 variants across are detailed in **Supplementary Tables**
9-10.
ASI = arterial stiffness index, CAD = coronary artery disease