Title: Genetic Association of Arterial Stiffness Index with Blood Pressure and Coronary Artery Disease

Brief Title: Arterial Stiffness Index Genetic Analyses

Seyedeh M. Zekavat BS (1-4), Mary Haas PhD (1), Krishna Aragam MD MS (1,4,5,6), Connor Emdin DPhil (1,5), Amit V. Khera MD MSc (1,4,5,6), Derek Klarin MD (1,4), Hongyu Zhao PhD (3,7), Pradeep Natarajan MD MMSc (1,4,5,6)

1. Program in Medical and Population Genetics, Broad Institute of MIT and Harvard, Cambridge, MA;
2. Yale School of Medicine, New Haven, CT;
3. Computational Biology & Bioinformatics Program, Yale University, New Haven, CT;
4. Center for Genomic Medicine, Massachusetts General Hospital, Boston, MA;
5. Harvard Medical School, Boston, MA;
6. Cardiovascular Research Center, Massachusetts General Hospital, Boston, MA;
7. Department of Biostatistics, Yale School of Public Health, New Haven, CT

Please address correspondence to:
 Pradeep Natarajan, MD MMSc
 pnatarajan@mgh.harvard.edu
 185 Cambridge Street, CPZN 3.184, Boston, MA 02114
 Tel: 617-724-3526
 Fax: 617-726-2203

Main text word count: 3,424

Funding: S.M.Z. and P.N. had full access to all the data in the study and take responsibility for the integrity of the data and accuracy of the data analysis. S.M.Z. and P.N. conceived of and designed the study. P.N. acquired the data. S.M.Z., M.H., and K.A. analyzed the data. S.M.Z., C.E., A.V.K., D.K., H.Z., and P.N. interpreted the data. S.M.Z. and P.N. drafted the manuscript. S.M.Z. is supported by the National Institutes of Health's Medical Scientist Training Program at the Yale School of Medicine and the Paul and Daisy Soros Fellowship. P.N. is supported by a Hassenfeld Scholar Award from the Massachusetts General Hospital, and K08 HL140203 from the National Heart, Lung, and Blood Institute.

Disclosures: P.N. reports investigator-initiated grants from Amgen and Boston Scientific, and consulting income from Apple.

Acknowledgements: The authors would like to acknowledge and thank the participants and staff of the UK Biobank and of the CARDIOGramplusC4D consortium cohorts.

Abstract

Background: Arterial stiffness index (ASI) is independently associated with blood pressure and coronary artery disease (CAD) in epidemiologic studies. However, it is unknown whether these associations represent causal relationships.

Objectives: Here, we assess whether genetic predisposition to increased ASI is associated with elevated blood pressure and CAD risk.

Methods: Genome-wide association analysis (GWAS) of finger photoplethysmography-derived ASI was performed in 131,686 participants from the UK Biobank. Across UK Biobank participants not in the ASI GWAS, a 6-variant ASI polygenic risk score was calculated. The ASI polygenic score was associated with systolic and diastolic blood pressures (SBP, DBP, N=208,897), and with incident CAD over 10 years follow-up (N=223,061; 7,534 cases). The lack of CAD association observed was replicated among 184,305 participants (60,810 cases) from the Coronary Artery Disease Genetics Consortium (CARDIOGRAMplusC4D).

Results: We replicated prior reports of the epidemiologic association of ASI with SBP (Beta 0.55mmHg, [95% CI, 0.45-0.65], $P=5.77 \times 10^{-24}$), DBP (Beta 1.05mmHg, [95% CI, 0.99-1.11], $P=7.27 \times 10^{-272}$), and incident CAD (HR 1.08 [95% CI, 1.04-1.11], $P=1.5 \times 10^{-6}$) in multivariable models. While each SD increase in genetic predisposition to elevated ASI was highly associated with SBP (Beta 4.63 mmHg [95% CI, 2.1-7.2]; $P=3.37 \times 10^{-4}$), and DBP (Beta 2.61 mmHg [95% CI, 1.2-4.0]; $P=2.85 \times 10^{-4}$), no association was observed with incident CAD in UK Biobank (HR 1.12 [95% CI, 0.55-2.3]; $P=0.75$), or with prevalent CAD in CARDIOGRAMplusC4D (OR 0.56 [95% CI, 0.26-1.24]; $P=0.15$).

Conclusions: A genetic predisposition to higher ASI was associated with elevated blood pressure but not with increased risk of developing CAD.

Condensed Abstract: Arterial stiffness index (ASI) is proposed by some as a surrogate of blood pressure and coronary artery disease (CAD) risk based on epidemiologic analyses. We tested whether genetic predisposition to increased ASI is associated with elevated blood pressure and CAD risk to assess whether these represent causal relationships. We find that a genetic predisposition to higher ASI is associated with elevated systolic (Beta 4.63 mmHg [95% CI, 2.1-7.2]) and diastolic blood pressures (Beta 2.61 mmHg [95% CI, 1.2-4.0]) in the UK Biobank, but not associated with incident CAD in the UK Biobank ($P=0.75$) or with prevalent CAD in CARDIOGRAMplusC4D ($P=0.15$). These data support a causal relationship of ASI with blood pressure but do not support the notion that ASI is a suitable surrogate for CAD risk.

Keywords: Arterial Stiffness, Blood Pressure, Coronary Artery Disease, Genetic Epidemiology, Mendelian Randomization, Population Genetics

Abbreviations: ASI=arterial stiffness index, CAD=coronary artery disease, CARDIOGRAMplusC4D=Coronary Artery Disease Genetics Consortium, DBP=diastolic blood pressure, GRS=genetic risk score, PPG=photoplethysmography, SBP=systolic blood pressure

Introduction:

Arterial stiffness index (ASI), as measured non-invasively via pulse wave analysis, is independently associated with cardiovascular disease risk in multiple epidemiological studies (1-9). Increased vascular resistance and diminished viscoelasticity are key features of vascular aging which were previously associated with systolic hypertension (5), coronary artery disease (CAD) (2,4,7), and all-cause mortality (10). Arterial stiffness may be influenced by variations in collagen, elastin, smooth muscle tone, and endothelial dysfunction, in addition to other factors (11-17). Carotid-femoral (aortic) pulse wave velocity is the 'gold-standard' approach for assessing arterial stiffness. ASI measurement using finger infrared analysis is a scalable, non-invasive approach to assess ASI and is correlated with aortic pulse wave velocity (18-20).

While arterial stiffness measures are associated with cardiovascular diseases (1-8), whether the associations are causal is not clear. For example, non-causal risk factors, such as high-density lipoprotein cholesterol for CAD, are good risk predictors but are disappointing therapeutic targets (21-26). Lifestyle factors are separately linked to arterial stiffness and cardiovascular diseases, potentially confounding the observed relationships (27). Furthermore, reverse causality could lead to a statistically robust but non-causal relationship. For example, individuals with increased arterial stiffness might develop cardiovascular disease because of reduced exercise (28).

Some propose that ASI should be considered a non-invasive surrogate end point for cardiovascular events largely based on robust epidemiological associations (29-

31)·(2,32-38). Understanding whether ASI causally mediates CAD, independent of blood

pressure, may help determine whether ASI is a suitable surrogate end point for CAD

separate from its utility as a risk predictor. Mendelian randomization uses human

genetics for causal inference by leveraging the random assortment of genetic variants

during meiosis at conception, which diminishes susceptibility to confounding or reverse

causality (39). Here, we used Mendelian randomization to determine whether a genetic

predisposition to increased ASI is associated with elevated blood pressure and

increased risk for incident CAD.

Methods:

UK Biobank study participants and phenotypes

Individual-level genomic data and longitudinal phenotypic data from the UK Biobank, a

large-scale population-based dataset consisting of genotype and phenotype data in

approximately 500,000 volunteer participants collected from 2007-2017, was used.

Clinical disease definitions are detailed in **Supplementary Table 1**. In summary, the

main outcome, CAD, was defined by billing codes for heart attack, angina pectoris,

unstable angina, myocardial infarction, coronary atherosclerosis, coronary artery

revascularization, and other acute, subacute, and chronic forms of ischemic heart

disease, or with self-reported angina, heart attack/myocardial infarction, coronary

angioplasty +/- stent, or coronary artery bypass graft (CABG) surgery. We also

assessed systolic and diastolic blood pressures, and adjusted for blood pressure

medications by adding 15 and 10 mmHg to systolic and diastolic blood pressures,

respectively (40).

Arterial stiffness index measurement

ASI was previously measured in the UK Biobank using the PulseTrace PCA2 (CareFusion, San Diego, CA), which uses finger photoplethysmography over a 10- to 15-second timeframe to obtain the pulse waveform from an infrared sensor clipped to the end of the index finger. ASI (in m/s) was calculated by dividing standing height by the time between the systolic and diastolic peaks of the pulse waveform. ASI by this approach was previously correlated with aortic pulse wave velocity, which is regarded as the gold standard (18). ASI was inverse rank normalized for analysis (with mean = 0, SD = 1).

Genotyping and imputation

Genome-wide genotyping was previously performed in the UK Biobank using two genotyping arrays sharing 95% of marker content: Applied Biosystems UK BiLEVE Axiom Array (807,411 markers in 49,950 participants) and Applied Biosystems UK Biobank Axiom Array (825,927 markers in 438,427 participants) both by Affymetrix (Santa Clara, CA) (41). Variants used in the present analysis include those also imputed using the Haplotype Reference Consortium reference panel of up to 39M SNPs (42,43).

Quality control and variant annotation

Poor quality variants and genotypes were filtered as previously described (41). We further filtered out individuals from both genetic and epidemiological analyses using the following genetic criteria: non-white or not of British ancestry, gender mismatch between reported and genotypic genders, sex chromosome aneuploidy, or one from each pair of 1st or 2nd degree relatives (**Supplementary Table 2**). Non-consenting individuals with

prevalent peripheral arterial disease, aortic valve disease, or CAD were excluded, as

were extreme outliers for any of the arterial pulse wave phenotypes listed in

Supplementary Table 3. Extreme outliers were determined by adjusting the traditional

box and whisker upper and lower bounds and accounting for skewness in the

phenotypic data identified using the Robustbase package in R (setting range=3)

(https://cran.r-project.org/web/packages/robustbase/robustbase.pdf).

After filtering samples, variants were further filtered by the following criteria: not in

Hardy-Weinberg Equilibrium ($P<1\text{x}10^{-10}$), low imputation quality (INFO score < 0.3), call

rate < 95%, and minor allele frequency < 0.05% (minor allele count < 66).

Variant consequences were annotated using with Ensembl's Variant Effect Predictor

(VEP), ascribing the most severe consequence and associated gene among the

canonical transcripts present for each variant[44]. The Hail v0.1 software (https://hail.is)

was used to perform quality control and variant annotation (45).

Epidemiological association analyses with arterial stiffness index

Epidemiological association of ASI with blood pressure phenotypes and incident CAD

was performed using linear regression and Cox proportional hazards model,

respectively, in R (version 3.3, R Foundation, Vienna, Austria). For CAD, adjustment

was performed for age, sex, ever smoking status, heart rate at pulse wave analysis,

prevalent hypertension, prevalent hypercholesterolemia, prevalent diabetes, alcohol

intake (self-reported alcohol intake of at least once per month), exercise (self-reported

exercise of at least 3x per week), and vegetable intake (self-reported intake of at least 6

tablespoons of vegetable intake per day). The same adjustment variables were used for

SBP and DBP, except prevalent hypertension was not included as a covariate.

Analyses were performed using a Cox proportional hazards model for incident CAD,

and linear regression for the blood pressure traits. The threshold for significance for the

three primary phenotypes was assigned as alpha = 0.05/3 tests = 0.017.

Genome-wide association analysis of arterial stiffness

A genome-wide association of ASI was performed using individual-level data from

131,686 individuals of European descent from the UK Biobank, collected from 2007 to

2017. Each variant was individually associated with ASI in an additive linear regression

model and adjusted for sex, age, smoking status, genotyping array type, and the first

ten principal components of ancestry (46). Only variants with minor allele frequency >

0.05% (minor allele count > 66) were considered. $P < 5 \times 10^{-8}$ was considered to be

significant. The Hail software version 0.1 (https://hail.is) was used for genome-wide

association analysis (45).

Mendelian randomization

An additive genetic risk score (GRS) was calculated as $\sum_{i=1}^{m} ln(OR_i) \times SNP_{ij}$ were m is

the number of SNPs, $ln(OR_i)$ is the weight for SNP_i from the discovery sample, SNP_{ij} is

the number of alleles (i.e., 0, 1, or 2) for SNP_i in person j in the validation sample. Six

independent variants (linkage disequilibrium $r^2 < 0.25$ within 500kb windows)

demonstrating at least suggestive association with ASI ($P<5 \times 10^{-7}$) were included in the

GRS. The raw GRS was calculated for each individual using PLINK (47), inverse rank

normalized, then re-scaled such that one unit increase in the GRS was equivalent to a

one standard deviation (SD) increase in ASI.

To confirm that the GRS for ASI was a strong instrument for ASI, an F-statistic for the

instrument was calculated in the UK Biobank. An F-statistic is a measure of the

significance of an instrument (the GRS) for prediction of the exposure (ASI), controlling

for additional covariates (age, sex, ever smoked, 10 principal components of ancestry,

and genotyping array type). An F-statistic greater than 10 is evidence of a strong

instrument. Furthermore, sensitivity analyses were performed to evaluate for

associations between the ASI GRS and potential environmental confounders including

sex, ever smoking status, diet (alcohol intake, vegetable intake), and exercise frequency

among individuals not in the ASI genome-wide association analyses.

A linear regression model was used to associate the ASI GRS with systolic and diastolic

blood pressures. A Cox proportional hazards model was used to associate ASI GRS

with incident CAD. For CAD, adjustment was performed for age, sex, ever smoking

status, heart rate at blood pressure measurement, prevalent hypertension, prevalent

hypercholesterolemia, prevalent diabetes, alcohol intake (self-reported alcohol intake of

at least once per month), exercise (self-reported exercise of at least 3x per week), and

vegetable intake (self-reported intake of at least 6 tablespoons of vegetable intake per

day), where indicated. The same adjustment variables were used for SBP and DBP,

except for prevalent hypertension.

2-Sample Mendelian randomization with coronary artery disease

To address potential power limitations from the lack of association between ASI and

CAD, we also pursued 2-sample Mendelian randomization using variant-level summary

statistics from prior genome-wide association analyses of CAD from several

independent case-control studies, specifically 184,305 individuals from the Coronary

Artery Disease Genetics Consortium (CARDIOGRAMplusC4D) (48). The effect

estimates and standard errors for the six GRS variants for ASI (from UK Biobank) and

for CAD (from CARDIOGramplusC4D) were used to perform robust, penalized inverse

variance weighted (IVW) 2-sample Mendelian randomization using the

MendelianRandomization package in R (49,50). IVW 2-sample Mendelian

randomization uses a weighted linear regression of the ratio of the SNP effects on the

outcomes to the SNP effects on the risk factor, without using an intercept term. The

threshold for significance was defined as alpha = 0.05.

Additionally, analyses were performed to evaluate the reverse association, of CAD

causally impacting ASI. 77 known, independent, genome-wide significant CAD locus

variants were identified across several published sources (48,51-53) (**Supplementary**

Table 9). These 77 CAD locus variants were used as an instrument in 2-sample

Mendelian randomization to evaluate whether CAD causally affects ASI.

<u>**Results:**</u>

Baseline characteristics

A total of 131,686 individuals in the UK Biobank had ASI measured, genotype data available, and passed quality control (**Supplementary Table 2**). Among these individuals, median age was 59 (IQR 51-63) years, 53.8% were female, 4.6% had diabetes, 27.1% had hypertension, and 12.9% had hypercholesterolemia. Median SBP was 139 (IQR 127-153) mmHg, median DBP was 82 (IQR 75-89) mmHg. 44.1% of individuals were prior or current smokers, and 10.1% of individuals were on antihypertensive medications (**Table 1**). The median ASI was 9 (IQR 7-11) m/s (**Supplementary Table 3**).

Epidemiological associations of ASI

Univariate association of cardiovascular risk factors with ASI showed the following associations with at least nominal significance ($P<0.05$): for age (0.024 SD/year, $P<1 \times 10^{-300}$), sex (0.40 SD higher in males, $P<1 \times 10^{-300}$), blood pressure medication (0.34 SD, $P=1.4 \times 10^{-317}$), hypertension (0.21 SD, $P=1.4 \times 10^{-269}$), hypercholesterolemia (0.20 SD, $P=4.1 \times 10^{-137}$), diabetes (0.20 SD, $P=9.1 \times 10^{-54}$), ever smoking (0.18 SD, $P=3.0 \times 10^{-250}$), exercise $\geq$3x/wk (-0.16 SD, $P=2.9 \times 10^{-66}$), alcohol intake $\geq$1x/mo (0.05 SD, $P=3.3 \times 10^{-20}$), and $\geq$6 tablespoons vegetable intake per day (-0.063 SD, $P=3.1 \times 10^{-4}$) (**Supplementary Table 4**).

For the associations of ASI with SBP and DBP, both univariable and multivariable, adjusting for age, sex, smoking status, prevalent hypercholesterolemia, prevalent diabetes, vegetable intake, alcohol intake, and exercise, analyses showed strong associations (**Figure 1A**). Each SD of ASI was associated with elevated SBP by 0.55

mmHg ([95% CI, 0.45-0.65], P=5.77x10^{-24}) and DBP by 1.05 mmHg ([95% CI, 0.99-

1.11], P=7.27x10^{-272}).

ASI was also significantly independently associated with incident CAD, adjusting for

age, sex, ever smoking status, heart rate, prevalent hypertension, prevalent

hypercholesterolemia, prevalent diabetes, vegetable intake, alcohol intake, and exercise

(HR 1.08 per SD ASI [95% CI, 1.04-1.11], P=7.67x10^{-6}) (**Figure 2A**).

Genome-wide association analysis of ASI

A genome-wide association analysis of ASI was performed among 131,686 individuals

and 13,995,214 variants in the UK Biobank. A quantile-quantile plot of the genome-wide

association statistics did not show substantial genomic inflation ($\lambda = 1.05$)

(**Supplementary Figure 1**). Two genome-wide significant loci were identified (P<5x10^{-}

^{8}), the top variants of which were in second intron of *TEX41* (rs1006923, -0.025 SD,

P=3.7x10^{-10}, minor allele frequency (MAF)=0.32), and first intron of *FOXO1* (rs7331212,

-0.024 SD, P=9.3x10^{-9}, MAF=0.26). Three additional suggestive loci (P<5x10^{-7}) were

also identified, of which the top variants are intronic variants in *COL4A2* (rs872588, -

0.020 SD, P=2.3x10^{-7}, MAF=0.42), *RNF126* (rs1009628, -0.027 SD, P=1.2x10^{-7},

MAF=0.15), and *TCF20* (rs55906806, -0.024 SD, P=2.4x10^{-7}, MAF=0.20). Through

chromatin conformational changes, intronic variants at *TEX41* and *COL4A2* may

influence gene expression at nearby enhancers **Supplementary Results,**

Supplementary Figure 2). Interrogation of disruptive protein-coding variants yielded

moderate association for *HFE* p.Cys282Tyr (MAF 0.076), the most common variant

implicated in hereditary hemochromatosis (**Supplementary Results**, **Supplementary**

Table 4).

Mendelian randomization in the UK Biobank

Six independent and at least suggestive ($P<5\times10^{-7}$) variants were used towards an ASI

genetic risk score (GRS) (**Supplementary Table 5**). The raw ASI GRS was associated

with a 0.85 SD increase in ASI (SE: 0.072; $P=8.0\times10^{-32}$). The F-statistic of the GRS was

123 (recommended F-statistic > 10), suggesting high instrument strength. The inverse-

rank normalized GRS was re-scaled such that each unit reflected one SD in ASI for

comparison with the phenotypic associations (**Supplementary Figure 3**). Sensitivity

analysis was performed to evaluate for associations between the ASI GRS and potential

environmental confounders including sex, ever smoking status, diet (alcohol intake,

vegetable intake), and exercise frequency. No significant associations between the ASI

GRS and environmental confounders were observed (**Supplementary Table 6**).

A 1-SD increase in genetically-mediated ASI was significantly associated with elevated

SBP (Beta 4.63 mmHg [95% CI, 2.1-7.2]; $P=3.37\times10^{-4}$), and DBP (Beta 2.61 mmHg

[95% CI, 1.2-4.0]; $P=2.85\times10^{-4}$), independent of cardiometabolic risk factors (age, sex,

and smoking status, prevalent hypercholesterolemia, prevalent diabetes, heart rate,

vegetable intake, alcohol intake, and exercise frequency) (**Figure 1B**).

The ASI GRS, however, was not associated with incident CAD in UK Biobank in an

unadjusted model (HR 1.3 [95% CI, 0.64-2.6]; $P=0.47$) or an adjusted model including

age, sex, smoking status, prevalent hypertension, prevalent hypercholesterolemia,

prevalent diabetes, heart rate, vegetable intake, alcohol intake, and exercise frequency

as covariates (HR 1.12 [95% CI, 0.55-2.3]; *P*=0.75) (**Figure 2B**).

2-Sample Mendelian randomization with coronary artery disease

To address potential power limitations impeding association of ASI GRS with incident

CAD in the UK Biobank, we also pursued 2-sample Mendelian randomization between

ASI and prevalent CAD using variant-level summary statistics from 184,305 separate

individuals in the Coronary Artery Disease Genetics Consortium

(CARDIOGRAMplusC4D) (48). Robust, penalized inverse-variance weighted 2-sample

Mendelian randomization similarly did not demonstrate an association between

genetically-elevated ASI and CAD (OR 0.56 [95% CI, 0.26-1.24], *P*=0.15) (**Figure 3**).

Furthermore, the six variants showing suggestive association with ASI did not

demonstrate a significant positive association with CAD across several different 2-

sample Mendelian randomization methods (**Supplementary Table 8**).

We also developed an expanded ASI polygenic score using 321 independent variants

(P<1x10^{-4}, LD r^2< 0.25) to capture additional genetic variation of ASI. The expanded ASI

polygenic score explained 3.3% of ASI variance conferring >80% power to detect the

CAD effect estimate observed in epidemiologic analyses (i.e., OR=1.08) with alpha =

0.05. With this approach, we again confirmed no significant association in inverse-

variance weighted 2-sample Mendelian randomization (OR 0.95 [95% CI, 0.89-1.02],

P=0.13).

77 genome-wide significant CAD loci from prior GWAS (48,52,53) were identified, and

CAD risk effect estimates prior studies and ASI effect estimates from this study were

catalogued (**Figure 4**). While 3 of 77 previously-associated CAD loci showed evidence

of association with ASI ($P<0.05/77=6.5 \times 10^{-4}$), effect directions were inconsistent

between ASI and CAD. For example, the variant rs9349379-A, an intronic variant in

PHACTR1, was associated with increased ASI (0.015 SD, $P=4.5 \times 10^{-5}$) but decreased

risk for CAD (OR= 0.87, $P=1.8 \times 10^{-42}$). Similarly, ASI-raising alleles at the *ZEB2-TEX41*

and *ABO* loci decrease CAD risk, while ASI-raising alleles at *CYP17A1-CNNM2-NT4C2*

and *SH2B3* increase CAD risk. Detailed variant-level summary statistics for these 77

CAD locus variants are provided in **Supplementary Tables 9-10**. These 77 CAD locus

variants were also used as an instrument in 2-sample Mendelian randomization for a

putative reverse association – whether a genetic susceptibility to CAD increases ASI.

No significant associations were observed across various 2-sample Mendelian

randomization methods for the reverse association (**Supplementary Table 11**).

Discussion:

We performed the largest genome-wide association analysis to-date of a measure of

vascular aging, ASI, in 131,686 individuals, and leveraged these observations to

perform causal inference analyses with blood pressure and risk of CAD in up to 407,366

separate individuals. In our genome-wide association analyses, we discover the first

genome-wide variants associated with ASI. We replicate the epidemiologic associations

of ASI with blood pressure and CAD, and find that genetic analyses do indeed support a

causal relationship between ASI and blood pressure. However, our genetic analyses do

not support a causal relationship between ASI and CAD.

These results permit several conclusions. First, we observe strong epidemiologic and

genetic association between ASI and blood pressure. These data indicate that non-

invasive photoplethysmography, employed by a finger probe or potentially

commercially-available wearable monitors that measure heart rate (54), may be used to

impute continuous blood pressure, and that changes will track with blood pressure

changes. However, given independent clinical effects and imperfect correlation, ASI

measurement may complement blood pressure assessments. Second, there is a long-

standing debate whether ASI precedes elevated blood pressure or vice versa (55).

Compared to its phenotypic effect, the effect conferred by genetically-elevated ASI is

8.4-fold higher for SBP (4.63 mmHg for ASI GRS versus 0.55 mmHg for ASI phenotype)

and 2.5-fold higher for DBP (2.61 mmHg for ASI GRS versus 1.05 mmHg for ASI

phenotype), potentially representing the effects of life-long exposure to elevated arterial

stiffness on blood pressure. This supports the notion that arterial stiffness may predate

the onset of elevated blood pressure indicating that ASI may identify individuals at

heightened risk for future blood pressure elevations.

Third, our epidemiological and genetic analyses indicate that ASI is an independent,

non-causal risk factor for CAD. Arterial stiffness may be a parallel disrupted pathway in

the setting of CAD, as opposed to an upstream causal mediating factor. Thus, while ASI

monitoring may still serve as a good proxy for blood pressure, therapeutic modulation of

ASI in isolation may not have a meaningful impact on CAD outcomes. Similarly, a

recent study of twins showed that while carotid-femoral pulse wave velocity was

heritable, it did not associate with 5-year progression of carotid intima media thickness

(56). The lack of significance between genetically-elevated ASI and CAD is also

consistent with prior mixed results in experimental models. Fragmentation of elastin

fibers and deposition of collagen fibers are features of vascular aging implicated in

arterial stiffness (57). However, murine models lacking elastin do not have endothelial

damage, thrombosis, or inflammation which typically occur with atherosclerosis (58).

Furthermore, we found that while few variants associated with CAD show apparent

association with ASI, our data indicate that ASI may not be mediating the apparent CAD

risk. We observed generally inconsistent genetic effects between ASI and CAD risk. In

particular, an intronic variant within *PHACTR1* (rs9349379-A), which was recently

shown to influence endothelin-1 expression in the vasculature, is associated with

decreased risk for CAD (59), increased blood pressure (60), and increased ASI. For this

variant, the divergent directionalities of effect on CAD and blood pressure may be due

to the differential expression of *EDNRA* versus *EDNRB* in the coronary arteries

compared to peripheral vasculature (59). Additionally, genetic variants disrupting nitric

oxide signaling at the *NOS3* and *GUCY13* loci influence both blood pressure and risk of

CAD (61-63). Notably, in our study, risk variants at these loci were not strongly

associated with ASI. Extensive prior experimental work linked nitric oxide signaling and

endothelin-1 with endothelial function and vascular tone (64-68). Our data suggests

that increased risk of CAD through these pathways is unlikely to be through changes in

photoplethymsography-detected ASI but potentially through alternative vascular

mechanisms.

While our study has several strengths, some limitations should be considered. First,

lack of ASI genetic risk score association with CAD may be due to limited statistical

power. Our replication of the lack of association using 2-sample Mendelian

randomization including with an expanded polygenic score, combined with our analysis

showing inconsistent effects of individual variants between CAD and ASI suggests that

this is less likely. Second, our imputation of untreated blood pressure among those with

prescribed hypertensives assumes a homogenous blood pressure effect across the

population. Without prescription data in the UK Biobank, we are unable to account for

different medication regimens and adherence.

<u>Conclusion:</u>

A genetic predisposition to higher ASI was associated with increased blood pressure,

but not increased risk of CAD. Our data support the conclusion that finger

photoplethysmography-derived ASI is an independent, causal risk factor for blood

pressure and an independent, non-causal risk factor for CAD.

Clinical Perspectives:

Core Clinical Competencies: A genetic predisposition to higher ASI was associated with elevated blood pressure, but not with elevated risk for CAD.

Translational Outlook: Further research should be conducted to determine whether photoplethysmography-derived ASI may be used in wearables as a continuous proxy for blood pressure phenotypes for prevention and monitoring.

Translational Outlook: Further research is required to understand whether the novel genes identified and implicated in ASI are suitable novel targets for blood pressure-lowering.

References:

1. Ikonomidis I, Ntai K, Kadoglou NP et al. The evaluation of pulse wave velocity using Arteriograph and Complior apparatus across multiple cohorts of cardiovascular-related diseases. Int J Cardiol 2013;168:4890-2.
2. Mattace-Raso FU, van der Cammen TJ, Hofman A et al. Arterial stiffness and risk of coronary heart disease and stroke: the Rotterdam Study. Circulation 2006;113:657-63.
3. Ben-Shlomo Y, Spears M, Boustred C et al. Aortic pulse wave velocity improves cardiovascular event prediction: an individual participant meta-analysis of prospective observational data from 17,635 subjects. J Am Coll Cardiol 2014;63:636-46.
4. Ikonomidis I, Makavos G, Lekakis J. Arterial stiffness and coronary artery disease. Curr Opin Cardiol 2015;30:422-31.
5. Franklin SS, Gustin Wt, Wong ND et al. Hemodynamic patterns of age-related changes in blood pressure. The Framingham Heart Study. Circulation 1997;96:308-15.
6. Chirinos JA, Kips JG, Jacobs DR, Jr. et al. Arterial wave reflections and incident cardiovascular events and heart failure: MESA (Multiethnic Study of Atherosclerosis). J Am Coll Cardiol 2012;60:2170-7.
7. Sutton-Tyrrell K, Najjar SS, Boudreau RM et al. Elevated aortic pulse wave velocity, a marker of arterial stiffness, predicts cardiovascular events in well-functioning older adults. Circulation 2005;111:3384-90.
8. Tsao CW, Lyass A, Larson MG et al. Relation of Central Arterial Stiffness to Incident Heart Failure in the Community. J Am Heart Assoc 2015;4.
9. Said MA, Eppinga RN, Lipsic E, Verweij N, van der Harst P. Relationship of Arterial Stiffness Index and Pulse Pressure With Cardiovascular Disease and Mortality. J Am Heart Assoc 2018;7.
10. Vlachopoulos C, Aznaouridis K, Stefanadis C. Prediction of cardiovascular events and all-cause mortality with arterial stiffness: a systematic review and meta-analysis. J Am Coll Cardiol 2010;55:1318-27.
11. Bank AJ, Kaiser DR. Smooth muscle relaxation: effects on arterial compliance, distensibility, elastic modulus, and pulse wave velocity. Hypertension 1998;32:356-9.
12. Anderson TJ. Arterial stiffness or endothelial dysfunction as a surrogate marker of vascular risk. Can J Cardiol 2006;22 Suppl B:72B-80B.
13. Ikonomidis I, Lekakis J, Papadopoulos C et al. Incremental value of pulse wave velocity in the determination of coronary microcirculatory dysfunction in never-treated patients with essential hypertension. Am J Hypertens 2008;21:806-13.
14. McNulty M, Mahmud A, Spiers P, Feely J. Collagen type-I degradation is related to arterial stiffness in hypertensive and normotensive subjects. J Hum Hypertens 2006;20:867-73.
15. Nigam A, Mitchell GF, Lambert J, Tardif JC. Relation between conduit vessel stiffness (assessed by tonometry) and endothelial function (assessed by flow-mediated dilatation) in patients with and without coronary heart disease. Am J Cardiol 2003;92:395-9.

16. Wagenseil JE, Mecham RP. Elastin in large artery stiffness and hypertension. J Cardiovasc Transl Res 2012;5:264-73.

17. Zieman SJ, Melenovsky V, Kass DA. Mechanisms, pathophysiology, and therapy of arterial stiffness. Arterioscler Thromb Vasc Biol 2005;25:932-43.

18. Woodman RJ, Kingwell BA, Beilin LJ, Hamilton SE, Dart AM, Watts GF. Assessment of central and peripheral arterial stiffness: studies indicating the need to use a combination of techniques. Am J Hypertens 2005;18:249-60.

19. Millasseau SC, Kelly RP, Ritter JM, Chowienczyk PJ. Determination of age-related increases in large artery stiffness by digital pulse contour analysis. Clin Sci (Lond) 2002;103:371-7.

20. Alty SR, Angarita-Jaimes N, Millasseau SC, Chowienczyk PJ. Predicting arterial stiffness from the digital volume pulse waveform. IEEE Trans Biomed Eng 2007;54:2268-75.

21. Brousseau ME, Schaefer EJ, Wolfe ML et al. Effects of an inhibitor of cholesteryl ester transfer protein on HDL cholesterol. N Engl J Med 2004;350:1505-15.

22. Emerging Risk Factors C, Di Angelantonio E, Sarwar N et al. Major lipids, apolipoproteins, and risk of vascular disease. JAMA 2009;302:1993-2000.

23. Group HTC, Landray MJ, Haynes R et al. Effects of extended-release niacin with laropiprant in high-risk patients. N Engl J Med 2014;371:203-12.

24. Investigators A-H, Boden WE, Probstfield JL et al. Niacin in patients with low HDL cholesterol levels receiving intensive statin therapy. N Engl J Med 2011;365:2255-67.

25. Schwartz GG, Olsson AG, Abt M et al. Effects of dalcetrapib in patients with a recent acute coronary syndrome. N Engl J Med 2012;367:2089-99.

26. Voight BF, Peloso GM, Orho-Melander M et al. Plasma HDL cholesterol and risk of myocardial infarction: a mendelian randomisation study. Lancet 2012;380:572-80.

27. Tanaka H, Safar ME. Influence of lifestyle modification on arterial stiffness and wave reflections. Am J Hypertens 2005;18:137-44.

28. Sacre JW, Jennings GL, Kingwell BA. Exercise and dietary influences on arterial stiffness in cardiometabolic disease. Hypertension 2014;63:888-93.

29. Laurent S, Briet M, Boutouyrie P. Arterial stiffness as surrogate end point: needed clinical trials. Hypertension 2012;60:518-22.

30. Laurent S. Arterial stiffness: intermediate or surrogate endpoint for cardiovascular events? Eur Heart J 2005;26:1152-4.

31. Cohn JN, Quyyumi AA, Hollenberg NK, Jamerson KA. Surrogate markers for cardiovascular disease: functional markers. Circulation 2004;109:IV31-46.

32. Blacher J, Guerin AP, Pannier B, Marchais SJ, Safar ME, London GM. Impact of aortic stiffness on survival in end-stage renal disease. Circulation 1999;99:2434-9.

33. Laurent S, Boutouyrie P, Asmar R et al. Aortic stiffness is an independent predictor of all-cause and cardiovascular mortality in hypertensive patients. Hypertension 2001;37:1236-41.

34. London GM, Blacher J, Pannier B, Guerin AP, Marchais SJ, Safar ME. Arterial wave reflections and survival in end-stage renal failure. Hypertension 2001;38:434-8.

35. Mitchell GF, Moye LA, Braunwald E et al. Sphygmomanometrically determined pulse pressure is a powerful independent predictor of recurrent events after myocardial infarction in patients with impaired left ventricular function. SAVE investigators. Survival and Ventricular Enlargement. Circulation 1997;96:4254-60.

36. Safar ME. Systolic blood pressure, pulse pressure and arterial stiffness as cardiovascular risk factors. Curr Opin Nephrol Hypertens 2001;10:257-61.

37. van Popele NM, Grobbee DE, Bots ML et al. Association between arterial stiffness and atherosclerosis: the Rotterdam Study. Stroke 2001;32:454-60.

38. Weber T, Auer J, O'Rourke MF et al. Arterial stiffness, wave reflections, and the risk of coronary artery disease. Circulation 2004;109:184-9.

39. Smith GD, Ebrahim S. 'Mendelian randomization': can genetic epidemiology contribute to understanding environmental determinants of disease? Int J Epidemiol 2003;32:1-22.

40. Warren HR, Evangelou E, Cabrera CP et al. Genome-wide association analysis identifies novel blood pressure loci and offers biological insights into cardiovascular risk. Nat Genet 2017;49:403-415.

41. Bycroft C, Freeman C, Petkova D et al. The UK Biobank resource with deep phenotyping and genomic data. Nature 2018;562:203-209.

42. McCarthy S, Das S, Kretzschmar W et al. A reference panel of 64,976 haplotypes for genotype imputation. Nat Genet 2016;48:1279-83.

43. Bycroft C, Freeman C, Petkova D et al. Genome-wide genetic data on ~500,000 UK Biobank participants. bioRxiv 2017.

44. McLaren W, Pritchard B, Rios D, Chen Y, Flicek P, Cunningham F. Deriving the consequences of genomic variants with the Ensembl API and SNP Effect Predictor. Bioinformatics 2010;26:2069-70.

45. Ganna A, Genovese G, Howrigan DP et al. Ultra-rare disruptive and damaging mutations influence educational attainment in the general population. Nat Neurosci 2016;19:1563-1565.

46. Hoggart CJ, Parra EJ, Shriver MD et al. Control of confounding of genetic associations in stratified populations. Am J Hum Genet 2003;72:1492-1504.

47. Purcell S, Neale B, Todd-Brown K et al. PLINK: a tool set for whole-genome association and population-based linkage analyses. Am J Hum Genet 2007;81:559-75.

48. Nikpay M, Goel A, Won HH et al. A comprehensive 1,000 Genomes-based genome-wide association meta-analysis of coronary artery disease. Nat Genet 2015;47:1121-1130.

49. Yavorska OO, Burgess S. MendelianRandomization: an R package for performing Mendelian randomization analyses using summarized data. Int J Epidemiol 2017;46:1734-1739.

50. Burgess S, Bowden J, Dudbridge F, Thompson SG. Robust instrumental variable methods using multiple candidate instruments with application to Mendelian randomization. ArXiv e-prints, 2016.

51. Webb TR, Erdmann J, Stirrups KE et al. Systematic Evaluation of Pleiotropy Identifies 6 Further Loci Associated With Coronary Artery Disease. J Am Coll Cardiol 2017;69:823-836.

52. Klarin D, Zhu QM, Emdin CA et al. Genetic analysis in UK Biobank links insulin resistance and transendothelial migration pathways to coronary artery disease. Nat Genet 2017;49:1392-1397.

53. Consortium CAD, Deloukas P, Kanoni S et al. Large-scale association analysis identifies new risk loci for coronary artery disease. Nat Genet 2013;45:25-33.

54. Georgiou K, Larentzakis AV, Khamis NN, Alsuhaibani GI, Alaska YA, Giallafos EJ. Can Wearable Devices Accurately Measure Heart Rate Variability? A Systematic Review. Folia Med (Plovdiv) 2018;60:7-20.

55. Mitchell GF. Arterial stiffness and hypertension: chicken or egg? Hypertension 2014;64:210-4.

56. Cecelja M, Jiang B, Keehn L et al. Arterial stiffening is a heritable trait associated with arterial dilation but not wall thickening: a longitudinal study in the twins UK cohort. Eur Heart J 2018;39:2282-2288.

57. Xu J, Shi GP. Vascular wall extracellular matrix proteins and vascular diseases. Biochim Biophys Acta 2014;1842:2106-2119.

58. Li DY, Brooke B, Davis EC et al. Elastin is an essential determinant of arterial morphogenesis. Nature 1998;393:276-80.

59. Gupta RM, Hadaya J, Trehan A et al. A Genetic Variant Associated with Five Vascular Diseases Is a Distal Regulator of Endothelin-1 Gene Expression. Cell 2017;170:522-533 e15.

60. Surendran P, Drenos F, Young R et al. Trans-ancestry meta-analyses identify rare and common variants associated with blood pressure and hypertension. Nat Genet 2016;48:1151-1161.

61. Emdin CA, Khera AV, Klarin D et al. Phenotypic Consequences of a Genetic Predisposition to Enhanced Nitric Oxide Signaling. Circulation 2018;137:222-232.

62. Kessler T, Wobst J, Wolf B et al. Functional Characterization of the GUCY1A3 Coronary Artery Disease Risk Locus. Circulation 2017;136:476-489.

63. Erdmann J, Stark K, Esslinger UB et al. Dysfunctional nitric oxide signalling increases risk of myocardial infarction. Nature 2013;504:432-6.

64. Wilkinson IB, Qasem A, McEniery CM, Webb DJ, Avolio AP, Cockcroft JR. Nitric oxide regulates local arterial distensibility in vivo. Circulation 2002;105:213-7.

65. Stewart AD, Millasseau SC, Kearney MT, Ritter JM, Chowienczyk PJ. Effects of inhibition of basal nitric oxide synthesis on carotid-femoral pulse wave velocity and augmentation index in humans. Hypertension 2003;42:915-8.

66. McEniery CM, Qasem A, Schmitt M, Avolio AP, Cockcroft JR, Wilkinson IB. Endothelin-1 regulates arterial pulse wave velocity in vivo. J Am Coll Cardiol 2003;42:1975-81.

67. Li JS, Lariviere R, Schiffrin EL. Effect of a nonselective endothelin antagonist on vascular remodeling in deoxycorticosterone acetate-salt hypertensive rats. Evidence for a role of endothelin in vascular hypertrophy. Hypertension 1994;24:183-8.

68. Amiri F, Virdis A, Neves MF et al. Endothelium-restricted overexpression of human endothelin-1 causes vascular remodeling and endothelial dysfunction. Circulation 2004;110:2233-40.

Central Illustration: Epidemiologic and genetic associations of arterial stiffness
with blood pressure and coronary artery disease.
Association between phenotypic ASI, and separately, genotypic ASI (from the ASI
GRS), with systolic and diastolic blood pressures, as well as with incident and prevalent
CAD. Blood pressure results are adjusted by age, sex, smoking status, prevalent
hypercholesterolemia, prevalent diabetes, heart rate, vegetable intake, alcohol intake,
and exercise frequency. Blood pressure effect estimates represent mmHg increase in
blood pressure resulting from (A) 1 SD increase in ASI phenotype, and (B) 1SD
increase in genetically-mediated ASI from the ASI GRS.
Incident CAD associations in the UK Biobank, as well as prevalent CAD associations
using the CARDIOGRAMplusC4D consortium, with phenotypic ASI and the ASI GRS
are provided. Incident CAD results were derived using individual-level data from the UK
Biobank and adjusting by cardiometabolic risk factors (age, sex, smoking status,
prevalent hypertension, prevalent hypercholesterolemia, prevalent diabetes, heart rate,
vegetable intake, alcohol intake, and exercise frequency). Prevalent CAD results were
derived from summary-level genome-wide association data from the
CARDIOGRAMplusC4D consortium using robust, penalized inverse-variance weighted
2-sample Mendelian randomization.
ASI = Arterial stiffness index, CAD = coronary artery disease, DBP = diastolic blood
pressure, GRS = genetic risk score, SBP = systolic blood pressure, SD = standard
deviation
22
23
Figure 1: Epidemiologic and genetic associations of arterial stiffness index with
blood pressure.
Association between (A) phenotypic ASI, and, (B) genotypic ASI (ie: the ASI GRS), with
systolic and diastolic blood pressures in the UK Biobank. Results are presented as both
unadjusted and, separately, adjusted by age, sex, smoking status, prevalent
hypercholesterolemia, prevalent diabetes, heart rate, vegetable intake, alcohol intake,
and exercise frequency. For the ASI GRS instrument, analysis was performed excluding
individuals used in the ASI genome-wide association study. Effect estimates represent
mmHg increase in blood pressure resulting from (A) 1 SD increase in ASI phenotype,
and (B) 1SD increase in genetically-mediated ASI from the ASI GRS.
ASI = Arterial stiffness index, DBP = diastolic blood pressure, GRS = genetic risk score,
SBP = systolic blood pressure, SD = standard deviation
36
37 **Figure 2: Epidemiologic and genetic associations of arterial stiffness index with**
38 **incident coronary artery disease.**
39 Association between (A) phenotypic ASI, and, (B) the ASI GRS, with incident coronary
40 artery disease in the UK Biobank. Results are presented as both unadjusted (cyan) and

adjusted (purple) by age, sex, smoking status, prevalent hypertension, prevalent hypercholesterolemia, prevalent diabetes, heart rate, vegetable intake, alcohol intake, and exercise frequency. For the ASI GRS instrument, analysis was performed excluding individuals used in the ASI genome-wide association study. Hazard ratios represent increased risk of incident CAD resulting from (A) 1 SD increase in ASI phenotype, and (B) 1SD increase in genetically-mediated ASI from the ASI GRS. Sample sizes for (A) the phenotypic association are 3,692 cases, 126,615 controls, and for (B) the genotypic association are 7,534 cases, 215,527 controls.

ASI = Arterial stiffness index, CAD = coronary artery disease, GRS = genetic risk score, HR = hazard ratio, SD = standard deviation.

Figure 3: One- and two-sample Mendelian randomization analyses of arterial stiffness index with coronary artery disease.

Association between the ASI GRS and incident CAD in the UK Biobank, as well as prevalent CAD in the CARDIOGRAMplusC4D consortium. Incident CAD results were derived using individual-level data from the UK Biobank and adjusting by cardiometabolic risk factors (age, sex, smoking status, prevalent hypertension, prevalent hypercholesterolemia, prevalent diabetes, heart rate, vegetable intake, alcohol intake, and exercise frequency). Individuals used in the ASI genome-wide association study were excluded in the analyses. Prevalent CAD results were derived from summary-level genome-wide association data from the CARDIOGRAMplusC4D consortium using robust, penalized inverse-variance weighted 2-sample Mendelian randomization. For the ASI GRS instrument, analysis was performed excluding individuals used in the ASI genome-wide association study.

ASI = Arterial stiffness index, CAD = coronary artery disease, GRS = genetic risk score, HR = hazard ratio, OR = odds ratio

Figure 4: Comparison of variant level-effects with arterial stiffness index and with coronary artery disease shows inconsistency.

Variant-level effect estimates (from CARDIOGRAMplusC4D) from variants at 77 independent known CAD loci, were compared to their ASI associations. Highlighted are 5 out of the 77 variants with at least suggestive significance with ASI (*P*<0.005), showing that ASI-raising alleles have inconsistent effects on CAD risk. The variant-level summary statistics for these 77 variants across are detailed in **Supplementary Tables 9-10**.

ASI = arterial stiffness index, CAD = coronary artery disease

Table 1: Baseline characteristics of analyzed participants with arterial stiffness index and genotypes

Category	Phenotype†	
Demographic phenotypes	Age (Median; Q1-Q3 (N))	59; 51-63 (131,686)
	Sex (% Female)	70,847 (53.8%)
Prevalent Disease (Cases/Controls)	Prevalent Diabetes	6019/125667 (4.6%)
	Prevalent Hypertension	35639/96047 (27.1%)
	Prevalent Hypercholesterolemia	17056/114630 (12.9%)
	Prevalent Atrial Fibrillation or Atrial Flutter	1830/129856 (1.4%)
	Prevalent Heart Failure	305/131381 (0.23%)
Blood Pressure (Median; Q1-Q3 (N))	SBP	139; 127-153 (131,084)
	DBP	82; 75-89 (131,086)
Lifestyle factors & Medications N (%)	Previous or Current Smoker	57,974 (44.1%)
	Antihypertensive Medication	13,296 (10.1%)

†these values reflect the 131,686 samples with all pulse wave analysis phenotypes and genotype data present used in the genome-wide association analysis; sample outliers for quantitative phenotypes were removed as described in the methods. SBP=systolic blood pressure, DBP=diastolic blood pressure.

Arterial Stiffness Index

Systolic Blood Pressure

ASI Instrument:	Beta	95% CI
Phenotype	0.55	[0.45; 0.65]
Genetic	4.63	[2.10; 7.16]

0   2   4   6   8

mmHg per SD ASI

Diastolic Blood Pressure

ASI Instrument:	Beta	95% CI
Phenotype	1.05	[0.99; 1.1]
Genetic	2.61	[1.20; 4.0]

0   2   4   6   8

mmHg per SD ASI

Coronary Artery Disease

ASI Instrument:		95% CI
UK Biobank		
Phenotype	HR=1.08	[1.04; 1.1]
Genetic	HR=1.12	[0.55; 2.3]
CARDIOGRAMplusC4D		
Genetic	OR=0.56	[0.26; 1.2]

0.3 0.5   1   2   3

risk per SD ASI

A.

Effect of Phenotypic ASI on Blood Pressure
(mmHg per SD ASI Phenotype) **Beta** **95% CI** **P** **N**

SBP
| Unadjusted | 2.96 | [2.84; 3.08] | <1e−300 | 137,858 |
| Adjusted | 0.55 | [0.45; 0.65] | 5.77e−24 | 137,858 |

DBP
| Unadjusted | 2.07 | [2.01; 2.13] | <1e−300 | 137,862 |
| Adjusted | 1.05 | [0.99; 1.11] | 7.27e−272 | 137,862 |

0 2 4 6 8 10

B.

Effect of Genotypic ASI on Blood Pressure
(mmHg per SD ASI GRS) **Beta** **95% CI** **P** **N**

SBP
| Unadjusted | 6.37 | [3.6; 9.2] | 8.13e−06 | 208,897 |
| Adjusted | 4.63 | [2.1; 7.2] | 3.37e−04 | 208,897 |

DBP
| Unadjusted | 3.48 | [1.9; 5.0] | 8.72e−06 | 208,894 |
| Adjusted | 2.61 | [1.2; 4.0] | 2.85e−04 | 208,894 |

0 2 4 6 8 10