Identification of *Slit3* as a locus affecting nicotine preference in zebrafish and human smoking behaviour.

Judit García-González[1*], Alistair J. Brock[1*], Matthew O. Parker[2], Riva Riley[1], David Jolliffe[3], Ari Sudwarts[1], Muy-Teck Teh[4], Elisabeth M. Busch-Nentwich[5,6], Derek L. Stemple[5], Adrian R. Martineau[3], Jaakko Kaprio[7,8], Teemu Palviainen[7], Valerie Kuan[9], Robert T. Walton[3*], Caroline H. Brennan[1*]

Affiliations: [1] School of Biological and Chemical Sciences, Queen Mary, University of London, London E1 4NS. [2] Brain and Behaviour Laboratory, School of Pharmacy and Biomedical Science, University of Portsmouth, PO1 2DT, UK. [3] Barts and The London School of Medicine and Dentistry, Blizard Institute, 58 Turner St, Whitechapel, London E1 2AB. [4] Centre for Immunobiology and Regenerative Medicine, Institute of Dentistry, Barts and The London School of Medicine and Dentistry, Blizard Building, 4 Newark St, Whitechapel, London E1 2AT. [5] Wellcome Trust Sanger Institute, [6] Cambridge Institute of Therapeutic Immunology & Infectious Disease (CITIID), Jeffrey Cheah Biomedical Centre, University of Cambridge, Puddicombe Way, Cambridge, CB2 0AW [7] Institute for Molecular Medicine FIMM, HiLIFE. PO Box 20 (Tukholmankatu 8). 00014 Helsinki, Finland. [8] Department of Public Health, Faculty of Medicine, University of Helsinki. Helsinki, Finland. [9] Institute of Cardiovascular Science, University College London, London, UK.

Keywords: Slit3, acoustic startle, smoking, conditioned place preference, gene association

*Equally contributed

Number of words in the abstract: 149 words

Number of words in the main text: 8406

Number of figures: 10

Number of tables: 2

Number of Figure –figure supplements: 2

Number of supplementary tables: 8

ABSTRACT

To facilitate smoking genetics research we determined whether a screen of mutagenized zebrafish for nicotine preference could predict loci affecting smoking behaviour. Of 30 ENU mutagenized families screened, two showed increased or decreased nicotine preference. Out of 25 inactivating mutations in the families, one in the *slit3* gene segregated with increased nicotine preference in heterozygous individuals. Focussed SNP analysis of the human *SLIT3* locus in cohorts from UK (n=863) and Finland (n=1715) identified two variants that predict cigarette consumption and likelihood of cessation. Characterisation of *slit3* mutant larvae and adult fish revealed decreased sensitivity to the dopaminergic and serotonergic antagonist amisulpride, known to affect startle reflex that is correlated with addiction in humans, and increased *htr1aa* mRNA expression in mutant larvae. No effect on neuronal pathfinding was detected. These findings reveal a role for SLIT3 in development of pathways affecting responses to nicotine in zebrafish and smoking in humans.

INTRODUCTION

Tobacco smoking is the leading preventable cause of death worldwide placing a

heavy social and financial burden on society (1–3). It is well established that aspects

of smoking behaviour have a strong genetic component (4–7). However, identifying

causal genetic factors and exploring the mechanisms by which they act is challenging

in human studies: the field has been characterized by small effect sizes and lack of

replication such that there are remarkably few genes and loci that can be confidently

linked to smoking. The strongest evidence for causal effects is for functional variants

in *CHRNA5* (8) and *CYP2A6* (4), affecting amount smoked and nicotine metabolism,

respectively. Recent large studies have identified numerous new association loci, but

their significance is yet to be biologically characterised (6,7).

As approaches to identify genetic risk are difficult in humans, research has been

facilitated by studies in animal models, with a focus on genomic analysis of inbred

and selectively-bred, naturally occurring genetic strains (9). This type of study

produces quantitative trait loci (QTL) maps of multiple loci, each with a small impact

on the phenotype. However, as with human studies, it is inherently difficult to identify

relevant genes from QTL maps, as the overall phenotype cannot be predicted by

individual genotypes. Mutagenesis studies in animal model systems can overcome

these limitations: e.g. N-ethyl-N-nitrosourea (ENU) mutagenesis introduces thousands

of point mutations into the genome with the potential to generate much stronger

phenotypes than those occurring in a natural population thereby facilitating

identification of causal mutations. Examination of phenotypic variation in ENU

mutagenized model species could be applied to identify novel, naturally occurring

variants influencing human addictive behaviour by identifying key genes and
pathways affecting conserved behavioural phenotypes.

Drug-induced reinforcement of behaviour, that reflects the hedonic value of drugs of
abuse including nicotine, is highly conserved in both mammalian and non-mammalian
species (10–13). Conditioned place preference (CPP), where drug exposure is paired
with specific environmental cues, is commonly used as a measure of drug-induced
reward or reinforcement (14). ENU Mutagenesis screens for cocaine or amphetamine-
induced CPP have been undertaken in zebrafish (9,15), however, despite successful
isolation of lines with altered reinforcement responses to these drugs, the causal
mutations have not been identified and the predictive validity of these screens for
human behaviour has not been established. Larval locomotor response to nicotine has
also been used to explore nicotine response genetics (16) but the relevance of larval
locomotion to addiction is somewhat less clear.

Here, we conducted a forward genetic screen of families of ENU-mutagenized
zebrafish for nicotine-induced CPP. Zebrafish express homologues of all 16 members
of the nicotinic acetylcholine receptor family present in mammals (17–19) with
similar binding characteristics (20,21). However, as a result of a local gene
duplication event in the ray fish lineage and a teleost genome tetraploidation event,
zebrafish have duplicate copies of nicotinic receptor α2, α4, α7, α9, α10, β1, β3 and
β5. In addition, zebrafish have additional receptors (α8 and α11) that have been lost in
humans (19). Zebrafish show robust CPP to nicotine (21–24). Nicotinic receptor
partial agonists, that modulate striatal dopamine release in response to nicotine in
mammalian systems, also inhibit nicotine-induced CPP in zebrafish (21). Further, on
prolonged exposure to nicotine or ethanol, adult zebrafish show conserved adaptive

changes in gene expression and develop dependence-related behaviours, such as

persistent drug seeking despite adverse stimuli or reinstatement of drug seeking

following periods of abstinence (24). These data demonstrate the existence of a

conserved nicotine-responsive reward pathway and support the suitability of zebrafish

to examine the genetic and molecular mechanisms underlying behavioural responses

to nicotine.

To evaluate the use of a behavioural CPP screen in zebrafish to predict loci affecting

human smoking behaviour we initially assessed 1) the ability of varenicline and

bupropion, pharmacological agents used to treat human nicotine addiction, to reduce

zebrafish nicotine-induced place preference and 2) the heritability of nicotine

responses in ENU-mutagenized fish. We then screened 30 families of ENU-

mutagenized fish to identify families with increased/decreased CPP for nicotine. For

two families with altered CPP response, the phenotype was confirmed following

independent replication with a larger number of fish. Exome sequence information

was used to generate a list of coding, loss of function candidate mutations affecting

the phenotype. One family with a mutation co-segregating with increased nicotine

CPP was selected for further study. Firstly, the effect of the identified gene on

nicotine-induced CPP was confirmed using an independent line carrying a similar

predicted loss of function mutation in the same gene. We then characterized the

mutation using gene expression analysis, immunohistochemical analysis of neuronal

pathways and behavioural responses to acoustic startle; a response known to be

modulated by serotonergic and dopaminergic signalling and, in humans, associated

with vulnerability to addiction (25–27). Finally, we used focused SNP analysis of

human cohorts to assess the predictive validity of findings in fish for human smoking

behaviour.

In agreement with previous studies zebrafish showed a robust CPP to nicotine.

Nicotine-induced CPP was abolished by varenicline and bupropion and found to be

heritable in fish. The screening of ENU mutagenized families identified mutations in

the *slit3* gene influencing sensitivity to rewarding effects of nicotine. *Slit3* mutant

larvae and adult fish showed reduced behavioural sensitivity to amisulpride and larvae

showed increased *ht1aa* receptor expression. No effect on neuronal pathfinding was

detected. Analysis of the *SLIT3* locus in two independent human cohorts identified

two genetic markers that predict level of cigarette consumption and likelihood of

cessation. This proof of principle study demonstrates that screening of zebrafish is

able to predict loci affecting complex human behavioural phenotypes and suggests a

role for SLIT3 signalling in the development of dopaminergic and serotonergic

pathways affecting behaviours associated with nicotine sensitivity.

RESULTS

Nicotine CPP in zebrafish is inhibited by varenicline and buproprion

The hedonic value of drugs of abuse, that gives rise to reinforced behaviour, is

commonly assessed using either self-administration protocols or CPP. The ability of

compounds used as therapeutics in humans to prevent rodent nicotine self-

administration is used to support the translational relevance of nicotine-self

administration in that model (28–30). As our aim was to use nicotine-CPP to predict

genes affecting smoking behaviour, we assessed the ability of the nicotine

therapeutics varenicline and bupropion to inhibit nicotine induced CPP in zebrafish.

As seen previously (22,24,31), 10μM nicotine induced a robust 15-20% change in

place preference. Consistent with previous results (21), pre-incubation in varenicline

or bupropion dose-dependently inhibited the nicotine CPP response (Figure 1).

97
Figure 1. 10μM nicotine induced place preference in zebrafish is sensitive to

inhibition by therapeutics effective in humans. A) varenicline (nicotine partial

agonist) and B) bupropion (norepinephrine and dopamine reuptake inhibitor with

nicotine antagonist properties when metabolised). Bars represent mean and error bars

represent +SEM. Asterisk (*) represents significance at p<0.05.

Nicotine CPP is heritable in zebrafish

We selected a nicotine concentration predicted to induce a minimal detectable CPP in

wild types (5μM) (22,23), to enable us to detect both increased and decreased

response to nicotine in mutants. To ensure that this strategy could detect genetic

factors affecting response to nicotine, we assessed the heritability of the CPP response

in ENU mutagenized fish using a selective breeding approach over three generations.

Figure 2A shows our assessment strategy where fish showing the highest and lowest

CPP response are selected for further breeding. In the first generation, the CPP change

score phenotype was normally distributed (Shapiro-Wilks p = 0.83) and there was a

mean CPP change score of 0.11 to the drug paired side. CPP change scores ranged

from - 0.4 to 0.6.

An increasing difference in nicotine preference between offspring of fish from the

upper vs lower extremes of the distribution (Shift of Cohen's d = 0.89 in Second

generation CPP to d = 1.64 in Third generation CPP) indicates that nicotine CPP

behaviour is heritable in zebrafish (Figure 2B), and that our CPP strategy is able to

identify heritable differences in both extremes of the distribution. Phenotypes in the

second and third generation screen may result from selecting for multiple co-

segregating mutations that strengthened the phenotypes, or from selecting against

other contrary mutations that weaken effects.

Figure 2. A) Breeding and selection to assess heritability of nicotine-induced place preference in ENU-mutagenized zebrafish. To test whether nicotine preference is heritable, fish in the upper (blue circle) and lower (red circle) 10% of the change in preference distribution curve were inbred and screened for CPP (Second generation CPP assay). A similar approach was used for the third generation CPP assay. **B) CPP for nicotine is heritable**. Mean preference change is increasingly distinct for the second and third generation CPP analysis. Plot represents mean and ±SEM. First generation (corresponding to the F_3 families used for the screen) (n=120): mean=0.11; SD=0.17. Second generation: Offspring of fish from *upper* 10% of the first generation screen (n=92): mean=0.17; SD=0.14. Offspring of fish from *lower* 10% of the first generation screen (n=64): mean=0.08; SD=0.15. Third generation. Offspring of fish from *upper* 10% of the second generation screen (n=69): mean=0.21; SD=0.10. Offspring of fish from *lower* 10% of the second generation screen (n=67): mean=0.01; SD=0.09.

Identification of *slit3* mutations affecting nicotine place preference in zebrafish

Out of 30 families screened, individuals from nine families were in the top 5% of the change in preference distribution, and individuals from five families in the bottom

5%. To identify candidate mutations affecting nicotine preference in fish, we focussed

on families where all individuals included in the screen clustered at one or other

extreme of the distribution curve. Two families (called AJBQM1 and AJBQM2 after

the researcher who conducted the screen), which clustered at the top (AJBQM1) and

bottom (AJBQM2) of the nicotine preference distribution, were selected for further

study. We first assessed nicotine CPP in the remaining siblings not initially included

in the screen. As shown in Figure 3, the phenotypes were conserved when remaining

siblings were assessed. Exome sequencing of fish (32) used to generate AJBQM1 and

AJBQM2 identified 25 nonsense and essential splice site mutations. We genotyped

fish at these 25 loci and determined the co-segregation with nicotine preference.

Figure 3. AJBQM1 and AJBQM2 families show increased and decreased

nicotine place preference. AJBQM1 and AJBQM2 siblings, not included in the

screen (n=10 for AJBQM1; n=14 for AJBQM2), AJBQM1 significantly differed from

the parental strain, Tupfel longfin (TLF) wild type (w/t) saline control (n=17) and

wild type nicotine exposed fish (n=7). AJBQM2 differed from wild type nicotine

exposed fish but not wild type saline controls. Different superscript letters indicate

significant difference between groups (p<0.05), same superscript letters indicate no

significant differences between groups. Bars indicate Mean ±SEM.

Of the 25 coding, predicted loss of function mutations in AJBQM1 and AJBQM2

(Listed in Supplementary Table 1), only $slit3^{sa1569/+}$ (exon 7 splice acceptor site

disruption at amino acid position 176), segregated with nicotine preference (Figure

4A & Supplementary Table 5A). None of the coding, predicted loss of function

mutations in AJBQM2 segregated with nicotine preference and this line was not

examined further (Supplementary Table 5B).

To confirm that loss of $slit3$ function was related to nicotine seeking behaviour we

obtained an independent family of fish, $slit3^{sa202}$, carrying a G>T transversion

producing a premature stop codon at amino acid position 163 in the SLIT3 protein

from the Sanger Institute. Although not as marked as in AJBQM1 mutants (hereafter

called $slit3^{sa1569}$), heterozygous $slit3^{sa202}$ fish showed enhanced nicotine CPP (p=0.03)

compared to wild type siblings (Figure 4C & 4D). The $slit3^{sa1569}$ allele affects slicing

and $slit3^{sa202}$ introduces a premature stop codon. Both alleles reside before the second

leucine rich repeat (LRR) domain in the encoded protein (Figure 4B), which is

essential for interaction with ROBO receptor proteins (33).

Figure 4. **A. Segregation of *slit3^{sa1569}* mutation with nicotine seeking.** CPP change

scores for individual un-mutagenized TLF wild type fish (n=7) and AJBQM1 fish

(n=10). Following CPP analysis, fish were genotyped for 25 loss of function

mutations contained within the family. Black dots indicate *slit3^{sa1569/+}* heterozygous

mutant fish. White dots indicate *slit3^{sa1569+/+}* fish. Heterozygosity for *slit3^{sa1569}*

segregates with increased nicotine seeking behaviour. **B. Position of ENU-induced**

mutations in zebrafish Slit3 protein. *slit3^{sa1569}* (A>G transition) disrupts a splice site

in intron 7 affecting translation at amino acid 176. *slit3^{sa202}* (G>T transversion)

introduces a stop codon at amino acid 163. Both mutations truncate the protein before

the leucine rich repeat domain 2 (D2), which interacts with membrane bound ROBO

during SLIT-ROBO signalling. **C: Nicotine preference of *slit3^{sa202}* line.** *slit3^{sa202/+}*

fish (n=18) show increased nicotine preference compared to wild type TLF controls

(n=8) (p = 0.001) and wild type siblings $slit3^{+/+}$ (n=14) (p<0.05). Bars indicate mean

+SEM. **D: Segregation of $slit3^{sa202}$ allele with nicotine seeking**. CPP change scores

for individual un-mutagenised TLF wild type parent strain fish (n=8) and $slit3^{sa202}$ fish

(n=21). Black dots indicate $slit3^{sa202/+}$ heterozygous mutant fish, white dots indicate

$slit3^{sa202+/+}$ fish. Mutations in $slit3^{sa202}$ co-segregate with nicotine preference.

Heterozygous $slit3^{+/sa202}$ present increased place preference compared to $slit3^{sa202+/+}$

siblings (n=11).

Characterisation of $slit3^{sa1569}$ mutants

SLIT3 is a member of a family of proteins with established axon guidance properties

and previously suggested to be involved in dopaminergic and serotonergic pathfinding

(34). Therefore, we performed immunostaining in three-day-old zebrafish larvae and

examined the number of cell bodies and axonal projections of serotonergic (5-HT)

and catecholaminergic neurons in the brain using anti-5-HT and anti-tyrosine

hydroxylase (TH) antibodies (Figure 5).

No differences between $slit3^{sa1569}$ mutant and wild type larvae were observed in the

number of cells labelled by anti-5HT antibody in the raphe nucleus (Mean ± SEM:

$slit3^{+/+}$: 40.2±2.6 vs. $slit3^{sa1569/sa1569}$: 47.0±4.7, p=0.23) rostral hypothalamus ($slit3^{+/+}$:

15.4±2.2 vs. $slit3^{sa1569/sa1569}$: 19.4±3.8, p=0.38) or inferior hypothalamic lobes

($slit3^{+/+}$: 76.8±13.5 vs. $slit3^{sa1569/sa1569}$: 94.8±12.9, p=0.36), nor in the number of cells

labelled by anti-TH antibody in the diencephalic dopaminergic cluster ($slit3^{+/+}$:

209±18 vs. $slit3^{sa1569/sa1569}$: 140±35, p=0.12) and medulla oblongata interfascicular

zone and vagal area ($slit3^{+/+}$: 196±40 vs. $slit3^{sa1569/sa1569}$: 124±27, p=0.17). Similarly,

we observed no significant differences in the number of anti-TH labelled axon tracts

projecting to the midline across the 3 planes examined ($slit3^{+/+}$: 2.6±2.3 vs

$slit3^{sa1569/sa1569}$: 4.3±1, p=0.53) (Figure 5).

We also looked at the expression patterns using anti-acetylated tubulin antibody along

the midline in the ventral forebrain, where $slit3$ is known to be expressed (35).

However, no obvious differences were observed. Staining of $slbp^{ty77e/ty77e}$ mutant

larvae, known to have fewer neurons and axonal defects (16) were used as positive

control (Figure 5-figure supplement 1).

Figure 5: Fluorescent immunohistochemistry in three-day old wild type *slit3*^{sa1569+/+} **and homozygous mutant** *slit3*^{sa1569/sa1569}. A-D) anti-5-HT, E-M) anti TH:

A) 5-HT-labelled neurons in wild type zebrafish brain. Circles indicate regions used

for quantification of cell number in rostral hypothalamus (RH), inferior hypothalamic

lobes (HL-R, HL-L) and raphe nucleus (RN). B) anti-5-HT labelled cells in *slit3* wild

type brain, C) anti-5-HT labelled cells in *slit3^{sa1569}* homozygous mutant brain. D)

Quantification of anti-5HT labelled cell number in wild type and *slit3* mutant brains

No significant differences were observed between wild type and *slit3* mutant larvae.

E) Unprocessed maximum intensity projection of anti-TH-labelled whole mounted

wild type zebrafish brain. Circles indicate areas used for quantification, or in the case

of LC-R and LC-L, landmarks used as reference to determine the extension of the

medial longitudinal catecholaminergic tract (MLC) used when quantifying the

number of anti-TH labelled projections to the midline (panels L, M), F) Cell

quantification for diencephalic dopaminergic cluster (DDC). No significant

differences were observed between wild type and *slit3* mutant larvae. G) Cell

quantification for medulla oblongata interfascicular zone and vagal area, and area

postrema (MO, AP). No significant differences were observed between wild type and

slit3 mutant larvae. H-K) Anti-TH labelled wild types and *slit3^{sa1569}*. Zoomed-in

visualization of diencephalic dopaminergic cluster (H-I) and medulla oblongata

interfascicular zone and vagal area (J-K). L-M) Quantification of catecholaminergic

projections projecting to the midline. Examples of projections are indicated with

yellow arrows. Projections were assessed from posterior to anterior using the locus

coerulus and posterior extent of the raphe nucleus as landmarks (Panel L, yellow line)

and from dorsal to ventral (Panel M, stacks 1-3). Figure 5-figure supplement 1 shows

individual planes. n=5 samples per genotype group.

Although we did not observe differences between wild type and *slit3^{sa1569}* mutants,

subtle effects on circuit formation may not have been detected by our antibody

staining. To further characterise the *slit3* phenotype and explore potential functional

differences in catecholamine circuitry, we examined the response and habituation to

acoustic startle stimuli in wild type and mutant fish. Habituation to acoustic startle is

known to involve catecholamine signalling and to be sensitive to

dopaminergic/serotonergic antagonists such as amisulpride (36) and, in humans, is

associated with vulnerability to addiction (25–27). Five day old larvae were subjected

to 10 sound/vibration stimuli over a total of 20 seconds (2 second interval between

each stimulus) in the presence of 0, 0.05 mg/L, 0.1 mg/L or 0.5 mg/L amisulpride in

0.05% dimethyl sulfoxide (DMSO). The distance travelled one second after each

stimulus was recorded for each fish.

Response and habituation to the stimuli was quantified as the percentage of fish

moving more than 4.6 mm, which corresponds to 85% of the mean distance travelled

one second after the first startle (Figure 6A). Using this criterion – and in line with the

habituation response paradigm (37) – a lower percentage of fish responded as the

number of stimuli increased: 68% or wild type non drug-treated individuals responded

to the first stimulus, 81% to the first and/or second stimulus, whereas 16% responded

to the last stimulus (Figure 6B).

In drug-free conditions, there were no differences across *slit3*^{sa1569} genotype groups

(Figure 6C). However, when larvae were treated with amisulpride, the habituation to

startle responses across taps was different across genotypes. Amisulpride caused a

biphasic dose dependent effect on stimulus response in wild types such that 0.05mg/L

caused an increase in responders across all 10 stimuli, and 0.5mg/L caused a decrease

(Effect of amisulpride dose p<0.001). A similar pattern was observed for

heterozygous *slit3*^{sa1569}, but the effect of amisulpride was not significant (p=0.083).

Amisulpride dose had no significant effect on stimulus response in homozygous

slit3^{*sa1569*}, that showed an increase in response to low doses but were less sensitive to

inhibition at high doses (Figure 6D-F). The presence of a *slit3*^{*sa1569*} genotype by

amisulpride dose interaction across taps was confirmed by a three-way interaction in

the regression models ($p=0.04$). The interaction between dose and stimulus event

number was a significant predictor of response in wild type larvae ($p=0.044$) and

heterozygous larvae ($p=0.02$) but not in homozygous larvae ($p=0.16$).

There were no significant differences in locomotion in the 15 seconds before the first

startle, in magnitude of the response to the first tap stimulus, nor in total distance

moved across all tap stimuli across experimental groups (Figure 6-figure supplement

1) indicating that differences in behaviour were not confounded by differences in

locomotion per se.

Figure 6. Habituation response in the presence and absence of amisulpride. A-B:
Response and habituation to 10 stimuli with two seconds interval between stimuli
in wild type, drug free zebrafish. Mean distances travelled were measured in one
second time bins. Line indicates 4.6 mm, which corresponds to 85% of the population
mean distance travelled one second after the first stimulus and was used to define
respondents. The percentage of fish responding to the stimuli decreases with
stimulus/tap number (Main effects of tap number $p<0.05$) 67% respond to the first
tap; 16% respond to the last tap. Respondents are defined as fish moving more than
4.6 mm. **C: proportion of responders across the ten stimuli in drug free**
individuals from each genotype: there was no significant effect of genotype on
response across taps ($p=0.340$) or responsiveness ($p=0.352$) in drug free fish. **D:**

Mean percentage of responders across the ten stimuli (±SEM). Data are stratified

by *slit3*^{sa1569} genotype and amisulpride dose normalised to response in absence of

drug. The effect of amisulpride on habituation varies by genotype. **E/F: proportion**

of individuals responding in each amisulpride dose condition in wild type and

homozygous mutant fish, respectively. The interaction between amisulpride dose

and stimulus event number had a significant effect on the proportion of responsive

individuals in wild type individuals (p<0.05) but not homozygous mutants (p=0.160).

Adult *slit3*$^{sa1569/ sa1569}$ mutant zebrafish showed a qualitatively different response to

inhibition of CPP by amisulpride compared to wild type siblings, consistent with a

persistent difference in sensitivity to this drug. The minimal CPP induced by 5µM

nicotine in wild type fish was prevented by pre-exposure to 0.5mg/L amisulpride.

Nicotine-induced CPP in *slit3*^{sa1569} homozygous mutants was not affected (Figure 7).

Figure 7. CPP induced by 5μM nicotine is blocked by 0.5mg/L

dopamine/serotonin antagonist amisulpride in wild type *slit3^sa1569+/+* fish but not

in *slit3^sa1569* homozygous mutants. Bars represent mean (+SEM). (n=11-14 fish per

group). *Two-way ANOVA followed by post-hoc Tukey tests (p < 0.05).

As *slit3^sa1569* homozygous mutant fish showed altered sensitivity to nicotine and

amisulpride, we examined whether expression of genes previously associated with

nicotine dependence was dysregulated in *slit3^sa1569* mutant larvae using quantitative

real-time PCR (qPCR). We included the nicotinic receptors and genes from the

dopamine receptor family -*drd1* (40–42)(38), *drd2* (39) and *drd3* (40)-, and the

dopamine transporter *dat* (41)). Genes from the adreno-receptor families (*adra1* and

adra2) were also included due to their links with nicotine addiction and use as

putative targets for smoking treatment (43–47)(42–44). Finally, expression of

serotonin receptor genes was included, again due to their well-established links with

nicotine addiction (45–49).

For several genes (i.e. *drd3, chrnb3, htr4, adra2b*), up-regulation of gene expression

in mutant larvae showed nominal significance (Supplementary Table 8). However,

only *htr1aa* ($[F(2,6)=44]$, $p=0.0003$) showed a significant difference across genotypes

after correcting for multiple testing (Figure 8).

Figure 8. Quantitative real-time PCR analysis of five day old wild type *slit3*[sa1569+/+]**,** *slit3*[sa1569/+] **heterozygous and** *slit3*[sa1569/sa1569] **homozygous mutant larvae** (Total n=30, 3 samples per experimental group with n=10 embryos per sample). Only *htr1aa* ([F(2,6)=44], p=0.0003) showed a significant difference across

genotypes after correcting for multiple testing. *Two-way ANOVA followed by post-

hoc Tukey test ($p < 0.05$).

Variations at the *SLIT3* locus predict smoking behaviour in human samples

We next examined associations between 19 single nucleotide polymorphisms (SNPs)

in the human *SLIT3* gene and smoking behaviour in two London cohorts. Two SNPs,

rs12654448 and rs17734503 in high linkage disequilibrium (Figure 9) were associated

with level of cigarette consumption ($p=0.00125$ and $p=0.00227$). We repeated the

analysis on heavy smokers: rs12654448 ($p=0.0003397$) and rs17734503

($p=0.0008575$) were again associated with cigarette consumption together with

rs11742567 ($p=0.004715$). The SNP rs11742567 was associated with cigarette

consumption in light smokers (<20 cigarettes per day, $p=0.003909$)) and with

quitting. Associations are reported in Table 1. No other *SLIT3* polymorphisms were

associated with smoking initiation, persistent smoking or cessation (Supplementary

Tables 6 & 7).

Figure 9: Linkage disequilibrium (LD) plot of *SLIT3* SNPs in human smoking association analysis. Numbers within each square indicate D' values (white: D' < 1, LOD < 2; blue: D' = 1, LOD < 2; pink: D' < 1, LOD ≥ 2; and bright red: D' = 1, LOD ≥ 2). Top part of the figure shows domain organization of the SLIT protein based on the UCSC Genome Browser (http://genome.ucsc.edu/) in relation to the SNP location. LRR: leucin-rich repeats. EGF: epidermal growth factor domains. LamG; Laminin G domain. Some intron numbers were added for reference.

Table 1. Associations of *SLIT3* SNPs with level of tobacco consumption for the London study groups (n=863). Regression coefficients,

confidence intervals and p-values from linear regression of cigarettes smoked per day (CPD) on minor allele count for smokers from COPD,

asthma and general cohorts, adjusted for age, sex and cohort. β coefficient represents effect of each additional minor allele. Benjamini-Hochberg

cut-off at q-value 0.1 = 0.01053. **Associations of *SLIT3* SNPs with tobacco consumption in a subset of heavy smokers (≥20 cigs/day).**

Adjusted for age, sex and cohort. (q-value 0.1 = 0.01579). **Associations of *SLIT3* SNPs in a subset of light smokers (<20 cigs/day).** Adjusted

for age, sex and cohort (q-value 0.1 = 0.00526). **Association analysis of *SLIT3* SNPs with smoking cessation.** Logistic regression of current

smokers *vs* ever smokers controlling for age, sex and cohort. Odds ratio >1 indicates minor allele increases odds of persistent smoking relative to

major allele. P: p-value, SE: standard error, L95: lower limit of 95% confidence interval, U95: upper limit. For all panels, associations in bold

remained significant after adjustment for multiple comparisons using a Benjamini-Hochberg procedure to control false discovery rate at 10%.

	Tobacco consumption				Tobacco consumption - heavy smokers (≥20 cigs/day)				Tobacco consumption - light smokers (<20 cigs/day)				Smoking cessation				
SNP	P value	β	SE	95% CI	P value	β	SE	95% CI	P value	β	SE	95% CI	OR	SE	L95	U95	P value
rs10036727	0.629	-0.388	0.802	(-1.960, 1.183)	0.448	-0.653	0.860	(-2.337, 1.032)	0.940	-0.051	0.686	(-1.396, 1.293)	0.947	0.160	0.693	1.295	0.734
rs11134527	0.218	1.014	0.822	(-0.596, 2.625)	0.327	-0.867	0.883	(-2.599, 0.864)	0.261	0.795	0.705	(-0.586, 2.176)	0.665	0.165	0.482	0.918	0.013
rs11742567	0.135	-1.166	0.779	(-2.691, 0.361)	**0.005**	**-2.346**	**0.825**	**(-3.962, -0.730)**	**0.004**	**1.888**	**0.644**	**(0.6258, 3.151)**	**1.586**	**0.163**	**1.153**	**2.183**	**0.005**
rs11749001	0.059	1.972	1.044	(-0.074, 4.018)	0.873	0.177	1.103	(-1.985, 2.338)	0.206	1.200	0.944	(-0.651, 3.051)	0.953	0.206	0.637	1.426	0.817
rs12515725	0.592	-0.406	0.756	(-1.888, 1.076)	0.488	-0.565	0.813	(-2.159, 1.029)	0.278	-0.688	0.631	(-1.925, 0.550)	1.028	0.151	0.765	1.381	0.855
rs12521041	0.904	-0.105	0.865	(-1.801, 1.591)	0.996	-0.005	0.942	(-1.851, 1.841)	0.059	-1.354	0.710	(-2.746, 0.038)	1.554	0.178	1.096	2.205	0.013
rs12654448	**0.001**	**-4.241**	**1.307**	**(-6.803, -1.680)**	**0.0003**	**-4.830**	**1.334**	**(-7.444, -2.216)**	0.410	-1.034	1.251	(-3.486, 1.417)	1.625	0.279	0.941	2.808	0.082
rs1345588	0.240	-1.268	1.078	(-3.380, 0.845)	0.253	-1.334	1.164	(-3.616, 0.948)	0.869	-0.150	0.907	(-1.927, 1.627)	1.417	0.222	0.918	2.189	0.116
rs1421763	0.272	-0.982	0.894	(-2.735, 0.770)	0.978	-0.027	0.959	(-1.908, 1.853)	0.162	-1.074	0.764	(-2.571, 0.424)	0.917	0.176	0.649	1.294	0.622
rs1559051	0.961	-0.040	0.819	(-1.644, 1.564)	0.458	0.656	0.882	(-1.073, 2.384)	0.507	0.455	0.685	(-0.880, 1.797)	0.919	0.163	0.668	1.265	0.606
rs17665158	0.131	1.338	0.884	(-0.394, 3.070)	0.236	1.114	0.939	(-0.727, 2.955)	0.034	1.620	0.758	(0.1354, 3.106)	0.723	0.172	0.516	1.013	0.060
rs17734503	**0.002**	**-3.987**	**1.299**	**(-6.534, -1.441)**	**0.001**	**-4.458**	**1.325**	**(-7.055, -1.861)**	0.410	-1.034	1.251	(-3.486, 1.417)	1.616	0.275	0.942	2.773	0.081
rs2938774	0.140	1.101	0.745	(-0.359, 2.562)	0.528	0.496	0.786	(-1.044, 2.036)	0.015	-1.655	0.674	(-2.976, -0.333)	0.753	0.148	0.563	1.007	0.056
rs295994	0.714	0.283	0.770	(-1.227, 1.793)	0.643	0.378	0.813	(-1.215, 1.971)	0.238	-0.796	0.672	(-2.114, 0.521)	0.799	0.154	0.591	1.082	0.147
rs297886	0.620	0.442	0.890	(-1.303, 2.187)	0.961	-0.048	0.986	(-1.979, 1.884)	0.489	0.488	0.704	(-0.891, 1.867)	1.108	0.177	0.784	1.568	0.561
rs3733975	0.909	-0.099	0.860	(-1.784, 1.587)	0.982	0.022	0.934	(-1.809, 1.852)	0.059	-1.354	0.710	(-2.746, 0.038)	1.488	0.176	1.054	2.101	0.024
rs4282339	0.669	-0.434	1.013	(-2.419, 1.552)	0.942	-0.080	1.103	(-2.241, 2.081)	0.238	-1.006	0.849	(-2.670, 0.658)	0.984	0.203	0.661	1.464	0.936
rs7728604	0.701	0.286	0.744	(-1.173, 1.745)	0.321	0.827	0.832	(-0.803, 2.457)	0.654	0.262	0.583	(-0.880, 1.404)	0.935	0.149	0.698	1.253	0.653
rs9688032	0.948	-0.050	0.766	(-1.551, 1.451)	0.770	0.246	0.839	(-1.398, 1.890)	0.080	-1.076	0.610	(-2.272, 0.119)	1.066	0.156	0.786	1.446	0.680

We subsequently investigated associations with more detailed smoking phenotypes in

the Finnish twins cohort (50) (Table 2). Associations were observed between

rs17734503 and DSM-IV nicotine dependence symptoms (p=0.0322) and age at onset

of weekly smoking (p=0.00116) and between rs12654448 and age at onset of weekly

smoking (p=0.00105). Associations were seen elsewhere between *SLIT3* markers and

Fagerström Test for Nicotine Dependence (FTND), cigarettes smoked each day,

sensation felt after smoking first cigarette and time to first cigarette in the morning. In

keeping with the London studies the minor allele was associated with a lower degree

of dependence and decreased cigarette consumption.

The SNPs rs12654448 and rs17734503 are in non-coding domains, therefore it was

not possible to predict loss or gain of function of SLIT3 from the SNP location. No

evidence of affecting gene expression was found as per GTEx database

(https://gtexportal.org/home/).

Table 2: Associations between detailed nicotine dependence phenotypes and SLIT3 genotype in a Finnish twin cohort (n=1715).

Associations of *SLIT3* SNPs with DSM-IV nicotine dependence symptoms, Fagerström scores (FTND), cigarettes smoked each day (CPD), and

sensation felt after smoking first cigarette and time to first cigarette in the morning. The three SNPs that were linked to smoking behaviour in the

London cohorts are shown in bold.

SNP	DSM-IV ND diagnosis			DSM-IV ND symptoms			FTND (≥4)			FTND score		
	β	SE	P value	β	SE	P value	β	SE	P value	β	SE	P value
rs12654448	-0.0343	0.0262	0.190975	-0.1839	0.0964	0.056728	0.0526	0.0287	0.066509	0.075	0.1365	0.58286
rs17734503	-0.0354	0.0259	0.171821	-0.2044	0.0954	**0.032199**	0.0474	0.0283	0.094383	0.0443	0.135	0.743052
rs11742567	0.0006	0.0163	0.97262	-0.0359	0.0601	0.55086	0.0134	0.0179	0.45384	0.0449	0.0851	0.597682
rs17665158	0.0117	0.019	0.538639	0.1536	0.0696	0.027544	0.0178	0.0207	0.391096	0.0935	0.0988	0.344157
rs1345588	-0.0031	0.0222	0.889847	-0.0389	0.0817	0.634184	0.0578	0.0242	0.01708	0.1901	0.1157	0.100729
rs7728604	-0.0049	0.0162	0.761485	-0.0442	0.0597	0.459743	0.0004	0.0177	0.980706	-0.0261	0.0846	0.757849
rs11134527	0.0296	0.0171	0.084576	0.0927	0.063	0.141369	0.0324	0.0187	0.083498	0.1376	0.0891	0.122807
rs10036727	0.0067	0.0165	0.68406	0.0207	0.0605	0.732266	0.0022	0.018	0.903865	0.046	0.0857	0.591583
rs1559051	0.0249	0.0193	0.198647	0.0492	0.0703	0.484353	-0.0433	0.0208	0.037736	-0.1011	0.0995	0.309836
rs12515725	0.0072	0.0159	0.6502	0.0277	0.0584	0.635739	0.0586	0.0172	0.000696	0.2482	0.0824	0.002637
rs2938774	0.0042	0.0173	0.80717	0.0096	0.0642	0.881054	-0.0157	0.0191	0.41163	-0.0173	0.0907	0.848978
rs295994	-0.014	0.0171	0.410864	-0.0144	0.0622	0.816397	-0.016	0.0184	0.38542	-0.0985	0.0879	0.262584
rs9688032	-0.0174	0.0173	0.31299	-0.0347	0.0636	0.585081	0.0234	0.0189	0.216144	0.0626	0.09	0.4869

SNP	β	SE	P value	β	SE	P value	β	SE	P value	β	SE	P value
rs11749001	0.0178	0.0235	0.448278	-0.0062	0.0865	0.942516	0.0224	0.0257	0.383552	0.0067	0.1222	0.956097
rs4282339	0.0118	0.0201	0.557544	0.0526	0.0739	0.476216	-0.0077	0.0219	0.724058	0.1623	0.1045	0.120641
rs297886	-0.0216	0.0171	0.207835	-0.045	0.0634	0.478517	-0.0256	0.0188	0.173314	-0.1354	0.0897	0.131469
rs1421763	0.0079	0.0187	0.671624	0.0178	0.0687	0.795522	0.0641	0.0203	0.001641	0.2582	0.0971	0.007892
rs3733975	-0.013	0.0167	0.436903	-0.0798	0.0613	0.192755	-0.0384	0.0181	0.034371	-0.2083	0.0866	0.016295
rs12521041	-0.0098	0.0167	0.559173	-0.0669	0.0613	0.275274	-0.0365	0.0182	0.044962	-0.1905	0.0868	0.028295

	CPD			max CPD			Age of onset of weekly smoking			First time sensation			FTND time to first cigarette		
SNP	β	SE	P value	β	SE	P value	β	SE	P value	β	SE	P value	β	SE	P value
rs12654448	-0.3509	0.5669	0.536029	-1.0602	0.7743	0.171106	0.7826	0.2384	**0.001051**	-0.0861	0.1423	0.545206	0.0047	0.0802	0.953291
rs17734503	-0.479	0.5608	0.393086	-1.2329	0.7657	0.107544	0.7689	0.2362	**0.001156**	-0.1039	0.1406	0.460344	0.0188	0.0795	0.812682
rs11742567	0.0179	0.3532	0.959588	-0.4621	0.4823	0.338159	0.0965	0.1493	0.518066	-0.1003	0.0884	0.256427	-0.0216	0.05	0.665265
rs17665158	0.8135	0.4096	0.047191	1.5424	0.5587	0.005828	0.0562	0.1732	0.745385	0.2476	0.1027	0.01603	-0.0884	0.058	0.12787
rs1345588	0.294	0.4805	0.54066	0.3968	0.6562	0.54542	0.0989	0.2031	0.626431	-0.0618	0.1204	0.607606	-0.1303	0.068	0.055678
rs7728604	-0.0772	0.3511	0.825888	0.0691	0.4795	0.885363	-0.0261	0.1486	0.860643	-0.029	0.0875	0.740682	-0.0193	0.0497	0.6978
rs11134527	0.1831	0.3705	0.621187	0.7441	0.5057	0.141392	-0.2347	0.1563	0.133517	0.1142	0.093	0.21965	-0.1089	0.0523	0.037681
rs10036727	0.1482	0.3557	0.67697	0.4246	0.4858	0.382161	-0.0456	0.1507	0.762197	0.0289	0.0896	0.74711	-0.0639	0.0504	0.205061
rs1559051	-0.4816	0.413	0.243693	-0.4779	0.5641	0.397066	0.1437	0.175	0.411533	-0.0289	0.1045	0.782381	0.0731	0.0586	0.212174
rs12515725	0.5491	0.3429	0.10948	0.7708	0.4684	0.100032	-0.1629	0.1452	0.26192	-0.0368	0.0865	0.670165	-0.1385	0.0485	0.00434
rs2938774	-0.2796	0.377	0.45835	0.1945	0.5149	0.70567	-0.0575	0.1598	0.718862	0.043	0.0933	0.645221	-0.0136	0.0533	0.797909
rs295994	-0.2793	0.3651	0.444276	-0.2585	0.4988	0.604451	0.1543	0.1548	0.318928	0.0625	0.0926	0.499881	0.0527	0.0517	0.307869
rs9688032	0.2452	0.3738	0.511921	0.4211	0.5107	0.409766	-0.1142	0.1584	0.471283	-0.2227	0.0937	0.01755	-0.0517	0.0531	0.329867

rs11749001	-0.0301	0.5078	0.952789	0.0574	0.6939	0.934054	-0.1994	0.2147	0.353064	0.1497	0.1274	0.240255	0.0075	0.0718	0.916823
rs4282339	0.4086	0.434	0.346592	0.3083	0.593	0.603197	0.0952	0.1836	0.603958	-0.0394	0.1084	0.716204	-0.0524	0.0614	0.394221
rs297886	-0.1375	0.3727	0.712273	-0.4782	0.5091	0.347622	0.1262	0.1575	0.423104	0.0519	0.0928	0.576255	0.0632	0.0527	0.230861
rs1421763	0.5585	0.4037	0.166723	0.6702	0.5515	0.224417	-0.1481	0.1706	0.385475	-0.0799	0.1018	0.432497	-0.1269	0.0571	0.026442
rs3733975	-0.7784	0.3597	0.030606	-1.0555	0.4911	0.031758	0.0902	0.1521	0.553335	-0.2932	0.0896	0.001085	0.1373	0.0509	0.007035
rs12521041	-0.7312	0.3602	0.042534	-0.8864	0.492	0.071805	0.0522	0.1523	0.731943	-0.3129	0.0897	0.000499	0.1257	0.051	0.01373

DISCUSSION

The aim of this study was to use forward genetic screening in zebrafish to identify

loci affecting human smoking behaviour. Among 30 mutant families screened for

CPP, we identified one family showing increased nicotine preference compared to

wild types. Out of the 13 pre-identified loss-of-function mutations in that family, only

one in the *slit3* gene co-segregated with the behaviour. We confirmed the association

between *slit3* loss of function and increased nicotine preference using an independent

zebrafish mutant line with a different loss of function mutation in *slit3*. Next, we

established the relevance in humans by identifying two markers in *SLIT3* where the

presence of the minor allele was associated with fewer cigarettes smoked each day

and with smoking cessation. Studies in a separate twin cohort showed that these same

alleles were associated with DSM-IV nicotine dependence symptoms and age at onset

of weekly smoking. Taken together these findings suggest that zebrafish can be used

to identify genes associated with smoking behaviour in humans and that variants in

the *slit3* gene are linked in humans with a disruption of SLIT3 function that may

affect propensity to develop tobacco dependence.

Consistent with previous findings in zebrafish and other species (21), varenicline and

bupropion inhibited nicotine induced place preference in zebrafish. The pattern of

inhibition differed such that varenicline (partial agonist) showed increasing inhibition

at increasing concentrations, whereas inhibition by bupropion (re-uptake inhibitor)

was maximal at 1μM, decreasing as concentration increased. The difference in

response profile presumably reflects differences in the modes of action of these two

compounds.

Our screening of ENU-mutagenized zebrafish families followed by sibling re-screen

is a proof of principle study that indicates the relevance of zebrafish for human studies

and emphasizes the advantage of first using a screen with a low number of individuals

per family to increase efficiency. The classic three generation forward genetic screen

examines phenotypes in groups of 20 or more individuals from each family (51).

Logistical considerations make it difficult to apply such an approach to adult

behavioural screens. Our approach increases efficiency by initially screening a small

number of individuals from a large number of families and only selecting those

families that occur at the extremes of the distribution for further analysis. Although in

this study we were able to confirm phenotypes using a relatively small population of

siblings, re-screening of a larger number would increase the power of the analysis and

allow more subtle phenotypes to be identified.

One limitation of our forward genetic approach is that a large number of genes are

duplicated in zebrafish due to the teleost tetraploidization; ENU-mutagenesis in one

copy may not be sufficient to produce behavioural changes due to genetic

compensation from the other copy of the same gene. *Slit3* is present as a single copy

in the zebrafish genome which may have facilitated our ability to identify its role in

responses to nicotine. However, arguably the most significant consequence of the

teleost tetraploidization is the temporal and spatial specific expression of the gene

duplicates. Spatial and/or temporal differences in gene expression patterns may offset

concerns regarding compensation, and offer great potential to study region-specific

functionality.

We identified a loss of function mutation in the zebrafish *slit3* gene associated with

increased nicotine place preference and confirmed the phenotype in an independent

line. SLIT molecules bind to ROBO receptors through a highly conserved leucine-rich

repeat (LRR) domain (52). In the AJBQM1 (*slit3^{sa1569}*) line the loss of function

mutation causes a truncation at amino acid 176 and in the *slit3^{sa202}* line at amino acid

163. These are immediately adjacent to the LRR2 domain responsible for SLIT3's

functional interaction with ROBO proteins (52) and would therefore be predicted to

lead to formation of non-functional proteins. Initially identified as a family of axon

guidance molecules, SLIT proteins are known to be expressed in a range of tissues

and, by regulating cell polarity, to play major roles in many developmental process

including cell migration, proliferation, adhesion, neuronal topographic map formation

and dendritic spine remodelling (53). *In vitro* SLIT proteins bind promiscuously to

ROBO receptors suggesting that the proteins may co-operate *in vivo* in areas in which

they overlap. However, their restricted spatial distributions, particularly of SLIT3 in

the central nervous system (54) suggest the individual proteins play subtly different

roles in vivo.

Despite its neuronal expression, the most prominent phenotype seen in SLIT3

deficient mice is postnatal diaphragmatic hernia (55,56) with no obvious neuronal or

axon pathfinding defects having been reported. Similarly, we did not detect any major

differences in axon pathfinding nor in number of serotonergic and catecholaminergic

cells in *slit3* mutant zebrafish larvae. As suggested previously (57) it may be that

overlap of expression with other SLIT molecules compensates for loss of SLIT3 in

the brain preventing gross neuronal pathfinding defects. However, subtle differences

in circuit formation and/or axon branching may have escaped our analysis.

As our antibody staining may not have detected subtle, functionally important

differences in dopaminergic and/or serotonergic circuit formation we examined the

impact of the dopaminergic and serotonergic antagonist amisulpride on the acoustic

startle response, a behaviour associated with vulnerability to addiction and known to

be sensitive to modulation by dopaminergic antagonists in zebrafish as well as

mammals (36,58,59). Although the binding affinity of amisulpride in zebrafish has

not been examined, reported behavioural effects (60) are consistent with binding

characteristics and behaviours seen in mammalian species. In mammals, amisulpride

binds to D2/3 pre-and post- synaptic receptors with greater affinity at presynaptic than

post-synaptic receptors (61,62). Presynaptic D2 receptors in mammals act as

autoreceptors and inhibit the synthesis and subsequent release of dopamine. D2/3

postsynaptic receptors are coupled to Gi/o G proteins mediating inhibitory

neurotransmission (63). Treatment of rodents with D2/3 receptor antagonists leads to

biphasic effects on locomotion such that low doses inhibit locomotion via pre-

synaptic D2/3 autoreceptors, and high doses increase locomotion via post-synaptic

D2/3 receptors (64). Treatment of adult zebrafish with amisulpride has a similar

biphasic dose-dependent effect on locomotion such that low doses inhibit locomotion

and high doses increase locomotion (60). These findings suggest that amisulpride has

similar binding affinities at D2/3 receptors in fish as in rodents and imply the

existence of pre-and post- synaptic D2/3 receptors with similar functional properties.

Our finding that high concentrations of amisulpride increased habituation to acoustic

startle in wild type fish is in agreement with the effect of amisulpride in humans (36).

A biphasic dose-dependent effect of amisulpride on habituation in wild type larvae

suggests the involvement of both pre- and post- synaptic dopamine receptors.

Inhibition of presynaptic receptors at low dose leading to increased responsiveness

(reduced habituation), and inhibition of postsynaptic receptors at high doses causing

reduced responsiveness (increased habituation). In contrast to results in wild type fish,

$slit3^{sa1569}$ mutant larvae showed decreased habituation in the presence of both high

and low dose amisulpride, suggesting a reduction of sensitivity to amisulpride at post-

synaptic sites, possibly related to the marginal increase in dopamine D3 receptors in

slit3 mutants. Differential sensitivity to amisulpride was also seen at adult stages,

where amisulpride inhibited nicotine-induced CPP in adult wild type fish but not in

slit3[sa1569] mutants. These findings are consistent with a disrupted dopaminergic

system, and/or disrupted interactions between dopaminergic and serotonergic systems

caused by *slit3* loss of function.

Although our results are consistent with differences in dopaminergic signalling,

differential sensitivity to actions of amisulpride at serotonergic receptors cannot be

ruled out: amisulpride also acts as an antagonist at *Htr7* and *Htr2b* receptors with

affinities approximately four times lower than at dopaminergic receptors. In mice,

acoustic startle is sensitive to inhibition at *Htr2b* sites, and genetic ablation of the

Htr2b receptors induced a reduction in startle amplitude and a deficit in prepulse

inhibition of the startle reflex in loss of function mice (65). Loss of function of *Htr7*

has no effect on acoustic startle or pre-pulse inhibition of acoustic startle in mice (66).

Although gene expression analyses revealed subtle up-regulation in several receptors -

including *drd3, chrnb3, adra2b and htr4* - in *slit3* mutants, significant difference was

only seen for the *ht1aa* receptor subtype. Zebrafish possess two homologues of the

HTR1A gene, *htr1aa* and *htr1ab*, with overlapping expression domains (67). The

observation of increased *htr1aa* expression in *slit3* mutants is of interest: Serotonergic

signalling has been previously linked to drug reward processes including nicotine use

and dependence (68,69). Manipulations which decrease brain serotonin

neurotransmission (e.g., a neurotoxic serotonin depletion or a lasting serotonin

synthesis inhibition) elevate self-administration of several different drugs including

nicotine in rats (69–71). Compounds that facilitate serotonin neurotransmission, such

as selective serotonin reuptake inhibitors, decrease nicotine intake (72) whereas the

HTR1A specific antagonist WAY100635 has also been reported to block nicotine

enhancement of cocaine and methamphetamine self-administration in adolescent rats

(73). Nicotine increases serotonin release in the striatum, hippocampus, cortex, dorsal

raphe′ nucleus (DRN), spinal cord and hypothalamus (74). The effects in the cortex,

hippocampus, and DRN involve stimulation of *Htr1a* receptors, and in the striatum,

Htr3 receptors. In the DRN, *Htr1a* receptors play a role in mediating the anxiolytic

effects of nicotine. In contrast, in the dorsal hippocampus and lateral septum, these

same receptors mediate its anxiogenic effects. Further, pharmacological studies in

rodents have shown that the *Htr1a* receptor antagonists WAY100635 and LY426965

alleviate the anxiety-related behavioural responses induced by nicotine withdrawal

(75–77). Although it is possible that an anxiolytic effect of nicotine contributed to the

increased nicotine-induced place preference, preliminary assessment of anxiety-like

responses (tank diving) in *slit3* mutants, where mutants show decreased anxiety-like

behaviour (n.s), argue against this.

It is perhaps of particular interest that it is a homologue of the HTR1A receptor that is

up-regulated in *slit3* mutants. HTR1A is the major inhibitory serotonergic receptor in

mammalian systems. In mammals, it is present as autoreceptors on cell bodies and

dendrites in the raphe nucleus and as post-synaptic heteroreceptors in brain regions

implicated in mood and anxiety such as prefrontal cortex, hippocampus and

amygdala. Projections from raphe release serotonin throughout the entire forebrain.

and brainstem and modulate a range of activities with additional raphe nuclei also

providing innervation to the midbrain (see (78) for review). Interaction between

HTR1A receptor and dopaminergic signalling are well established. Systemic

administration of HTR1A receptor agonists leads to increased dopamine transmission

in the nigrostrial pathway, ventral tegmental area and frontal cortex (78,79). Although

the mechanism underlying this increased dopamine transmission is not clear in all

areas, the likely mechanism is by action on autoreceptors in the raphe so inhibiting

serotonergic projections and disinhibiting dopaminergic transmission (79). Within the

nucleus accumbens, a key area in drug reward, HTR1A agonists have little effect on

dopamine release under baseline conditions, but inhibit amphetamine induced release

(80). Further interactions between dopaminergic D2 receptor systems and HTR1A

receptor signalling come from studies of atypical neuroleptics for the treatment of

schizophrenia. Considerable evidence suggests that the balance between the properties

of D2 receptors and HTR1A receptors influences the profile of action of these drugs

in preclinical models (81,82). Whilst synergistic effects of atypical neuroleptics may

enhance dopamine release via inhibition of D2 autoreceptors and secondarily via

disinhibition of projections from the raphe, it has been suggested (83) that association

of D2 receptors and HTR1A in functional heterodimers that exhibit properties distinct

to either G protein coupled receptor may also be involved. Such an interaction is seen

between D2 receptors and adenosine A2a receptors both in vitro and in vivo and has

been reported for D2 receptors and HTR1A in vitro (83). Whilst speculative, it is also

potentially of relevance that HTR1A protein expression is upregulated in the brains of

schizophrenics (84) and variants in both *HTR1A* and *SLIT3* are associated with

psychiatric disorders (85–87), known to involve dopaminergic and serotoninergic

pathways.

The mechanism by which loss of function in *slit3* leads to increased *htr1aa* expression

and disrupted dopaminergic signalling in zebrafish mutants is yet to be established.

However, *Htr1a* is up-regulated in conditions of reduced serotonergic signalling in

other systems (88). Loss of serotonergic signalling from the DRN affects dopaminergic axonal outgrowth to the rat medial prefrontal cortex at developmental stages (89) such that lesioning of the DRN at neonatal stages results in significant increase in dopaminergic fibres. These effects are stage specific, raising the possibility that, although we did not detect any differences in catecholamine axon projections at three days post fertilisation, more detailed analysis at later stages of development would reveal significant differences. The observation that *slit3* is strongly expressed in the posterior raphe nucleus overlapping with *htr1aa* expression supports interaction between these two systems. Although detailed co-expression studies have not been performed, each of the other genes found to have marginal changes in expression also show overlapping expression with *slit3* (35,67,90,91).

Thus, our findings of an increase in *ht1aa* expression, altered sensitivity to amisulpride and altered nicotine CPP support a role for *slit3* signalling in the formation of dopaminergic and serotonergic pathways involved in responses to nicotine.

There are limitations to our findings: we used zebrafish of various ages in different experiments. While this confirms that loss-of-function in *slit3* alters behaviour from early life to adulthood, suggesting a developmental role, mechanisms underlying the two behavioural phenotypes may differ. We used whole embryos for the qPCR study so changes in expression in non-neuronal tissue may contribute to the observed differences, further, expression of genes in one tissues may mask changes of expression in another. In addition, we only examined a limited number of receptors and transporters for key neurotransmitter pathways. Important differences in other transmitter pathways and neurotransmitter metabolism may have been missed. We

confirmed the translational effects of *SLIT3* gene variants in a human study, and the

association was validated in a second, independent cohort. The sample sizes used in

the human studies are small in comparison with those used in human discovery

studies, however we used the two human studies to validate the findings in fish and

analysed only a small set of SNPs focussed on just one gene which minimises the

chance of type 1 error. Analyses were also corrected for multiple comparisons. The

Finnish cohort was not used as a formal replication of the findings in the London

cohort but instead to explore the effects of genetic variation on richer and more

informative smoking phenotypes.

Associations between *SLIT3* and aspects of smoking phenotype have also been found

in previous GWAS (https://atlas.ctglab.nl) (92). However, larger studies would be

necessary to obtain greater precision on estimates of the effect size. Further studies

are also required to determine the effects of genetic variation in *SLIT3* on anatomical

pathways in the human brain and their functioning with view to identifying people

who are at high risk of developing dependence. This could be achieved by using

imaging techniques to study brain activation in response to smoking related cues in

smokers who have the *SLIT3* polymorphisms linked to smoking (particularly

rs12654448).

To our knowledge, this is the first report of a forward behavioural genetic screen in

adult zebrafish successfully predicting a novel human coding genetic region involved

in a complex human behavioural trait. Taken together, these results provide evidence

for a role for *SLIT3* in regulating smoking behaviour in humans and confirm adult

zebrafish as a translationally relevant animal model for exploration of addiction-

related behaviours. Further work analysing the cellular processes affected as a result

of the *Slit3* mutation may provide useful targets when designing tailored treatments to

aid smoking cessation.

 MATERIALS AND METHODS

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
Genetic reagent (*danio rerio*)	slit3	Sanger Institute	sa202	Generated by ENU mutagenesis. Soon to be available from ZIRC zebrafish resource centre
Genetic reagent (danio rerio)	slit3	Sanger Institute	sa1569	Generated by ENU mutagenesis. Soon to be available from ZIRC zebrafish resource centre
Strain (danio rerio)	Tupfel	Sanger Institute	Tubingen longfin	Wild type strain, now available from the ZIRC zebrafish resource centre
antibody	anti-5-HT (Rabbit polyclonal)	Sigma	Cat#S5545,	IHC (1/200)
antibody	anti-tyrosine hydroxylase (mouse monoclonal)	Abcam	Cat# AB152	IHC (1/1000))
Chemical compound	Amisulpride	Tocris	C2132	0.05-0.5mg/L

Chemical compound	varenicline	sigma	PZ0004	10-20micromolar
Chemical compound	bupropion	Sigma	B1277	1-10 micromolar
Chemical compound	Nicotine hemisulphate	Sigma	N1019	5-10 micromolar

Animals: All in vivo experimental work was carried out following consultation of the ARRIVE guidelines (NC3Rs, UK). Required sample size was estimated following pilot studies to determine effect sizes, and power calculations (beta = 0.8, alpha = 0.05). All animals were selected at random from groups of conspecifics for testing.

Generation of F3 families of ENU-mutagenised fish: Wild type and ENU-mutagenized Tupfel longfin (TLF) fish were obtained from the Sanger Institute, as part of the Zebrafish Mutation Project which aimed to create a knockout allele in every zebrafish protein-coding gene [https://www.sanger.ac.uk/resources/zebrafish/zmp/]. At the Sanger, ENU-mutagenized TLF F_0 males were outcrossed to create a population of F_1 fish heterozygous for ENU-induced mutations. Due to the high ENU mutation rate (1/300 kb in F1 fish) and homologous recombination when F_1 gametes are generated, all F_2 were heterozygous for multiple mutations. F_2 families, each generated from a separate F_1 fish, were imported from the Sanger Institute to Queen Mary University of London (QMUL).

At QMUL a single male and female fish from each F_2 family were inbred to generate

30 F_3 families that would be 25% wild type, 50% heterozygous and 25% homozygous

mutant for any single mutation, assuming Mendelian genetics. Based on exome

sequencing data from the F_1 generation performed at the Sanger, each F_3 family

contained 10-20 known predicted loss of function exonic mutations, approximately

100 non-synonymous coding mutations and approximately 1500 unknown mutations

in non-coding domains across the entire genome (32). Breeding scheme is detailed in

Figure 10.

Figure 10. Zebrafish breeding scheme to generate F3 families. F_2 ENU-

mutagenized zebrafish, heterozygous for multiple mutations across the entire genome

were obtained from the Wellcome Sanger Institute as part of the Zebrafish Mutation

Project. At Queen Mary University of London, heterozygous F_2 fish were incrossed to

generate 30 F_3 families, each containing 10-20 nonsense or essential splice site

mutations and about 1500 additional exonic and intronic point mutations. F_3 Families

were arbitrarily numbered 1-30. Expected mutation rate and type of mutations in

coding regions are specified on the right hand side.

Fish maintenance:

Fish were housed in a recirculating system (Tecniplast, UK) on a 14h:10h light:dark

cycle (0830–2230). The housing and testing rooms were maintained at ~25–28ºC.

Fish were maintained in aquarium-treated water and fed three times daily with live

artemia (twice daily) and flake food (once). All procedures were carried out under

license in accordance with the Animals (Scientific Procedures) Act, 1986 and under

guidance from the local animal welfare and ethical review board at Queen Mary

University of London.

Conditioned place preference (CPP): All fish were age and weight matched for all

behavioural analysis and were approximately 5 months old, weighing 0.2-0.25g at the

start of testing. Following habituation and determination of basal preference, animals

were conditioned to 5μM nicotine (Sigma, Gillingham, UK Catalogue number:

N1019) over three consecutive days and assessed for a change in place preference the

following day. 5μM nicotine was used because it was predicted to induce a minimum

detectable change in place preference based on results of previous studies (22,24).

This minimal effective dose was used to avoid possible ceiling effects if using a

higher concentration. CPP was assessed as described previously (22,23,93): The

testing apparatus was an opaque 3 L rectangular tank that could be divided in half

with a Perspex divider. Each end of the tank had distinct visual cues (1.5cm diameter

black spots versus vertical 0.5cm wide black and white stripes, matched for

luminosity). After habituation to the apparatus and handling, we determined the basal

preference for each fish: individual fish were placed in the tank for 10 min and the

time spent at either end determined using a ceiling mounted camera and Ethovision

tracking software (Noldus, Wageningen, NL). Any fish showing >70% preference for

either end was excluded from further analysis (between 10 and 20% of fish). Fish

were then conditioned with nicotine in the least preferred environment for 20 min, on

3 consecutive days: Each day each fish was restricted first to its preferred side for 20

min in fish water and then to its least preferred side with nicotine or, if a control fish,

vehicle (fish water) added, for another 20 min. After 20 min in the nicotine (or

vehicle)-paired environment each fish was returned to its home tank. After 3 days of

conditioning, on the following day, fish were subject to a probe trial whereby each

fish was placed in the conditioning tank in the absence of divider and the time spent at

either end of the tank over a 10 min period was determined as for assessment of basal

preference. The change in place preference was determined as the proportion time

spent in the nicotine-paired zone during the probe trial minus the proportion time

spent in the nicotine-paired zone during basal testing. The CPP procedure has been

used and validated previously with nicotine as well as other drugs (22,23,93) .

Data analysis: Change in preference scores were calculated as proportion time spent

in drug paired stimulus after conditioning minus proportion time spent in drug paired

stimulus before conditioning. Population means between generations were compared

using independent two-sample t-tests, and effect-sizes ascertained using Cohen's d

(94). For the rescreen of outlier sibling families and $slit3^{sa202}$ line, mutant lines were

compared with wild type controls using an independent two-sample t-test.

***CPP in the presence or absence of antagonists*:** To assess the ability of compounds

(varenicline (Sigma, Gillingham, UK, PZ0004), buproprion (Sigma, Gillingham, UK,

B1277) or amisulpride (Tocris, Bristol, UK, C2132) to inhibit subjective effects of

nicotine a modified version of the CPP procedure was used (20). In this modified

version, following habituation and establishment of basal preference, each day each

fish was restricted first to its preferred side for 20 min in fish water and then removed

from the conditioning tank and transferred to a tank containing the appropriate

concentration of test compound or fish water (plus carrier where required) for 10min.

After 10min the fish was returned to its least preferred side in the conditioning tank

with nicotine or, if a control fish, vehicle (fish water) for another 20 min. After 20

min in the nicotine (or vehicle)-paired environment, each fish was returned to its

home tank. After three days of conditioning to nicotine in the presence or absence of

test compound, on the following day, fish were subject to a probe trial whereby each

fish was placed in the conditioning tank in the absence of divider and the time spent at

either end of the tank over a 10 min period determined. To assess the ability of

varenicline (0-20μM) or buproprion (0-10μM) to inhibit subjective effects of nicotine

in wild type fish, fish were incubated in the presence and absence of increasing doses

of test compound (or vehicle) for 10 min before conditioning to 10μM nicotine.

Statistical analysis was performed using a univariate analysis of variance (ANOVA),

followed by Tukey's post hoc test.

To test the effect of amisulpride on nicotine-induced CPP in wild type and *Slit3^{sa1569}*

mutant fish, fish were incubated in the presence or absence of 0.5mg/L amisulpride

for 10 min before conditioning to 5μM nicotine. We selected 0.5mg/L amisulpride as

this concentration has been shown to impact behaviour in adults previously (60) and

wild type and homozygous mutants were found to be differentially sensitive to the

effect of this concentration on acoustic startle response. Although previous studies

with amisulpride in adult zebrafish used a pre-incubation period of 30min, to be

consistent with our analysis of inhibitory effects of varenicline and bupropion, and to

reduce the possibility of prolonged confinement in a small volume of drug affecting

stress levels which may confound results, we used a preincubation period of 10 min.

Two-way ANOVA was performed with genotype ($slit3^{+/+}$ and $slit3^{sa1569/sa1569}$) and

treatment (control, nicotine, nicotine+amisulpride) as independent variables. Values

of $p<0.05$ were considered significant.

Breeding and selection to assess heritability of nicotine-induced place preference:

To test whether nicotine preference is heritable, fish falling in the upper and lower

deciles of the 'change in place preference' distribution were kept for analysis and

further breeding. Individuals were bred (in-cross of fish from the upper decile and in-

cross of fish from the lower decile done separately) and their offspring screened for

CPP (Second Generation CPP analysis). The same approach was repeated again: fish

at the extremes of the Second Generation CPP distribution curve were selected and in-

crossed, and their offspring were used to perform a Third Generation CPP analysis.

Identification of ENU-induced mutations influencing nicotine place preference: To

investigate whether ENU-induced mutations affect fish sensitivity to the rewarding

effects of nicotine, candidate families were selected when the 3-4 fish from a family

tested clustered together at one or other extreme of the change in preference

distribution curve. To confirm the genetic effect on the CPP phenotype in candidate

families, all remaining siblings of that family were assessed for nicotine induced CPP,

along with non-mutagenized TLF control fish, to confirm the genetic effect on the

phenotype. The experimental team and technical staff were blind to the fish genotype.

Candidate mutations, obtained from exome sequencing on F_1 fish, were assessed for

co-segregation with behaviour using site specific PCR (95) (See Supplementary

Methods). Once a co-segregating candidate mutation was identified, larvae from an

independent line carrying a predicted loss of function allele in the same gene were

obtained from the Sanger Institute (*slit3*^{sa202}) to confirm the association. Heterozygous

slit3$^{sa202/+}$ and sibling *slit3*$^{+/+}$ larvae were reared to adulthood and assessed for

nicotine-induced CPP as described above. All fish were fin clipped and genotyped

following CPP.

Characterization of larvae:

Antibody staining: In order to visualize axonal pathways, fluorescent

immunohistochemistry was carried out in three day old embryos from wild type

slit3$^{+/+}$, and homozygous mutant *slit3*$^{sa1569/sa1569}$ in-crosses. To prevent skin

pigmentation, embryos were incubated in 0.2mM of 1-phenyl 2-thiourea (Sigma,

Gillingham, UK) from 24 hours after fertilization. At three days, they were fixed in

4% paraformaldehyde (Sigma, Gillingham, UK) to avoid tissue degradation. For the

immunostaining, rabbit polyclonal anti-tyrosine hydroxylase primary antibody (1:200;

Sigma, Gillingham, AB152), rabbit polyclonal anti-serotonin (5HT) antibody (1:100,

Sigma, Gillingham, S5545) and mouse anti-acetylated tubulin monoclonal antibody

(1:1000; Sigma Gillingham, UK, T6793) were used. The three primary antibodies

were detected with Alexa 546-conjugated secondary antibodies (1:400; Fisher

Scientific, Loughborough, UK A11010). Whole-mount immunohistochemistry and

mounting was performed as described previously (96).

Confocal microscopy imaging and analysis: Images were acquired using a Leica SP5

confocal microscope. Confocal z-stacks were recorded under the same conditions

using diode laser and images were processed under ImageJ environment. Areas of

interest for quantification were isolated, making sure that for all the individuals the

same number of Z stacks, (covering the same D/V distance) were included. The

number of cells was quantified using the ImageJ plugin "3D Objects counter"

(https://biii.eu/3d-objects-counter). To quantify the number of catecholaminergic

axons crossing the midline, a line was drawn from the Medulla oblongata

interfascicular zone and vagal area to the locus coeruleus (Figure 5-figure

supplement1). Every 10 stacks (~7.6 microns), the number of intensity peaks (defined

as grey value intensity >20) was measured. Unpaired t-tests were calculated to assess

genotype differences in the number of cells and intensity peaks.

Startle response in the presence or absence of amisulpride: Five day old larvae,

generated from adult *slit3* wild type and homozygous mutant (*slit3$^{sa1569/sa1569}$*) fish as

for quantitative PCR, were individually placed in 24 well plates. In the drug-free

condition, each well contained 300μl system water and 0.05% of dimethyl sulfoxide

(DMSO, Sigma, Gillingham, UK). In the pharmacological conditions, serial dilutions

of the dopaminergic and serotonergic antagonist amisulpride (Tocris, Bristol, UK,

71675-86-9) were prepared to give final concentrations of 0.05 mg/L, 0.1 mg/L or 0.5

mg/L amisulpride in 0.05% DMSO. Amisulpride concentrations were chosen based

on previous studies in zebrafish (60) and correspond to 50, 100 and 500 times its Ki

value for the D2 receptor in mammals (61,62). To ensure that larvae were exposed to

the drug for the same amount of time, amisulpride was added 15 minutes before

undertaking the experiment. Care was taken regarding the distribution of

concentrations and genotypes to ensure that experimental groups were randomly

distributed in the plates. Plates were placed in a custom-made filming tower with a

tapping device that applied 10 sound/vibration stimuli with two seconds interval

between them. The setup for this device has been described elsewhere (97). Larval movement was recorded using Ethovision XT software (Noldus Information Technology, Wageningen, NL) and data were outputted in one second time-bins. Three technical replicates were performed (three different days) with three 24 well plates assayed each day.

For each fish, distance travelled (mm) during one second after each tap was recorded. All videos used to evaluate responses were checked for tracking errors. Any points where tracking errors were detected were removed. In total 105 points (31 wild type, 33 heterozygous, 41 homozygous) out of 5,460 were removed due to tracking errors. We also excluded individuals that failed to respond to all 10 taps using the criteria defined below. 39 wild type, 38 heterozygous, and 39 homozygous individuals were removed (a total of 115 individuals out of 546).

To evaluate how amisulpride dose and genotype affect habituation to the startle stimulus, we defined a threshold of distance moved: if a fish moved more than this threshold distance, we defined it as a 'response' (binary variable). We established our threshold by calculating the maximum distance each individual moved over all 10 taps. We ascertained that genotype was not a significant predictor of maximum distance through a linear model with maximum distance each individual moved as the response variable and genotype as the explanatory variable (ANOVA, $F_{2,543} = 0.10$, p=0.902). We therefore calculated the mean of the population maximum distance and defined our threshold as 70% of the mean maximum distance.

The percentage of fish responding to stimulus together with amisulpride dose, stimulus event number and genotype group were modelled in a beta regression conducted using the R package "betareg". The proportion of individuals responding

was the response variable and the three-way interaction between amisulpride dose,

genotype, and stimulus event number was the explanatory variable. To determine

whether this interaction was a significant predictor of individual responsiveness

(indicating that genotypes varied in how amisulpride dose affected their habituation to

repeated stimuli), likelihood ratio tests for nested regression models were performed.

Results of all statistical analyses were reported with respect to a type-1 error rate of

$\alpha=0.05$.

Real-time quantitative PCR: Adult *slit3* wild type and *slit3^{sa1569}* homozygous mutant

fish, generated from a *slit3$^{sa1569/+}$* heterozygous in cross, were bred to generate

homozygous wild type, heterozygous mutant and homozygous mutant larvae.

Embryos were carefully staged at 1, 24 and 48 hour and at five day post fertilisation

to ensure, based on morphological criteria, there were no differences in development

between groups. mRNA from 3 samples of five day old embryos (n=10 pooled

embryos per sample) for each genotype was isolated using the phenol-chloroform

method. cDNA was generated using the ProtoScript® II First Strand cDNA Synthesis

Kit (NEB (UK Ltd.), Hitchen, UK). Relative qPCR assays were performed using the

LightCycler 480 qPCR system from Roche Diagnostics, Ltd. with all reactions carried

out in triplicates. Reference genes for all the qPCR analyses were β-actin, ef1α and

rpl13α based on previous studies (93,98,99). Accession numbers and primer

sequences for the genes can be found in Supplementary Table 2.

Relative mRNA expression in qPCR was calculated against reference gene cycle-

threshold (Ct) values, and then subjected to one-way ANOVA. To account for

multiple testing a Bonferroni correction was applied, and significance was declared at

a threshold of 0.001.

Human Cohorts: In London human subjects were recruited from three clinical

groups: patients with chronic obstructive pulmonary disease (COPD) (Cohort 1;

n=272); patients with asthma (Cohort 2; n=293); and residents and carers in sheltered

accommodation, with neither condition (Cohort 3; n=298). The methods used for

recruitment and definition of phenotypes are reported elsewhere (100–102). The

studies were approved by East London and The City Research Ethics Committee 1

(09/H0703/67, 09/H0703/76 and 09/H0703/112). Written informed consent was

obtained from all participants.

Details of the Finnish twin cohort are reported elsewhere (103–105). In brief, twin

pairs concordant for moderate to heavy smoking were identified from the population-

based Finnish Twin Cohort survey responders. The twin pairs and their siblings were

invited to a computer-assisted, telephone-based, structured, psychiatric interview

(SSAGA) (103), to yield detailed information on smoking behaviour and nicotine

dependence as defined by Fagerström Test for Nicotine Dependence (FTND) and

DSM-IV diagnoses. Human phenotypes to be investigated in relation to zebrafish

nicotine seeking behaviour were determined by consensus *a priori*.

Sample characteristics of the human cohorts and detailed definitions of both London

and Finnish phenotypes can be found in the supplementary material and

Supplementary Table 3.

Human genotyping: For the London cohorts, DNA from participants was extracted

from whole blood using the salting-out method (106) and normalized to 5ng/µl. 10ng

DNA was used as template for 2 µl TaqMan assays (Applied Biosystems, Foster City,

CA, USA) performed on the ABI 7900HT platform in 384-well format and analysed

with Autocaller software. Pre-developed assays were used to type all SNPs. See

Supplementary Table 4 for primer and reporter sequences. Typing for two SNP

(rs6127118 and rs11574010) failed. For the Finnish cohort, DNA was extracted from

whole blood and genotyping was performed at the Wellcome Trust Sanger Institute

(Hinxton, UK) on the Human670-QuadCustom Illumina BeadChip (Illumina, Inc.,

San Diego, CA, USA), as previously described (103–105). Genotyping and

imputation for the Finnish cohort at the Wellcome Trust Sanger Centre have been

described previously (107).

Human association analyses: We attempted to replicate the zebrafish findings

initially in a cohort from London using a narrow set of SNPs in *SLIT3*. We then used

the same set of SNPS to evaluate effects on more detailed smoking phenotypes in a

Finnish twin cohort.

London cohort association analysis was performed using PLINK v1.07 (108). *SLIT3*

SNPs that had been previously associated with disease phenotype were identified and

the 20 with low linkage disequilibrium score selected for analysis. Of twenty *SLIT3*

SNPs, one departed from Hardy-Weinberg equilibrium (rs13183458) and was

excluded. Linear regression was performed on average number of cigarettes smoked

per day, controlling for age, sex and cohort. Since physiological and genetic

mechanisms may be different in heavy (more dependent) and light (less dependent)

smokers we repeated on heavy smokers ($\geq$ 20 cigarettes per day) and light smokers ($<$

20 cigarettes per day). Smoking cessation (current *vs* ever smokers) was analysed

using logistic regression controlling for age, sex and cohort. All analyses were

performed under the additive genetic model and multiple testing was taken into

account using the Benjamini-Hochberg adjustment. Only individuals from European

ancestry were included in analyses.

Association analyses for the Finnish Twin Cohort were performed using GEMMA

v0.94 (109) with linear mixed model against the allelic dosages controlling for age

and sex. Sample relatedness and population stratification were taken into account by

using genetic relatedness matrix as random effect of the model.

Acknowledgements

Funding: MRC UK, G1000403 (CHB/RW). NC3Rs G1000053 (CHB). BBSRC BB/M007863 (CHB) NIH U01 DA044400-03 (CHB) NIHR PGfAR RP-PG-0609-10181 (RW). NIHR PGfAR, RP-PG-0407-10398 (ARM). CHB is a member of the Royal Society Industry Fellows' College. RW is an NIHR Senior investigator (NF-SI-0515-10076). VK holds a Wellcome Trust Clinical Research Fellowship. JK is supported by the Academy of Finland (grants 308248, 312073). CHB conceived the project and directed the research. RW, AM and JK oversaw the human studies. AJB, JGG, DJ, MTT, AS, MOP, CHB conducted experiments. TP, VK analysed the human data. EMB and DS supplied the mutant lines. Some of the data in this manuscript has been published previously as a preprint: bioRxiv 453928: doi: https://doi.org/10.1101/453928

Competing interests

The authors of this manuscript certify that they have NO affiliations with or involvement in any organization or entity with any financial interest or non-financial interest in the subject matter or materials discussed in this manuscript

References

1. World Health Organization. WHO report on the global tobacco epidemic, 2017: monitoring tobacco use and prevention policies. 2017.

2. National Center for Chronic Disease Prevention and Health Promotion (US) Office on Smoking and Health. The Health Consequences of Smoking—50 Years of Progress: A Report of the Surgeon General [Internet]. Atlanta (GA): Centers for Disease Control and Prevention (US); 2014. (Reports of the Surgeon General).

3. Xu X, Bishop EE, Kennedy SM, Simpson SA, Pechacek TF. Annual healthcare spending attributable to cigarette smoking: an update. Am J Prev Med. 2015 Mar;48(3):326–33.

4. Munafò M, Clark T, Johnstone E, Murphy M, Walton R. The genetic basis for smoking behavior: a systematic review and meta-analysis. Nicotine Tob Res. 2004 Aug;6(4):583–97.

5. Batra V, Patkar AA, Berrettini WH, Weinstein SP, Leone FT. The genetic determinants of smoking. Chest. 2003 May;123(5):1730–9.

6. Liu M, Jiang Y, Wedow R, Li Y, Brazel DM, Chen F, et al. Association studies of up to 1.2 million individuals yield new insights into the genetic etiology of tobacco and alcohol use. Nat Genet. 2019 Jan 14;

7. Erzurumluoglu AM, Liu M, Jackson VE, Barnes DR, Datta G, Melbourne CA, et al. Meta-analysis of up to 622,409 individuals identifies 40 novel smoking behaviour associated genetic loci. Mol Psychiatry. 2019 Jan 7;

8. Chen L-S, Hung RJ, Baker T, Horton A, Culverhouse R, Saccone N, et al. CHRNA5 Risk Variant Predicts Delayed Smoking Cessation and Earlier Lung Cancer Diagnosis—A Meta-Analysis. J Natl Cancer Inst. 2015 Apr 13;107(5).

9. Darland T, Dowling JE. Behavioral screening for cocaine sensitivity in mutagenized zebrafish. Proc Natl Acad Sci USA. 2001 Sep 25;98(20):11691–6.

10. Brennan CH, Parker MO. Zebrafish (Danio rerio) models of substance abuse: Harnessing the capabilities. Behaviour. 2012;149(10–12):1037–62.

11. Engleman EA, Katner SN, Neal-Beliveau BS. Caenorhabditis elegans as a Model to Study the Molecular and Genetic Mechanisms of Drug Addiction. Prog Mol Biol Transl Sci. 2016;137:229–52.

12. Shipley AT, Imeh-Nathaniel A, Orfanakos VB, Wormack LN, Huber R, Nathaniel TI. The Sensitivity of the Crayfish Reward System to Mammalian Drugs of Abuse. Front Physiol. 2017;8.

13. Berridge KC, Kringelbach ML. Affective neuroscience of pleasure: reward in humans and animals. Psychopharmacology (Berl). 2008 Aug;199(3):457–80.

14. Tzschentke TM. Measuring reward with the conditioned place preference paradigm: a comprehensive review of drug effects, recent progress and new issues. Prog Neurobiol. 1998 Dec;56(6):613–72.

15. Ninkovic J, Folchert A, Makhankov YV, Neuhauss SCF, Sillaber I, Straehle U, et al. Genetic identification of AChE as a positive modulator of addiction to the psychostimulant D-amphetamine in zebrafish. Journal of Neurobiology. 2006 Apr;66(5):463–75.

16. Petzold AM, Balciunas D, Sivasubbu S, Clark KJ, Bedell VM, Westcot SE, et al. Nicotine response genetics in the zebrafish. PNAS. 2009 Nov 3;106(44):18662–7.

17. Zirger JM, Beattie CE, McKay DB, Thomas Boyd R. Cloning and expression of zebrafish neuronal nicotinic acetylcholine receptors. Gene Expression Patterns. 2003 Dec 1;3(6):747–54.

18. Ackerman KM, Nakkula R, Zirger JM, Beattie CE, Boyd RT. Cloning and spatiotemporal expression of zebrafish neuronal nicotinic acetylcholine receptor alpha 6 and alpha 4 subunit RNAs. Dev Dyn. 2009 Apr;238(4):980–92.

19. Pedersen JE, Bergqvist CA, Larhammar D. Evolution of vertebrate nicotinic acetylcholine receptors. BMC Evolutionary Biology. 2019 Jan 30;19(1):38.

20. Papke RL, Ono F, Stokes C, Urban J, Boyd RT. The nicotinic acetylcholine receptors of zebrafish and an evaluation of pharmacological tools used for their study. Biochem Pharmacol. 2012 Aug 1;84(3):352–65.

21. Ponzoni L, Braida D, Pucci L, Andrea D, Fasoli F, Manfredi I, et al. The cytisine derivatives, CC4 and CC26, reduce nicotine-induced conditioned place preference in zebrafish by acting on heteromeric neuronal nicotinic acetylcholine receptors. Psychopharmacology (Berl). 2014 Dec;231(24):4681–93.

22. Kily LJM, Cowe YCM, Hussain O, Patel S, McElwaine S, Cotter FE, et al. Gene expression changes in a zebrafish model of drug dependency suggest conservation of neuro-adaptation pathways. J Exp Biol. 2008 May;211(Pt 10):1623–34.

23. Brock AJ, Goody SMG, Mead AN, Sudwarts A, Parker MO, Brennan CH. Assessing the Value of the Zebrafish Conditioned Place Preference Model for Predicting Human Abuse Potential. J Pharmacol Exp Ther. 2017 Oct;363(1):66–79.

24. Kedikian X, Faillace MP, Bernabeu R. Behavioral and Molecular Analysis of Nicotine-Conditioned Place Preference in Zebrafish. PLOS ONE. 2013 Jul 24;8(7):e69453.

25. Loeber S, Croissant B, Nakovics H, Zimmer A, Georgi A, Klein S, et al. The startle reflex in alcohol-dependent patients: changes after cognitive-behavioral therapy and predictive validity for drinking behavior. A pilot study. Psychother Psychosom. 2007;76(6):385–90.

26. Vrana SR, Calhoun PS, Dennis MF, Kirby AC, Beckham JC. Acoustic startle and prepulse inhibition predict smoking lapse in posttraumatic stress disorder. J Psychopharmacol. 2015 Oct 1;29(10):1070–6.

27. Kumari V, Gray JA. Smoking withdrawal, nicotine dependence and prepulse inhibition of the acoustic startle reflex. Psychopharmacology. 1999 Jan 1;141(1):11–5.

28. Hall BJ, Slade S, Wells C, Rose JE, Levin ED. Bupropion–varenicline interactions and nicotine self-administration behavior in rats. Pharmacology Biochemistry and Behavior. 2015 Mar 1;130:84–9.

29. O'Connor EC, Parker D, Rollema H, Mead AN. The α4β2 nicotinic acetylcholine-receptor partial agonist varenicline inhibits both nicotine self-administration following repeated dosing and reinstatement of nicotine seeking in rats. Psychopharmacology. 2010 Feb 1;208(3):365–76.

30. Le Foll B, Chakraborty-Chatterjee M, Lev-Ran S, Barnes C, Pushparaj A, Gamaleddin I, et al. Varenicline decreases nicotine self-administration and cue-induced reinstatement of nicotine-seeking behaviour in rats when a long pretreatment time is used. Int J Neuropsychopharmacol. 2012 Oct 1;15(9):1265–74.

31. Brennan CH, Parmar A, Kily LKM, Ananthathevan A, Doshi A, Patel S. Conditioned Place Preference Models of Drug Dependence and Relapse to Drug Seeking: Studies with Nicotine and Ethanol. In: Kalueff AV, Cachat JM, editors. Zebrafish Models in Neurobehavioral Research. Totowa, NJ: Humana Press; 2011. p. 163–79. (Neuromethods).

32. Kettleborough RNW, Busch-Nentwich EM, Harvey SA, Dooley CM, Bruijn E de, Eeden F van, et al. A systematic genome-wide analysis of zebrafish protein-coding gene function. Nature. 2013 Apr;496(7446):494–7.

33. Brose K, Bland KS, Wang KH, Arnott D, Henzel W, Goodman CS, et al. Slit Proteins Bind Robo Receptors and Have an Evolutionarily Conserved Role in Repulsive Axon Guidance. Cell. 1999 Mar 19;96(6):795–806.

34. Smidt MP, Burbach JPH. How to make a mesodiencephalic dopaminergic neuron. Nat Rev Neurosci. 2007 Jan;8(1):21–32.

35. Miyasaka N, Sato Y, Yeo S-Y, Hutson LD, Chien C-B, Okamoto H, et al. Robo2 is required for establishment of a precise glomerular map in the zebrafish olfactory system. Development. 2005 Mar 15;132(6):1283–93.

36. Quednow BB, Wagner M, Westheide J, Beckmann K, Bliesener N, Maier W, et al. Sensorimotor gating and habituation of the startle response in schizophrenic patients randomly treated with amisulpride or olanzapine. Biol Psychiatry. 2006 Mar 15;59(6):536–45.

37. Rankin CH, Abrams T, Barry RJ, Bhatnagar S, Clayton D, Colombo J, et al. Habituation Revisited: An Updated and Revised Description of the Behavioral Characteristics of Habituation. Neurobiol Learn Mem. 2009 Sep;92(2):135–8.

38. Huang W, Ma JZ, Payne TJ, Beuten J, Dupont RT, Li MD. Significant association of DRD1 with nicotine dependence. Hum Genet. 2008 Mar;123(2):133–40.

39. Swan GE, Valdes AM, Ring HZ, Khroyan TV, Jack LM, Ton CC, et al. Dopamine receptor DRD2 genotype and smoking cessation outcome following treatment with bupropion SR. Pharmacogenomics J. 2005;5(1):21–9.

40. Huang W, Payne TJ, Ma JZ, Li MD. A functional polymorphism, rs6280, in DRD3 is significantly associated with nicotine dependence in European-American smokers. Am J Med Genet B Neuropsychiatr Genet. 2008 Oct 5;147B(7):1109–15.

41. Sieminska A, Buczkowski K, Jassem E, Niedoszytko M, Tkacz E. Influences of polymorphic variants of DRD2 and SLC6A3 genes, and their combinations on smoking in Polish population. BMC Med Genet. 2009 Sep 17;10:92.

42. Forget B, Wertheim C, Mascia P, Pushparaj A, Goldberg SR, Le Foll B. Noradrenergic alpha1 receptors as a novel target for the treatment of nicotine addiction. Neuropsychopharmacology. 2010 Jul;35(8):1751–60.

43. Kotagale NR, Taksande BG, Gahane AY, Ugale RR, Chopde CT. Repeated agmatine treatment attenuates nicotine sensitization in mice: modulation by alpha2-adrenoceptors. Behav Brain Res. 2010 Dec 1;213(2):161–74.

44. Swan GE, Hops H, Wilhelmsen KC, Lessov-Schlaggar CN, Cheng LS-C, Hudmon KS, et al. A genome-wide screen for nicotine dependence susceptibility loci. Am J Med Genet B Neuropsychiatr Genet. 2006 Jun 5;141B(4):354–60.

45. Kenny PJ, File SE, Rattray M. Nicotine regulates 5-HT(1A) receptor gene expression in the cerebral cortex and dorsal hippocampus. Eur J Neurosci. 2001 Mar;13(6):1267–71.

46. Levin ED, Slade S, Johnson M, Petro A, Horton K, Williams P, et al. Ketanserin, a 5-HT2 receptor antagonist, decreases nicotine self-administration in rats. Eur J Pharmacol. 2008 Dec 14;600(1–3):93–7.

47. de Bruin NMWJ, McCreary AC, van Loevezijn A, de Vries TJ, Venhorst J, van Drimmelen M, et al. A novel highly selective 5-HT6 receptor antagonist attenuates ethanol and nicotine seeking but does not affect inhibitory response control in Wistar rats. Behav Brain Res. 2013 Jan 1;236(1):157–65.

48. Hauser SR, Hedlund PB, Roberts AJ, Sari Y, Bell RL, Engleman EA. The 5-HT7 receptor as a potential target for treating drug and alcohol abuse. Front Neurosci. 2014;8:448.

49. Bloch B, Reshef A, Cohen T, Tafla A, Gathas S, Israel S, et al. Preliminary effects of bupropion and the promoter region (HTTLPR) serotonin transporter (SLC6A4)

polymorphism on smoking behavior in schizophrenia. Psychiatry Res. 2010 Jan 30;175(1–2):38–42.

50. Kaprio J. Twin Studies in Finland 2006. Twin Research and Human Genetics. 2006 Dec;9(6):772–7.

51. Lawson ND, Wolfe SA. Forward and Reverse Genetic Approaches for the Analysis of Vertebrate Development in the Zebrafish. Developmental Cell. 2011 Jul 19;21(1):48–64.

52. Morlot C, Thielens NM, Ravelli RBG, Hemrika W, Romijn RA, Gros P, et al. Structural insights into the Slit-Robo complex. PNAS. 2007 Sep 18;104(38):14923–8.

53. Blockus H, Chédotal A. The multifaceted roles of Slits and Robos in cortical circuits: from proliferation to axon guidance and neurological diseases. Current Opinion in Neurobiology. 2014 Aug 1;27:82–8.

54. Marillat V, Cases O, Nguyenf-Ba-Charvet KT, Tessier-Lavigne M, Sotelo C, Chédotal A. Spatiotemporal expression patterns of slit and robo genes in the rat brain. Journal of Comparative Neurology. 2002 Jan 7;442(2):130–55.

55. Yuan W, Rao Y, Babiuk RP, Greer J, Wu JY, Ornitz DM. A genetic model for a central (septum transversum) congenital diaphragmatic hernia in mice lacking Slit3. PNAS. 2003 Apr 29;100(9):5217–22.

56. Liu J, Zhang L, Wang D, Shen H, Jiang M, Mei P, et al. Congenital diaphragmatic hernia, kidney agenesis and cardiac defects associated with Slit3-deficiency in mice. Mechanisms of Development. 2003 Sep 1;120(9):1059–70.

57. Long H, Sabatier C, Le Ma, Plump A, Yuan W, Ornitz DM, et al. Conserved Roles for Slit and Robo Proteins in Midline Commissural Axon Guidance. Neuron. 2004 Apr 22;42(2):213–23.

58. Halberstadt AL, Geyer MA. Habituation and sensitization of acoustic startle: opposite influences of dopamine D1 and D2-family receptors. Neurobiol Learn Mem. 2009 Sep;92(2):243–8.

59. Burgess HA, Granato M. Sensorimotor gating in larval zebrafish. J Neurosci. 2007 May 2;27(18):4984–94.

60. Tran S, Nowicki M, Muraleetharan A, Gerlai R. Differential effects of dopamine D1 and D 2/3 receptor antagonism on motor responses. Psychopharmacology (Berl). 2015 Feb;232(4):795–806.

61. Schoemaker H, Claustre Y, Fage D, Rouquier L, Chergui K, Curet O, et al. Neurochemical characteristics of amisulpride, an atypical dopamine D2/D3 receptor antagonist with both presynaptic and limbic selectivity. J Pharmacol Exp Ther. 1997 Jan;280(1):83–97.

62. Perrault G, Depoortere R, Morel E, Sanger DJ, Scatton B. Psychopharmacological profile of amisulpride: an antipsychotic drug with presynaptic D2/D3 dopamine receptor antagonist activity and limbic selectivity. J Pharmacol Exp Ther. 1997 Jan;280(1):73–82.

63. Beaulieu J-M, Gainetdinov RR. The physiology, signaling, and pharmacology of dopamine receptors. Pharmacol Rev. 2011 Mar;63(1):182–217.

64. Millan MJ, Seguin L, Gobert A, Cussac D, Brocco M. The role of dopamine D3 compared with D2 receptors in the control of locomotor activity: a combined behavioural and neurochemical analysis with novel, selective antagonists in rats. Psychopharmacology (Berl). 2004 Jul;174(3):341–57.

65. Pitychoutis PM, Belmer A, Moutkine I, Adrien J, Maroteaux L. Mice Lacking the Serotonin Htr2B Receptor Gene Present an Antipsychotic-Sensitive Schizophrenic-Like Phenotype. Neuropsychopharmacology. 2015 Nov;40(12):2764–73.

66. Semenova S, Geyer MA, Sutcliffe JG, Markou A, Hedlund PB. Inactivation of the 5-HT7 Receptor Partially Blocks Phencyclidine-Induced Disruption of Prepulse Inhibition. Biological Psychiatry. 2008 Jan 1;63(1):98–105.

67. Norton WHJ, Folchert A, Bally-Cuif L. Comparative analysis of serotonin receptor (HTR1A/HTR1B families) and transporter (slc6a4a/b) gene expression in the zebrafish brain. J Comp Neurol. 2008 Dec 1;511(4):521–42.

68. Fletcher PJ, Lê AD, Higgins GA. Serotonin receptors as potential targets for modulation of nicotine use and dependence. In: Di Giovann G, Di Matteo V, Esposito E, editors. Progress in Brain Research. 2008. vol. 172p. 361–83.

69. Olausson P, Engel JA, Söderpalm B. Involvement of serotonin in nicotine dependence: Processes relevant to positive and negative regulation of drug intake. Pharmacology Biochemistry and Behavior. 2002 Apr 1;71(4):757–71.

70. Roberts DCS, Loh EA, Baker GB, Vickers G. Lesions of central serotonin systems affect responding on a progressive ratio schedule reinforced either by intravenous cocaine or by food. Pharmacology Biochemistry and Behavior. 1994 Sep 1;49(1):177–82.

71. LeMarquand D, Pihl RO, Benkelfat C. Serotonin and alcohol intake, abuse, and dependence: Findings of animal studies. Biological Psychiatry. 1994 Sep 15;36(6):395–421.

72. Opitz K, Weischer ML. Volitional oral intake of nicotine in tupaias: drug-induced alterations. Drug Alcohol Depend. 1988 May;21(2):99–104.

73. Dao JM, McQuown SC, Loughlin SE, Belluzzi JD, Leslie FM. Nicotine alters limbic function in adolescent rat by a 5-HT1A receptor mechanism. Neuropsychopharmacology. 2011 Jun;36(7):1319–31.

74. Seth P, Cheeta S, Tucci S, File SE. Nicotinic–serotonergic interactions in brain and behaviour. Pharmacology Biochemistry and Behavior. 2002 Apr 1;71(4):795–805.

75. Rasmussen K, Kallman MJ, Helton DR. Serotonin-1A antagonists attenuate the effects of nicotine withdrawal on the auditory startle response. Synapse. 1997 Oct;27(2):145–52.

76. Rasmussen K, Calligaro DO, Czachura JF, Dreshfield-Ahmad LJ, Evans DC, Hemrick-Luecke SK, et al. The novel 5-Hydroxytryptamine(1A) antagonist LY426965: effects on nicotine withdrawal and interactions with fluoxetine. J Pharmacol Exp Ther. 2000 Aug;294(2):688–700.

77. Harrison AA, Liem YT, Markou A. Fluoxetine combined with a serotonin-1A receptor antagonist reversed reward deficits observed during nicotine and amphetamine withdrawal in rats. Neuropsychopharmacology. 2001 Jul;25(1):55–71.

78. Garcia-Garcia AL, Newman-Tancredi A, Leonardo ED. 5-HT(1A) [corrected] receptors in mood and anxiety: recent insights into autoreceptor versus heteroreceptor function. Psychopharmacology (Berl). 2014 Feb;231(4):623–36.

79. Bantick RA, Deakin JF, Grasby PM. The 5-HT1A receptor in schizophrenia: a promising target for novel atypical neuroleptics? J Psychopharmacol (Oxford). 2001 Mar;15(1):37–46.

80. Ichikawa J, Kuroki T, Kitchen MT, Meltzer HY. R(+)-8-OH-DPAT, a 5-HT1A receptor agonist, inhibits amphetamine-induced dopamine release in rat striatum and nucleus accumbens. Eur J Pharmacol. 1995 Dec 12;287(2):179–84.

81. Newman-Tancredi A. The importance of 5-HT1A receptor agonism in antipsychotic drug action: rationale and perspectives. Curr Opin Investig Drugs. 2010 Jul;11(7):802–12.

82. Newman-Tancredi A, Kleven MS. Comparative pharmacology of antipsychotics possessing combined dopamine D2 and serotonin 5-HT1A receptor properties. Psychopharmacology (Berl). 2011 Aug;216(4):451–73.

83. Łukasiewicz S, Błasiak E, Szafran-Pilch K, Dziedzicka-Wasylewska M. Dopamine D2 and serotonin 5-HT1A receptor interaction in the context of the effects of antipsychotics - in vitro studies. J Neurochem. 2016;137(4):549–60.

84. Carrard A, Salzmann A, Malafosse A, Karege F. Increased DNA methylation status of the serotonin receptor 5HTR1A gene promoter in schizophrenia and bipolar disorder. J Affect Disord. 2011 Aug;132(3):450–3.

85. Shi Y, Zhao X, Yu L, Tao R, Tang J, La Y, et al. Genetic Structure Adds Power to Detect Schizophrenia Susceptibility at SLIT3 in the Chinese Han Population. Genome Res. 2004 Jul;14(7):1345–9.

86. Glessner JT, Wang K, Sleiman PMA, Zhang H, Kim CE, Flory JH, et al. Duplication of the SLIT3 locus on 5q35.1 predisposes to major depressive disorder. PLoS ONE. 2010 Dec 1;5(12):e15463.

87. Cukier HN, Dueker ND, Slifer SH, Lee JM, Whitehead PL, Lalanne E, et al. Exome sequencing of extended families with autism reveals genes shared across neurodevelopmental and neuropsychiatric disorders. Mol Autism. 2014 Jan 10;5(1):1.

88. Chen CP, Alder JT, Bray L, Kingsbury AE, Francis PT, Foster OJ. Post-synaptic 5-HT1A and 5-HT2A receptors are increased in Parkinson's disease neocortex. Ann N Y Acad Sci. 1998 Dec 15;861:288–9.

89. Benes FM, Taylor JB, Cunningham MC. Convergence and plasticity of monoaminergic systems in the medial prefrontal cortex during the postnatal period: implications for the development of psychopathology. Cereb Cortex. 2000 Oct;10(10):1014–27.

90. Boehmler W, Obrecht-Pflumio S, Canfield V, Thisse C, Thisse B, Levenson R. Evolution and expression of D2 and D3 dopamine receptor genes in zebrafish. Developmental Dynamics. 2004 Jul 1;230(3):481–93.

91. Ackerman KM, Boyd RT. Analysis of Nicotinic Acetylcholine Receptor (nAChR) Gene Expression in Zebrafish (Danio rerio) by In Situ Hybridization and PCR. In: Li MD, editor. Nicotinic Acetylcholine Receptor Technologies. New York, NY: Springer; 2016. p. 1–31. (Neuromethods).

92. Watanabe K, Stringer S, Frei O, Umićević Mirkov M, de Leeuw C, Polderman TJC, et al. A global overview of pleiotropy and genetic architecture in complex traits. Nat Genet. 2019;51(9):1339–48.

93. Parker MO, Evans AM-D, Brock AJ, Combe FJ, Teh M-T, Brennan CH. Moderate alcohol exposure during early brain development increases stimulus-response habits in adulthood. Addict Biol. 2016 Jan;21(1):49–60.

94. Cohen J. A power primer. Psychol Bull. 1992 Jul;112(1):155–9.

95. Hamajima N, Saito T, Matsuo K, Kozaki K, Takahashi T, Tajima K. Polymerase chain reaction with confronting two-pair primers for polymorphism genotyping. Jpn J Cancer Res. 2000 Sep;91(9):865–8.

96. Driever W, Rath M, Nitschke R, Filippi A, Ronneberger O, Driever W. Generation of high quality multi-view confocal 3D datasets of zebrafish larval brains suitable for analysis using Virtual Brain Explorer (ViBE-Z) software. Protocol Exchange. 2012 Jun 22; Available from: http://www.nature.com/protocolexchange/protocols/2408

97. Brennan CH, Parker MO. Translational Pharmacology of a Putative Measure of Motor Impulsivity in Larval Zebrafish. Current Psychopharmacology. 2016. Available from: http://www.eurekaselect.com/142514/article

98. Tang R, Dodd A, Lai D, McNabb WC, Love DR. Validation of zebrafish (Danio rerio) reference genes for quantitative real-time RT-PCR normalization. Acta Biochim Biophys Sin (Shanghai). 2007 May;39(5):384–90.

99. Collier AD, Echevarria DJ. The utility of the zebrafish model in conditioned place preference to assess the rewarding effects of drugs. Behavioural Pharmacology. 2013 Sep;24(5 and 6):375.

100. Martineau AR, James WY, Hooper RL, Barnes NC, Jolliffe DA, Greiller CL, et al. Vitamin D3 supplementation in patients with chronic obstructive pulmonary disease (ViDiCO): a multicentre, double-blind, randomised controlled trial. Lancet Respir Med. 2015 Feb;3(2):120–30.

101. Martineau AR, MacLaughlin BD, Hooper RL, Barnes NC, Jolliffe DA, Greiller CL, et al. Double-blind randomised placebo-controlled trial of bolus-dose vitamin D3 supplementation in adults with asthma (ViDiAs). Thorax. 2015 May;70(5):451–7.

102. Martineau AR, Hanifa Y, Witt KD, Barnes NC, Hooper RL, Patel M, et al. Double-blind randomised controlled trial of vitamin D3 supplementation for the prevention of acute respiratory infection in older adults and their carers (ViDiFlu). Thorax. 2015 Oct;70(10):953–60.

103. Loukola A, Wedenoja J, Keskitalo-Vuokko K, Broms U, Korhonen T, Ripatti S, et al. Genome-wide association study on detailed profiles of smoking behavior and nicotine dependence in a twin sample. Mol Psychiatry. 2014 May;19(5):615–24.

104. Loukola A, Broms U, Maunu H, Widén E, Heikkilä K, Siivola M, et al. Linkage of nicotine dependence and smoking behavior on 10q, 7q and 11p in twins with homogeneous genetic background. Pharmacogenomics J. 2008 Jun;8(3):209–19.

105. Broms U, Wedenoja J, Largeau MR, Korhonen T, Pitkäniemi J, Keskitalo-Vuokko K, et al. Analysis of detailed phenotype profiles reveals CHRNA5-CHRNA3-CHRNB4 gene cluster association with several nicotine dependence traits. Nicotine Tob Res. 2012 Jun;14(6):720–33.

106. Miller SA, Dykes DD, Polesky HF. A simple salting out procedure for extracting DNA from human nucleated cells. Nucleic Acids Res. 1988 Feb 11;16(3):1215.

107. Loukola A, Buchwald J, Gupta R, Palviainen T, Hällfors J, Tikkanen E, et al. A Genome-Wide Association Study of a Biomarker of Nicotine Metabolism. PLOS Genetics. 2015 Sep 25;11(9):e1005498.

108. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, et al. PLINK: a tool set for whole-genome association and population-based linkage analyses. Am J Hum Genet. 2007 Sep;81(3):559–75.

109. Zhou X, Stephens M. Genome-wide Efficient Mixed Model Analysis for Association Studies. Nat Genet. 2012 Jun 17;44(7):821–4.

110. Turner KJ, Hoyle J, Valdivia LE, Cerveny KL, Hart W, Mangoli M, et al. Abrogation of Stem Loop Binding Protein (Slbp) function leads to a failure of cells to transition from proliferation to differentiation, retinal coloboma and midline axon guidance deficits. PLOS ONE. 2019 Jan 29;14(1):e0211073.

<u>**Supplementary Methods**</u>

Zebrafish genomic DNA extraction: Genomic DNA was extracted from fin-clips using

QIAGEN DNeasy® Blood and Tissue Kit (Qiagen, Manchester, UK) according to

manufacturer's instructions. Samples were eluted into distilled water and stored at -20°C until

later use.

Site Specific Polymerase Chain Reaction: Allele-specific PCR single nucleotide

polymorphism (SNP) assays were used for genotyping F3 individuals for mutations known to

be present in the ENU-mutagenized F1 generation. Four primer pairs were designed to carry

out PCR genotyping as previously described (95). The list of loss-of-function mutations in

the AJBQM1 and AJBQM2 lines is detailed in Supplementary Table 1. For each line, a

primer was designed with 3' complementary to the ENU-SNP with a second primer ~100bp

downstream. The second pair had one primer with 3' complementary to the wild type base

with a second primer ~200bp upstream. The resulting PCR results in a 300bp fragment that

spans the region and acts as an internal control for the PCR plus one 100bp fragment if

homozygous for the mutation, 2 bands of 100bp and 200bp if heterozygous, and one 200bp

fragment if homozygous wild type. The 4-primer groups were designed with melting

temperatures as close as possible using the NCBI primer design tool and were ordered from

Eurofins, MWG operon (Ebersberg, DE).

Supplementary Table 1: List of loss-of-function mutations in the AJBQM1 (A) and

AJBQM2 (B) lines. List was derived from exome sequencing and provided by the Wellcome

Sanger Trust, Hinxton, Cambridge.

A)

SNP Name	Allele Number	Location	Description
cacna1ba (Cacna1b)	sa1562	Zv9:5:31016641	voltage-dependent N-type calcium channel subunit alpha-1B
vcana (VCAN)	sa1563	Zv9:5:48057817	novel protein similar to vertebrate chondroitin sulfate proteoglycan 2
si:ch211-157f15.1 (EVPL)	sa1564	Zv9:6:21645941	envoplakin
mobkl2a (MOBKL2A)	sa1565	Zv9:8:20954361	mps one binder kinase activator-like 2A
ENSDARG00000068026 (PRKG1)	sa1566	Zv9:8:53199402	protein kinase, cGMP-dependent, type I
glis3 (GLIS3)	sa1567	Zv9:10:663606	zinc finger protein GLIS3
si:dkey-220f10.4 (TULP2)	sa1568	Zv9:12:21973687	novel tub family member protein
slit3 (SLIT3)	sa1569	Zv9:14:25591202	slit homolog 3 protein
dchs1 (DCHS1)	sa1570	Zv9:15:31441900	dachsous 1
flad1 (FLAD1)	sa1571	Zv9:16:25049338	Molybdenum cofactor biosynthesis protein-like region FAD synthase region
si:ch211-199m3.2 (AKD1)	sa1572	Zv9:20:33741430	adenylate kinase domain containing 1
si:dkey-4c23.3 (???)	sa1573	Zv9:22:25367694	novel protein similar to vitellogenin 1 (Vg1)
magi2 (MAGI2)	sa1574	Zv9:25:21478784	membrane associated guanylate kinase, WW and PDZ domain containing 2
zgc:101050 (TRIMM55)	sa158	Zv9:23:17631394	hypothetical protein LOC445187 (tripartite motif-containing 55)

26 **B)**

SNP Name	Allele Number	Location	Description
capn3 (CAPN3)	sa150	Zv9:17:45493037	calpain-3
chrna9 (CHRNA9)	sa575	Zv9:1:22190803	cholinergic receptor, nicotinic, alpha 9
snmp70 (SNRNP70)	sa576	Zv9:3:32068963	U1 small nuclear ribonucleoprotein 70 kDa
zgc:153677 (SV2B)	sa577	Zv9:7:18060163	synaptic vesicle protein 2B homolog
wu:fa56e12 (AC103686.1)	sa578	Zv9:7:44124381	DNA-dependent protein kinase catalytic subunit
kctd4 (KCTD4)	sa580	Zv9:9:19499015	potassium channel tetramerisation domain containing 4
LOC557854 (SLC19A3)	sa581	Zv9:15:34443534	solute carrier family 19, member 3
tspan3a (TSPAN3)	sa584	Zv9:18:26858396	tetraspanin 3
rapsn (RAPSN)	sa585	Zv9:18:20289930	43 kDa receptor-associated protein of the synapse
si:ch211-132b12.1 (SLC6A11)	sa586	Zv9:18:38859333	hypothetical protein LOC100034467
pkhd1l1 (PKHD1L1)	sa587	Zv9:19:23349482	polycystic kidney and hepatic disease 1 (autosomal recessive)-like 1
klf11a (KLF-11)	sa588	Zv9:20:29529553	kruppel-like factor 11a

29 **Supplementary Table 2**. Gene ID and primer sequences used in gene expression analysis.

Gene name	Transcript ID	Sequence (5' -> 3')	Fragment size (bp)
Serotonergic pathway			
htr1aa	NM_001123321.1	Forward: TTCTACATCCCGCTCATCCTCA	180
		Reverse: CCTCCAAGTTTTACCCACCTCTC	
htr1ab	NM_001145766.1	Forward: AAACACCGAGGCGAAGAGGAA	99
		Reverse: GGCAGCCAACACAGAATGAAAGT	
htr2a	XM_684208.9	Forward: TACGGTGGCTGGGAACATTTTAG	187
		Reverse: GGGACACAGTGATGCAGGGAAA	
htr5a	NM_001126410.2	Forward: TGGATCAAAGAGGACCAACACC	118
		Reverse: CTGAAACGTCACCGTGGCAT	
htr2cl1	NM_001129893.1	Forward: AACTTCTTCCTCCGCTCACTCG	179
		Reverse: ATGGCACACAGGTGCATGATGG	
htr2cl2	XM_001339004.7	Forward: CACAACCCCACCAACTTCTTCC	152
		Reverse: ACGTCCAGAAAGATCCACAGCG	
htr4	XM_021481160.1 XM_009291062.3	Forward: GTTTCTTTCCAAGCGCCTC	168
		Reverse: ACTTCTTCCATCTCAGGCATC	
htr6	XM_009297078.3	Forward: ACTACAGTCATCAGGAGCCACC	147
		Reverse: GCCAGGCACTGAAGAATAGTCC	
htr7	XM_003199584.5	Forward: TGGATGTGATGTGCTGTACCGC	118
		Reverse: GCCATGCACTTTCCACTCTGTC	
slc6a4/ SERT	NM_001039972.1	Forward: ACCAGGGGCGAAGCCAAGCA	117
		Reverse: GCCACAGGCCCCGCTGTTA	
htr1b	NM_001128709.1	Forward: CCTTGTCGTCAGTTCTGGGT	112
		Reverse: ATCAGAAAGTTCGCCGGTGT	
Nicotinic pathway			
chrna2	NC_007128.7	Forward: TGGCTGCAGATCAGTCAAAGAC	271
		Reverse: CCCTCTAACTGTCCCTTCACAA	
chrna3	Not found in BLAST	Forward: TGTACATCCGCCGATTACCGCT	?
		Reverse: TCCGCAGTCGGAGGGCAGTA	
chrna4b	ENSDART00000018614.7	Forward: TTACAAGAGGTTTGGGCGCT	90
		Reverse: ACAGACCAGTAGATCATCACTCC	
chrna5	NM_001017885.1	Forward: GGCTCCCAGGTCGACATTCTC	103
		Reverse: AACCCCGGTTACCAGTGGCCT	
chrna6	NM_001042684	Forward: AGGCTCTTTCGTCGTTTATTC	156
		Reverse: TCTCAGCCAAAGGTTTGTTTC	
chrna7	ENSDART00000171463.3	Forward: ACCGTGTCACATTGTTCATTCTC	105

	and ENSDART00000166391.2	Reverse: ACAGGTCTCTCCAGTGGGTTA	
chrnb2b	ENSDART00000041625.7 and ENSDART00000185728.1	Forward: CACAAAGTCACGCTCCGATAC	160
		Reverse: CCGTCGCTCTGAGCAGATAA	
chrnb3b	ENSDARG00000038508.5	Forward: CAGGAGTCAACCTCCGCTTT	106
		Reverse: TGAATCTGAACGCACTGGCT	
chrnb4	NC_007129.7	Forward: ATGTGAATGAATGGCGGTGTGTG	203
		Reverse: ATGCGCGTGTCAGATTTACCC	

Dopaminergic pathway

drd1b	NM_001135976.2	Forward: TGGTTCCTTTCTGCAACCCA	100
		Reverse: AGTGATGAGTTCGCCCAACC	
drd2a	XM_009291617.3 and XM_005157501.4	Forward: TCCACAAAATCAGGAAAAGCGT	106
		Reverse: CAGCCAATGTAAACCGGCAA	
drd3	XM_021470111.1 and XM_005162673.4	Forward: ATCGAGTTTCGCAGAGCCTT	95
		Reverse: TCCACAGTGTCTGAAAGCCG	
slc6a3	NM_131755.1	Forward: GCCTGGTTTTACGGAGTGGA	66
		Reverse: GGAGGATTGAAGGTGGCGAA	

Adrenergic pathway

adra1aa	NM_001324454.1	Forward: AAGAAGGCCGCCAAGACTTT	114
		Reverse: GTCCGAGGGTCTGTACGTTG	
adra1d	XM_691951.6	Forward: AAGCTGCTAAAACCCTCGCC	103
		Reverse: GGCTTCAGAGCTGGGAAGAAT	
adra2b	NM_207638.1	Forward: AAAAGCCAGGCCTCCAACTT	92
		Reverse: GGGCTTGCAGAAGGTTGTTG	
*adra2*c	NM_207639.1	Forward: CGCCGTTTTAACGAGCAGAG	87
		Reverse: AGTGTGGCCACCAGAATGTC	
adra2da	NM_194364.2	Forward: CATCATCCTCGTGGTGTCCC	188
		Reverse: ATCCCATGATCTCGTTGGCG	
adra2db	NM_194365.1	Forward: TGCCACTTTGGTCATTCCGT	88
		Reverse: AGCCAGGTAGAAAGCACACC	

30
31

Human Association Studies

Supplementary Table 3: Sample characteristics of the human cohorts. Detailed inclusion and exclusion criteria for the London cohorts can be found in https://clinicaltrials.gov. Further details about recruitment, power calculation and definition of medical phenotypes can be found elsewhere (100–102). Genotype and phenotype Finnish twins are deposited in the Biobank of the National Institute for Health and Welfare (https://thl.fi/en/web/thl-biobank/for-researchers). **Differences in sample size for Finnish cohorts was due to hard-call genotype probability threshold. DSM-IV nicotine dependence symptoms, Fagerström scores and cigarettes smoked each day (N = 1715). Sensation felt after smoking first cigarette (N = 1915). Time to first cigarette in the morning (N= 1726).

Cohort name	N	Country	Cohort description	ClinicalTrials. gov ID	Mean age (years)	% female	Smoking phenotypes investigated	N heavy smokers
ViDiCO	272	UK	Subjects with mild, moderate or severe chronic obstructive pulmonary disease (COPD) treated with the same bi-monthly 3mg vitamin D3 intervention.	NCT00977873	64.6	40	Tobacco consumption; smoking cessation (current vs ever smokers)	249
ViDiAs	293	UK	Adult patients with asthma treated with inhaled corticosteroids treated with a bi-monthly 3mg vitamin D3 intervention	NCT00978315	47	56	Tobacco consumption; smoking cessation (current vs ever smokers)	17
ViDiFLU	298	UK	Adults in sheltered accommodation given 10 mcg vitamin D3 daily as well as bi-monthly 3mg vitamin D3 interventions	NCT01069874	66.8	66	Tobacco consumption; smoking cessation (current vs ever smokers)	66

| Finnish Twins | 1915, 1715, 1726* | Finland | Study sample ascertained from the Finnish Twin Cohort study (N=35834 adult twins) concordant for moderate to heavy smoking | NA | 55 | 48 | DSM-IV nicotine dependence symptoms; Fagerström scores; cigarettes smoked each day; sensation felt after smoking first cigarette and time to first cigarette in the morning | NA |

Phenotype definitions for the London cohorts: _Amount smoked_ was defined as the average number of cigarettes smoked per day (CPD) for each participant. Participants met criteria for _smoking cessation_ if they reported being 'ever smokers' and reported **not** smoking currently. The percentage of current smokers in the cohort was 42%, 7% and 18% for ViDiCO, ViDiAs and ViDiFLU, respectively.

Phenotype definition for the Finnish twin cohort study: Definitions of the phenotypes were adapted from Broms et al (105).

Amount smoked

Cigarettes per day (CPD) constitutes of eight categories: 1-2, 3-5, 6-10, 11-15, 16-19, 20-25, 26-39, ≥40 CPD. In the statistical analyses of the CPD variables, original categorical observations were replaced with class means of CPD (1.5, 3.5, 8, 13, 17.5, 22.5, 32.5, and 45 cigarettes per day, respectively). Regression coefficients can therefore be interpreted as the average change in number of cigarettes smoked per day when the number of minor allele is increased by one.

- **CPD**: Number of cigarettes smoked per day during month of heaviest smoking. Values ranged from 1 to >40 with mean=19.8 cigarettes per day.

Maximum CPD: Maximum number of cigarettes ever smoked during one day (24h

period). Values ranged from 2 to 98 with mean=30 cigarettes per day.

Smoking initiation

Age of onset of weekly smoking: Age (years) when started to smoke weekly ("How

old were you when you first smoked a cigarette at least once a week for at least two

months in a row?"). Values ranged from 6 to 54, mean=17.3 years.

First time sensation. Sensation felt after smoking the first cigarette or first puffs.

Sensation measured as: "While smoking your very first cigarettes, did you (1) like the

taste or smell of the cigarette, (2) cough, (3) feel dizzy or light-headed, (4) feel more

relaxed, (5) get a headache, (6) feel a pleasurable rush or buzz, (7) feel your heart

racing, (8) feel nauseated, like vomiting, (9) feel your muscles tremble or become

jittery, (10) feel burning in your throat"). Sum score of 10 questions (items #1, #4,

and #6 were reverse-scored before summation): 0 points if answered "No", 1 = "A

little bit", 2="Some", 3= "Quite a bit", 4="A great deal". Cronbach's alpha = 0.70.

Values ranged from 3.6 to 15.8. Mean =10.2.

Nicotine dependence

DSM-IV ND diagnosis: Nicotine dependence by DSM-IV diagnosis ($\geq$3 symptoms

out of 7 occurring within a year). Prevalence = 53.5%.

DSM-IV ND symptoms: Number of DSM-IV ND symptoms from 0 to 7. Mean=3

FTND ($\geq$4): Nicotine dependent if $\geq$4 out of 10 points in Fagerström Test for

Nicotine Dependence. Prevalence = 50.4%

FTND score: Fagerström Test for Nicotine Dependence (FTND) score: 0 to 10

points. Mean=3.7.

▪ ***FTND time to first cigarette (TTF):*** Time to first cigarette in the morning (one item

of the FTND scale). Five categories: 0-5 min, 6-15 min 6, 16-30 min 6, 31-60 min,

>60 min. Categorization differs from original four categories (3), i.e., 6-30 minutes is

split into 6-15 min and 16-30 min. In our data set 46% of smokers belong to the group

of 6-30 min, and from the smoking behaviour point of view there is a significant

difference whether one smokes the first cigarette within 6 minutes or 30 minutes from

waking up. In this data set 22% of smokers belong to the 6-15 min and 24% to the 16-

30 min group. Values ranged from 1 to 5 with a mean=3.1.

Supplementary Table 4. Primer and reporter sequences used for human genotyping.

GENE	SNP	Sequence name	Sequence
CYP3A4	rs2740574		
		Forward	CCAGGCATAGGTAAAGATCTGTAGGT
		Reverse	CTCAAGTGGAGCCATTGGCATA
		Reporters	ACAAGGGCAAGAGAG and
			ACAAGGGCAGGAGAG
CUBN	rs3740165		
		Forward	GCAATGAGATTAAATCTTCAGGAAACACA
		Reverse	CTGGAGGTATAGGAAGCAGTGAAG
		Reporters	CCGCCATATGGCCTG and CGCCATACGGCCTG
RXRA	rs7861779		
		Forward	TGGCCCATGCACGAGTAG
		Reverse	ACCGAGACAGGCCAAACTC
		Reporters	CAGCAGAGGTGGCCGA and
			CAGCAGAGATGGCCGA

<u>**Supplementary results**</u>

Supplementary Table 5: Site specific PCR genotyping of AJBQM1 (A) and AJBQM2

(B) outlier siblings. The siblings were genotyped at each of the candidate loci using site

specific PCR and results compared with each individual place preference change scores. Each

row represents one fish, with their corresponding CPP change score, and genotype for each

locus assessed (13 loci for AJBQM1, 12 loci for AJBQM2). P-values result from independent

two-sample t-tests comparing the CPP change scores between genotype groups at each locus.

(A)

Fish	CPP Change Score	Gene name												
		slit3	*cacne*	*vcan*	*evpl*	*mob3a*	*prkg1*	*glis3*	*tulp2*	*dchs1*	*flad1*	*akd1*	*magI2*	*trimm55*
1	**0**	**WT**	WT	HET	WT	HET	HET	WT	HOM	WT	HOM	HOM	HET	WT
2	**0.01**	**WT**	HET	HET	WT	HET	HET	HET	HOM	WT	HOM	HOM	WT	HET
3	**0.07**	**WT**	WT	WT	WT	HET	HET	HET	WT	WT	WT	WT	HET	WT
4	**0.15**	**WT**	WT	HET	WT	HOM	HET	HET	HET	WT	HET	HET	WT	HET
5	**0.32**	**HET**	HOM	HOM	WT	HOM	HET	WT	WT	WT	WT	WT	HET	WT
6	**0.43**	**HET**	HET	WT	WT	HET	HET	WT	HET	WT	WT	HET	HOM	WT
7	**0.44**	**HET**	WT	HET	WT	WT	HET	WT	HET	WT	WT	HET	WT	WT
8	**0.47**	**HET**	WT	HET	WT	HET	HET	HET	HET	WT	HET	HET	WT	WT
9	**0.51**	**HET**	WT	WT	WT	WT	HOM	WT	WT	WT	HET	WT	HOM	HET
10	**0.6**	**HET**	WT	WT	WT	HET	HOM	HET	WT	WT	HET	WT	HET	WT
	P-value	**7.66×10^{-5}**	0.691	0.259	-	0.236	-	0.602	0.481	-	0.981	0.418	0.73	0.51

14   **(B)**

Fish number	CPP Change Scores	Gene name											
		tspan3a	*raspn*	*a9*	*capn3*	*klf11a*	*kctd4*	*slc6a11*	*pkhd1l1*	*slc19a3*	*sv2b*	*snrnp70*	*ac10103686*
1	-0.38	HET	WT	WT	HET	WT	HET	HET	WT	WT	WT	WT	WT
2	-0.28	HET	HOM	WT	HET	WT	WT	HET	WT	WT	WT	WT	WT
3	-0.27	HET	WT	HET	HET	WT	WT	HET	WT	WT	HET	WT	WT
4	-0.23	WT	WT	WT	WT	HET	HET	WT	HET	WT	WT	WT	WT
5	-0.21	WT	WT	WT	HET	WT	WT	HET	HOM	WT	WT	WT	WT
6	-0.18	WT	HOM	WT	WT	WT	WT	WT	HOM	WT	WT	WT	WT
7	-0.17	HET	HOM	HET	WT	WT	HET	HET	HET	WT	WT	WT	WT
8	-0.17	HET	HOM	WT	HET	HET	WT	HET	WT	WT	HET	WT	WT
9	-0.12	WT	WT	HET	WT	WT	HET	WT	WT	WT	HOM	WT	WT
10	-0.09	WT	WT	HET	HET	WT	HET	WT	WT	WT	HET	WT	WT
11	-0.09	WT	WT	HET	WT	HET	HET	WT	WT	WT	HOM	WT	WT
12	-0.07	HET	HOM	HET	WT	WT	HET	WT	WT	WT	HOM	WT	WT
13	0.04	HET	WT	HET	WT	HET	WT	HET	WT	WT	HET	WT	WT
14	0.07	HET	HOM	HET	WT	WT	WT	WT	HOM	WT	HOM	WT	WT
	P-value	0.583	0.792	0.339	0.911	0.318	0.252	0.697	0.499	-	0.269	-	-

Supplementary Table 6: Results of association analysis of *SLIT3* SNPs on **smoking**

initiation. Logistic regression of initiation vs non-initiation on additive genotype, controlling

for age, sex and cohort. OR: Odds ratio. >1 value indicates that the minor allele increases

odds of persistent smoking relative to the major allele, SE: standard error, L95: lower limit of

95% confidence interval, U95: upper limit of 95% confidence interval. Benjamini Hochberg

cut off at 0.1 = 0.00526.

SNP	OR	SE	L95	U95	P value
rs2938774	0.7253	0.1418	0.5493	0.9578	0.02357
rs11742567	1.328	0.1538	0.9825	1.796	0.06496
rs4282339	0.7277	0.1815	0.5099	1.039	0.07991
rs297886	1.328	0.176	0.9405	1.875	0.1071
rs9688032	1.269	0.1495	0.9467	1.701	0.111
rs7728604	1.198	0.1446	0.9024	1.591	0.2112
rs1345588	0.7788	0.2046	0.5215	1.163	0.2218
rs17734503	0.7362	0.2632	0.4395	1.233	0.2445
rs12515725	0.8612	0.1458	0.6471	1.146	0.3052
rs11749001	0.8698	0.1951	0.5933	1.275	0.4746
rs3733975	1.116	0.1641	0.8092	1.54	0.5029
rs12521041	1.116	0.1641	0.8092	1.54	0.5029
rs11134527	0.9212	0.155	0.6798	1.248	0.5964
rs12654448	0.8718	0.2654	0.5182	1.467	0.6052
rs1559051	0.9257	0.1575	0.6799	1.26	0.6241
rs17665158	0.9304	0.171	0.6654	1.301	0.673
rs1421763	0.9423	0.1704	0.6748	1.316	0.7271
rs295994	0.9732	0.1404	0.7391	1.281	0.8464
rs10036727	1.01	0.1522	0.7491	1.361	0.9503

Supplementary Table 7: Results of association analysis of *SLIT3* SNPs on **persistent**

smoking. Logistic regression of initiation vs non-initiation on additive genotype, controlling

for age, sex and cohort. OR: Odds ratio. >1 value indicates that the minor allele increases

odds of persistent smoking relative to the major allele, SE: standard error, L95: lower limit of

95% confidence interval, U95: upper limit of 95% confidence interval. Benjamini Hochberg

cut off at 0.1 = 0.00526.

SNP	OR	SE	L95	U95	P value
rs11134527	1.428	0.1573	1.049	1.943	0.02359
rs12521041	0.6871	0.1736	0.489	0.9655	0.03061
rs11742567	0.7288	0.1547	0.5382	0.987	0.04089
rs3733975	0.7165	0.1712	0.5123	1.002	0.05146
rs17734503	0.6142	0.2631	0.3667	1.029	0.06398
rs1345588	0.6786	0.2145	0.4457	1.033	0.07068
rs17665158	1.338	0.163	0.9722	1.842	0.07393
rs12654448	0.6225	0.2671	0.3688	1.051	0.07597
rs2938774	1.232	0.1394	0.9373	1.619	0.1348
rs295994	1.214	0.1443	0.9147	1.61	0.1796
rs7728604	1.152	0.1434	0.8699	1.526	0.3232
rs1559051	1.115	0.1549	0.8231	1.511	0.4821
rs12515725	0.9108	0.1456	0.6846	1.212	0.521
rs297886	0.9342	0.1726	0.6661	1.31	0.6932
rs4282339	0.9391	0.19	0.6471	1.363	0.7411
rs10036727	0.9596	0.1516	0.713	1.292	0.7857
rs1421763	1.029	0.1671	0.7417	1.428	0.8633
rs9688032	1.013	0.1511	0.7531	1.362	0.9328
rs11749001	0.9943	0.1969	0.676	1.463	0.977

34
Figure 5-figure supplement 1. Fluorescent immunohistochemistry in three-day old wild

type and homozygous mutant *slit3*^{sa1569} labelled with tyrosine hydroxylase (A) and tubulin

(B-F). A) Example of three Z-planes used to quantify catecholaminergic projections crossing

the midline. (Stacks 1-3). B) Anti-tubulin staining of *slbp*$^{ty77e/ty77e}$ mutant larvae, known to

have fewer neurons and axonal defects (110). C-D) Anti-tubulin staining of wild type and

homozygous mutant *slit3*^{sa1569}. E-F) Zoom-in of C-D for midline in the ventral forebrain,

where *slit3* is known to be expressed (35).

Figure 6 –figure supplement 1. Average distance moved before (Figure 1A) and during

startle stimuli (Figure 1B) in wild type and *slit3*[sa1569] mutant five day old zebrafish larvae. **A)**

Distance moved as function of amisulpride dose, fish *slit3* genotype and their interaction. The

effect of dose and genotype was tested in a linear mixed model. Timepoint, well where the

fish were placed and plate were also included as fixed factors and the fish ID as random

factor. **B)** Distance moved during taps as function of amisulpride dose and fish *slit3*

genotype. Drug and genotype effects were examined in a linear mixed model including

stimulus number, well, plate used and distance moved before stimuli as fixed factors and Fish

ID as random factor. Zebrafish larvae did not differ in the distance travelled before or during

startle stimuli as a function of amisulpride dose nor genotype ($p > 0.05$). Bars represent

estimated marginal means $\pm$ SEM (n=42-48 fish per group).

Supplementary Table 8. Full results for quantitative real-time PCR analysis of five day old

wild type *slit3$^{sa1569+/+}$*, *slit3$^{sa1569/+}$* heterozygous and *slit3$^{sa1569/sa1569}$* homozygous mutant

larvae (Total n=30, 3 samples per experimental group with n=10 embryos per sample). Only

htr1aa ([F(2,6)=44], p=0.0003) showed a significant difference across genotypes after

correcting for multiple testing (p < 0.001). DFn and DFd represents degree of freedom for the

numerator and denominator of the F ratio.

Gene name	Mean (SD) Slit3$^{+/+}$	Mean (SD) Slit3$^{sa1569/-}$	Mean (SD) Slit3$^{sa1569/sa1569}$	F (DFn, DFd)	p
Serotonergic pathway					
htr1aa	**5.30×10^{-4}**	**1.43×10^{-3}**	**1.27×10^{-3}**	**F (2, 6) = 44**	**0.0003**
	(2.60×10^{-5})	(1.53×10^{-4})	(1.53×10^{-4})		
htr1ab	8.63×10^{-4}	9.30×10^{-4}	9.23×10^{-4}	F (2, 6) = 0.2594	0.7797
	(1.25×10^{-4})	(1.54×10^{-4})	(8.62×10^{-5})		
htr2a	5.87×10^{-4}	7.43×10^{-4}	5.73×10^{-4}	F (2, 6) = 0.8987	0.4556
	(1.05×10^{-4})	(2.35×10^{-4})	(1.53×10^{-4})		
htr5a	2.90×10^{-5}	5.20×10^{-5}	4.73×10^{-5}	F (2, 6) = 0.4694	0.6465
	(2.20×10^{-5})	(4.22×10^{-5})	(2.41×10^{-5})		
htr2cl1	1.13×10^{-3}	1.43×10^{-3}	1.33×10^{-3}	F (2, 6) = 1.500	0.2963
	(1.16×10^{-4})	(3.22×10^{-4})	(1.53×10^{-4})		
htr2cl2	3.87×10^{-5}	6.50×10^{-5}	5.17×10^{-5}	F (2, 6) = 0.9413	0.4410
	(5.13×10^{-6})	(3.21×10^{-5})	(2.46×10^{-5})		
htr4	1.02×10^{-3}	1.40×10^{-3}	1.43×10^{-3}	F (2, 6) = 7.654	0.0223
	(1.33×10^{-4})	(2.00×10^{-4})	(5.77×10^{-5})		
htr6	1.50×10^{-5}	6.17×10^{-5}	4.23×10^{-5}	F (2, 6) = 4.184	0.0728
	(2.00×10^{-6})	(3.26×10^{-5})	(1.07×10^{-5})		
htr7	1.23×10^{-4}	1.12×10^{-4}	1.73×10^{-4}	F (2, 6) = 0.8930	0.4576
	(3.11×10^{-5})	(5.35×10^{-5})	(8.39×10^{-5})		
slc6a4/ sert	1.40×10^{-3}	2.00×10^{-3}	1.83×10^{-3}	F (2, 6) = 3.408	0.1026
	(1.00×10^{-4})	(4.36×10^{-4})	(2.31×10^{-4})		
htr1b	9.53×10^{-4}	1.33×10^{-3}	1.13×10^{-3}	F (2, 6) = 1.617	0.2744
	(4.51×10^{-5})	(4.16×10^{-4})	(1.62×10^{-4})		
Overall serotonergic	*6.09×10^{-4}*	*8.69×10^{-4}*	*8.00×10^{-4}*	*F (2, 30) = 0.5137*	*0.6034*
	(5.01×10^{-4})	*(7.05×10^{-4})*	*(6.51×10^{-4})*		
nicotinic pathway					
chrna2	1.49×10^{-5}	1.60×10^{-5}	3.80×10^{-7}	F (2, 6) = 0.8431	0.4757
	(6.47×10^{-6})	(2.77×10^{-5})	(2.82×10^{-7})		
chrna3	3.53×10^{-4}	4.13×10^{-4}	3.70×10^{-4}	F (2, 6) = 1.044	0.4081
	(5.03×10^{-5})	(4.62×10^{-5})	(6.00×10^{-5})		
chrna4b	4.13×10^{-4}	6.10×10^{-4}	5.73×10^{-4}	F (2, 6) = 5.836	0.0391
	(4.51×10^{-5})	(1.05×10^{-4})	(6.11×10^{-5})		
chrna5	1.43×10^{-3}	2.00×10^{-3}	1.90×10^{-3}	F (2, 6) = 2.906	0.1311
	(1.53×10^{-4})	(3.61×10^{-4})	(3.61×10^{-4})		

chrna6	2.13×10^{-3}	2.63×10^{-3}	2.23×10^{-3}	$F(2, 6) = 2.333$	0.178
	(5.77×10^{-5})	(5.03×10^{-4})	(1.16×10^{-4})		
chrna7	5.75×10^{-5}	8.53×10^{-5}	7.50×10^{-5}	$F(2, 6) = 0.4101$	0.6808
	(5.04×10^{-5})	(4.15×10^{-5})	(8.72×10^{-6})		
chrnb2b	5.20×10^{-4}	9.67×10^{-4}	8.40×10^{-4}	$F(2, 6) = 18.18$	0.0028
	(2.00×10^{-5})	(1.53×10^{-4})	(5.00×10^{-5})		
chrnb3b	6.27×10^{-6}	8.73×10^{-5}	1.45×10^{-4}	$F(2, 6) = 5.870$	0.0387
	(3.51×10^{-6})	(2.47×10^{-5})	(8.26×10^{-5})		
chrnb4	4.70×10^{-5}	7.57×10^{-5}	5.93×10^{-5}	$F(2, 6) = 0.8718$	0.4652
	(9.17×10^{-6})	(3.84×10^{-5})	(2.40×10^{-5})		
Overall nicotinic	5.53×10^{-4}	7.65×10^{-4}	6.89×10^{-4}	$F(2, 24) = 0.1455$	0.8654
	(7.44×10^{-4})	(9.46×10^{-4})	(8.32×10^{-4})		

dopaminergic pathway

drd1b	1.65×10^{-3}	2.15×10^{-3}	2.01×10^{-3}	$F(2, 6) = 4.333$	0.0685
	(2.46×10^{-4})	(2.66×10^{-4})	(7.00×10^{-5})		
drd2a	6.42×10^{-4}	8.72×10^{-4}	8.29×10^{-4}	$F(2, 6) = 1.484$	0.2995
	(1.25×10^{-4})	(2.51×10^{-4})	(1.11×10^{-4})		
drd3	1.30×10^{-3}	1.53×10^{-3}	1.60×10^{-3}	$F(2, 6) = 5.154$	0.0498
	(1.00×10^{-4})	(1.53×10^{-4})	(1.00×10^{-4})		
slc6a3	5.43×10^{-4}	5.80×10^{-4}	6.60×10^{-4}	$F(2, 6) = 4.089$	0.0758
	(3.51×10^{-5})	(6.25×10^{-5})	(5.20×10^{-5})		
Overall dopaminergic	1.04×10^{-3}	1.28×10^{-3}	1.28×10^{-3}	$F(2, 9) = 0.2025$	0.8203
	(5.32×10^{-4})	(7.02×10^{-4})	(6.39×10^{-4})		

adrenergic pathway

adra1aa	1.20×10^{-3}	1.70×10^{-3}	1.57×10^{-3}	$F(2, 6) = 4.892$	0.0549
	(1.73×10^{-4})	(2.65×10^{-4})	(1.53×10^{-4})		
adra1d	8.30×10^{-4}	9.43×10^{-4}	8.63×10^{-4}	$F(2, 6) = 0.2954$	0.7545
	(8.72×10^{-5})	(2.89×10^{-4})	(1.10×10^{-4})		
adra2b	3.93×10^{-3}	5.23×10^{-3}	4.43×10^{-3}	$F(2, 6) = 12.90$	0.0067
	(2.52×10^{-4})	(3.79×10^{-4})	(3.06×10^{-4})		
adra2c	1.30×10^{-3}	1.60×10^{-3}	1.50×10^{-3}	$F(2, 6) = 3.500$	0.0983
	(1.00×10^{-4})	(2.00×10^{-4})	(1.00×10^{-4})		
adra2da	3.27×10^{-4}	4.17×10^{-4}	4.00×10^{-4}	$F(2, 6) = 3.005$	0.1247
	(2.31×10^{-5})	(7.51×10^{-5})	(2.65×10^{-5})		
adra2db	2.90×10^{-4}	2.87×10^{-4}	3.57×10^{-4}	$F(2, 6) = 0.6630$	0.5494
	(9.54×10^{-5})	(8.02×10^{-5})	(7.51×10^{-5})		
Overall adrenergic	1.31×10^{-3}	1.70×10^{-3}	1.52×10^{-3}	$F(2, 15) = 0.08865$	0.9156
	(1.35×10^{-3})	(1.83×10^{-3})	(1.52×10^{-3})		