Highly Predictive Transdiagnostic Features Shared across Schizophrenia, Bipolar Disorder, and ADHD Identified Using a Machine Learning Based Approach

Yuelu Liu[1], Monika S. Mellem[1], Humberto Gonzalez[1], Matthew Kollada[1], Atul R.
Mahableshwarkar[1], Annette Madrid[1], William J. Martin[1], Parvez Ahammad[1]*

[1]Blackthorn Therapeutics, Inc., San Francisco, CA, USA

* Correspondence:
Parvez Ahammad, PhD
parvez.ahammad@blackthornrx.com

Keywords: transdiagnostic, schizophrenia, bipolar disorder, ADHD, machine learning.

Abstract

The Diagnostic and Statistical Manual of Mental Disorders (DSM) is the standard for diagnosing
psychiatric disorders in the United States. However, evidence has suggested that symptoms in
psychiatric disorders are not restricted to the boundaries between DSM categories, implying an
underlying latent transdiagnostic structure of psychopathology. Here, we applied an importance-
guided machine learning technique for model selection to item-level data from self-reported
instruments contained within the Consortium for Neuropsychiatric Phenomics dataset. From 578
questionnaire items, we identified a set of features which consisted of 85 items that were shared
across diagnoses of schizophrenia (SCZ), bipolar disorder (BD), and attention deficit/hyperactivity
disorder (ADHD). A classifier trained on the transdiagnostic features reliably distinguished the
patient group as a whole from healthy controls (classification AUC = 0.95) and only 10 items were
needed to attain the performance level of AUC being 0.90. A sum score created from the items
produced high separability between patients and healthy controls (Cohen's d = 2.85), and it
outperformed predefined sum scores and sub-scores within the instruments (Cohen's d ranging
between 0.13 and 1.21). The transdiagnostic features comprised both symptom domains (e.g.
dysregulated mood, attention deficit, and anhedonia) and personality traits (e.g. neuroticism,
impulsivity, and extraversion). Moreover, by comparing the features that were common across the
three patient groups with those that were most predictive of a single patient category, we can describe
the unique features for each patient group superimposed on the transdiagnostic feature structure.
Overall, our results reveal a latent transdiagnostic symptom/behavioral phenotypic structure shared
across SCZ, BD, and ADHD and present a new perspective to understand insights offered by self-
report psychiatric instruments.

Number of words: 6463
Number of words in abstract: 267
Number of figures: 4

1   Introduction

The Diagnostic and Statistical Manual of Mental Disorders (DSM) provides a symptom-based
taxonomy which serves to help clinicians classify various clusters of symptoms and abnormal
behaviors into distinct categories of disorders. The uniformity of diagnostic criteria in DSM serves to
effectively index psychiatric disorders but does not provide a data-driven framework within which to
understand the shared and unique features across disorders. For example, dimensionality and
comorbidity are pervasive in terms of symptoms across different DSM categories (Kessler et al.,
2005; Markon, 2009; Krueger and Markon, 2011). Such dimensionality manifests as heterogeneity in
symptom clusters within disease categories defined by the DSM and is exemplified across DSM
categories (Kessler et al., 2007). In the area of anxiety and mood disorders, more than 50% of
individuals are diagnosed as having more than a single category of disorders according to the DSM at
a given time (Grisanzio et al., 2017). Similarly, about 50% of bipolar disorder patients exhibit
schizophrenia-like psychotic symptoms during illness episodes (Coryell et al., 2001; Keck et al.,
2003). The presence of such psychotic symptoms can be mood-incongruent (Pacheco et al., 2010)
and can occur outside of illness episodes (Pope and Lipinski, 1978; Abrams and Taylor, 1981). These
observations highlight the likelihood of a latent trans-diagnostic dimensional structure that spans
multiple disorders (Krueger and Markon, 2006) and underscore the importance of understanding
patients at the symptom-level, rather than simply at a diagnostic level, to create more effective
treatments.

Studies have attempted to uncover the latent structure of psychopathology, between or within
categories, through multimodal assessments that measure symptoms, behavior, physiology, imaging,
and genetics. One such example is the large-scale study conducted by the UCLA Consortium for
Neuropsychiatric Phenomics (CNP), which seeks to identify links among phenotypic data, imaging,
and genetics (Poldrack et al., 2016). Overall, genetic studies have pointed to the heritability of major
neuropsychiatric disorders (Cross-Disorder Group of the Psychiatric Genomics Consortium, 2013;
Hamshere et al., 2013; Larsson et al., 2013; The Brainstorm Consortium et al., 2017; Bipolar
Disorder and Schizophrenia Working Group of the Psychiatric Genomics Consortium et al., 2018;
Gandal et al., 2018) as well as the genetic commonality amongst disorders (Purcell et al., 2009; Lotan
et al., 2014) such as schizophrenia (SCZ), bipolar disorder (BD), and attention deficit/hyperactivity
disorder (ADHD). Recent data-driven studies based on symptom and behavior have focused on
classifying and subtyping patients within a single diagnostic category (Lamers et al., 2012; van Loo
et al., 2012; Georgiades et al., 2013; Doshi-Velez et al., 2014; van Hulst et al., 2014; Costa Dias et
al., 2015; Geisler et al., 2015; Sun et al., 2015; Drysdale et al., 2016; Gheiratmand et al., 2017).
Several of these studies identified important shared abnormal features associated with the latent
transdiagnostic structure across major psychiatric disorders.

Despite recent advancements, several unresolved issues still remain in the field. First, the clinical
utility of using the features identified in the above-mentioned studies to reliably classify patients
remains an open question. Emerging studies have used unsupervised machine learning approaches,
such as clustering and dimensionality reduction algorithms, to uncover the transdiagnostic structure
across disorders (Grisanzio et al., 2017; Xia et al., 2018). However, the lack of ground truth on how
patients should be assigned to an identified cluster/subtype limits the application of these insights.
Second, despite recent genetic studies documenting shared risk factors among SCZ, BD, and ADHD
(Cross-Disorder Group of the Psychiatric Genomics Consortium, 2013; Larsson et al., 2013), a trans-
diagnostic dimensional structure shared across the three disorders has not been discovered in other
modalities such as neuroimaging and clinical characteristics. While a substantial body of
neuroimaging studies have focused on investigating shared etiology between SCZ and BD (see e.g.,

Rashid et al., 2014), studies that further incorporated ADHD are scarce. Third, how well different
feature modalities can be used as markers to reliably identify psychiatric patients in a clinical setting
has not been systematically compared in prior literature. Studies typically reported systematic
deviations within a single feature modality among psychiatric patients (Buckholtz and Meyer-
Lindenberg, 2012; Goodkind et al., 2015; Sha et al., 2018) and the relative predictive power of
various feature modalities in transdiagnostic scenarios remains unknown.

In the current study, we addressed the above issues by taking a patient-focused approach to identify
transdiagnostic features that are shared across SCZ, BD, and ADHD. Following the definition used in
the recent literature, we used the term "transdiagnostic" to denote features that extend beyond a
single DSM category. Using an importance-guided model selection approach, the supervised
machine learning framework used in this study allowed us to evaluate the performance of the
transdiagnostic features and hence to iteratively identify the optimal set of features required to
distinguish the patient group from healthy controls (HCs). Based on the CNP dataset, we used
multiple data modalities including the behavioral/symptom phenotypes (from here on referred to as
phenotypes) defined in self-reported instruments and neuroimaging data (sMRI and fMRI) to obtain
the optimal transdiagnostic features. Because the self-reported instruments were administered to
acquire a rich set of phenotypic information rather than providing diagnoses, our study also sought to
establish the clinical utility of these phenotypic features in distinguishing patients from HCs. In
addition, the clinical utility of the markers identified in each feature modality was systematically
evaluated by comparing the performance of models trained on each modality. We then report these
shared features and discuss the identified latent psychopathological structure across these psychiatric
disorders.

2   Materials and Methods

2.1   The CNP dataset

We utilized the openly available dataset from the CNP LA5c Study conducted at the University of
California, Los Angeles (the CNP dataset: https://openneuro.org/datasets/ds000030/versions/00016).
Detailed information on the CNP study/dataset can be found in (Poldrack et al., 2016). The CNP
dataset contains a variety of data modalities. In this study, we focused on identifying shared
transdiagnostic features based on the item-level data from self-reported instruments and
neuroimaging data (including both sMRI and resting-state fMRI). The dataset in this study includes
272 subjects, of which 50 are diagnosed with schizophrenia (SCZ), 49 with bipolar disorder (BD),
and 43 with attention deficit/hyperactivity disorder (ADHD). The remaining 130 subjects are age-
matched healthy controls (HC). The diagnoses were given by following the Diagnostic and Statistical
Manual of Mental Disorders, Fourth Edition, Text Revision (DSM-IV-TR; American Psychiatric
Association, 2000) and were based on the Structured Clinical Interview for DSM-IV (First et al.,
2002). To better characterize ADHD related symptoms, the Adult ADHD Interview (Kaufman et al.,
2000) was further used as a supplement. Out of all subjects, 1 had incomplete phenotype data from
the instruments used in this study, 10 had missing structural MRI (sMRI) data, and 10 had missing
resting-state functional MRI (fMRI) data. Fifty-five (55) subjects had an aliasing artifact in their
sMRI data potentially caused by the headset used in the scanner, whereas 22 subjects had errors in
the structural-functional alignment step during MRI preprocessing. These subjects were excluded
from the corresponding modeling analyses performed in this study. The final number of subjects
included in each modeling analysis is as follows: phenotype data, 271; sMRI, 206; fMRI, 229;

sMRI+fMRI, 178; phenotype+sMRI, 205; phenotype+fMRI, 228; phenotype+sMRI+fMRI: 177. The
demographic information from the subjects are given in **Table 1**.

2.2 Phenotype data

It should be noted that we used the term phenotype to broadly refer to behavioral and symptom
measures characterized by the clinical instruments. A total of 20 clinical instruments were
administered in the CNP dataset to capture a wide range of phenotype data including specific
behavioral traits and symptom dimensions (Poldrack et al., 2016). These instruments are either
clinician-rated or self-reported. While the clinician-rated questionnaires only covered relevant patient
groups, 13 self-reported clinical scales were given to all three patient groups as well as the heathy
controls. We therefore selected to only use subjects' answers to each of the individual questions
coming from these 13 self-reported scales as input features to our models because these scales
captured phenotypic features across all diagnostic groups as well as the HCs. It should be noted that
these 13 self-reported scales were not used to provide the official diagnosis in the CNP data since
they are not designed for such purposes. Specifically, the 13 self-reported scales used in this study
are: Chapman Social Anhedonia Scale, Chapman Physical Anhedonia Scale, Chapman Perceptual
Aberrations Scale, Hypomanic Personality Scale, Hopkins Symptom Checklist, Temperament and
Character Inventory, Adult ADHD Self-Report Scale v1.1 Screener, Barratt Impulsiveness Scale,
Dickman Functional and Dysfunctional Impulsivity Scale, Multidimensional Personality
Questionnaire – Control Subscale, Eysenck's Impulsivity Inventory, Scale for Traits that Increase
Risk for Bipolar II Disorder, and Golden and Meehl's Seven MMPI Items Selected by Taxonomic
Method. Together, these self-reported scales cover domains including symptoms, personality traits,
positive and negative affect, cognition, as well as sensory and social processing. The scores for
known sum and sub-scores within these self-reported instruments for each patient group as well as
the HCs are reported in **Supplementary Table 1**.

2.3 MRI data acquisition parameters

MRI data were acquired on one of two 3T Siemens Trio scanners both housed at the University of
California, Los Angeles. The sMRI data used in this study are T1-weighted and were acquired using
a magnetization-prepared rapid gradient-echo (MPRAGE) sequence with the following acquisition
parameters: TR = 1.9 s, TE = 2.26 ms, FOV = 250 mm, matrix = 256 x 256, 176 1-mm thick slices
oriented along the sagittal plane. The resting-state fMRI data contain a single run lasting 304 s. The
scan was acquired using a T2*-weighted echoplanar imaging (EPI) sequence using the following
parameters: 34 oblique slices, slice thickness = 4 mm, TR = 2 s, TE = 30 ms, flip angle = 90°, matrix
size 64 x 64, FOV = 192 mm. During the resting-state scan, subjects remained still and relaxed inside
the scanner, and kept their eyes open. No specific stimulus or task was presented to them.

2.4 MRI preprocessing

2.4.1 sMRI

Structural MRI preprocessing was implemented using Freesurfer's *recon-all* processing pipeline
(http://surfer.nmr.mgh.harvard.edu/). Briefly, the T1-weighted structural image from each subject
was intensity normalized and skull-stripped. The subcortical structures, white matter, and ventricles
were segmented and labeled according to the algorithm described in (Fischl et al., 2002). The pial
and white matter surfaces were then extracted and tessellated (Fischl et al., 2001), and cortical
parcellation was obtained on the surfaces according to a gyral-based anatomical atlas which partitions
each hemisphere into 34 regions (Desikan et al., 2006).

2.4.2 Resting-state fMRI

Resting-state fMRI preprocessing was implemented in AFNI (http://afni.nimh.nih.gov/afni).
Specifically, the first 3 volumes in the data were discarded to remove any transient magnetization
effects in the data. Spikes in the resting-state fMRI data were then removed and all volumes were
spatially registered with the 4th volume to correct for any head motion. The T1w structural image was
deobliqued and uniformized to remove shading artifacts before skull-stripping. The skull-stripped
structural image was then spatially registered with motion corrected fMRI data. The fMRI data were
further spatially smoothed using a 6-mm FWHM Gaussian kernel and converted to percent signal
change. Separately, the Freesurfer-generated aparc+aseg image from sMRI preprocessing was also
spatially registered with and resampled to have the same spatial resolution of the BOLD image.
Based on this, eroded white matter and ventricle masks were created, from which nuisance tissue
regressors were built based on non-spatially smoothed fMRI data to model and remove variances that
are not part of the BOLD signal. Specifically, we used the ANATICOR procedure (Jo et al., 2010),
where a locally averaged signal from the eroded white matter mask within a 25-mm radius spherical
region of interest (ROI) centered at each gray matter voxel was used to create a voxel-wise local
estimate of the white matter nuisance signal. This local estimate of the white matter nuisance signal,
along with the estimated head motions and average signal from the ventricles were detrended with a
4th order polynomial and then regressed out from the fMRI data. Finally, the clean resting-state fMRI
data was spatially normalized to the MNI template and resampled to have 2 mm isotropic voxels.
Note that in our preprocessing pipeline, the spatial normalization was performed after regressing out
nuisance signals. This allowed nuisance tissue regression to be performed in each subject's native
space to achieve a more accurate removal of these signals.

2.5 Feature extraction

We extracted measures from 3 data modalities as features: phenotype data from self-reported
instruments, measures derived from the sMRI data, and functional correlations based on resting-state
fMRI data. For phenotype features from self-reported instruments, we directly used subjects'
responses from a total of 578 questions from the above listed 13 instruments. Responses from non-
True/False type questions were normalized to have a range of between 0 and 1 to match those from
True/False type questions. For sMRI features, we specifically used 1) the volume of subcortical
structures generated by Freesurfer's subcortical volumetric segmentation, and 2) the area, thickness,
and volume of cortical brain regions estimated from Freesurfer's surface-based analysis pipeline. For
resting-state fMRI features, we first parceled the brain into 264 regions according to the atlas
proposed in (Power et al., 2011). A 5-mm radius spherical ROI was seeded according to the MNI
coordinates of each brain region specified in the atlas. Second, the clean resting-state BOLD time
series from all voxels within a given 5-mm radius spherical ROI were averaged to create the
representative time series for the brain region. Third, functional connectivity between ROIs was
estimated via the Pearson's correlation coefficient between the average time series from all pairs of
brain regions. This produced a 264-by-264 correlation matrix, from which 34,716 are unique
correlations between two distinct ROIs and were used as input features to the models.

2.6 Model fitting and feature importance weighting

The primary goals of machine learning analyses in this study were two-fold: 1) to identify important
features shared across a transdiagnostic patient group and 2) to evaluate the clinical utility of the
transdiagnostic features via classifiers that can reliably separate the patient group as a whole from
healthy controls. To achieve these goals, we built classifiers based on the logistic regression model as
implemented in the *scikit-learn* toolbox to classify patients from HCs. To identify predictive

transdiagnostic features embedded within each feature modality, separate logistic regression models
were independently trained using each of the above extracted feature modalities (i.e., item-level
phenotype data, sMRI measures, and resting-state fMRI correlations) as inputs and their
performances were evaluated in each of the transdiagnostic scenarios. Combinations of 2 and 3
feature modalities were also used as classifiers' inputs and their performances were evaluated in the
same fashion.

Because the number of features we extracted was relatively large compared to the sample size in
CNP data, an elastic net regularization term (Zou and Hastie, 2005) was added in all of our logistic
regression models to prevent overfitting. The use of elastic net regularization in our models also
enabled feature selection as the regularization induces sparse models via the grouping effect where
all the important features will be retained and the unimportant ones set to zero (Zou and Hastie, 2005;
Ryali et al., 2012). This allowed us to identify predictive features that are shared across multiple
patient categories.

We adopted the following procedure to determine the best regularization parameters. First, the input
data were randomly partitioned into a development set and an evaluation set. The development set
contains 80% of the data upon which a grid search with 3-fold cross validation procedure was
implemented to determine the best regularization parameters. Then the model with the best
regularization parameters was further tested on the remaining 20% of evaluation set. All features
were standardized to have zero mean and unit variance within the training data and the mean and
variance from the training data were used to standardize the corresponding test data. The entire
process was implemented 100 times. The following metrics were used to quantify the model
performances: area under the receiver operating characteristics curve (AUC), accuracy, sensitivity,
and specificity. The mean and standard deviation of the above metrics over the 100 evaluation sets
were reported.

From the above models, the predictive power of each feature is assessed via the weights of the
logistic regression model in our transdiagnostic classifiers. For each feature, we calculated its
corresponding standardized model weight (mean model weight divided by the standard deviation)
across the 100 model implementations as the proxy for feature importance. Features with large
importance values from our transdiagnostic classifiers are potentially symptoms, traits, and
neuropathological mechanisms shared across patient groups but are distinct from healthy controls.

To identify the set of most predictive transdiagnostic features within a given data modality, we used
the following feature importance-guided sequential model selection procedure. Specifically, we first
rank ordered the features in the classifiers according to their standardized model weights. Next, a
series of truncated models was built such that each model only takes the top k most predictive
features as inputs to perform the same classification tasks. The model weights and the best
regularization parameters for each truncated model were estimated via the 3-fold cross validation
procedure within the development set. We let k range from the top 1 most predictive feature to all
available features in steps of 1 for phenotype features, sMRI features, and the combination of the two
feature sets. For any feature or feature combinations involving fMRI correlations, because of the
significantly increased feature dimension, the k's were chosen from a geometric sequence with a
common ratio of 2 (i.e., 1, 2, 4, 8, 16, …). The sequential model selection procedure was
implemented 10 times for each feature set. Model performances based on the evaluation set were
obtained for each truncated model and were evaluated as a function of the number of top features (k)
included in each truncated model to determine the optimal feature set.

2.7   Statistical analyses

To statistically examine whether the models' performances are significantly above chance level, we
performed a random permutation test where labels in the training data (e.g., HC vs. Patients) were
shuffled 100 times and truncated models based on the best set of features were trained on these label-
shuffled data using exactly the same approach as described above (Ojala and Garriga, 2009). The
performances from the 100 models were used to construct the empirical null distribution against
which the performance of the best truncated models based on the actual unshuffled data was then
compared. This random permutation test procedure also helped us to determine whether overfitting
occurred during training.

To evaluate differences in sum scores obtained from the top features between HC and patients, we
used two sample t-tests on the sample means since the sum scores are quasi normally distributed.
Effect sizes were measured using Cohen's d, which captures the shift in mean scaled by the data's
standard deviation. Tests on the difference in AUC between the full model and the best truncated
model were carried out via the Wilcoxon's rank-sum test.

3   Results

In total, we trained classifiers to distinguish HCs from the patient group as a whole based on 7 sets of
features by either using each individual feature modality (self-reported instruments, sMRI, and fMRI)
or combinations of 2 or 3 feature modalities (e.g., instruments+sMRI+fMRI). The classifiers'
performances using each of the 7 feature sets for the HC vs. Patients classifier are reported in **Table**
2. Overall, classifiers trained on feature sets involving phenotypical data from self-reported
instruments (i.e., scales and scales + MRI feature sets) outperformed those only trained on MRI
features (sMRI, fMRI, and sMRI+fMRI). For classifiers using features involving these instruments,
the mean AUC ranged from 0.83 to 0.89 (mean accuracy: 0.77 – 0.91), whereas the mean AUC
ranged from 0.56 to 0.59 (mean accuracy: 0.58 – 0.61) for MRI feature sets.

Next, to identify the optimal set of transdiagnostic features shared across the CNP patient population
that are highly distinct from HC, we examined the performance measures from the best truncated
classification models during sequential model selection (**Figure 1** and **Table 2;** see **Supplementary**
Fig. 1 for AUC as a function of input feature dimensions). Significantly improved performances were
obtained from the best truncated classification models compared with the corresponding models
using the full sets of features (all p's < 0.05, FDR corrected, as assessed by the rank-sum test; **Table**
2). The AUCs from all feature sets were also significantly above chance level as assessed via the
random permutation test (all p's < 0.01, FDR corrected; **Supplementary Fig. 2**). Additionally, the
computational time for the importance guided sequential model selection method grew linearly as the
number of features increased, which is highly efficient compared to the brute force feature selection
procedure (exponential time complexity; **Supplementary Fig. 3**).

The truncated classification model involving data from the self-reported instruments alone had high
performance of distinguishing patients from HCs with the mean AUC being 0.95 (accuracy: 0.88;
sensitivity: 0.87; specificity: 0.88; **Figure 1**; **Table 2**). This truncated model selected 85 items as the
most predictive features from the total of 578 items contained in the 13 self-reported instruments.
Moreover, only 10 items were needed to achieve an AUC of 0.90 (accuracy: 0.81; sensitivity: 0.79;
specificity: 0.84), suggesting that a concise scale can be constructed potentially for screening
purposes. The model involving data from self-reported instruments performed better compared to

those using feature sets based solely on MRI (mean AUC ranging from 0.77 to 0.87; mean accuracy
ranging from 0.71 to 0.85). Combining MRI features with data from instruments only slightly
improved the model performance (mean AUC being 0.96 – 0.98) (**Figure 1**; **Table 2**). Taken
together, this indicates that the phenotypical data captured by the 13 self-reported instruments contain
a set of transdiagnostic features common across the patient populations studied while distinguishing
them from the healthy controls. We hence focused on discussing these transdiagnostic phenotypical
features below.

It should be noted that small yet systematic differences exist between patients and HCs in the CNP
dataset despite the effort to fully counterbalance subjects' demographic information during
recruiting. From the demographic information listed in **Table 1**, a statistical test on age showed that
the median age for patients were significantly higher than HCs (Mann-Whitney's U test: $p = 0.001$).
The gender ratio on the other hand did not significantly differ between patients and HCs (Z-test on
proportions: $p = 0.14$). Patients had significantly lower years of education compared with HCs
(Mann-Whitney's U test: $p < 0.001$). The racial distribution also showed a significant difference
between patients and HCs (χ^2-test: $p = 0.004$). To ensure that the results were not mainly driven by
differences in demographics, we built classifiers based solely on these demographic variables. The
mean test AUC on the evaluation set for the best model was 0.71 ($SD = 0.06$), which was
substantially lower than the performance of both the full model and the best truncated model based
on the self-reported instruments alone (mean AUCs being 0.83 and 0.95, respectively). Additionally,
despite mixing the demographic variables with the self-reported instrument items slightly improved
model performance compared to those obtained from instruments alone (mean AUC for truncated
model using demographics and instruments: 0.98), the highest-ranking demographic variable ranked
only 43[rd] among all instrument items and demographic features. Taken together, these results suggest
that our models were unlikely to be mainly driven by demographic differences.

A simple sum score constructed by adding up an individual's responses to the 85 most predictive
items (with item responses having negative model coefficients reversed) demonstrated high
separability between healthy controls and patients (Cohen's $d = 2.85$; test on the difference in sample
mean: $t = 10.27$, $p < 0.001$; **Figure 2A**). The separability based on the top 10 most predictive items
(corresponding AUC = 0.90) was also very high (Cohen's $d = 2.16$; test on the difference in sample
mean: $t = 18.13$, $p < 0.001$). The sum scores had higher separability than other known sum
scores/sub-scores of the self-reported instruments (**Figure 2C & 2D; Supplementary Table 2**),
indicating the items selected by the sequential model selection procedure do not adhere to known
dimensional structures within the instruments. Calculating the sum score for each individual patient
category showed that all 3 patient categories had elevated sum scores compared to healthy controls (t
> 5.671, $p < 0.001$); yet, the difference between the patient categories was insignificant ($t < 1.940$, p
> 0.056; **Figure 2B**). This suggests that the 85 items captured transdiagnostic phenotypic features
shared across the patient groups as a whole rather than driven by a single patient category.

Figure 3 illustrates the proportion of questionnaire items selected from each instrument that were
included in transdiagnostic set of phenotypic features of the best truncated model (i.e. the one with
the highest AUC). Items from all 13 instruments were selected to be among the top features by the
classifiers. Overall, these instruments measure a wide range of phenotype and symptom domains
encompassing personality traits, positive and negative affect (reward/anhedonia, fear, and anxiety),
cognition (attention, response inhibition), sensory processing (perceptual disturbances), and social
processing. While all items included among the set of transdiagnostic phenotypic features jointly
formed a highly predictive set to distinguish patients from healthy controls, the Temperament and
Character Inventory (TCI) contributed the largest proportion of items in the set of transdiagnostic

features. The proportion of TCI items selected among the 85 most predictive items (32.9%)
significantly exceeded the proportion of all TCI items among all 578 items from the 13 instruments
(24.0%; p = 0.04 as assessed via the Z-test). The disproportionately high number of TCI items
indicates that certain personality traits are strong predictors of shared psychopathology regarding
SCZ, BD, and ADHD.

To better understand which features strongly predicted psychopathology, we focused on the top 20
items that contributed the largest magnitude of model weights. We identified the specific
behavioral/symptom phenotypes characterized by each item and then grouped the items accordingly.
Among personality traits, the top transdiagnostic phenotypes included neuroticism, extraversion, and
impulsivity (**Figure 4A**); whereas the top symptom domains consisted of mood dysregulation,
inattention, hyperactivity/agitation, and social anhedonia and apathy. In addition, the importance of
religion was also a shared feature across patients (**Figure 4A**; see **Supplementary Table 4** for item
grouping). We next compared the most predictive transdiagnostic features with those most predictive
of a single patient category from healthy controls to identify category-specific differences (**Figure
4B-D**; see **Supplementary Table 3 and Supplementary Fig. 4** for classification results between HC
and each patient category). SCZ patients exhibited additional features including perceptual
aberration, physical anhedonia, and psychological distress that are not among the top transdiagnostic
features. On the other hand, extraversion, impulsivity, inattention, and religion which were present in
the transdiagnostic features set were not among the most predictive features for SCZ (**Figure 4B** and
Supplementary Table 5). By contrast, transdiagnostic features overlapped with BD patient-specific
features. Nonetheless, BD patients exhibited additional features of increased energy, psychological
distress and physical anhedonia; yet psychomotor agitation and neuroticism contributed little
predictive value (**Figure 4C**, **Supplementary Table 6**). ADHD patients were effectively classified
by additional features such as indecision and physical anhedonia, with little predictive contribution
from apathy, neuroticism, and religion (**Figure 4D**, **Supplementary Table 7**).

4   Discussion

In this study, using self-reported instruments provided in the CNP dataset, we generated predictive
models to identify a set of transdiagnostic phenotypic features that were shared across SCZ, BD, and
ADHD. These models were quantified for performance (e.g. accuracy, sensitivity and specificity) and
were interpretable along dimensions of personality traits and symptom domains. We found the set of
85 items is highly predictive of the patient group as a whole from HCs. To our surprise, a compact
model of only 10 items is sufficient to achieve a performance AUC value of 0.90. Further, we
demonstrated that a simple sum score can be calculated to enable high separability between patients
and HCs. Our importance-guided sequential model selection approach revealed which phenotypical
features were shared across transdiagnostic patient groups. Within each patient population, we also
show which abnormal psychopathological personality traits and symptom domains deviated from the
transdiagnostic classifier. Importantly, many of these features are consistent with established clinical
intuition. Taken together, this study offers new perspectives on the shared psychopathology across
SCZ, BD, and ADHD and underscores the potential of creating a short transdiagnostic screening
scale based on the selected items.

The application of machine learning to systematically search for consistent patterns in clinical data
across disease categories defined in DSM is an emerging trend in the field of computational
psychiatry (Bzdok and Meyer-Lindenberg, 2017). Nonetheless, our approach to identifying

transdiagnostic features in psychiatric disorders differs both conceptually and methodologically from
previous studies. Numerous investigators have focused on patient subtyping within a given disorder
(Rhebergen et al., 2011; Lamers et al., 2012; van Loo et al., 2012, 2014; Georgiades et al., 2013;
Brodersen et al., 2014; Doshi-Velez et al., 2014; Lewandowski et al., 2014; van Hulst et al., 2014;
Veatch et al., 2014; Costa Dias et al., 2015; Geisler et al., 2015; Sun et al., 2015; Clementz et al.,
2016; Drysdale et al., 2016; Mostert et al., 2018) or have mined transdiagnostic symptom dimensions
underlying various psychiatric disorders (Grisanzio et al., 2017; Elliott et al., 2018; Xia et al., 2018a,
b). Among studies examining the transdiagnostic symptom dimensions, most adopted an
unsupervised machine learning predictive framework. However, the differences in distance/similarity
metrics used, coupled with the lack of ground truth in the unsupervised machine learning algorithms
used to detect the transdiagnostic structure, make it difficult to validate the clinical utility of the
identified features. We designed our study to overcome these limitations. To our best knowledge, our
study is the first to use feature importance to guide forward model selection under a supervised
machine learning framework to identify transdiagnostic psychopathological features across multiple
DSM categories. The high performance of our truncated models selected via the model selection
approach demonstrate the potential clinical utility of the identified transdiagnostic features.

Though we built models with different modalities as inputs (e.g. personality traits, symptoms and
neuroimaging), we found high performance models could be obtained without significant
contribution of the imaging modalities. This finding contrasts with what would be predicted from the
published literature. For example, a recent meta-analysis of studies on psychiatric disorders involving
structural magnetic resonance imaging (sMRI) identified shared abnormalities in certain brain
regions underlying common psychiatric disorders (Goodkind et al., 2015). In addition, studies using
functional MRI (fMRI) found altered functional connectivity patterns shared across multiple
categories of disorders such as SCZ, BD, and major depressive disorder (MDD) (Buckholtz and
Meyer-Lindenberg, 2012; Wei et al., 2018). Similarly, another recent study focusing on MDD, post-
traumatic stress disorder, and panic disorder identified 6 distinct subtypes based on 3 orthogonal
symptom dimensions shared across the DSM diagnoses and their corresponding biomarkers in
electroencephalogram (EEG) beta activity (Grisanzio et al., 2017). Although these studies did not
systematically compare the predictability in each data modality, it is possible that the sample size in
CNP or other methodological differences (e.g., parcellation used during sMRI and fMRI feature
extraction) limited the weighted importance of structural or functional measures in our models.

A broad set of behavioral phenotypic features from the self-report instruments were identified by our
sequential model selection procedure to be shared across the three patient groups. The phenotypes are
distributed across all 13 self-reported instruments and covers symptom domains encompassing
personality and traits, positive and negative affect, cognition, sensory and social processing. It should
be noted that these 13 self-reported instruments are not designed to yield diagnoses. Therefore, a set
of phenotypic features that can be identified algorithmically and used to distinguish patients from
HCs with an accuracy level close to a clinician's performance demonstrates the potential clinical
utility of the phenotypic features and the sequential model selection approach. For the top 20 most
predictive features, mood dysregulation, impulsivity, inattention, neuroticism, social anhedonia and
apathy weighted prominently in the transdiagnostic model. This high level of shared symptom
domains across SCZ, BD, and ADHD is in line with recent genetic studies reporting significantly
correlated risk factors for heritability among these three disorders (Larsson et al., 2013; The
Brainstorm Consortium et al., 2017; Bipolar Disorder and Schizophrenia Working Group of the
Psychiatric Genomics Consortium et al., 2018). For SCZ and BD, previous studies have identified
shared features both in terms of symptoms and the underlying psychopathology and biology
(Pearlson, 2015). Similarly, studies have identified shared symptoms and biology between SCZ and

ADHD (Peralta et al., 2011; Park et al., 2018) and have found high levels of comorbidity between
BD and ADHD along with the shared features between the two disorders (Nierenberg et al., 2005;
Klassen et al., 2010; van Hulzen et al., 2017; Wang et al., 2017). Despite these prior studies, the three
diagnostic categories have not been considered together in a single study. Consistent with the
findings reported in these studies, our study provides an important data-driven confirmation on the
shared behavioral and symptoms features across the three disease categories.

Since the TCI is less commonly used in clinical practice and historically a greater emphasis has been
placed on symptoms than personality traits, we were surprised by the finding that the TCI contributed
the largest proportion of questions among the set of 85 most predictive items determined by the
transdiagnostic classifier. Prior studies have found that the personality traits and characters defined in
the TCI are associated with various mood disorders (Cloninger et al., 1998; Grucza et al., 2003).
Specifically, for disorders in the CNP dataset, studies have found positive association between
personality dimensions characterized in TCI and overall ADHD symptom (Lynn et al., 2005;
Anckarsäter et al., 2006) as well as subtypes of ADHD (Salgado et al., 2009). For SCZ, studies have
identified links between positive and negative symptom dimensions and TCI factors (Guillem et al.,
2002; Hori et al., 2008). Among BD patients, (Hajirezaei et al., 2017) identified personality profiles
that are distinct from healthy controls and these profiles were further found to be shared with MDD.
Since these studies associated disease symptoms solely with the known factor scores in the TCI, the
contribution of the nuanced personality profiles captured in individual items in the TCI could not be
determined. In the current study, the fact that we identified items in the TCI that corresponded to
shared symptoms such as apathy, anhedonia, and distress extends prior literature and is consistent
with studies documenting the relationship between TCI factor scores and symptoms such as
anhedonia (Martinotti et al., 2008), as well as depression and anxiety (Jylhä and Isometsä, 2006).
Additionally, prior studies only examined TCI's association with symptoms without simultaneously
including other instruments as covariates in the model. Such an approach cannot evaluate the relative
importance of the personality traits in TCI against the broader set of phenotypical features defined in
other instruments. In this regard, our study established the usefulness of personality traits as a set of
reliable transdiagnostic features among all features defined in the self-reported instruments in the
CNP data.

The sum score of the 85 most predictive transdiagnostic items achieved much higher separability
between HC and patients than known sub-scores and sum scores in the instruments that were
specifically designed to assess diagnosis-specific symptom domains. This is true even for the subset
of top 10 most predictive transdiagnostic items, which indicates that the shared phenotypic features
across patient groups do not fully adhere to known dimensional structures in the instruments. Thus,
using the total score and/or the sub-scores according to pre-defined subscales of a given instrument
cannot identify the optimal set of transdiagnostic features. One explanation for this phenomenon is
that because the patients share a broad range of phenotypic features, the pre-defined subscales and
sum scores become insufficient in capturing the full dimensional structure since most of the
instruments are designed to measure a limited set of constructs targeting a specific patient population
(Avila et al., 2015). This further demonstrates the advantage of our importance-guided sequential
model selection approach in identifying potentially clinically relevant transdiagnostic features across
a large set of instruments.

While patients shared a broad set of phenotypic features, our results showed deviations from this
transdiagnostic structure within the most predictive features for each patient group. These differences
may in particular reflect clustered personality traits and symptom domains that are most unique for
each patient population. For SCZ, the unique features of perceptual aberration and psychological

distress, along with other features that are consistent with the transdiagnostic structure, largely
conform to the positive and negative symptom dimensions associative with SCZ patients. For BD,
the unique features of increased energy, physical anhedonia, and psychological distress serve to
shape the symptom structure along the manic and depressive dimensions. For ADHD, the increased
representation in inattention, hyperactivity, and impulsivity is consistent with the overall
symptomatology of ADHD patients. Overall, the concurrent existence of shared and category-
specific phenotypic features across the CNP patient groups is consistent with recent studies reporting
both shared and distinct properties in functional brain networks (Grisanzio et al., 2017; Xia et al.,
2018a, b) and genetic neuropathology (The Brainstorm Consortium et al., 2017; Gandal et al., 2018)
across major psychiatric disorders. Our results raise the possibility of exploring the relationship
between the predictive phenotypic features and the underlying genetics of the individuals or groups
that present with these features.

In conclusion, we identified a set of transdiagnostic phenotypic features shared across SCZ, BD, and
ADHD. This set of features distinguished the patient group from HC with high accuracy and a
compact transdiagnostic screening scale can be derived from the corresponding top 10 most
predictive questionnaire items. The feature importance guided sequential model selection provides a
data-driven method to identify shared features under a supervised machine learning framework, in
which the performance of the identified feature sets is evaluated on unseen data. This is an advantage
over unsupervised machine learning methods. Moreover, the importance guided sequential model
selection can be generalized to identify clinically-useful transdiagnostic features across categories
defined in DSM-5 and ICD-10, or alternatively to identify the neural correlates of symptom severity
across psychiatric disorders (Mellem et al., 2018). It should be noted that the medication status in the
CNP dataset is not controlled. This suggests that although reliable transdiagnostic features could be
identified across patient groups, the underlying cause of the observed symptom structure could
potentially be confounded by the uncontrolled medication and symptom status. Future studies should
further validate the transdiagnostic features identified in this study on other datasets with similar
patient populations and with better controlled medication status. Including these additional datasets
as out-of-sample validations can demonstrate the generalizability of the current results and
methodology to the wider population. Additionally, because the phenotypic features largely reflected
behaviors and symptoms through self-reports, the high performance of our models based on these
features may be reflecting the close mapping between the identified features and symptom-based
diagnostic criteria in DSM. Nevertheless, our findings represent an important step in the ongoing
effort to characterize clinically useful transdiagnostic phenotypes. Future studies could potentially
investigate the underlying brain circuits associated with these clinically-relevant phenotypic features.

5　Conflict of Interest

The authors are employees of BlackThorn Therapeutics, Inc, and are compensated financially by
BlackThorn Therapeutics, Inc.

6　Author Contributions

YL, MM, HG, WM, PA designed the study; YL, MM, HG, MK, ARM, AM performed the analysis;
YL, MM, HG, MK, ARM, WM, PA wrote the manuscript.

7   Funding

525   This work is funded by BlackThorn Therapeutics, Inc.

8   Acknowledgments

528   The authors would like to thank Clark Gao, Lori Jean Van Orden, Martine Meyer, and Simone
529   Krupka for their help and discussions in shaping this work.

9   References

Abrams R., & Taylor M. (1981). Importance of schizophrenic symptoms in the diagnosis of mania.
Am J Psychiatry, 138(5):658–661.

American Psychiatric Association. (2000). Diagnostic and statistical manual of mental disorders. (4th
ed.). Washington, DC: American Psychiatric Press Inc.

Anckarsäter H., Stahlberg O., Larson T., Hakansson C., Jutblad S.B., Niklasson L., Nydén A., Wentz
E., Westergren S., Cloninger C.R., Gillberg C., & Rastam M. (2006). The Impact of ADHD and
autism spectrum disorders on temperament, character, and personality development. Am J
Psychiatry, 163(7):1239–1244.

Avila M., Stinson J., Kiss A., Brandão L.R., Uleryk E., & Feldman B.M. (2015). A critical review of
scoring options for clinical measurement tools. BMC Res Notes 8:612.

Bipolar Disorder and Schizophrenia Working Group of the Psychiatric Genomics Consortium.
(2018). Genomic Dissection of Bipolar Disorder and Schizophrenia, Including 28 Subphenotypes.
Cell 173(7):1705-1715.e16.

Brodersen K.H., Deserno L., Schlagenhauf F., Lin Z., Penny W.D., Buhmann J.M., & Stephan K.E.
(2014). Dissecting psychiatric spectrum disorders by generative embedding. Neuroimage Clin 4:98–
111.

Buckholtz J.W., & Meyer-Lindenberg A. (2012). Psychopathology and the human connectome:
toward a transdiagnostic model of risk for mental illness. Neuron 74(6):990–1004.

Bzdok D., & Meyer-Lindenberg A. (2017). Machine Learning for Precision Psychiatry:
Opportunities and Challenges. Biol Psychiatry Cogn Neurosci Neuroimaging 3(3):223-230.

Clementz B.A., Sweeney J.A., Hamm JP, Ivleva EI, Ethridge LE, Pearlson GD, Keshavan MS, &
Tamminga CA (2016) Identification of Distinct Psychosis Biotypes Using Brain-Based Biomarkers.
Am J Psychiatry 173(4):373–384.

Cloninger CR, Bayon C, & Svrakic DM (1998) Measurement of temperament and character in mood
disorders: a model of fundamental states as personality types. J Affect Disorder 51(1):21–32.

Coryell W, Leon AC, Turvey C, Akiskal HS, Mueller T, & Endicott J (2001) The significance of
psychotic features in manic episodes: a report from the NIMH collaborative study. J Affect Disord
67(1-3):79–88.

Cross-Disorder Group of the Psychiatric Genomics Consortium (2013) Identification of risk loci with
shared effects on five major psychiatric disorders: a genome-wide analysis. Lancet 381(9875):1371–
1379.

Desikan RS, Ségonne F, Fischl B, Quinn BT, Dickerson BC, Blacker D, Buckner RL, Dale AM,
Maguire PR, Hyman BT, Albert MS, & Killiany RJ (2006) An automated labeling system for
subdividing the human cerebral cortex on MRI scans into gyral based regions of interest.
Neuroimage 31(3):968–980.

Costa Dias TG, Iyer SP, Carpenter SD, Cary RP, Wilson VB, Mitchell SH, Nigg JT, & Fair DA
(2015) Characterizing heterogeneity in children with and without ADHD based on reward system
connectivity. Dev Cogn Neurosci 11:155–174.

Doshi-Velez F, Ge Y, & Kohane I (2014) Comorbidity clusters in autism spectrum disorders: an
electronic health record time-series analysis. Pediatrics 133(1):e54–63.

Drysdale AT, Grosenick L, Downar J, Dunlop K, Mansouri F, Meng Y, … Liston C (2016) Resting-
state connectivity biomarkers define neurophysiological subtypes of depression. Nat Med 23(1):28-
38.

Elliott ML, Romer AL, Knodt AR, & Hariri AR (2018) A Connectome-Wide Functional Signature of
Transdiagnostic Risk for Mental Illness. Biol Psychiat 84(6):452-459.

First, M. B., Spitzer, R. L., Gibbon, M. & Williams, J. B.W. Structured Clinical Interview for DSM-
IV-TR Axis I Disorders, Research Version, Patient Edition. (SCID-I/P) (2002).

Fischl B, Liu A, Dale AM (2001) Automated Manifold Surgery: Constructing Geometrically
Accurate and Topologically Correct Models of the Human Cerebral Cortex. IEEE Trans Med
Imaging 20:70.

Fischl B, Salat DH, Busa E, Albert M, Dieterich M, Haselgrove C, van der Kouwe A, Killiany R,
Kennedy D, Klaveness S, Montillo A, Makris N, Rosen B, Dale AM (2002) Whole Brain
Segmentation Automated Labeling of Neuroanatomical Structures in the Human Brain. Neuron
33:341–355.

Gandal MJ, Haney JR, Parikshak NN, Leppa V, Ramaswami G, Hartl C, Schork AJ, Appadurai V,
Buil A, Werge TM, Liu C, White KP, Consortium C, Consortium P, Group iPSYCH-B, Horvath S,
Geschwind DH (2018) Shared molecular neuropathology across major psychiatric disorders parallels
polygenic overlap. Science 359:693–697.

Geisler D, Walton E, Naylor M, Roessner V, Lim KO, Schulz CS, Gollub RL, Calhoun VD,
Sponheim SR, Ehrlich S (2015) Brain structure and function correlates of cognitive subtypes in
schizophrenia. Psychiatry Res Neuroimaging 234:74–83.

Georgiades S, Szatmari P, Boyle M, Hanna S, Duku E, Zwaigenbaum L, Bryson S, Fombonne E,
Volden J, Mirenda P, Smith I, Roberts W, Vaillancourt T, Waddell C, Bennett T, Thompson A, in
Team P (2013) Investigating phenotypic heterogeneity in children with autism spectrum disorder: a
factor mixture modeling approach. J Child Psychol Psyc 54:206–215.

Gheiratmand M, Rish I, Cecchi GA, Brown MR, Greiner R, Polosecki PI, Bashivan P, Greenshaw
AJ, Ramasubbu R, Dursun SM (2017) Learning stable and predictive network-based patterns of
schizophrenia and its clinical symptoms. Npj Schizophrenia 3:22.

Goodkind M, Eickhoff SB, Oathes DJ, Jiang Y, Chang A, Jones-Hagata LB, Ortega BN, Zaiko YV,
Roach EL, Korgaonkar MS, Grieve SM, Galatzer-Levy I, Fox PT, Etkin A (2015) Identification of a
Common Neurobiological Substrate for Mental Illness. JAMA Psychiatry 72:305–315.

Grisanzio KA, Goldstein-Piekarski AN, Wang M, Ahmed AP, Samara Z, Williams LM (2017)
Transdiagnostic Symptom Clusters and Associations With Brain, Behavior, and Daily Function in
Mood, Anxiety, and Trauma Disorders. JAMA Psychiatry 75(2):201-209.

Grucza RA, Przybeck TR, Spitznagel EL, Cloninger CR (2003) Personality and depressive
symptoms: a multi-dimensional analysis. J Affect Disorders 74:123–130.

Guillem F, Bicu M, Semkovska M, Debruille JB (2002) The dimensional symptom structure of
schizophrenia and its association with temperament and character. Schizophr Res 56:137–147.

Hajirezaei S, Mohammadi A, Soleimani M, Rahiminezhad F, Mohammadi M, Cloninger RC (2017)
Comparing the Profile of Temperament and Character Dimensions in Patients with Major Depressive
Disorder and Bipolar Mood Disorder with a Control Group. Iranian J Psychiatry 12:147–153.

Hamshere ML, Stergiakouli E, Langley K, Martin J, Holmans P, Kent L, Owen MJ, Gill M, Thapar
A, O'Donovan M, Craddock N (2013) Shared polygenic contribution between childhood attention-
deficit hyperactivity disorder and adult schizophrenia. Br J Psychiatry 203:107–111.

Hori H, Noguchi H, Hashimoto R, Nakabayashi T, Saitoh O, Murray RM, Okabe S, Kunugi H (2008)
Personality in schizophrenia assessed with the Temperament and Character Inventory (TCI). Psychiat
Res 160:175–183.

Jo H, Saad ZS, Simmons KW, Milbury LA, Cox RW (2010) Mapping sources of correlation in
resting state FMRI, with artifact detection and removal. Neuroimage 52:571–582.

Jylhä P, Isometsä E (2006) Temperament, character and symptoms of anxiety and depression in the
general population. Eur Psychiat 21(6):389-395.

Kaufman, J., Birmaher, B., Brent, D. A., Ryan, N. D. & Rao, U. (2000). K-SADS-PL. J Am Acad
Child Adolesc Psychiatry 39(10): 1208.

Keck PE, McElroy SL, Havens J, Altshuler LL, Nolen WA, Frye MA, Suppes T, Denicoff KD,
Kupka R, Leverich GS, Rush AJ, Post RM (2003) Psychosis in bipolar disorder: phenomenology and
impact on morbidity and course of illness. Compr Psychiat 44:263–269.

Kessler R, Gruber M, Hettema J, Hwang I, Sampson N, Yonkers K (2007) Co-morbid major
depression and generalized anxiety disorders in the National Comorbidity Survey follow-up. Psychol
Med 38:365–374.

Kessler RC, Demler O, Frank RG, Olfson M, Pincus H, Walters EE, Wang P, Wells KB, Zaslavsky
AM (2005) Prevalence and Treatment of Mental Disorders, 1990 to 2003. New Engl J Medicine
352:2515–2523.

Klassen LJ, Katzman MA, Chokka P (2010) Adult ADHD and its comorbidities, with a focus on
bipolar disorder. J Affect Disorders 124:1–8.

Krueger RF, Markon KE (2006) Reinterpreting Comorbidity: A Model-Based Approach to
Understanding and Classifying Psychopathology. Annu Rev Clin Psycho 2:111–133.

Krueger RF, Markon KE (2011) A Dimensional-Spectrum Model of Psychopathology: Progress and
Opportunities. Arch Gen Psychiat 68:10–11.

Lamers F, Burstein M, He J, Avenevoli S, Angst J, Merikangas KR (2012) Structure of major
depressive disorder in adolescents and adults in the US general population. Br J Psychiatry 201:143–
150.

Larsson H, Rydén E, Boman M, Långström N, Lichtenstein P, Landén M (2013) Risk of bipolar
disorder and schizophrenia in relatives of people with attention-deficit hyperactivity disorder. Br J
Psychiatry 203:103–106.

Lewandowski K, Sperry S, Cohen B, Öngür D (2014) Cognitive variability in psychotic disorders: a
cross-diagnostic cluster analysis. Psychol Med 44:3239–3248.

Lotan A, Michaela Fenckova , Janita Bralten, Aet Alttoa, Luanna Dixson, Robert W. Williams and
Monique van der Voet (2014) Neuroinformatic analyses of common and distinct genetic components
associated with major neuropsychiatric disorders. Frontiers in Neuroscience 8: 1-15.

Lynn DE, Lubke G, Yang M, McCracken JT, McGough JJ, Ishii J, Loo SK, Nelson SF, Smalley SL
(2005) Temperament and Character Profiles and the Dopamine D4 Receptor Gene in ADHD. Am J
Psychiat 162:906–913.

Markon K (2009) Modeling psychopathology structure: a symptom-level analysis of Axis I and II
disorders. Psychol Med 40:273–288.

Martinotti G, Cloninger CR, Janiri L (2009) Temperament and character inventory dimensions and
anhedonia in detoxified substance-dependent subjects. Am J Drug Alcohol Abus 34(2):177-183.

Mellem SM, Liu Y, Gonzalez H, Kollada M, Martin WJ, Ahammad P (2018) Machine learning
models identify multimodal measurements highly predictive of transdiagnostic symptom severity for
mood, anhedonia, and anxiety. BioRxiv, doi: https://doi.org/10.1101/414037

Mostert JC, Hoogman M, Onnink MA, van Rooij D, von Rhein D, van Hulzen KJ, Dammers J, Kan
CC, Buitelaar JK, Norris DG, Franke B (2018) Similar Subgroups Based on Cognitive Performance
Parse Heterogeneity in Adults with ADHD and Healthy Controls. J Atten Disord 22:281–292.

Nierenberg AA, Miyahara S, Spencer T, Wisniewski SR, Otto MW, Simon N, Pollack MH, Ostacher
MJ, Yan L, Siegel R, Sachs GS, Investigators S-B (2005) Clinical and Diagnostic Implications of
Lifetime Attention-Deficit/Hyperactivity Disorder Comorbidity in Adults with Bipolar Disorder:
Data from the First 1000 STEP-BD Participants. Biol Psychiat 57:1467–1473.

Ojala M, Garriga GC (2009) Permutation Tests for Studying Classifier Performance. 2009 Ninth Ieee
Int Conf Data Min:908–913.

Pacheco A, Barguil M, Contreras J, Montero P, Dassori A, Escamilla MA, Raventós H (2010) Social
and clinical comparison between schizophrenia and bipolar disorder type I with psychosis in Costa
Rica. Soc Psych Psych Epid 45:675–680.

Park MM, Raznahan A, Shaw P, Gogtay N, Lerch JP, Chakravarty MM (2018) Neuroanatomical
phenotypes in mental illness: identifying convergent and divergent cortical phenotypes across autism,
ADHD and schizophrenia. J Psychiatry Neurosci Jpn 43:201–212.

Pearlson GD (2015) Etiologic, Phenomenologic, and Endophenotypic Overlap of Schizophrenia and
Bipolar Disorder. Annu Rev Clin Psycho 11:1–31.

Peralta V, de Jalón E, Campos MS, Zandio M, Sanchez-Torres A, Cuesta MJ (2011) The meaning of
childhood attention-deficit hyperactivity symptoms in patients with a first-episode of schizophrenia-
spectrum psychosis. Schizophr Res 126:28–35.

Poldrack RA, Congdon E, Triplett W, Gorgolewski KJ, Karlsgodt KH, Mumford JA, Sabb FW,
Freimer NB, London ED, Cannon TD, Bilder RM (2016) A phenome-wide examination of neural
and cognitive function. Sci Data 3:sdata2016110.

Pope HG, Lipinski JF (1978) Diagnosis in Schizophrenia and Manic-Depressive Illness: A
Reassessment of the Specificity of "Schizophrenic" Symptoms in the Light of Current Research.
Arch Gen Psychiat 35:811–828.

Power JD, Cohen AL, Nelson SM, Wig GS, Barnes K, Church JA, Vogel AC, Laumann TO, Miezin
FM, Schlaggar BL, Petersen SE (2011) Functional Network Organization of the Human Brain.
Neuron 72:665–678.

Purcell SM et al. (2009) Common polygenic variation contributes to risk of schizophrenia and
bipolar disorder. Nature 460:748.

Rashid B, Damaraju E, Pearlson GD, Calhoun VD (2014) Dynamic connectivity states estimated
from resting fMRI Identify differences among Schizophrenia, bipolar disorder, and healthy control
subjects. Front Hum Neurosci 8:897.

Rhebergen D, Lamers F, Spijker J, de Graaf R, Beekman A, Penninx B (2011) Course trajectories of
unipolar depressive disorders identified by latent class growth analysis. Psychol Med 42:1383–1396.

Ryali S, Chen T, Supekar K, Menon V (2012) Estimation of functional connectivity in fMRI data
using stability selection-based sparse partial correlation with elastic net penalty. Neuroimage
59:3852–3861.

Salgado CAI, Bau CHD, Grevet EH, Fischer AG, Victor MM, Kalil KLS, Sousa NO, Garcia CR,
Belmonte-de-Abreu P (2009) Inattention and Hyperactivity Dimensions of ADHD Are Associated
with Different Personality Profiles. Psychopathology 42:108–112.

Sha Z, Wager TD, Mechelli A, He Y (2018) Common Dysfunction of Large-Scale Neurocognitive
Networks across Psychiatric Disorders. Biol Psychiat.

Sun H, Lui S, Yao L, Deng W, Xiao Y, Zhang W, Huang X, Hu J, Bi F, Li T, Sweeney JA, Gong Q
(2015) Two Patterns of White Matter Abnormalities in Medication-Naive Patients with First-Episode
Schizophrenia Revealed by Diffusion Tensor Imaging and Cluster Analysis. JAMA Psychiatry
72:678–686.

The Brainstorm Consortium et al. (2017). Analysis of shared heritability in common disorders of the
brain. Science, 360 (6395):eaap8757.

van Hulst B, de Zeeuw P, Durston S (2014) Distinct neuropsychological profiles within ADHD: a
latent class analysis of cognitive control, reward sensitivity and timing. Psychol Med 45:735–745.

van Hulzen K, Scholz CJ, Franke B, Ripke S, Klein M, McQuillin A, Sonuga-Barke EJ, Group P,
Kelsoe JR, Landén M, Andreassen OA, Group P, Lesch K-P, Weber H, Faraone SV, Arias-Vasquez
A, Reif A (2017) Genetic Overlap Between Attention-Deficit/Hyperactivity Disorder and Bipolar
Disorder: Evidence from Genome-wide Association Study Meta-analysis. Biol Psychiat 82:634–641.

van Loo HM et al. (2014) Major depressive disorder subtypes to predict long-term course. Depress
Anxiety 31:765–777.

van Loo HM, de Jonge P, Romeijn J-W, Kessler RC, Schoevers RA (2012) Data-driven subtypes of
major depressive disorder: a systematic review. Bmc Med 10:1–12.

Veatch O, Veenstra□VanderWeele J, Potter M, Pericak□Vance M, Haines J (2014) Genetically
meaningful phenotypic subgroups in autism spectrum disorders. Genes Brain Behav 13:276–285.

Wang H, Jung Y-E, Chung S-K, Hong J, Kang N, Kim M-D, Bahk W-M (2017) Prevalence and
correlates of bipolar spectrum disorder comorbid with ADHD features in nonclinical young adults. J
Affect Disorders 207:175–180.

Wei Y, Chang M, Womer FY, Zhou Q, Yin Z, Wei S, Zhou Y, Jiang X, Yao X, Duan J, Xu K, Zuo
X-N, Tang Y, Wang F (2018) Local functional connectivity alterations in schizophrenia, bipolar
disorder, and major depressive disorder. J Affect Disorders 236:266–273.

Xia C.H. et al. (2018a) Linked dimensions of psychopathology and connectivity in functional brain
networks. Nat Commun 9(1):3003.

Xia M, Womer FY, Chang M, Zhu Y, Zhou Q, Edmiston EK, Jiang X, Wei S, Duan J, Xu K, Tang
Y, He Y, Wang F (2018b) Shared and Distinct Functional Architectures of Brain Networks Across
Psychiatric Disorders. Schizophr Bull. doi: 10.1093/schbul/sby046

Zou H, Hastie T (2005) Regularization and variable selection via the elastic net. J Royal Statistical
Soc Ser B Statistical Methodol 67:301–320.

Table 1. Demographic Information*

	HC	SCZ	BD	ADHD	Total
No. of subjects	130	50	49	43	272
With complete phenotype data	130	50	48	43	271
With sMRI data**	98	30	44	34	206
With fMRI data[†]	104	47	41	37	229
Age					
Mean age	31.26	36.46	35.15	33.09	
SD age	8.74	8.88	9.07	10.76	
Range age	21-50	22-49	21-50	21-50	
Gender					
No. of female subjects	62	12	21	22	
Percent female subjects	47.69%	24.00%	42.86%	51.16%	
Race					
American Indian or Alaskan Native	19.23%	22.00%	6.25%	0%	
Asian	1.54%	2.00%	0%	2.33%	
Black/African American	0.77%	4.00%	2.08%	2.33%	
White	78.46%	66.00%	77.08%	88.37%	
More than one race	0%	2.00%	14.58%	6.98%	
Education					
No high school	1.54%	18.00%	2.08%	0%	
High school	12.31%	44.00%	29.17%	23.26%	
Some college	20.77%	18.00%	25.00%	30.23%	
Associate's degree	7.69%	4.00%	6.25%	6.98%	
Bachelor's degree	50.00%	10.00%	29.17%	32.56%	
Graduate degree	6.92%	0%	4.17%	2.33%	
Other	0.77%	4.00%	4.17%	4.65%	

* Demographic information is based on initial number of subjects
** Excluding subjects with aliasing artifacts
[†] Excluding subjects with misaligned structural-function imaging data

Table 2. Performances of the transdiagnostic models on each feature set*

Performance of the full model:

	Scales	sMRI	fMRI	s+fMRI	Scales+sMRI	Scales+fMRI	Scales+s+fMRI
AUC	0.83(0.04)	0.56(0.05)	0.59(0.04)	0.57(0.05)	0.89(0.07)	0.86(0.06)	0.86(0.05)
Accuracy	0.77(0.05)	0.58(0.08)	0.60(0.06)	0.61(0.07)	0.91(0.04)	0.87(0.05)	0.86(0.05)
Sensitivity	0.77(0.07)	0.74(0.11)	0.60(0.11)	0.62(0.11)	0.86(0.08)	0.82(0.12)	0.86(0.08)
Specificity	0.82(0.07)	0.38(0.11)	0.56(0.12)	0.49(0.17)	0.80(0.15)	0.61(0.30)	0.48(0.26)

Performance of the best truncated model:

	Scales	sMRI	fMRI	s+fMRI	Scales+sMRI	Scales+fMRI	Scales+s+fMRI
AUC	0.95(0.02)	0.78(0.06)	0.87(0.08)	0.77(0.06)	0.96(0.03)	0.98(0.02)	0.96(0.03)
Accuracy	0.88(0.04)	0.71(0.06)	0.85(0.07)	0.77(0.06)	0.87(0.05)	0.92(0.04)	0.90(0.04)
Sensitivity	0.87(0.08)	0.81(0.09)	0.86(0.09)	0.77(0.08)	0.93(0.07)	0.91(0.06)	0.94(0.05)
Specificity	0.88(0.04)	0.60(0.16)	0.84(0.18)	0.76(0.07)	0.80(0.15)	0.92(0.04)	0.85(0.09)
No. of features	85	131	8192	16384	238	32	64

Test on AUCs between the full and the truncated model**:

	Scales	sMRI	fMRI	s+fMRI	Scales+sMRI	Scales+fMRI	Scales+s+fMRI
Test statistic	100	100	100	99.5	82.5	100	95
p-value[†]	< 0.001	< 0.001	< 0.001	< 0.001	0.01537	< 0.001	< 0.001

* The mean performance measures across 10 implementations are reported here with the standard deviation shown in parentheses
** Wilcoxon's rank-sum test
† FDR-corrected

g

Figure Legends

Figure 1. The performances of the best transdiagnostic models selected via the feature importance-guided sequential model selection procedure. **A)** The receiver operating characteristic (ROC) curve for the best truncated models based on each feature set. Area under the ROC curve (AUC) for each model is listed in the legend. **B)** Box plots showing the AUC of the best truncated model for each feature set measured across 10 implementations of sequential model selection procedure.

Figure 2. Distributions and effect sizes of the model's derived scores vs. the existing scale scores. **A)** Sum score calculated from the identified 85 most predictive items showing high separability in terms of Cohen's d between HC and Patients. **B)** All three patient categories showed elevated sum scores relative to HC (p < 0.001). **C)** The 4 temperament sub-scores in TCI included in the CNP dataset showing only medium effect sizes between HC and Patients. **D)** Box plot showing significantly higher effect size from the identified 85 items (asterisk) compared to all predefined sum and subscores in self-reported instruments in CNP data. The asterisk represents the Cohen's d between HC and Patients from the top 85 items, whereas the box plot shows the effect sizes from all predefined sum and subscores (also see Supplementary Table 2).

Figure 3. The percentage of items from each of the 13 self-reported instruments among the set of 85 most predictive transdiagnostic items.

Figure 4. The grouping of items into specific phenotypic domains for the top 20 most predictive items from **A)** the HC vs. All Patients transdiagnostic model and **B) – D)** the 3 HC vs. a single patient category classifiers. The radius in the spider plots represents item counts.