Article

Reporter assays for Ebola virus nucleoprotein oligomerization, virion-like particle budding, and minigenome activity reveal the importance of nucleoprotein amino acid position 111

Aaron E. Lin [1,2,3,12,*], **William E. Diehl** [4,12], **Yingyun Cai** [5], **Courtney L. Finch** [5], **Chidiebere Akusobi** [6], **Robert N. Kirchdoerfer** [7], **Laura Bollinger** [5], **Stephen F. Schaffner** [2,3], **Elizabeth A. Brown** [2,3], **Erica Ollmann Saphire** [8], **Kristian G. Andersen** [9,10], **Jens H. Kuhn** [5,13], **Jeremy Luban** [4,13], **Pardis C. Sabeti** [1,2,3,11,13,*]

[1] Harvard Program in Virology, Harvard Medical School, Boston, MA 02115, USA
[2] FAS Center for Systems Biology, Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, MA 02138, USA
[3] Broad Institute of MIT and Harvard, Cambridge, MA 02142, USA
[4] Program in Molecular Medicine, University of Massachusetts Medical School, Worcester, MA 01605, USA
[5] Integrated Research Facility at Fort Detrick, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Frederick, MD 21702, USA
[6] Department of Immunology and Infectious Diseases, Harvard T.H. Chan School of Public Health, Boston, MA 02120, USA
[7] Department of Biochemistry, University of Wisconsin-Madison, Madison, WI 53706, USA
[8] La Jolla Institute for Immunology, La Jolla, CA 92037, USA
[9] Department of Immunology and Microbial Sciences, The Scripps Research Institute, La Jolla, CA 92037, USA
[10] Scripps Translational Science Institute, La Jolla, CA 92037, USA
[11] Howard Hughes Medical Institute, Chevy Chase, MD 20815, USA
[12] Co-first author
[13] Co-last author

* Correspondence: alin@broadinstitute.org (A.E.L.), pardis@broadinstitute.org (P.C.S.)

Abstract: For highly pathogenic viruses, reporter assays that can be rapidly performed are critically needed to identify potentially functional mutations for further study under maximal containment (e.g., biosafety level 4 [BSL-4]). The Ebola virus nucleoprotein (NP) plays multiple essential roles during the viral life cycle, yet few tools exist to study the protein under BSL-2 or equivalent containment. Therefore, we adapted reporter assays to measure NP oligomerization and virion-like particle (VLP) production in live cells and further measure transcription and replication using established minigenome assays. As a proof-of-concept, we examined the NP-R111C substitution, which emerged during the 2013–2016 Western African Ebola virus disease epidemic and rose to high frequency. NP-R111C slightly increased NP oligomerization and VLP budding but slightly decreased transcription and replication. By contrast, a synthetic charge-reversal mutant, NP-R111E, greatly increased oligomerization but abrogated transcription and replication. These results are intriguing in light of recent structures of NP oligomers, which reveal that the neighboring residue, K110, forms a salt bridge with E349 on adjacent NP molecules. By developing and utilizing multiple reporter assays, we find that the NP-111 position mediates a complex interplay between NP's roles in protein structure, virion budding, and transcription and replication.

Keywords: Ebola virus; nucleoprotein; budding; oligomerization; reporter assays; viral evolution

1. Introduction

Ebola virus (EBOV) remains a serious hazard to human heath; however, studying live virus requires maximum (biosafety level 4 [BSL-4]) containment, restricting experiments to a handful of laboratories around the globe. Therefore, reporter assays can complement live virus experiments in two ways. First, reporter assays can selectively study one or a handful of viral phenotypes for better understanding of specific molecular mechanisms. Second, many reporter assays can be performed at BSL-1 or -2 containment, opening up performance of EBOV studies in many more laboratories. Therefore, reporter assays can be a first-line tool to assess how rapidly evolving viral genotypes affect phenotype before time- and resource-intensive BSL-4 studies are conducted.

Given the past and current public health threats caused by Ebola virus disease (EVD) outbreaks, rapidly evaluating whether EBOV genomic mutations change viral phenotypes is critically important. As an RNA virus, EBOV generates many mutations over the course of an outbreak. The vast majority of these mutations likely will not be adaptive and will instead have negligible or negative effects on EBOV viability and replication [1]. Yet, changes in the EBOV genome over time can have important implications for clinical patient care, epidemiological modeling, and vaccine development, and thus influence prospective outbreak prediction and response.

The need to better understand EBOV evolution became clear during the 2013–2016 EVD epidemic in Western Africa caused by the EBOV Makona variant. This epidemic was the largest EVD epidemic on record with over 28,000 infections and more than 11,000 deaths [2]. EBOV replication generated thousands of mutations over numerous rounds of human-to-human transmission [3–14], but only a handful of mutations became common enough to have had a sizeable impact on the epidemic [3–9]. One key mutation, C6283U, results in an A82V substitution in the EBOV glycoprotein (GP-A82V) and has been studied extensively through well-established BSL-2 surrogate model systems and live virus BSL-4 studies. GP-A82V increases EBOV infectivity for human and other primate cell types *in vitro*, suggesting that the mutation confers a selective advantage to EBOV [15–20]; however increased EBOV infectivity has not yet been demonstrated clearly *in vivo* [21,22].

A second high-frequency mutation that emerged amidst the epidemic, C800U, results in an R111C substitution in the EBOV nucleoprotein (NP-R111C). Both the GP-A82V and NP-R111C substitutions arose on the same viral lineage and rose to >90% frequency. In contrast to GP-A82V, NP-R111C has not been thoroughly studied. Though NP has many functions, a limited number of assays are available for studying these functions. This limitation presents an important challenge and opportunity to develop assays to gain mechanistic insight into NP biology and variation.

The N-terminal domain of EBOV NP, which contains the R111 residue, is necessary and sufficient to drive oligomerization of NP monomers into long, flexible helices. These oligomers help shape virions and coat viral RNA during the viral life cycle. Many studies have determined NP structures and modeled how NP interacts with itself and RNA [23–29]. In addition to a flexible oligomerization domain (OD) at the very N terminus of NP that is essential for oligomerization, numerous other residues appear to be involved in NP-NP interactions as visualized in cryo-electron microscopy (cryo-EM) of helical NP oligomers [26–30]. Between vertically adjacent NP proteins, NP-K110 on one monomer appears to form a salt bridge with NP-E349 on another monomer [28,29]. No naturally occurring mutations have been shown to affect EBOV NP oligomerization, though the proximity of NP-R111 to the NP-K110 residue suggests that the NP-R111C substitution could affect oligomerization. Although many biochemical methods are available for assessing NP oligomerization *in vitro*, few assays exist to quantitatively measure oligomerization in live cells.

The ability of NP to oligomerize likely affects its role in virion assembly and budding. During the viral life cycle, the EBOV matrix protein VP40 induces curvature of the cell's plasma membrane and engages with EBOV nucleocapsids (composed of NP oligomers, polymerase cofactor VP35, and minor capsid VP24) to form EBOV particles that exit the cell by budding [31]. Expression of VP40 alone generates similarly shaped particles, dubbed virion-like particles (VLPs) [32–34]. Co-expression of NP with VP40 significantly increases the number of VLPs released from cells [35],

but the mechanism by which NP increases VLP production is not fully elucidated. For example, NP
oligomers could physically stabilize VLP structure, and/or NP could engage with host proteins to
promote VLP assembly and budding.
EBOV NP oligomers also play an essential role in viral transcription and replication. By
directly interacting with EBOV RNA, VP35, and transcription regulator VP30, NP recruits the
RNA-directed RNA polymerase L to enact both of these essential functions [36]. NP structural data,
homology modeling to other viral nucleoproteins, and site-directed mutagenesis experiments have
identified key EBOV NP residues that interact with EBOV RNA [23–25], VP35 [24,25], and VP30
[37]. The NP residue 111 lies outside of any of these annotated interaction surfaces, suggesting that
NP-R111C most likely does not directly affect binding affinity to RNA. However, only NP
oligomers, and not NP monomers, bind RNA [24,25,27], so mutations that affect NP
oligomerization propensity could still impact RNA binding avidity and thereby influence
transcription and replication.
For this study, we adapted and modified BSL-2 methods to study key functions of EBOV NP in
cell culture – oligomerization and VLP budding – and used established minigenome systems that
measure viral transcription and replication. Such tools are critical for rapidly characterizing
unknown or emerging mutants since studying live EBOV requires limited BSL-4 facilities.
Moreover, existing recombinant live virus systems typically use a genetic backbone that is different
from the EBOV Makona variant isolate C-15 sequence [38–41], and generating new recombinant
systems remains logistically and financially challenging due to restrictions on their use and
associated synthesis costs. Using these straightforward, modular reporter assays, we found that the
NP-R111C substitution slightly increased EBOV NP oligomerization and VLP budding and slightly
decreased transcription and replication. Charge-reversal at this position, NP-R111E, significantly
altered NP oligomerization, supporting the K110-E349 salt bridge in cryo-EM structures of NP
oligomers. Unexpectedly, NP-R111E dramatically ablated viral transcription and replication. Our
findings demonstrate the interconnectedness of multiple viral phenotypes controlled by the
NP-R111 residue, and support the possibility that NP-R111C affects replication of live virus.

2. Materials and Methods

Key Resources Table

We attached a Key Resources Table, including oligonucleotide sequences, as File S1.

2.1. Ebola virus Genome Sequences and Phylogenetic Analysis

We obtained Ebola virus (EBOV) genomes from the US National Institute of Allergy and
Infectious Diseases (NIAID) Virus Pathogen Database and Analysis Resource (ViPR) through the web
site at http://www.viprbrc.org/ [42] on October 2017. We removed short sequences, sequences from
tissue-cultured EBOV isolates, duplicate sequences from the same clinical EVD case, and sequences
with >0.2% ambiguous or missing nucleotide calls. The final dataset consisted of 1,823 EBOV complete
or near-complete genomes.
We aligned these genomes with MAFFT v6.902b [43] using the parameters (L-INS-i): --localpair
--maxiterate 1000 --reorder --ep 0.123. We trimmed the alignment using trimAl v1.4 [44] with
-automated1. Lastly, we generated a maximum likelihood tree with RAxML v7.3.0 [45] under a
generalized time-reversible (GTRγ) nucleotide substitution model with 100 bootstrap
pseudoreplicates (Figure 1).
For cloning and functional characterization, we used the genome sequence of Ebola
virus/H.sapiens-wt/GIN/2014/Makona-C15 (EBOV/Mak-C15; GenBank #KJ660346.2; *Filoviridae: Zaire
ebolavirus*) as the EBOV Makona variant reference sequence for NP, VP40, VP35, and GP, unless
otherwise noted. The structural analysis of EBOV NP was based on the Ebola
virus/H.sapiens-tc/COD/1976/Yambuku-Mayinga NP (EBOV/Yam-May) crystal structure under
Protein Data Bank (PDB) #4YPI [24], with manual annotation of key residues based on other studies
[24,25]. The structure of EBOV/Mak-C15 NP has not yet been elucidated, but the amino acid sequence

is 98% identical (14 mutations / 739 residues) to EBOV/Yam-May NP, and the N-terminal 450 amino
acids of the two isolates are 99.3% identical (3 mutations / 450 residues). The EBOV/Yam-May crystal
structure #4YPI was visualized using PyMOL (Schrödinger, New York City, NY) (Figure 2) [46].

2.2. Constructs and Cloning

We performed most assays with the same mammalian expression vector for EBOV/Mak-C15 NP
and its mutants, except where indicated. We synthesized EBOV NP-R111 in 2 dsDNA gBlocks
(Integrated DNA Technologies [IDT], Coralville, IA) and cloned these gBlocks into pGL4.23-CMV [15]
modified with a C-terminal V5 peptide tag. To generate all NP mutants, we performed a modified
site-directed mutagenesis (SDM) protocol, as described in [15]. For many assays, we expressed
enhanced green fluorescent protein (eGFP) in cells in place of NP as a negative control. We generated
the corresponding vector by cloning eGFP into pcDNA3.3-CMV (Thermo Fisher Scientific, Waltham,
MA) modified by an in-frame C-terminal V5 peptide tag.

For co-immunoprecipitation (co-IP) and oligomerization studies, we generated NP with different
C-terminal tags in the pGL4.23-CMV backbone. For the traditional dual-tag co-IP-western blot (WB)
strategy in Figure S1, we generated myc-tagged NP (pGL4.23-CMV/NP-myc). For the
bioluminescence resonance energy transfer (BRET) oligomerization assay (Figure 3), we replaced the
C-terminal V5 tag with either HaloTag or NanoLuc (NLuc) from the NanoBRET Nano-Glo Detection
System (Promega, Madison, WI). As a negative control, we removed NP amino acid residues 20–38,
abrogating the oligomerization domain (NP-ΔOD) [25], by SDM.

For the BRET experiment to study the NP-VP35 interaction (Figure 3C), we modified the
pcDNA3.3 backbone with a woodchuck hepatitis virus post-transcriptional regulatory element
(WPRE) to increase insert expression, using pcDNA3.3/KLF4 (a gift from Derrick Rossi; Addgene,
Cambridge, MA; plasmid #26815) [47] and pLV-WPRE/mCherry (a gift from Pantelis Tsoulfas;
Addgene; plasmid #36084) as source material. We then cloned in eGFP, porcine teschovirus 1 2A
'self-cleaving' peptide (P2A) [48], and EBOV/Mak-C15 VP35 amino acids 1–80 containing the
NP-binding peptide (NPBP) [25] from a gBlock (IDT) into a single open reading frame
(eGFP-P2A-VP35[NPBP]) upstream of the WPRE, resulting in
pcDNA3.3-WPRE/eGFP-P2A-VP35[NPBP]. As a negative control, we cloned a V5-tagged blue
fluorescent protein mTagBFP2 into the pcDNA3.3-WPRE backbone, using mTagBFP2-pBAD (a gift
from Michael Davidson; Addgene; plasmid #54572) [49] as source material.

For the VLP budding assay (Figures 4 and S3), we constructed a plasmid to express NLuc fused
to EBOV/Mak-C15 VP40 (NLuc-VP40). To create this plasmid, we obtained a pcDNA3.1(+)-based
vector expressing β-lactamase (Bla) fused to EBOV/Yam-May (Bla-VP40) from the US National
Institutes of Health (NIH)/NIAID Biodefense and Emerging Infections Research Resources Repository
(BEI Resources, Manassas, VA; #NR-19813) [50]. We replaced the Bla sequence with the gene encoding
NLuc from pNL1.1 (Promega) and replaced the EBOV/Yam-May VP40 sequence with that of
EBOV/Mak-C15 from a gBlock (IDT). pNL1.1, which expresses NLuc alone without VP40, was used as
a negative control. As an additional negative control, we performed SDM to introduce a L117R
substitution into NLuc-VP40 to generate loss-of-function (LOF) [51]. For electron microscopy (Figure
S4), we additionally co-expressed EBOV glycoprotein (GP) from a pGL4.23-CMV vector [15].

For monocistronic (1MG) minigenome experiments (Figure 5B), plasmids are described in [52]. In
this system, EBOV RNA-directed RNA polymerase (L), viral cofactor proteins (VP30 and VP35), and
NP were derived from the EBOV/Yam-May isolate and expressed from a pCAGGS vector. We
replaced EBOV/Yam-May NP with EBOV/Mak-C15 NP and its mutants before measuring
minigenome activity encoded by a firefly luciferase (FLuc) reporter gene.

For tetracistronic (4MG) minigenome experiments (Figures 5C and 5D), we additionally cloned L
and VP35 from EBOV/Mak-C15 into a pCAGGS vector. Since no amino acid differences are present
between EBOV/Yam-May and EBOV/Mak-C15 VP30, we were able to express EBOV/Mak-C15
sequences of all ribonucleoprotein (RNP) complex members (L, VP35, VP30, NP). We additionally
cloned the 4MG minigenome plasmid from EBOV/Mak-C15, expressing a *Renilla* luciferase (RLuc)
reporter gene, VP40, GP, and VP24.

2.3. Cell Culture and Plasmid Transfections

Unless otherwise specified, we grew human embryonic kidney (HEK) 293FT cells (Thermo
Fisher Scientific; #R70007) in Dulbecco's modified Eagle medium (DMEM) containing 10% fetal
bovine serum (FBS), 100 U/mL penicillin/streptomycin, non-essential amino acids, and sodium
pyruvate (Thermo Fisher Scientific), at 37 °C with 5% CO_2.
For most assays, we performed lipid-based reverse transfection using Lipofectamine 2000
(Thermo Fisher Scientific). For a 6-well plate, we incubated 2 µg of plasmid DNA with 125 µL of
Opti-MEM (Thermo Fisher Scientific) at room temperature for 5 minutes (min). We incubated this
mixture with 10 µL of Lipofectamine 2000 in 115 µL of Opti-MEM at room temperature for 45 min.
We added all 250 µL of the DNA:lipid mixture to a well of a 6-well plate and then added
trypsin-harvested cells. For smaller or larger plates, amounts were scaled accordingly. For BRET
experiments, we used Opti-MEM without phenol red (Thermo Fisher Scientific) to minimize
background fluorescence from culture media.
For the 1MG and 4MG minigenome assays and EM, we performed forward transfection by
incubating DNA with *Trans*IT-LT1 Transfection Reagent (Mirus Bio, Madison, WI) in a 1:3
DNA:reagent ratio in Opti-MEM for 15–20 min at room temperature, and then added the mixture
dropwise onto cells in 6- or 12-well plates.

2.4. Co-immunoprecipitation and western blot

We washed cells in 6-well plates with phosphate-buffered saline (PBS), harvested by scraping,
pelleted and resuspended cells in 30 µL of 1.2% (w/v) polyvinylpyrrolidine (PVP) in 20 mM K-HEPES
buffer pH 7.4, and snap-froze with liquid nitrogen. We lysed cells with 250–500 µL of pre-chilled lysis
buffer with end-over-end rotation at 4 °C for 30 min, cleared lysate of membranous debris by
centrifugation at 8000 x *g* at 4 °C for 10 min, and saved an aliquot as input. To capture the target
protein, we prepared a mixture of 25 µL each of Protein A and Protein G SureBeads Magnetic Beads
(Bio-Rad, Hercules, CA) and immobilized 1–2 µg of primary antibody on the beads by rotation at
room temperature for 20 min. We washed the bead-antibody complexes thrice with lysis buffer, and
then incubated with cleared cell lysate while rotating the mixture at 4 °C for 2 hours (h). After capture,
we washed beads six times with wash buffer followed by a final wash with PBS, and eluted proteins
by boiling the beads in 50 µL of Laemmli sample buffer (Bio-Rad) at 95 °C for 10 min.
We loaded the specified amount of input into denaturing 10% polyacrylamide gels, and
performed electrophoresis at 180 V until complete. We transferred proteins to Immun-Blot PVDF
Membranes (Bio-Rad) in a wet tank either at 200 mA for 1.5 h at 4 °C, or at 40 V overnight at 4 °C. We
blocked membranes by rocking in blocking buffer consisting of 5% non-fat dry milk (Santa Cruz
Biotechnology, Dallas, TX) dissolved in tris-buffered saline with 0.1% Tween 20 (TBS-T) for 1 h at
room temperature. We incubated membranes with primary antibody in blocking buffer for 45 min,
washed three times in TBS-T buffer, incubated the membrane with horseradish peroxidase
-conjugated secondary antibody in blocking buffer for 1 h, and washed three additional times. We
activated chemiluminesence with SuperSignal West Pico Chemiluminescent Substrate (Thermo Fisher
Scientific) and imaged with an AlphaInnotech ChemiImager (ProteinSimple, San Jose, CA) or
FluorChem E (ProteinSimple).
For the dual-tag co-IP-WB for NP oligomerization (Figure S1), we used RIPA buffer (50 mM Tris
pH 6.8, 150 mM NaCl, 0.5% (w/v) sodium deoxycholate, 1% (w/v) Triton X-100 (Sigma-Aldrich, St.
Louis, MO)) both as the lysis and wash buffer because the NP-NP interaction was very strong [53].
For mass spectrometry (Figure S5A) and reciprocal co-IP experiments (Figures S5B and S5C), we
used mild lysis and wash buffers, slightly modified from a previous study [54]. Mild lysis buffer
consisted of 20 mM K-HEPES buffer pH 7.4, 100 mM NaOAc, 2 mM $MgCl_2$, 0.1% (v/v) Tween 20, 250
mM NaCl, 0.5% (v/v) Triton X-100, 4 µg/mL DNase I (QIAgen, Hilden, Germany), 2 µg/mL RNase A
(QIAgen), 1/200 (v/v) each phosphatase inhibitor cocktails 2 and 3 (Sigma-Aldrich), and 1/100 (v/v)
protease inhibitor mixture (Sigma-Aldrich). We incubated cleared cell lysate with 1–2 µg of primary
antibody, rotated the mixture at 4 °C for 2–4 h, and then added 40 µL of Protein A/G PLUS-Agarose
beads (Santa Cruz Biotechnology), and rocked at 4 °C overnight. Wash buffer consisted of 20 mM

K-HEPES pH 7.4, 100 mM NaOAc, 2 mM MgCl$_2$, 0.1% (v/v) Tween 20, 500 mM NaCl, and 0.5% (v/v)
Triton X-100. We washed bead-antibody complexes four times with wash buffer, twice with PBS, and
eluted proteins as described above.

2.5. BRET NP Oligomerization Assay

We grew cells to near confluency, harvested by trypsinization, reverse-transfected, and plated
cells in poly-D-lysine-coated, 96-well black/clear flat-bottom plates (Corning, Corning, NY). We
reverse-transfected cells in each well with 10 ng of pGL4.23-CMV/NP-NLuc or pNL1.1/NLuc
(Promega) negative control and 100 ng of pGL4.23-CMV/NP-HaloTag or pcDNA3.3/eGFP negative
control. At the start of transfection, we also added HaloTag NanoBRET 618 ligand (NanoBRET
Nano-Glo Detection System, Promega) diluted in dimethylsulfoxide (DMSO) at a final concentration
of 100 nM or DMSO vehicle control (no HaloTag ligand) to cell culture media.
At 24 h post-transfection, we added 1:100 NanoBRET Nano-Glo Substrate, incubated cells in the
dark at room temperature for 45 min, and measured luminescence on a DTX880 Multimode Detector
(Beckman Coulter, Brea, CA) with emission filters of 625/35 nm (HaloTag ligand acceptor signal), and
then 465/35 nm (NLuc donor signal), both over 1-second (s) integrations. We calculated BRET signal
as the 625 nm / 465 nm ratio with HaloTag ligand subtracted by the same ratio for the corresponding
DMSO vehicle control, per manufacturer's protocol.
For the VP35 inhibition experiment (Figure 3C), we reverse-transfected cells in each well with 2
ng of pGL4.23-CMV/NP-NLuc and 10 ng of pGL4.23-CMV/NP-HaloTag. To test a range of
VP35[NPBP]   expression,   we   co-transfected   cells   with   decreasing   amounts   of
pcDNA3.3-WPRE/eGFP-P2A-VP35[NPBP] plasmid. To ensure that cells in each well received the
same total amount of DNA, we serially diluted pcDNA3.3-WPRE/eGFP-P2A-VP35(NPBP) in control
plasmid pcDNA3.3-WPRE/mTagBFP2, as described in the manufacturer's protocol. We performed the
remainder of the standard BRET protocol as described above.
For the donor saturation procedure (Figure 3D), we reverse-transfected cells in each well with 2
ng of pGL4.23-CMV/NP-NLuc. To test a range of NP-HaloTag expression, we co-transfected cells
with decreasing amounts (80, 20, 5, 0 ng) of pGL4.23-CMV/NP-HaloTag or pcDNA3.3/eGFP negative
control. To ensure that cells in each well received the same total amount of DNA, we serially diluted
pGL4.23-CMV/NP-HaloTag or pcDNA3.3/eGFP in control pcDNA3.3/eGFP plasmid, as described in
the manufacturer's protocol.

2.6. Virion-like Particle Budding Assay

We grew cells to near confluency, harvested by trypsinization, reverse-transfected, and plated
cells in 6-well poly-D-lysine-coated plates (Corning). We reverse-transfected cells in each well with 50
ng of pcDNA3.1/NLuc-VP40 or pNL1.1/NLuc negative control and 2000 ng of pGL4.23-CMV/NP-V5,
NP mutants, or pcDNA3.3/eGFP negative control.
At 16 h post-transfection, we removed supernatant, washed the cells with DMEM, and added 1.5
mL of fresh DMEM. At 40 h post-transfection (24 h later), we filtered culture supernatant through an
Acrodisc 0.45 µm low protein-binding filter (Pall Laboratory, Port Washington, NY). We underlaid 1
mL of filtered supernatant with 1 mL of 20% (w/v) sucrose (Sigma-Aldrich) in PBS and
ultracentrifuged at 222,000 x g at 4 °C for 2 h. We aspirated the supernatant, resuspended the
VLP-containing pellet in 170 µL of PBS, and rocked at room temperature for 1 h. We aliquoted
resuspended VLPs into 3 x 50 µL as technical triplicates, added 50 µL of Nano-Glo assay reagent
(Promega) to each, and incubated in 96-well non-binding-surface plates (Corning) in the dark at room
temperature for 10 min. We measured total luminescence on a SpectraMax L (Molecular Devices,
Sunnyvale, CA) over a 1-s integration. Technical triplicates were averaged and considered as a single
biological replicate.
For the thermal stability assay, we reverse transfected cells in each well with 50 ng of
pcDNA3.1/NLuc-VP40 per well as above. After filtration, we heated 1.2 mL of filtered supernatant at
4, 22, 37.1, 43.8, 60.2, or 95 °C, for 30 min on a Mastercycler pro S thermocycler (Eppendorf, Hamburg,
Germany). We saved 50 µL of heated supernatant for direct NLuc measurement. Subsequently, we

performed the remainder of the protocol described above (ultracentrifugation of 1 mL of heated
supernatant through sucrose to purify VLPs, and subsequent measurement of NLuc activity).

2.7. Electron Microscopy

We seeded 6 x 10^5 HEK 293 cells per well in 6-well plates. The following day, we transfected cells
in each well with 1250 ng of pcDNA3.1(+)-VP40 (untagged), 930 ng of pGL4.23-CMV/NP or
pGL4.23-CMV/NP-R111C (both untagged), and 310 ng of pGL4.23-CMV/GP-A82V [15] using 6.25 μL
of *Trans*IT-LT1 Transfection Reagent (Mirus Bio). We changed media the next morning. After 48 h, we
filtered culture supernatant through a 0.45 μm filter and overlaid it onto a 20% (w/v) sucrose in TNE
(10 mM Tris-Cl, 100 mM NaCl, 1 mM EDTA pH 7.5) cushion. VLPs were pelleted by
ultracentrifugation at 222,000 x g at 4 °C for 2 h. We aspirated the supernatant, washed the
VLP-containing pellet gently with 1 mL of ice-cold PBS, resuspended VLPs in 100 μL of 2% FBS in
PBS, and stored VLPs at 4 °C prior to EM.
We prepared samples for EM based on our previously described protocol [55]. Briefly, we
performed all spreads onto freshly prepared Carbon stabilized Formvar Support films on 200 mesh
copper grids. We adsorbed VLPs onto a carbon-coated Formvar support films for 30 s. We removed
excess liquid with filter paper, and negatively stained the samples immediately by running 6 drops of
1% uranyl acetate over the grid for contrast. We removed excess stain and air-dried the samples in a
controlled humidity chamber. We then examined the samples using a FEI Tecnai 12 Spirit BioTwin
transmission electron microscope (Thermo Fisher Scientific) using an accelerating voltage of 120 kV.
We captured micrographs at various magnifications to record the fine structure of VLPs and exported
micrographs into ImageJ [56] to measure the length and volume of individual particles.

2.8. Monocistronic Minigenome Assay

Monocistronic (1MG) minigenome plasmids and procedure were previously described [52]. We
seeded HEK 293T cells into 12-well plates, grew to 70% confluency, and transfected cells in each well
with 2 μg of pCAGGS/L, 0.25 μg of pCAGGS/VP30, 0.5 μg of pCAGGS/T7 RNA polymerase (T7pol),
0.5 μg of 1MG plasmid encoding FLuc, 0.1 μg of pCAGGS encoding RLuc, and 0.75 μg of
pCAGGS/NP-2A-VP35 for each NP mutant. After 2 days, we washed and lysed cells with 100 μL of
1X Passive Lysis Buffer (Dual Luciferase Assay Kit, Promega), freeze-thawed lysates, and cleared by
centrifugation. We incubated 10 μL of lysate with 50 μL of Luciferase Assay Reagent II, let the mixture
settle for 2 s, and integrated luminescence for 10 s on a Spark 10M microplate reader (Tecan, Zürich,
Switzerland) to measure FLuc activity. We then added 50 μL of Stop & Glo reagent and integrated
luminescence for 10 s to measure RLuc activity.

2.9. Tetracistronic Minigenome Assay

We followed an existing protocol for the tetracistronic (4MG) minigenome assay [57], with some
modifications. We first generated transcription- and replication-competent VLPs (trVLPs) using HEK
293T as producer (P0) cells. We seeded HEK 293T cells into collagen-coated 6-well plates, grew to 40%
confluency, and transfected cells in each well with the previously described plasmid ratio (125 ng of
pCAGGS/NP, 125 ng of pCAGGS/VP35, 75 ng of pCAGGS/VP30, 1000 ng of pCAGGS/L, 250 ng of
4MG plasmid encoding RLuc, 250 of ng pCAGGS/T7pol) [57,58], plus 250 ng of an FLuc-encoding
plasmid to normalize for transfection efficiency. We changed media 24 h post-transfection, collected
and clarified trVLP-containing P0 supernatant by centrifugation, and measured intracellular
luminescence in P0 cells 96 h post-transfection.
We used trVLP-containing supernatant to infect target Huh7 cells for a total of two trVLP
passages (P1 and P2 cells). We seeded Huh7 cells into collagen-coated 6-well plates as P1 cells, grew
to 40% confluency, and transfected cells in each well with the plasmid ratio above except without the
4MG plasmid or the pCAGGS/T7pol [57,58]. We inoculated P1 cells with 3 mL of trVLP-containing P0
supernatant 24 h post-transfection, changed media 24 h post-infection, and measured luminescence in
P1 cells and collected trVLP-containing P1 supernatant 96 h post-infection. We performed the

procedure again by transfecting fresh Huh7 cells, inoculating these P2 cells with P1 trVLPs, and
measuring luminescence.

To measure intracellular luminescence in P0, P1, and P2 cells, we lysed cells with 500 μL 1X
Passive Lysis Buffer (Dual Luciferase Assay Kit, Promega) at room temperature for 15 min,
freeze-thawed lysates, and detected FLuc and RLuc luminescence with the Dual Luciferase Assay Kit
as described above.

2.10. Tandem Mass Spectrometry

To assess protein-protein interactions of NP (Figure S5), we scaled up our co-IP protocol. We
grew two 15-cm^2 plates of cells to 40–60% confluency and transfected cells with 32 μg of
pGL4.23-CMV/NP-myc encoding either NP-R111 or NP-K109E/K110E/R111E. After 48 h, we
harvested cells by scraping in 1.2% (w/v) polyvinylpyrrolidine (PVP) in 20 mM K-HEPES buffer pH
7.4, snap-froze with liquid nitrogen, and then lysed in 2.5 mL of mild lysis buffer, described above.

We performed co-IP of myc-tagged NP complexes using 25 μg of mouse α-myc IgG or irrelevant
normal mouse IgG at 4 °C overnight, and bound complexes to 250 μL of Protein A/G PLUS-Agarose
beads at 4 °C for 2 h. We washed beads as described above and eluted proteins in 120 μL of Laemmli
sample buffer at 95 °C for 10 min. We separated proteins by SDS-PAGE, visualized with PageBlue
Protein Staining Solution (Thermo Fisher Scientific), and excised lanes excluding IgG chains.

We cut gel bands into approximately 1-mm^3 pieces and performed a modified in-gel trypsin
digestion procedure [59]. We dehydrated pieces with acetonitrile for 10 min, dried them completely in
a speed-vac pump, and rehydrated with 50 mM ammonium bicarbonate solution containing 12.5
ng/μL of sequencing-grade modified trypsin (Promega) at 4 °C for 45 min. To extract peptides, we
replaced the solution with 50 mM trypsin-free ammonium bicarbonate solution and incubated at 37
°C overnight. We washed peptides once with 50% acetonitrile and 1% formic acid, dried in a
speed-vac pump for ≈1 h and then stored at 4 °C. On the day of analysis, we reconstituted peptides in
5–10 μL of high-performance liquid chromatography (HPLC) solvent A (2.5% acetonitrile, 0.1% formic
acid). We packed nano-scale reverse-phase HPLC capillary columns with 2.6 μm C18 spherical silica
beads into fused silica capillary tubes (100 μm inner diameter x ≈25 cm length) using flame-drawn
tips [60]. After equilibrating the columns, we loaded each sample via a Famos autosampler (LC
Packings, San Francisco, CA). We eluted peptides with increasing concentrations of solvent B (97.5%
acetonitrile, 0.1% formic acid).

To detect peptides, we performed tandem mass spectrometry (MS/MS) on an LTQ Orbitrap
Velos Pro ion-trap mass spectrometer (Thermo Fisher Scientific). We matched MS/MS fragmentation
spectra to human forward protein databases and against reverse databases to a 1–2% false discovery
rate using the SEQUEST database search program (Thermo Fisher Scientific) [61]. We computed
unique and total peptide spectra matches (PSMs) for each identified protein.

To generate a list of putative NP interacting partners, we filtered proteins with at least 2 unique
PSMs in co-IPs of both NP-R111 and NP-K109E/K110E/R111E, and at least 2-fold greater-than-average
PSM enrichment of α-myc co-IP over both IgG controls combined. To eliminate abundant and 'sticky'
proteins, we normalized average PSM enrichment against PSMs identified in all 411 Contaminant
Repository for Affinity Purification (CRAPome) version 1.1 experiments [62], a collection of proteins
identified in negative control isolations. From each replicate, we used the top 10% proteins enriched
versus CRAPome experiments for Search Tool for the Retrieval of Interacting Genes/proteins
(STRING) version 10 analysis [63,64] and visualized interactions with Cytoscape [65]. See
Supplementary File S2 for raw and filtered peptide/protein PSM counts.

2.11. Statistical Analysis

We performed all hypothesis testing using Prism 7 (GraphPad Software, La Jolla, CA) and all
non-linear curve fitting using R [66] and the 'nlstools' package [67]. We generated most plots using the
'ggplot2' package in R [68].

For the VP35 inhibition experiment using BRET (Figure 3C), we expressed varying amounts of
VP35[NPBP] in cells in the presence of NP-NLuc and NP-HaloTag for 4 amounts of

pcDNA3.3-WPRE/eGFP-P2A-VP35[NPBP] plasmid (n = 3 biological replicates each). We fitted the
data to an inverse function using the 'nls' function in R:

$$\text{BRET} \sim \text{scale} / (\text{VP35[NPBP]} + \text{max}) + \text{min}, \tag{1}$$

in which VP35[NPBP] expression was the independent variable, BRET was the dependent variable,
and scale, max, and min were constants to be fitted. Using non-linear regression, scale = 1.9×10^5, max
= 3.4×10^4, and min = 0.49. In the absence of VP35 (VP35[NPBP] = 0), the maximum BRET signal
would be 5.80; with very high expression (VP35[NPBP] $\rightarrow \infty$), the minimum BRET signal would be
0.49. We used the 'nlstools' package to generate 999 bootstrap pseudoreplicates, inferred parameters
for each pseudoreplicate, and plotted the central 95% of values as a shaded region.
For donor saturation assay using BRET (Figure 3D), we performed the BRET protocol with
NLuc- and HaloTag-tagged NP-R111 or NP mutants or eGFP control for 4 amounts of
pGL4.23-CMV/NP-HaloTag plasmid (n = 6 biological replicates each). We fitted the data to saturation
binding curves using the 'nls' function in R:

$$\text{BRET} \sim \text{Max} * \text{NP-HaloTag} / (K_d + \text{NP-HaloTag}), \tag{2}$$

in which the amount of NP-HaloTag plasmid was the independent variable, BRET was the dependent
variable, and Max and K_d were constants to be fitted. For NP-R111 or each NP mutant or eGFP
control, we estimated Max and K_d and generated 95% confidence intervals using 'nlstools' as
described above. Data points from eGFP failed to generate an appropriate curve fit. To determine
whether the remaining curve fits were significantly different from each other, we performed analysis
of variance (ANOVA) with Dunnett's post-test in which NP-R111C, NP-R111E, and
NP-K109E/K110E/R111E were compared to NP-R111, and generated multiple hypothesis corrected
p-values using Prism 7.
To measure VLP production from NLuc-VP40, NLuc-VP40-L117R, or NLuc alone (Figure 4B), we
quantified raw NLuc intensities (n = 6 biological replicates each). To assess statistical significance, we
performed a repeated measures ANOVA (rANOVA) with Dunnett's post-test, in which each
condition was compared to NLuc-VP40 to generate corrected p-values.
To measure the impact of NP genotype on VLP production (Figure 4C), we co-expressed
NLuc-VP40 and NP-R111 or NP mutants (R111C, R111E, K109E/K110E/R111E, ΔC50 – a 50 amino
acid truncation of the NP C-terminus [35]) or eGFP control. rANOVA revealed significant day-to-day
(replicate-to-replicate) variability, so we normalized NLuc intensities for each NP mutant to NP-R111
for each replicate. We performed Dunnett's post-test with the NP-R111 group removed (since variance
and degrees of freedom of NP-R111 are both 0 after normalization) and compared each NP mutant or
eGFP versus 1 to generate corrected p-values.
To determine whether heating disrupted VLPs (Figure S3B), we expressed NLuc-VP40 in cells,
and heated cell culture supernatant to 4, 22, 37.1, 43.8, 60.2, or 95 °C either before or after purifying
VLPs via ultracentrifugation (n = 3 biological replicates in a repeated-measures design). We
normalized NLuc values for all temperatures to the 4 °C value for each replicate, log-transformed the
normalized values, and fitted the data to sigmoidal curves using the 'nls' function in R:

$$\log_{10}(\text{NLuc.Norm}) \sim \text{min} + \text{max}/(1 + e^{(\text{midpt - temp})}/\text{scale}), \tag{3}$$

in which temperature was the independent variable, NLuc.Norm was the dependent variable, and
min, max, midpt, and scale were all constants to be fitted. Additionally, we tested whether NLuc
luminescence differed following heating the supernatant to 60.2 °C either before or after purifying
VLPs with a paired t-test.
To measure the impact of NP genotype on 1MG minigenome activity (Figure 5B), we expressed
NP-R111 or NP mutants with the rest of the EBOV replication complex (VP35, VP30, L), the
FLuc-encoding 1MG minigenome, and an RLuc-encoding plasmid to control for transfection
efficiency (n = 3 biological replicates each). We normalized FLuc by RLuc luminescence for each
replicate, then normalized all values to the average FLuc/RLuc ratio of the NP-R111 replicates. We
performed ANOVA with Dunnett's post-test, in which each condition was compared to NP-R111 to
generate corrected p-values.

To measure the impact of NP genotype on 4MG minigenome activity (Figure 5C), we expressed
NP-R111 or NP mutants with the rest of the EBOV replication complex, the RLuc-encoding 4MG
minigenome, and an FLuc-encoding plasmid to control for transfection efficiency (n = 3–9 biological
replicates each). Due to significant day-to-day (replicate-to-replicate) variability, we normalized RLuc
by FLuc luminescence for each replicate, then normalized all values to the average RLuc/FLuc ratio of
the GP-A82/NP-R111 replicates prepared on that day. We performed ANOVA with Tukey's post-test,
in which each condition was compared to each other condition to generate corrected p-values. To
measure the change in minigenome activity between passages (Figure 5D), we passaged cell culture
supernatant from P0 to P1, and then from P1 to P2 cells, keeping track of each replicate. We
normalized P1 and P2 RLuc/FLuc values to their own P0 replicate, and then normalized all values to
the average value of the GP-A82/NP-R111 replicates prepared on that day. We assessed statistical
significance using ANOVA with Tukey's post-test to generate corrected p-values.

3. Results

3.1. EBOV NP-R111C Emerged Alongside the GP-A82V Substitution during the 2013–2016 Western
African Epidemic

Among the viral mutations that rose to dominate the EBOV population during the 2013–2016
Western African EVD epidemic, the NP-R111C substitution is of great interest because it shares
features with the GP-A82V substitution that enhanced viral infectivity *in vitro* [15–20]. GP-A82V
and NP-R111C were two major clade-defining substitutions that rose to high frequency during the
epidemic; other mutations did not affect the amino acid sequence of EBOV proteins [4,8]. Based on
phylogeny of EBOV genomes from clinical samples, the NP-R111C substitution (Figure 1A, blue)
occurred soon after the emergence of GP-A82V (Figure 1A, green) and temporally preceded the
inflection point of the epidemic (Figure 1B). Indeed, few EBOV Makona variant genomes encoded
GP-A82V in the absence of NP-R111C (23 cases, 1.26% of total), and the overwhelming majority of
genomes encoded both substitutions (1653 cases, 90.67% of total).

Figure 1. EBOV NP-R111C emerged alongside the GP-A82V substitution. (**a**) Phylogenetic analysis
of the 2013–2016 EVD epidemic. We constructed a maximum likelihood tree based on 1,823 EBOV
genome sequences, and colored branches based on GP-82 and NP-111 alleles. No GP-A82/NP-R111C
sequences were detected. Arrowheads point to the emergence of the GP-A82V (green) and
NP-R111C (blue) substitutions compared to genomes encoding the ancestral GP-A82/NP-R111
alleles (tan). Scale bar denotes substitutions/nucleotide; (**b**) Number of EVD cases over time,
stratified by genotype. Coloring is identical to Figure 1A.

3.2. The EBOV NP-R111 Residue Lies Outside Known Functional Regions of NP, but Could Impact NP-NP Interaction

To investigate the functional importance of the NP-R111 residue, we examined existing annotations and functions of NP. The NP-R111 residue lies outside of key sites known to interact with EBOV RNA and VP35 (Figure 2A). Moreover, in NP crystal structures [23,25], the R111 residue appeared on the same face of the protein as the NP oligomerization domain (Figure 2B, left), opposite the key VP35 and RNA interaction residues (Figure 2B, right). Electron microscopy (EM) subtomogram averaging indicated that R111 is proximally located to key NP oligomerization residues [26]. Interestingly, R111 lies amidst a conserved stretch of 3 basic residues, K109, K110, and R111, on the surface of the NP protein (Figure 2C, yellow). Recent cryo-EM structures identified K110, adjacent to R111, as a residue forming a key electrostatic interstrand NP-NP interaction [28,29] that is highly conserved (Figure 2C). Indeed, deuterium exchange mass spectrometry indicated that K110 and R111 residues were partially buried in wild-type NP compared to an oligomerization-incompetent NP [27]. Therefore, we focused on whether NP-R111C affects oligomerization during the EBOV life cycle, and further queried this residue by generating charge-reversed mutants (NP-R111E and NP-K109E/K110E/R111E).

Figure 2. The EBOV NP-R111 residue is unannotated, but could impact NP-NP interaction. (**a**) Schematic of NP. R111 (yellow) lies in an un-annotated region within the N-terminal lobe. Key residues for known NP interactions are highlighted; (**b**) Crystal structure (PDB #4YPI) of NP. Though the precise location of the oligomerization domain has yet to be determined by crystallography (orange dashed line), the R111 residue (yellow) is located on the same face as residues proximal to the oligomerization domain (orange: residues 39, 40), but opposite to the VP35 (magenta: residues 160, 171, 174) and RNA (red: residues 240, 248, 252) interaction interfaces; (**c**) Alignment of *Ebolavirus* species. The basic residues at 109, 110, and 111 (blue), and a recently identified electrostatic interaction between K110-E349 [28,29], are conserved in all known ebolaviruses except Sudan virus (SUDV, red).

3.3. EBOV NP Position 111 Significantly Affects Oligomerization of NP

To address whether substitution at NP-R111 could influence NP-NP interaction and thereby
NP oligomerization, we developed an assay to measure intracellular NP oligomerization using
bioluminescence resonance energy transfer (BRET). Traditional oligomerization assays in cell
culture involve tagging a protein separately with two different tags, co-expressing both tagged
proteins, and then targeting one tag with co-immunoprecipitation (co-IP) and detecting the other
tag by western blot (WB) [53,69,70]. However, WB often has linear dynamic range issues;
furthermore, co-IPs can introduce non-specific or spurious protein-protein interactions using
different cell lysis and binding buffers. To overcome these deficits of co-IPs and WBs, we used
BRET to study NP oligomerization in live cells. We tagged the NP C-terminus with either the
chemiluminescent enzyme NanoLuc (NLuc) or the HaloTag protein (which covalently binds to an
acceptor fluorophore). We co-expressed both tagged NPs in cells, and activated NP-NLuc with
substrate, resulting in emission of light at 465 nm. Spatial proximity of NP-NLuc to NP-HaloTag
due to NP oligomerization results in energy transfer and a second light emission at a longer
wavelength, 625 nm (Figure 3A) [71].

NP	Max	K_d
R111E	14.9	2.50
K109E/K110E/R111E	14.3	2.78
R111C	14.4	3.44
R111	13.9	3.89
ΔOD	9.3	73.68

$p < 0.031$

$p < 0.11$

Figure 3. NP position 111 significantly affects oligomerization of NP. (**a**) Schematic of the NP
oligomerization assay. We co-expressed NP fused to NanoLuc (NLuc, donor) and HaloTag
(acceptor) in HEK 293FT cells. NP-NP binding and oligomerization brought the tags into close
spatial proximity, producing bioluminescence resonance energy transfer (BRET) emission at 625 nm.
To calculate BRET signal in milliBRET units, we normalized 625 nm BRET luminescence against
NP-NLuc luminescence at 465 nm and subtracted spectral spillover from NP-NLuc into the 625 nm
channel; (**b**) BRET oligomerization assay controls. Absence of either tag, free NLuc, or deletion of
the NP oligomerization domain (ΔOD, residues 19–38) reduced BRET signal; (**c**) EBOV VP35
NP-binding peptide (NPBP) disrupted NP oligomerization. In addition to NP-NLuc and
NP-HaloTag, we co-expressed varying amounts of VP35[NPBP] in cells. To quantify VP35[NPBP]
expression, we fused it to enhanced green fluorescent protein (eGFP), separated by a 'self-cleaving'
porcine teschovirus 1 2A peptide (P2A). We fitted oligomerization versus eGFP fluorescence units
(FU) to an inverse function (n = 3 biological replicates per VP35[NPBP] plasmid amount). Shading
indicates 95% confidence intervals based on 999 bootstrap pseudoreplicates; (**d**) Donor saturation
assay with NP mutants. We expressed a constant amount of NP-NLuc (donor) and expressed
varying amounts of NP-HaloTag (acceptor) in cells to generate donor saturation curves (n = 6
biological replicates per NP-HaloTag plasmid amount). We fitted data to saturation binding curves,
calculated maximum oligomerization (Max) and amount of NP-HaloTag plasmid needed to reach
half Max (K_d) for each NP mutant. eGFP (black dots near x-axis) did not produce data suitable for
curve fitting. We assessed statistical significance of differences in K_d between NP mutants by
ANOVA with Dunnett's test to correct for multiple hypothesis testing. Shading indicates 95%
confidence intervals based on 999 bootstrap pseudoreplicates.

To verify that our assay was truly measuring NP oligomerization, we generated a
loss-of-function (LOF) mutant and disrupted oligomerization with EBOV VP35. We generated
NP-ΔOD (deletion of oligomerization domain, NP residues 20–38), a LOF mutant that biochemical
methods (size exclusion chromatography and multiangle light scattering) indicated to be defective
in oligomerization [25]. To replicate these previous results in cell culture, we expressed V5- and
myc-tagged NP-ΔOD in cells and confirmed that NP-ΔOD lacked oligomerization capability using
the traditional dual-tag co-IP-WB strategy (Figure S1). We then generated NLuc- and
HaloTag-tagged NP and NP-ΔOD and performed our BRET assay in live cells. As expected, the lack
of NP-NLuc or NP-HaloTag, expression of NP-ΔOD, or expression of free NLuc reduced BRET
signal appreciably (Figure 3B). To confirm our assay in a biologically relevant context, we expressed
the NP-binding peptide (NPBP) of EBOV VP35 in cells, which is known to disrupt NP
oligomerization [24,25]. To quantitatively detect VP35[NPBP] expression, we fused enhanced green
fluorescent protein (eGFP) to NPBP via a bridging porcine teschovirus 1 2A 'self-cleaving' peptide
[48] (eGFP-P2A-VP35[NPBP]) and co-transfected varying amounts of this plasmid. Increasing
expression of eGFP-P2A-VP35[NPBP] led to a quantitative decrease in BRET oligomerization signal,
fitting well to an inverse function (Figure 3C).
Next, we measured the propensity of NP-R111 mutants to oligomerize and found that
NP-R111C, and to an even greater extent NP-R111E and NP-K109E/K110E/R111E, increased NP
oligomerization. To quantify oligomerization, we expressed increasing amounts of acceptor
NP-HaloTag in cells to saturate the donor NP-NLuc signal. The resulting oligomerization curves fit
well to saturation binding curves, parameterized by Max (maximum oligomerization) and K_d
(amount of NP-HaloTag plasmid needed to reach half Max) (Figure 3D). As expected, control eGFP
substituted for NP-NLuc (black dots) resulted in no detectable BRET signal, and NP-ΔOD (gray)
resulted in background signal at high concentrations. Relative to NP-R111 (tan), NP-R111C (red)
slightly, but not statistically significantly, increased oligomerization (12% lower K_d; $p < 0.74$,
ANOVA with Dunnett's test to correct for multiple hypotheses), whereas the charge-reversed
NP-R111E (light blue; 36% lower K_d; $p < 0.031$, ANOVA-Dunnett's test) and
NP-K109E/K110E/R111E (dark blue; 28% lower K_d; $p < 0.11$, ANOVA-Dunnett's test) mutants
oligomerized at even lower NP concentrations (Figure 3D). We verified that different NP variants
were expressed at similar concentrations by comparing the luminescence of NP-NLuc mutants in

the absence of the HaloTag substrate (Figure S2). These results support our hypothesis that the NP
111 allele affects the K110-E349 NP-NP interaction (Figure 2C) suggested by cryo-EM [28,29].

3.4. EBOV NP-R111C Increases Budding of Virion-like Particles

To determine whether the different mutants impact NP's role in virion structure, we designed
and optimized a VLP budding assay. Traditionally, researchers assess viral budding efficiency by
harvesting cell culture supernatants, purifying VLPs by ultracentrifugation through sucrose, and
detecting VLPs by WB using antibodies to specific VLP components [35,51,72,73]. However, WBs
are often insensitive to modest changes in VLP numbers and can suffer from high technical
variability. By contrast, luminescence can be reproducibly detected over a larger linear dynamic
range. However, the size of firefly luciferase (FLuc; 60 kDa) can severely interfere with
incorporation into budding VLPs. Indeed, although the EBOV matrix protein VP40 (40 kDa) alone
is sufficient to bud VLPs [32–34], fusion of VP40 to FLuc decreased luciferase activity to
undetectable levels in a budding assay [72]. Here, we took advantage of the smaller size of NLuc (19
kDa) [74] and fused it to VP40. We expressed NLuc-VP40 in cells, purified VLPs following
established protocols, and measured NLuc reporter activity (Figure 4A).

Figure 4. EBOV NP-R111C increases budding of VLPs. (a) Schematic of the VLP budding assay.
We expressed NLuc fused to EBOV VP40 (NLuc-VP40) in HEK 293FT cells to form luminescent
VLPs, and co-expressed NP mutants to measure the impact of NP genotype on VLP budding; **(b)**
VLP budding assay control. NLuc-VP40 expression alone results in bright luminescence, expressed
in relative light units (RLU). VP40 loss-of-function (LOF) mutant L117R fails to form VLPs (n = 6
biological replicates). We assessed statistical significance by paired t-test. Error bars indicate mean ±
standard error of the mean (SEM); **(c)** VLP budding with NP variants. We measured and normalized
each NP mutant's VLP production to that of NP-R111 (n = 6 biological replicates) and assessed
statistical significance using repeated measures ANOVA with Dunnett's test to correct for multiple
hypothesis testing. Error bars indicate mean ± SEM.

One major challenge to VLP budding assays is that VP40 can be expelled from cells as a
monomer [75] perhaps via exosomes [76]. To distinguish between monomeric VP40 and VLPs, we
used a LOF mutant, VP40-L117R, which is defective in VLP budding as judged by
immunofluorescence microscopy and WB [51]. We expressed NLuc-VP40 or NLuc-(VP40-L117R) in
cells and collected unpurified supernatant or purified VLPs by ultracentrifugation through a 20%
(w/v) sucrose cushion. We added NLuc substrate and measured luminescence in unpurified
supernatant or the VLP-containing pellet. As expected, budding of VLPs containing
NLuc-(VP40-L117R) LOF mutant was impaired by >400-fold compared to NLuc-VP40 (p < 0.001;
paired t-test) (Figure 4B). We did not observe as large of a difference between NLuc-VP40 and
NLuc-(VP40-L117R) in unpurified culture supernatant (Figure S3A, left), suggesting that an

appreciable amount of monomeric VP40 was secreted from cells. Moreover, treating cells with Brefeldin A, an inhibitor of coat protein complex I (COPI)-mediated transport, did not affect luminescence in unpurified supernatant (Figure S3A), suggesting that secretion of monomeric VP40 was independent of this major transport mechanism, as reported previously [75]. To further verify that we were measuring luminescence from VLPs rather than monomeric VP40, we heated culture supernatant prior to ultracentrifugation. If VLPs were present, heating would dissociate VLPs into VP40 monomers [77], which would fail to pellet after ultracentrifugation through sucrose. Whereas heating supernatant to 60.2 °C does decrease NLuc activity [74], heating and subsequent ultracentrifugation reduced NLuc activity an additional 15-fold (p < 0.007; paired t-test), suggesting that NLuc-VP40 VLPs were denatured and thus were not pelleted and detected (Figure S3B).

We then further optimized our VLP assay to maximize the difference between NLuc-VP40 and the LOF mutant NLuc-(VP40-L117R). We first transfected cells with a range of NLuc-VP40 or NLuc-(VP40-L117R) plasmid amounts and measured luminescence in unpurified culture supernatant. A relatively small amount of NLuc-VP40 plasmid (2.22 ng) produced the greatest difference in luminescence between NLuc-VP40 and NLuc-(VP40-L117R) (Figure S3C). Though NLuc is a relatively small protein (19 kDa), fusion to VP40 (40 kDa) could still impair VP40 interactions and functions. Co-expressing increasing amounts of 'dark' untagged VP40 in cells did not increase the difference in luminescence between wild-type NLuc-VP40 and LOF mutant (Figure S3D), suggesting that the NLuc tag did not drastically interfere with VLP production. Lastly, we optimized the amount of NP plasmid to co-transfect with NLuc-VP40, and, consistent with NP's role in promoting VLP formation, increasing the amount of NP plasmid transfected always increased luminescence and greater difference in luminescence between NLuc-VP40 and NLuc-(VP40-L117R) (Figure S3E).

Finally, we tested how different NP mutants affected our VLP budding assay and found that only NP-R111C affected VLP production. Expression of viral nucleoproteins, including EBOV NP, is known to significantly increase matrix protein-induced VLP production [35]. We verified that ancestral NP-R111 (tan) increased NLuc-VP40 VLP production 1.93-fold compared to eGFP control (gray; p < 0.0002, repeated measures ANOVA with Dunnet's test to correct for multiple hypothesis testing) (Figure 4C). NP-R111C (red) significantly increased VLP production above NP-R111 (1.26-fold; p < 0.039, rANOVA-Dunnett's test), whereas the charge-reversed NP-R111E (light blue; 1.07-fold; p < 0.847, rANOVA-Dunnett's test) and NP-K109E/K110E/R111E (dark blue; 1.13-fold; p < 0.484, rANOVA-Dunnett's test) did not have reproducible effects. To determine whether increased luminescence was due to increased VLP size, we expressed unfused VP40, GP, and NP-R111 or NP-R111C, and visualized VLPs via EM (Figure S4A). However, we did not detect any changes in length or volume of VLPs due to NP-R111C substitution (Figure S4B).

3.5. EBOV NP Position 111 Influences Viral Transcription and Replication

The mechanism by which changes in NP's structural phenotypes (e.g., oligomerization, budding) affect viral transcription and replication is not obvious because NP is highly multi-functional. NP's ability to oligomerize can influence its ability to bind RNA and modulate transcription and replication, though the direction and size of this effect are not always clear [24,25,27]. We quantified viral transcription and replication using two minigenome reporter assays (Figure 5A) [52,57]. In each assay, we expressed the components of the EBOV ribonucleoprotein (RNP) complex (NP, VP35, VP30, and L) in cells in the presence of a minigenome encoding a luciferase reporter flanked by the EBOV promoter-like leader and trailer sequences. Transcription is essential for minimal luciferase activity; replication is further required to achieve maximum signal [78]. Whereas the first reporter system uses a monocistronic minigenome (1MG) to only assess transcription and replication [52], the second system uses a tetracistronic minigenome (4MG) to produce transcription- and replication-competent VLPs (trVLPs) (Figure 5A). These 4MG trVLPs can be further 'passaged' as a measure of ability to infect target cells and complete the viral life cycle [57]. As a positive control, we further tested the 4MG system in the presence of the GP-A82V mutation, which increases viral infectivity in cell culture [15–20].

Figure 5. EBOV NP position 111 influences viral transcription and replication. (**a**) Schematic of monocistronic minigenome (1MG) and tetracistronic minigenome (4MG) systems. Compared to live virus genome (top), 1MG only encodes the EBOV leader and trailer sequences (middle), whereas the 4MG encodes the structural proteins VP40, GP, and VP24 (bottom). In both cases, the replication complex proteins (NP, VP35, VP30, and L) are expressed from plasmids *in trans*. Tan line indicates the position of NP-R111, and the dashed line indicates the position of GP-A82 (encoded on 4MG but not 1MG); (**b**) 1MG assay. We expressed NP mutants or ancestral NP-R111 in HEK 293T cells in the presence of the EBOV replication complex (L, VP30, VP24), and measured transcription and replication (txn/rep) of the 1MG minigenome encoding firefly luciferase (FLuc) relative to a *Renilla* luciferase (RLuc) loading control (n = 3 biological replicates). We normalized values to NP-R111. Absence of L, VP30, or 1MG abolished FLuc signal. Both NP-R111E and NP-K109E/K110E/R111E charge-reversal mutants significantly decreased 1MG activity (ANOVA-Dunnett's test). Error bars indicate mean ± SEM; (**c**) P0 producer cells of 4MG assay. We expressed trVLPs harboring GP-A82 (left) or GP-A82V (right) and NP mutants in HEK 293T cells, and measured 4MG minigenome activity (RLuc) relative to an FLuc loading control (n = 3–9 biological replicates). We normalized values to GP-A82/NP-R111 and assessed statistical significance by ANOVA with Tukey's test (ANOVA-Tukey's test) to correct for multiple hypothesis testing. Error bars indicate mean ± SEM; (**d**) Target P1 and P2 cells of 4MG assay. We expressed the EBOV replication complex in P1 Huh7 cells, and then infected P1 cells with P0 trVLPs, and measured 4MG activity. We repeated the process by expressing the EBOV replication complex in P2 Huh7 cells, infecting P2 cells with P1 trVLPs, and measuring 4MG activity again. We normalized values for cells in each biological replicate to its own P0 activity and assessed statistical significance by ANOVA-Tukey's test. Solid

lines indicate GP-A82, whereas dashed lines indicate GP-A82V. Bold lines indicate mean value of 3–

9 biological replicates, shown as faded lines. Error bars indicate mean ± SEM.

Using the 1MG system, only the charge-reversal NP mutants affected transcription and
replication differently than the wild-type. As expected, the absence of EBOV VP30, L, or the
minigenome (gray) resulted in <5% normalized minigenome activity compared to cells expressing
the minigenome and the entire RNP complex with NP-R111 (tan, Figure 5B). Substitution of
NP-R111C (red) in place of NP-R111 yielded similar activity (99%). On the other hand, the
charge-reversal mutants NP-R111E (light blue; 23% reporter activity; $p < 0.003$; ANOVA-Dunnett's
test) and NP-K109E/K110E/R111E (dark blue; 44% activity; $p < 0.017$; ANOVA-Dunnett's test)
greatly attenuated transcription and replication.

In the 4MG trVLP system, we again observed that NP-R111E abrogated transcription and
replication in the producer (P0) cells, though NP-R111C modestly decreased activity as well. As
expected, we detected virtually no signal from any cells in the absence of EBOV L (gray). Paired
with the ancestral GP-A82 (Figure 5C, left), NP-R111E (light blue) dramatically reduced reporter
activity compared to NP-R111 (tan; 33% activity; $p < 0.0001$, ANOVA-Tukey's test to correct for
multiple hypothesis testing), and NP-R111C (red) modestly decreased activity as well (80% activity;
$p < 0.0001$, ANOVA-Tukey's test). Because the P0 cells are transfected with plasmid encoding the
4MG minigenome, these cells primarily account for transcription and replication without the
requirement for trVLP entry and spread; therefore, we do not expect the GP-A82V substitution,
which only increases viral entry [15–20], to dramatically affect transcription and replication in P0
cells. Indeed, compared to GP-A82, GP-A82V did not measurably increase reporter activity in the
context of NP-R111 (tan, left vs right; 102% activity; $p < 0.96$, ANOVA-Tukey's test), and marginally
enhanced the activity of NP-R111C (red, left vs right; 80% to 90% activity; $p < 0.006$,
ANOVA-Tukey's test) but not NP-R111E (light blue, left vs right; 33% to 35% activity; $p < 0.998$,
ANOVA-Tukey's test, Figure 5C).

We used trVLP-containing culture supernatant from P0 cells to inoculate a first round of target
cells (P1), and repeated the procedure using P1 trVLPs to inoculate a second round of target cells
(P2). P1 and P2 cells were not transfected with plasmid encoding 4MG; instead, these cells
contained 4MG minigenome if and only if a trVLP entered the cell. Since P1 and P2 minigenome
activity depended on transcription, replication, and viral entry and spread, we calculated
fold-change in reporter gene activity relative to P0 (Figure 5D). In the presence of ancestral GP-A82,
NP-R111C (solid red lines) modestly decreased reporter activity compared to NP-R111 in P1 (solid
tan lines; 77% of fold-change in activity; $p < 0.053$, ANOVA-Tukey's test) and further in P2 (59% of
fold-change in activity; $p < 0.35$, ANOVA-Tukey's test). NP-R111E (solid blue lines) appeared
completely unable to spread in P1 (1% of fold-change in activity; $p < 0.0001$, ANOVA-Tukey's test)
and P2 (0.05% of fold-change in activity; $p < 0.0003$, ANOVA-Tukey's test) cells. As expected,
GP-A82V (dashed lines) spread much better than ancestral GP-A82 (solid lines) in the context of
NP-R111 (tan; P1: 137% of fold-change in activity, $p < 0.0001$, ANOVA-Tukey's test; P2: 218% of
fold-change in activity, $p < 0.0001$, ANOVA-Tukey's test) and NP-R111C (red; P1: 77% to 104% of
fold-change in activity, $p < 0.014$; P2: 59% to 67% of fold-change in activity, $p < 0.999$), though
NP-R111E was completely defective regardless of GP-82 allele (dashed and solid light blue lines
overlap along the x-axis).

3.6. Probing EBOV NP's Interactome Reveals Interaction with the AP-1 Clathrin Adaptor Complex

As NP-NP interactions appeared to be affected by NP position 111, we sought to query
whether interactions with host proteins may explain how NP enhances VLP budding. Only a single
interactome study has been performed on EBOV NP fused to eGFP [79], which utilized the NP
amino acid sequence from the Mayinga isolate of the EBOV Yambuku variant (EBOV/Yam-May),
the first EBOV isolated in 1976. To build upon these previous results, we performed
co-immunoprecipitation tandem mass spectrometry (co-IP MS/MS) using myc-tagged NP from
EBOV/Makona bearing either R111 or K109E/K110E/R111E. Our approach yielded multiple

members of the adaptor related protein 1 (AP-1) complex as strong candidate host protein
interactors (Figure S5A, right), which were identified previously [79] but were not further
confirmed in that study. Here, we confirmed that both NP-R111 and NP-K109E/K110E/R111E
strongly interacted with AP-1 subunit M1 (AP1M1) and AP1G1 by reciprocal IP-WB (Figure S5B).
Yet, NP-R111C, NP-R11E, and NP-K109E/K110E/R111E all bound to the AP-1 complex with similar
affinity as NP-R111 (Figure S5C), suggesting that the AP-1 interaction did not explain differences
between the mutants in our assays.

4. Discussion

Here, we developed and modified BSL-2 assays to study in-depth a key EBOV NP substitution,
NP-R111C, which arose during the 2013–2016 Western African EVD epidemic. Though the NP-R111
residue has not been previously annotated as functional, the residue's proximity to a key NP-NP
electrostatic interaction led us to consider that substitution at this residue could affect multiple viral
phenotypes. Because EBOV NP plays many critical roles during the life cycle, we developed assays
for NP oligomerization and VLP budding in live cells with controls. These assays are biologically
relevant for EBOV and may also be appropriate for other viral nucleoproteins as well. Our data
reveal that NP residue 111 is importantly positioned to affect both phenotypes as well as viral
transcription and replication. NP-R111, the adjacent basic residues K109 and K110, and the
K109-E349 salt bridge identified by cryo-EM [28,29], are highly conserved among Ebolavirus
species, including in the newly described Bombali virus [80] (Figure 2C). This high degree of
conservation, despite significant evolutionary divergence between ebolaviruses, emphasizes the
importance of this highly basic region to NP functions.
Charge-reversal at these residues, NP-R111E and NP-K109E/K110E/R111E, produced
dramatic phenotypes, including significantly increased oligomerization (Figure 3D). This increase is
seemingly paradoxical, since charge-reversal should disrupt the K109-E349 electrostatic interaction
between NP monomers. However, even if this interaction is lost, a nearby residue, K352, may be
able to stabilize E349. Moreover, the interface between the two NP monomers is very hydrophilic
and likely filled with water. As such, modeling how NP-R111E would affect NP oligomer structure
is challenging. Nevertheless, this substitution increases BRET signal, suggesting that it either
increases NP oligomerization propensity or alters the oligomer structure to bring the NLuc and
HaloTag tags (at the NP C-terminus) in closer proximity.
It is intriguing that NP-R111E's increased oligomerization correlated with severely defective
minigenome transcription and replication (Figure 5B, 5C), to the degree that NP-R111E trVLPs were
unable to replicate over multiple passages (Figure 5D). NP oligomers coat viral RNA to defend
against host viral RNA sensors and nucleases; however, these oligomers also prevent the viral
polymerase L from accessing viral RNA. During viral transcription and replication, accumulation of
EBOV VP35 disrupts NP oligomers (Figure 3C), releasing free viral RNA as a template for
RNA-directed RNA synthesis [24,25]. One possible hypothesis is that NP-R111E stabilizes
oligomers too strongly, rendering viral RNA inaccessible to L, in turn abrogating minigenome
transcription and replication. Because NP-R111E trVLPs were extremely defective, charge-reversal,
unsurprisingly, has never been observed in EBOV or most other ebolavirus sequences (Figure 2C).
Surprisingly, Sudan virus (SUDV), a replication-competent ebolavirus that is lethal in humans, does
encode a charge-reversal substitution, NP-K110E (Figure 2C). The secondary structure of an SUDV
NP monomer is quite similar to that of EBOV NP [81], and residues 109–111 are positioned in the
same β-strand in both. However, SUDV NP oligomers have not yet been characterized, so it is
possible that one of the many other amino acid differences between SUDV and EBOV could
counteract the effect of NP-K110E in SUDV. Future study is required to better understand how
EBOV NP oligomerization affects viral RNA synthesis, and how SUDV NP promotes viral
transcription and replication despite a charge-reversal substitution.
By contrast, the epidemic-associated EBOV substitution, NP-R111C, produced more
intermediate phenotypes – slightly increased oligomerization (Figure 3D), moderately increased
VLP production (Figure 4C), and moderately reduced transcription and replication (Figure 5C, 5D).

To better understand how this substitution might influence VLP production, we performed co-IP
MS/MS and confirmed an interaction between NP and the clathrin adaptor AP-1 complex (Figure
S5B), which had been identified but not specifically validated previously [79]. In retroviruses, the
Gag protein facilitates budding by hijacking the AP-1 complex [82]. Because retroviral and EBOV
virions bud using the same cellular pathway and machinery [73], we hypothesize that EBOV NP
also co-opts the AP-1 complex for virion egress and trafficking. This interaction with the AP-1
complex could be a mechanism to explain how NP expression significantly enhances budding of
EBOV VLPs [35]. Yet both NP-R111 and NP-R111C bound to AP-1 with similar affinity (Figure
S5C). Thus, the interaction with AP-1 did not explain NP-R111C's increased ability to promote VLP
production. Much work remains to be done to better understand how NP enacts so many roles
during the viral life cycle.

Our data point to a complex interplay between NP oligomerization and NP's roles in structure
and replication, and highlight the importance of developing reporter assays for viruses, especially
pathogens of high consequence. Many viral proteins are highly multi-functional, making study of
individual mutations challenging without robust assays that are sensitive to subtle changes in viral
phenotype. Since luciferase-based reporter systems fit the aforementioned requirements, we took
advantage of these systems to develop a BRET assay for NP oligomerization and a VLP detection
assay, which offer a number of advantages over existing methods. Previous attempts to assess NP
oligomerization typically used biochemical methods with purified protein [24,25] or by
co-immunoprecipitation from cell lysates [53,69,70]. In our BRET assay, the NLuc substrate and
HaloTag ligand have minimal cytotoxicity and readily diffuse into live cells, key features that
minimized spurious interactions and allowed us to assess oligomerization over a range of
physiologically-relevant EBOV NP concentrations (Figure 3D). To measure VLP production,
previous attempts to fuse FLuc (60 kDa) to VP40 (40 kDa) resulted in near undetectable levels of
luminescent VLPs [72]. In our VLP budding assay, we took advantage of the brightness and small
size of NLuc (19 kDa) to show that NLuc fused to VP40 generated luminescent VLPs, and that
NP-R111C modestly increased VLP production (Figure 4C).

These BSL-2 assays are simple and flexible for testing viral mutations of interest like
NP-R111C, which emerged and quickly became a dominant substitution during the 2013–2016 EVD
epidemic. In particular, the BRET assay for NP oligomerization can be performed in
high-throughput. BRET signal is quite reproducible between replicates and plates because the assay
includes normalization based on NLuc luminescence to adjust for differences in transfection or
protein expression efficiency, and subtraction of background luminescence to compensate for
luminescence spillover from NLuc into the BRET channel. With more rigorous screening and
quantification of key metrics of variability like Z-factor, these assays could be used for
high-throughput screens of hundreds of EBOV NP mutants, interactions with host factors, or
antagonism by drug candidates. The flexibility of BSL-2 reporter assays makes them ideal as
first-line methods to probe mutations and mechanism prior to cumbersome BSL-4 studies with live
virus.

Testing whether emergent mutations like NP-R111C affect fitness of live virus brings
additional challenges because different viral stocks and cultured cells versus animal models can
cause discordant results. For example, the GP-A82V substitution has been shown numerous times
to enhance EBOV infectivity in cell culture [15–20], using multiple EBOV surrogate systems (e.g.,
EBOV VLPs, retroviral particles pseudotyped with EBOV GP, and recombinant live virus [17]) and
multiple cell types (e.g., human monocyte-derived dendritic cells [15]). However, a recent study
using immunocompromised laboratory mice and non-human primates [21] indicated that EBOV
Makona viral isolates encoding GP-A82V lead to modestly decreased viral load compared those
without the GP-A82V mutation. A second recent study found that GP-A82V may induce slightly
more morbidity and fatality in immunocompromised laboratory mice but not in domestic ferrets
[22]. The discrepancies between the *in vitro* and *in vivo* studies could be due to the use of clinical
EBOV isolates [21] that contain multiple additional mutations versus live EBOV generated from
recombinant DNA plasmids [17]. Limited recombinant live EBOV studies have been performed to

test the impact of the NP-R111C substitution specifically. One study showed that EBOV with
GP-A82V and NP-R111C outcompetes ancestral EBOV Makona in a head-to-head format in
cultured cells, but did not measure the impact of NP-R111C alone [17]. Another study found that
NP-R111C alone increases viral replication in cell culture, decreases morbidity and lethality in
immunocompromised laboratory mice, and does not differ from ancestral EBOV Makona in
domestic ferrets [22].

Revealing the importance of the NP 111 residue and establishing experimental systems
represent steps towards characterizing a key EBOV substitution, NP-R111C, that arose during the
2013–2016 Western African EVD epidemic. Given current limitations and restrictions of BSL-4
settings and the potential global threat of EVD epidemics, BSL-2 model systems can facilitate rapid
and broad exploration of any potentially consequential EBOV mutations. We demonstrate that
NP-R111C modestly increased VLP production while slightly decreasing viral transcription and
replication. On the other hand, a charge-reversal substitution at the same site, observed in SUDV
but not other EBOV or other ebolaviruses, caused drastically increased oligomerization and ablated
transcription and replication. These findings provide additional insight into the interplay between
the many functions of NP in oligomerization, viral assembly and budding, and transcription and
replication, and suggest that NP-R111C and other substitutions at the 111 residue merit further
study using live virus.

Supplementary Materials: The following are available online at www.mdpi.com/xxx/s1, Figure S1: Deletion of
the reported NP oligomerization domain (NP-ΔOD) prevented NP oligomerization in cell culture, Figure S2:
NP-NLuc variants expressed at similar concentrations, Figure S3: Optimization of NLuc-VP40 VLP assay,
Figure S4: Electron microscopy of NP-R111 and NP-R111C VLPs, Figure S5: EBOV NP interacts with the
adaptor protein 1 (AP-1) complex independent of the NP 111 allele, File S1: Key resources table, File S2: List of
all proteins identified by co-immunoprecipitation tandem mass spectrometry (co-IP-MS/MS), related to Figure
S5A.

Figure S1. Deletion of the reported NP oligomerization domain (NP-ΔOD) prevented NP
oligomerization in cell culture. We co-expressed myc- and V5-tagged NP or NP-ΔOD (deletion of
residues 20–38) in HEK 293FT cells, harvested and lysed cells, and performed
co-immunoprecipitation (co-IP) targeting either the myc (blue) or the V5 (orange) tag, or IgG isotype
controls. We used 2.5% of each input and 25% of each IP for detection of co-eluting proteins by WB
for each tag. Oligomerization is indicated by heterologous detection (NP-V5 band in the IP myc

fraction, and vice versa) with full-length NP, which did not occur with NP-ΔOD. Additional bands
are due to heavy (H) and light (L) chains of the co-IP antibody.

Figure S2. NP-NLuc variants expressed at similar concentrations. As part of the BRET assay, we
measured NLuc-driven luminescence of each NP genotype in the absence of HaloTag ligand in HEK
293FT cells. This luminescence solely reflects the amount of NP-NLuc expression since no resonant
energy transfer can occur without the HaloTag ligand. Coloring is the same as in Figure 3D.
NP-NLuc expression does not appear to correlate with oligomerization propensity. Error bars
indicate mean ± standard error of the mean (SEM).

Figure S3. Optimization of NLuc-VP40 VLP assay. (a) VP40 monomers are expelled from cells independent of coat protein complex I (COPI) transport. We expressed NLuc-VP40 wild-type (tan) or L117R substitution (gray) in HEK 293FT cells, and treated with either Brefeldin A (BrfA) or dimethyl sulfoxide (DMSO) vehicle control. We collected culture supernatant and measured luminescence directly, or purified VLPs through a 20% sucrose cushion and then measured luminescence. NLuc-(VP40-L117R) luminescence greatly decreased after ultracentrifugation, suggesting that VP40 in culture supernatant was monomeric. BrfA did not appear to affect NLuc-(VP40-L117R) luminescence, suggesting that this phenomenon is independent of COPI transport; (b) Heating supernatant prior to ultracentrifugation results in loss of VLP luminescence. We expressed NLuc-VP40 in cells, collected total supernatant, and heated at various temperatures.

We then either measured luminescence immediately to assess NLuc thermal stability (blue), or pelleted VLPs and then measured luminescence to assess VLP stability (tan) (n = 3 biological replicates). We normalized luminescence at each temperature to the 4 °C condition, log-transformed, and fitted to sigmoidal curves; (**c**) Lower amounts of NLuc-VP40 plasmid led to more VLP budding than monomer expulsion. We expressed NLuc-VP40 wild-type (tan) or L117R mutant (gray) in cells, collected culture supernatant, and measured NLuc luminescence (n = 4–6 biological replicates). We observed the largest difference between NLuc-VP40 and NLuc-(VP40-L117R) at 2.22 ng plasmid per well of a 96-well plate. Error bars indicate mean ± SEM; (**d**) Co-expressing untagged VP40 in cells does not improve NLuc-VP40 VLP budding. We co-expressed NLuc-VP40 wild-type (tan) or L117R mutant (gray) with 'dark' untagged VP40 in cells, and measured culture supernatant luminescence (n = 2 biological replicates). Increasing amounts of 'dark' VP40 did not increase the difference between the NLuc-VP40 and NLuc-(VP40-L117R). Error bars indicate mean; (**e**) Expression of NP increases VLP production. We co-expressed NLuc-VP40 wild-type (tan) or L117R (gray) with NP in cells, collected culture supernatant, and measured culture supernatant luminescence (n = 4 biological replicates). Increasing amounts of NP plasmid increased the difference between VP40 variants. Error bars indicate mean ± SEM.

Figure S4. Electron microscopy of NP-R111 and NP-R111C VLPs. (**a**) EM of VP40 VLPs created by with co-expression of VP40, GP-A82V, and NP-R111 or NP-R111C in HEK 293T cells. Scale bar (black line in bottom left) denotes 200 nm. Error bars indicate mean ± SEM; (**b**) Quantification of VLP size and volume for VLPs. VLPs bearing NP-R111 (tan) and NP-R111C (red) have similar size and volume. Error bars indicate mean ± SEM.

Author Contributions: conceptualization, A.E.L., W.E.D., J.L., and P.C.S.; methodology, A.E.L., W.E.D., Y.C.,
C.F., C.A., R.N.K., J.H.K., and J.L.; software, A.E.L. and K.G.A.; validation, W.E.D., Y.C., C.F., and R.N.K.;
formal analysis, A.E.L., W.E.D., R.N.K., and K.G.A.; investigation, A.E.L., W.E.D., Y.C., C.F., C.A., and R.N.K.;

data curation, A.E.L.; writing—original draft, A.E.L., W.E.D., S.F.S., E.A.B., K.G.A., J.L., and P.C.S.;
writing—review & editing, A.E.L., W.E.D., Y.C., C.F., C.A., R.N.K., L.B., S.F.S., E.A.B., E.O.S., K.G.A., J.H.K.,
J.L., and P.C.S.; visualization, A.E.L., W.E.D., L.B., and S.F.S.; supervision, E.O.S., K.G.A., J.H.K., J.L., and
P.C.S.; project administration, J.L. and P.C.S.

Funding: This material was based upon work supported by the Howard Hughes Medical Institute Investigator
Award (P.C.S) and the US National Institutes of Health (NIH) grants National Institute of Allergy and
Infectious Diseases (NIAID) U19AI110818 (P.C.S.), National Institute on Drug Abuse (NIDA) DP1DA034990
(J.L.), NIAID R01AI111809 (J.L.), and NIAID R01AI118016 (E.O.S.). A.E.L. was supported by National Science
Foundation (NSF) under Grant No. DGE 1144152. C.A. was supported by award Number T32GM007753 from
the NIH's National Institute of General Medical Sciences (NIGMS). E.O.S. was supported by an Investigators in
the Pathogenesis of Infectious Disease Award from the Burroughs Wellcome Fund, by the Skaggs Institute for
Chemical Biology, by the Skaggs Institute for Chemical Biology, and by the Shaffer Family Foundation. K.G.A.
was a Pew Biomedical Scholar and was supported by NIH National Center for Advancing Translational
Studies Clinical and Translational Science (NCATS) Award UL1TR001114, and NIAID contract
HHSN272201400048C. This work was also funded in part through Battelle Memorial Institute's prime contract
with the US NIAID under Contract No. HHSN272200700016I (Y.C., C.L.F., L.B., J.H.K.). The funders had no
role in study design, data collection and interpretation, or the decision to submit the work for publication.

Acknowledgments: We thank Ross Tomaino of the Harvard Taplin Mass Spectrometry Facility for sample and
instrument handling following immunoprecipitation of NP-containing protein complexes. We thank Gregory
Hendricks of the University of Massachusetts Medical School Core Electron Microscopy Facility for sample
and instrument handling following production of Ebola virion-like particles. Electron Microscopy (EM) was
supported by Award Number S10RR027897 from the National Center for Research Resources to the University
of Massachusetts Medical School Core EM Facility. Ebola virus phylogenetics was made possible by the Virus
Pathogen Database and Analysis Resource (ViPR) [42], which has been wholly funded with federal funds from
NIAID, the US National Institutes of Health, and the US Department of Health and Human Services (DHSS),
under Contract No. HHSN272201400028C. The authors are solely the responsibility for the content of this
paper and do not necessarily represent the official views of the National Center for Research Resources or of
US DHHS or of the institutions and companies affiliated with the authors.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Holmes, E.C. *The Evolution and Emergence of RNA Viruses*; Oxford University Press, 2009; ISBN
9780199211128.
2. World Health Organization (WHO) Ebola Situation Report - 30 March 2016. Available online:
http://apps.who.int/ebola/current-situation/ebola-situation-report-30-march-2016 (accessed on 9
December 2019).
3. Baize, S.; Pannetier, D.; Oestereich, L.; Rieger, T.; Koivogui, L.; Magassouba, N.; Soropogui, B.; Sow, M.S.;
Keïta, S.; De Clerck, H.; et al. Emergence of Zaire Ebola virus disease in Guinea. *N. Engl. J. Med.* **2014**, *371*,
1418–1425.
4. Gire, S.K.; Goba, A.; Andersen, K.G.; Sealfon, R.S.G.; Park, D.J.; Kanneh, L.; Jalloh, S.; Momoh, M.; Fullah,
M.; Dudas, G.; et al. Genomic surveillance elucidates Ebola virus origin and transmission during the 2014
outbreak. *Science* **2014**, *345*, 1369–1372.
5. Carroll, M.W.; Matthews, D.A.; Hiscox, J.A.; Elmore, M.J.; Pollakis, G.; Rambaut, A.; Hewson, R.;
García-Dorival, I.; Bore, J.A.; Koundouno, R.; et al. Temporal and spatial analysis of the 2014-2015 Ebola
virus outbreak in West Africa. *Nature* **2015**, *524*, 97–101.
6. Ladner, J.T.; Wiley, M.R.; Mate, S.; Dudas, G.; Prieto, K.; Lovett, S.; Nagle, E.R.; Beitzel, B.; Gilbert, M.L.;
Fakoli, L.; et al. Evolution and spread of Ebola virus in Liberia, 2014-2015. *Cell Host Microbe* **2015**, *18*, 659–
669.
7. Tong, Y.-G.; Shi, W.-F.; Liu, D.; Qian, J.; Liang, L.; Bo, X.-C.; Liu, J.; Ren, H.-G.; Fan, H.; Ni, M.; et al.
Genetic diversity and evolutionary dynamics of Ebola virus in Sierra Leone. *Nature* **2015**, *524*, 93–96.
8. Park, D.J.; Dudas, G.; Wohl, S.; Goba, A.; Whitmer, S.L.M.; Andersen, K.G.; Sealfon, R.S.; Ladner, J.T.;
Kugelman, J.R.; Matranga, C.B.; et al. Ebola virus epidemiology, transmission, and evolution during
seven months in Sierra Leone. *Cell* **2015**, *161*, 1516–1526.

9.   Simon-Loriere, E.; Faye, O.; Faye, O.; Koivogui, L.; Magassouba, N.; Keita, S.; Thiberge, J.-M.; Diancourt,
968   L.; Bouchier, C.; Vandenbogaert, M.; et al. Distinct lineages of Ebola virus in Guinea during the 2014 West
969   African epidemic. *Nature* **2015**, *524*, 102–104.

970   10.   Kugelman, J.R.; Sanchez-Lockhart, M.; Andersen, K.G.; Gire, S.; Park, D.J.; Sealfon, R.; Lin, A.E.; Wohl, S.;
971   Sabeti, P.C.; Kuhn, J.H.; et al. Evaluation of the potential impact of Ebola virus genomic drift on the
972   efficacy of sequence-based candidate therapeutics. *MBio* **2015**, *6*, e02227-14.

973   11.   Hoenen, T.; Safronetz, D.; Groseth, A.; Wollenberg, K.R.; Koita, O.A.; Diarra, B.; Fall, I.S.; Haidara, F.C.;
974   Diallo, F.; Sanogo, M.; et al. Virology. Mutation rate and genotype variation of Ebola virus from Mali case
975   sequences. *Science* **2015**, *348*, 117–119.

976   12.   Hoenen, T.; Groseth, A.; Rosenke, K.; Fischer, R.J.; Hoenen, A.; Judson, S.D.; Martellaro, C.; Falzarano, D.;
977   Marzi, A.; Squires, R.B.; et al. Nanopore sequencing as a rapidly deployable Ebola outbreak tool. *Emerg.*
978   *Infect. Dis.* **2016**, *22*, 331–334.

979   13.   Smits, S.L.; Pas, S.D.; Reusken, C.B.; Haagmans, B.L.; Pertile, P.; Cancedda, C.; Dierberg, K.; Wurie, I.;
980   Kamara, A.; Kargbo, D.; et al. Genotypic anomaly in Ebola virus strains circulating in Magazine Wharf
981   area, Freetown, Sierra Leone, 2015. *Euro Surveill.* **2015**, *20*, pii=30035.

982   14.   Quick, J.; Loman, N.J.; Duraffour, S.; Simpson, J.T.; Severi, E.; Cowley, L.; Bore, J.A.; Koundouno, R.;
983   Dudas, G.; Mikhail, A.; et al. Real-time, portable genome sequencing for Ebola surveillance. *Nature* **2016**,
984   *530*, 228–232.

985   15.   Diehl, W.E.; Lin, A.E.; Grubaugh, N.D.; Carvalho, L.M.; Kim, K.; Kyawe, P.P.; McCauley, S.M.; Donnard,
986   E.; Kucukural, A.; McDonel, P.; et al. Ebola virus glycoprotein with increased infectivity dominated the
987   2013-2016 epidemic. *Cell* **2016**, *167*, 1088–1098.e6.

988   16.   Urbanowicz, R.A.; Patrick McClure, C.; Sakuntabhai, A.; Sall, A.A.; Kobinger, G.; Müller, M.A.; Holmes,
989   E.C.; Rey, F.A.; Simon-Loriere, E.; Ball, J.K. Human adaptation of Ebola virus during the West African
990   outbreak. *Cell* **2016**, *167*, 1079–1087.e5.

991   17.   Dietzel, E.; Schudt, G.; Krähling, V.; Matrosovich, M.; Becker, S. Functional characterization of adaptive
992   mutations during the West African Ebola virus outbreak. *J. Virol.* **2017**, *91*, e01913-16.

993   18.   Ueda, M.T.; Kurosaki, Y.; Izumi, T.; Nakano, Y.; Oloniniyi, O.K.; Yasuda, J.; Koyanagi, Y.; Sato, K.;
994   Nakagawa, S. Functional mutations in spike glycoprotein of Zaire ebolavirus associated with an increase
995   in infection efficiency. *Genes Cells* **2017**, *22*, 148–159.

996   19.   Hoffmann, M.; Crone, L.; Dietzel, E.; Paijo, J.; González-Hernández, M.; Nehlmeier, I.; Kalinke, U.; Becker,
997   S.; Pöhlmann, S. A polymorphism within the internal fusion loop of the Ebola virus glycoprotein
998   modulates host cell entry. *J. Virol.* **2017**, *91*, e00177-17.

20.   Wang, M.K.; Lim, S.-Y.; Lee, S.M.; Cunningham, J.M. Biochemical basis for increased activity of Ebola
glycoprotein in the 2013-16 epidemic. *Cell Host Microbe* **2017**, *21*, 367–375.

21.   Marzi, A.; Chadinah, S.; Haddock, E.; Feldmann, F.; Arndt, N.; Martellaro, C.; Scott, D.P.; Hanley, P.W.;
Nyenswah, T.G.; Sow, S.; et al. Recently identified mutations in the Ebola virus-Makona genome do not
alter pathogenicity in animal models. *Cell Rep.* **2018**, *23*, 1806–1816.

22.   Wong, G.; He, S.; Leung, A.; Cao, W.; Bi, Y.; Zhang, Z.; Zhu, W.; Wang, L.; Zhao, Y.; Cheng, K.; et al.
Naturally-occurring single mutations in Ebola observably impact infectivity. *J. Virol.* **2018**, *93*, e01098-18.

23.   Dong, S.; Yang, P.; Li, G.; Liu, B.; Wang, W.; Liu, X.; Xia, B.; Yang, C.; Lou, Z.; Guo, Y.; et al. Insight into
the Ebola virus nucleocapsid assembly mechanism: crystal structure of Ebola virus nucleoprotein core
domain at 1.8 Å resolution. *Protein Cell* **2015**, *6*, 351–362.

24.   Leung, D.W.; Borek, D.; Luthra, P.; Binning, J.M.; Anantpadma, M.; Liu, G.; Harvey, I.B.; Su, Z.;
Endlich-Frazier, A.; Pan, J.; et al. An intrinsically disordered peptide from Ebola virus VP35 controls Viral
RNA synthesis by modulating nucleoprotein-RNA interactions. *Cell Rep.* **2015**, *11*, 376–389.

25.   Kirchdoerfer, R.N.; Abelson, D.M.; Li, S.; Wood, M.R.; Saphire, E.O. Assembly of the Ebola virus
nucleoprotein from a chaperoned VP35 complex. *Cell Rep.* **2015**, *12*, 140–149.

26.   Wan, W.; Kolesnikova, L.; Clarke, M.; Koehler, A.; Noda, T.; Becker, S.; Briggs, J.A.G. Structure and
assembly of the Ebola virus nucleocapsid. *Nature* **2017**, *551*, 394–397.

27.   Su, Z.; Wu, C.; Shi, L.; Luthra, P.; Pintilie, G.D.; Johnson, B.; Porter, J.R.; Ge, P.; Chen, M.; Liu, G.; et al.
Electron Cryo-microscopy Structure of Ebola virus nucleoprotein reveals a mechanism for
nucleocapsid-like assembly. *Cell* **2018**, *172*, 966–978.e12.

28.   Sugita, Y.; Matsunami, H.; Kawaoka, Y.; Noda, T.; Wolf, M. Cryo-EM structure of the Ebola virus
nucleoprotein–RNA complex at 3.6 Å resolution. *Nature* **2018**, *563*, 137–140.

29. Kirchdoerfer, R.N.; Saphire, E.O.; Ward, A.B. Cryo-EM structure of the Ebola virus nucleoprotein-RNA complex. *Acta Crystallogr. Sect. F Struct. Biol. Cryst. Commun.* **2019**, *75*, 340–347.

30. Kirchdoerfer, R.N.; Saphire, E.O.; Ward, A.B. Structure of the Ebola virus nucleoprotein - RNA complex.

31. Jasenosky, L.D.; Kawaoka, Y. Filovirus budding. *Virus Res.* **2004**, *106*, 181–188.

32. Harty, R.N.; Brown, M.E.; Wang, G.; Huibregtse, J.; Hayes, F.P. A PPxY motif within the VP40 protein of Ebola virus interacts physically and functionally with a ubiquitin ligase: implications for filovirus budding. *Proc. Natl. Acad. Sci. U. S. A.* **2000**, *97*, 13871–13876.

33. Jasenosky, L.D.; Neumann, G.; Lukashevich, I.; Kawaoka, Y. Ebola virus VP40-induced particle formation and association with the lipid bilayer. *J. Virol.* **2001**, *75*, 5205–5214.

34. Noda, T.; Sagara, H.; Suzuki, E.; Takada, A.; Kida, H.; Kawaoka, Y. Ebola virus VP40 drives the formation of virus-like filamentous particles along with GP. *J. Virol.* **2002**, *76*, 4855–4865.

35. Licata, J.M.; Johnson, R.F.; Han, Z.; Harty, R.N. Contribution of Ebola virus glycoprotein, nucleoprotein, and VP24 to budding of VP40 virus-like particles. *J. Virol.* **2004**, *78*, 7344–7351.

36. Groseth, A.; Charton, J.E.; Sauerborn, M.; Feldmann, F.; Jones, S.M.; Hoenen, T.; Feldmann, H. The Ebola virus ribonucleoprotein complex: A novel VP30–L interaction identified. *Virus Res.* **2009**, *140*, 8–14.

37. Kirchdoerfer, R.N.; Moyer, C.L.; Abelson, D.M.; Saphire, E.O. The Ebola virus VP30-NP interaction is a regulator of viral RNA synthesis. *PLoS Pathog.* **2016**, *12*, e1005937.

38. Volchkov, V.E.; Volchkova, V.A.; Mühlberger, E.; Kolesnikova, L.V.; Weik, M.; Dolnik, O.; Klenk, H.-D. Recovery of infectious Ebola virus from complementary DNA: RNA editing of the GP gene and viral cytotoxicity. *Science* **2001**, *291*, 1965–1969.

39. Neumann, G.; Feldmann, H.; Watanabe, S.; Lukashevich, I.; Kawaoka, Y. Reverse genetics demonstrates that proteolytic processing of the Ebola virus glycoprotein is not essential for replication in cell culture. *J. Virol.* **2002**, *76*, 406–410.

40. Towner, J.S.; Paragas, J.; Dover, J.E.; Gupta, M.; Goldsmith, C.S.; Huggins, J.W.; Nichol, S.T. Generation of eGFP expressing recombinant Zaire ebolavirus for analysis of early pathogenesis events and high-throughput antiviral drug screening. *Virology* **2005**, *332*, 20–27.

41. Hoenen, T.; Groseth, A.; Callison, J.; Takada, A.; Feldmann, H. A novel Ebola virus expressing luciferase allows for rapid and quantitative testing of antivirals. *Antiviral Res.* **2013**, *99*, 207–213.

42. Pickett, B.E.; Sadat, E.L.; Zhang, Y.; Noronha, J.M.; Squires, R.B.; Hunt, V.; Liu, M.; Kumar, S.; Zaremba, S.; Gu, Z.; et al. ViPR: an open bioinformatics database and analysis resource for virology research. *Nucleic Acids Res.* **2012**, *40*, D593–8.

43. Katoh, K.; Standley, D.M. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol. Biol. Evol.* **2013**, *30*, 772–780.

44. Capella-Gutiérrez, S.; Silla-Martínez, J.M.; Gabaldón, T. trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **2009**, *25*, 1972–1973.

45. Stamatakis, A.; Ludwig, T.; Meier, H. RAxML-III: a fast program for maximum likelihood-based inference of large phylogenetic trees. *Bioinformatics* **2005**, *21*, 456–463.

46. Schrödinger, LLC The PyMOL molecular graphics system, version 2.0. Available online: https://pymol.org/2 (accessed on 5 October 2017).

47. Warren, L.; Manos, P.D.; Ahfeldt, T.; Loh, Y.-H.; Li, H.; Lau, F.; Ebina, W.; Mandal, P.K.; Smith, Z.D.; Meissner, A.; et al. Highly efficient reprogramming to pluripotency and directed differentiation of human cells with synthetic modified mRNA. *Cell Stem Cell* **2010**, *7*, 618–630.

48. Kim, J.H.; Lee, S.-R.; Li, L.-H.; Park, H.-J.; Park, J.-H.; Lee, K.Y.; Kim, M.-K.; Shin, B.A.; Choi, S.-Y. High cleavage efficiency of a 2A peptide derived from porcine teschovirus-1 in human cell lines, zebrafish and mice. *PLoS One* **2011**, *6*, e18556.

49. Subach, O.M.; Cranfill, P.J.; Davidson, M.W.; Verkhusha, V.V. An enhanced monomeric blue fluorescent protein with the high chemical stability of the chromophore. *PLoS One* **2011**, *6*, e28674.

50. Manicassamy, B.; Rong, L. Expression of ebolavirus glycoprotein on the target cells enhances viral entry. *Virol. J.* **2009**, *6*, 75.

51. Bornholdt, Z.A.; Noda, T.; Abelson, D.M.; Halfmann, P.; Wood, M.R.; Kawaoka, Y.; Saphire, E.O. Structural rearrangement of Ebola virus VP40 begets multiple functions in the virus life cycle. *Cell* **2013**, *154*, 763–774.

52. Jasenosky, L.D.; Neumann, G.; Kawaoka, Y. Minigenome-based reporter system suitable for high-throughput screening of compounds able to inhibit ebolavirus replication and/or transcription. *Antimicrob. Agents Chemother.* **2010**, *54*, 3007–3010.

53. Watanabe, S.; Noda, T.; Kawaoka, Y. Functional mapping of the nucleoprotein of Ebola virus. *J. Virol.* **2006**, *80*, 3743–3751.

54. Lin, A.E.; Greco, T.M.; Döhner, K.; Sodeik, B.; Cristea, I.M. A proteomic perspective of inbuilt viral protein regulation: pUL46 tegument protein is targeted for degradation by ICP0 during herpes simplex virus type 1 infection. *Mol. Cell. Proteomics* **2013**, *12*, 3237–3252.

55. Gao, G.; Hendricks, G.M. Introducing genes into mammalian cells: viral vectors. Protocol 14: analysis of rAAv sample morphology using negative staining and high resolution electron microscopy. In *Molecular Cloning (A laboratory Manual)*; Green, M.R., Sambrook, J., Eds.; Cold Spring Harbor Press, 2012; Vol. 2, pp. 1301–1303 ISBN 9781936113422.

56. Schneider, C.A.; Rasband, W.S.; Eliceiri, K.W. NIH Image to ImageJ: 25 years of image analysis. *Nat. Methods* **2012**, *9*, 671–675.

57. Watt, A.; Moukambi, F.; Banadyga, L.; Groseth, A.; Callison, J.; Herwig, A.; Ebihara, H.; Feldmann, H.; Hoenen, T. A novel life cycle modeling system for Ebola virus shows a genome length-dependent role of VP24 in virus infectivity. *J. Virol.* **2014**, *88*, 10511–10524.

58. Watanabe, S.; Watanabe, T.; Noda, T.; Takada, A.; Feldmann, H.; Jasenosky, L.D.; Kawaoka, Y. Production of novel Ebola virus-like particles from cDNAs: an alternative to Ebola virus generation by reverse genetics. *J. Virol.* **2004**, *78*, 999–1005.

59. Shevchenko, A.; Wilm, M.; Vorm, O.; Mann, M. Mass spectrometric sequencing of proteins silver-stained polyacrylamide gels. *Anal. Chem.* **1996**, *68*, 850–858.

60. Peng, J.; Gygi, S.P. Proteomics: the move to mixtures. *J. Mass Spectrom.* **2001**, *36*, 1083–1091.

61. Eng, J.K.; McCormack, A.L.; Yates, J.R. An approach to correlate tandem mass spectral data of peptides with amino acid sequences in a protein database. *J. Am. Soc. Mass Spectrom.* **1994**, *5*, 976–989.

62. Mellacheruvu, D.; Wright, Z.; Couzens, A.L.; Lambert, J.-P.; St-Denis, N.A.; Li, T.; Miteva, Y.V.; Hauri, S.; Sardiu, M.E.; Low, T.Y.; et al. The CRAPome: a contaminant repository for affinity purification-mass spectrometry data. *Nat. Methods* **2013**, *10*, 730–736.

63. Snel, B. STRING: a web-server to retrieve and display the repeatedly occurring neighbourhood of a gene. *Nucleic Acids Res.* **2000**, *28*, 3442–3444.

64. Szklarczyk, D.; Franceschini, A.; Wyder, S.; Forslund, K.; Heller, D.; Huerta-Cepas, J.; Simonovic, M.; Roth, A.; Santos, A.; Tsafou, K.P.; et al. STRING v10: protein-protein interaction networks, integrated over the tree of life. *Nucleic Acids Res.* **2015**, *43*, D447–52.

65. Shannon, P.; Markiel, A.; Ozier, O.; Baliga, N.S.; Wang, J.T.; Ramage, D.; Amin, N.; Schwikowski, B.; Ideker, T. Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res.* **2003**, *13*, 2498–2504.

66. R Core Team R: A language and environment for statistical computing. *R Foundation for Statistical Computing, Vienna, Austria* **2016**. Available online: https://www.R-project.org.

67. Baty, F.; Ritz, C.; Charles, S.; Brutsche, M.; Flandrois, J.-P.; Delignette-Muller, M.-L. A toolbox for nonlinear regression in R: the package nlstools. *J. Stat. Softw.* **2015**, *66*.

68. Wickham, H. *ggplot2: Elegant Graphics for Data Analysis*; Springer, 2016; ISBN 9783319242774.

69. Ng, A.K.-L.; Lam, M.K.-H.; Zhang, H.; Liu, J.; Au, S.W.-N.; Chan, P.K.-S.; Wang, J.; Shaw, P.-C. Structural basis for RNA binding and homo-oligomer formation by influenza B virus nucleoprotein. *J. Virol.* **2012**, *86*, 6758–6767.

70. Ortiz-Riano, E.; Cheng, B.Y.H.; de la Torre, J.C.; Martinez-Sobrido, L. Self-association of lymphocytic choriomeningitis virus nucleoprotein is mediated by its N-terminal region and is not required for its anti-interferon function. *J. Virol.* **2012**, *86*, 3307–3317.

71. Machleidt, T.; Woodroofe, C.C.; Schwinn, M.K.; Méndez, J.; Robers, M.B.; Zimmerman, K.; Otto, P.; Daniels, D.L.; Kirkland, T.A.; Wood, K.V. NanoBRET--a novel BRET platform for the analysis of protein-protein interactions. *ACS Chem. Biol.* **2015**, *10*, 1797–1804.

72. McCarthy, S.E.; Licata, J.M.; Harty, R.N. A luciferase-based budding assay for Ebola virus. *J. Virol. Methods* **2006**, *137*, 115–119.

73. Liu, Y.; Cocka, L.; Okumura, A.; Zhang, Y.-A.; Sunyer, J.O.; Harty, R.N. Conserved motifs within Ebola and Marburg virus VP40 proteins are important for stability, localization, and subsequent budding of virus-like particles. *J. Virol.* **2010**, *84*, 2294–2303.

74. Hall, M.P.; Unch, J.; Binkowski, B.F.; Valley, M.P.; Butler, B.L.; Wood, M.G.; Otto, P.; Zimmerman, K.; Vidugiris, G.; Machleidt, T.; et al. Engineered luciferase reporter from a deep sea shrimp utilizing a novel imidazopyrazinone substrate. *ACS Chem. Biol.* **2012**, *7*, 1848–1857.

75. Reynard, O.; Reid, S.P.; Page, A.; Mateo, M.; Alazard-Dany, N.; Raoul, H.; Basler, C.F.; Volchkov, V.E. Unconventional secretion of Ebola virus matrix protein VP40. *J. Infect. Dis.* **2011**, *204 Suppl 3*, S833–9.

76. Pleet, M.L.; Mathiesen, A.; DeMarino, C.; Akpamagbo, Y.A.; Barclay, R.A.; Schwab, A.; Iordanskiy, S.; Sampey, G.C.; Lepene, B.; Nekhai, S.; et al. Ebola VP40 in exosomes can cause immune cell dysfunction. *Front. Microbiol.* **2016**, *7*, 1765.

77. Hu, L.; Trefethen, J.M.; Zeng, Y.; Yee, L.; Ohtake, S.; Lechuga-Ballesteros, D.; Warfield, K.L.; Aman, M.J.; Shulenin, S.; Unfer, R.; et al. Biophysical characterization and conformational stability of Ebola and Marburg virus-like particles. *J. Pharm. Sci.* **2011**, *100*, 5156–5173.

78. Hoenen, T.; Jung, S.; Herwig, A.; Groseth, A.; Becker, S. Both matrix proteins of Ebola virus contribute to the regulation of viral genome replication and transcription. *Virology* **2010**, *403*, 56–66.

79. García-Dorival, I.; Wu, W.; Armstrong, S.D.; Barr, J.N.; Carroll, M.W.; Hewson, R.; Hiscox, J.A. Elucidation of the cellular interactome of Ebola virus nucleoprotein and identification of therapeutic targets. *J. Proteome Res.* **2016**, *15*, 4290–4303.

80. Goldstein, T.; Anthony, S.J.; Gbakima, A.; Bird, B.H.; Bangura, J.; Tremeau-Bravard, A.; Belaganahalli, M.N.; Wells, H.L.; Dhanota, J.K.; Liang, E.; et al. The discovery of Bombali virus adds further support for bats as hosts of ebolaviruses. *Nature Microbiology* **2018**, *3*, 1084–1089.

81. Landeras-Bueno, S.; Oda, S.-I.; Norris, M.J.; Li Salie, Z.; Guenaga, J.; Wyatt, R.T.; Saphire, E.O. Sudan ebolavirus VP35-NP crystal structure reveals a potential target for pan-filovirus treatment. *MBio* **2019**, *10*, e00734-19.

82. Camus, G.; Segura-Morales, C.; Molle, D.; Lopez-Vergès, S.; Begon-Pescia, C.; Cazevieille, C.; Schu, P.; Bertrand, E.; Berlioz-Torrent, C.; Basyuk, E. The clathrin adaptor complex AP-1 binds HIV-1 and MLV Gag and facilitates their budding. *Mol. Biol. Cell* **2007**, *18*, 3193–3203.