1 **The convergent evolution of eusociality is based on a shared reproductive**
2 **groundplan plus lineage-specific sets of plastic genes**
3
4
5

6 Michael R. Warner[1], Lijun Qiu[2], Michael J. Holmes[2,3], Alexander S. Mikheyev[2,4], Timothy A.
7 Linksvayer[1]
8
9 [1]University of Pennsylvania, Philadelphia, Pennsylvania, USA
10 [2]Okinawa Institute of Science and Technology, Okinawa, Japan
11 [3]School of Life and Environmental Science, University of Sydney, Australia
12 [4]Research School of Biology, Australian National University, Canberra, Australia

13
14

Abstract

Eusociality has convergently evolved multiple times, but the genomic basis of caste-based

division of labor and degree to which independent origins of eusociality have utilized common

genes remains largely unknown. We characterized caste-specific transcriptomic profiles across

development and adult body segments from pharaoh ants (*Monomorium pharaonis*) and honey

bees (*Apis mellifera*), representing two independent origins of eusociality. We identified a

substantial shared core of genes upregulated in the abdomens of queen ants and honey bees that

also tends to be upregulated in mated female flies, suggesting that these genes are part of a

conserved insect reproductive groundplan. Outside of this shared groundplan, few genes are

differentially expressed in common. Instead, the majority of the thousands of caste-associated

genes are plastically-expressed, rapidly evolving, and relatively evolutionarily young. These

results emphasize that the recruitment of both highly conserved and lineage-specific genes

underlie the convergent evolution of novel traits such as eusociality.

Main Text

The degree to which convergent phenotypic evolution involves the same sets of genes or pathways is a major unanswered question[1]. Comparative genomic studies indicate that parallel adaptive changes in the protein-coding sequences of the same genes are frequently associated with the evolution of convergent phenotypes in closely related populations and species[2,3]. Decades of research in evolutionary developmental biology also emphasize that changes in the expression of a relatively small "toolkit" of deeply conserved genes are often associated with convergently evolved phenotypes in distantly related species[4]. Alternatively, convergent phenotypic evolution between lineages could involve distinct subsets of genes in each lineage, including taxonomically-restricted genes, genes which have no detectable orthology outside of a given lineage[5]. Taxonomically-restricted genes have been shown to be important for lineage-specific evolutionary novelties[6], but their relative contribution to the evolution of convergent phenotypes is unknown.

The evolution of eusociality in several insect lineages (e.g., ants, honey bees, vespid wasps, termites) provides a striking example of convergent phenotypic innovation[7]. Eusocial insect societies are founded upon a novel caste polyphenism, in which reproductive queen and non-reproductive worker female castes develop from the same genome, depending mainly on socially-regulated nutritional inputs[8,9]. Within the worker caste, further specialization often occurs as individuals age and progress through a series of tasks, including nursing and foraging[7].

Polyphenic traits are often thought to evolve from pre-existing developmental plasticity[10]. Leading hypotheses for the evolution of caste-based division of labor in social insects also stress the use and modification of highly conserved developmental and physiological mechanisms[11–15].

For example, the reproductive and non-reproductive phases of ancestral solitary insects are

thought to have been decoupled to produce reproductive and non-reproductive castes[11,16], and

worker division of labor is similarly thought to be derived from the decoupling of the ancestral

reproductive cycle[13,16,17]. Along the same lines, it has been suggested that the convergent

evolution of novel social behavior involves changes to the regulation of a core "toolkit" of genes

underlying highly conserved physiological processes such as metabolism[14,15].

Studies focused on candidate genes underlying the genetic basis of caste-based division

of labor within individual eusocial species have often found support for the importance of highly

conserved genes and pathways associated with reproduction and metabolism. For example,

worker division of labor in honey bees is regulated by interactions between juvenile hormone,

vitellogenin, and insulin/TOR signaling pathways[13,17,18]. Similar pathways also play key roles in

regulating division of labor between queen and worker castes in both ants and honey bees,

though the mechanistic details vary[19–22]. While comparative genomic and transcriptomic studies

have often similarly emphasized common general functions such as metabolism, such studies

have thus far only identified very small sets of specific genes associated with the convergent

evolution of caste, worker behavior, or eusociality in independent lineages[15,23–27]. Alternatively,

many transcriptomic studies have argued for the importance of taxonomically-restricted genes

for the evolution of caste-based division of labor[28–34]. It is unclear if the lack of common specific

genes is due to biological differences between the species or methodological details because

studies in each species were not designed, conducted, or analyzed in parallel.

Previous work has mainly focused on identifying whether there is significant overlap of

genes or gene pathways associated with caste-based division of labor between independent

lineages[23,26,27], but there has been little effort to quantify the relative importance of shared versus

unshared genes to the convergent evolution of caste-based division of labor. Most of these

studies have either focused on brain or whole body samples[15,22,24,26,33,35–37], although expression

bias between queens and workers has been shown to be dependent upon developmental stage and

tissue type[34,38–40]. Finally, the transcriptomic signatures of reproductive physiology are strongest

in the abdomen[34,41], the location of reproductive organs, but no past study has explicitly

compared caste bias in abdominal tissues in species from lineages representing independent

origins of eusociality.

Here, we present to date the most comprehensive developmental transcriptomic dataset

investigating gene expression associated with reproductive caste and age-based worker division

of labor in the pharaoh ant (*Monomorium pharaonis*) and the honey bee (*Apis mellifera*). We

chose these two study species because they represent two independent origins of eusociality in

the ant and corbiculate bee lineages[42] as well as two independent elaborations of eusociality,

each characterized by strong queen-worker dimorphism and age-based worker division of

labor[33,43]. We performed all sampling, sequencing, and analysis for the two species in parallel to

maximize compatibility between the data sets. We leverage this extensive dataset to quantify in

an unbiased manner the relative contribution of differential expression of shared versus distinct

genes at each life stage and tissue to the convergent evolution of caste-based division of labor.

Results

We constructed two large, parallel transcriptomic datasets in honey bees and pharaoh ants

spanning caste development as well as adult tissues separated by reproductive caste (queens

versus workers), behavior (nurse workers versus forager workers), and sex (queens and workers

versus males). In total we constructed 177 mRNA-sequencing libraries across 28 distinct sample

types for each species (Supplementary Table 1).

Differential expression between queens and workers

To identify genes associated with caste development and adult caste dimorphism, we performed

differential expression analysis between queens and workers at each developmental stage and

adult tissue, separately for each species. The number of differentially expressed genes (DEGs)

between queens and workers increased throughout development, peaking in the adult abdomen

(Fig. 1a). In all tissues and stages, the majority of caste-associated DEGs in one species were

either not differentially expressed or did not have an ortholog in the other species (Fig. 1a;

Supplementary Fig. 1a; Supplementary Table 2). The magnitude of gene-wise caste bias (as

measured by $\log_2$ fold-change between queen and worker samples) was weakly positively

correlated between ant and honey bee orthologs in all three adult tissues, with the strongest

correlation in the abdomen, but uncorrelated or negatively correlated in all larval and pupal

stages (Supplementary Fig. 2; $r_{head} = 0.089$; $r_{thorax} = 0.161$; $r_{abdomen} = 0.275$; N = 7460 1:1

orthologs; $P < 0.001$ in all cases). The top enriched Gene Ontology (GO) terms for caste-

associated DEGs in each species were dominated by metabolism, signaling and developmental

processes (Supplementary Tables 3-4).

Differential expression between nurses and foragers

Both honey bees[43] and pharaoh ants[33] exhibit age-based worker division of labor, in which

younger individuals tend to specialize on nursing and other within-nest activities and older

individuals specialize on foraging. To identify genes associated with age-based worker division

of labor, we performed differential expression analysis between nurses and foragers in each adult

tissue, separately for each species. In general, there were very few behavioral DEGs shared

between the two species (Fig. 1b; Supplementary Fig. 1b; Supplementary Table 5). Gene-wise

$\log_2$ fold-change between nurses and foragers was significantly but weakly correlated across ant

and honey bee orthologs (Supplementary Fig. 3; $r_{head} = 0.070$, $P_{head} < 0.001$; $r_{thorax} = 0.031$, P_{thorax}

$= 0.008$; $r_{abdomen} = 0.051$, $P_{abdomen} < 0.001$; N = 7460 1:1 orthologs). The top enriched GO terms

for behavioral DEGs in each species were dominated by metabolism and developmental

processes (Supplementary Tables 6-7).

Shared abdominal caste-bias in ancient genes

For the most part, our results indicate distinct genes are associated with caste and worker

division of labor in honey bees and ants. However, approximately one third of abdominal caste-

associated DEGs were common to both species (Fig. 1a; 1545 shared DEGs, comprising 35%

[1545/4395] of ant DEGs, and 29% [1545/5352] of honey bee DEGs). Most shared abdominal

differential expression was the result of shared queen-bias: 56% (858/1545 genes) of shared

abdominal caste-associated DEGs were upregulated in queen abdomens in both species,

compared to 22% (338/1545) that were worker-upregulated in both species and 23% (349/1545)

that reversed direction (i.e. were queen-biased in one species and worker-biased in the other).

Shared abdominal caste-associated DEGs were more likely to be identified as evolutionarily

ancient in comparison to non-biased genes (Fig. 1c; Supplementary Fig. 1c; Fisher Test; F =

3.41, P < 0.001). Furthermore, abdominal DEGs with shared queen bias were more likely to be

identified as ancient than DEGs with shared worker bias (Fig. 1c; Fisher Test; F = 2.51, P <

0.001). In general, the evolutionary age of genes was associated with expression bias between

castes, though the direction of the effect was not consistent across all tissues and stages

(Supplementary Fig. 4).

We next tried to put the seemingly large proportion of shared abdominal caste-associated

DEGs (35% for ants and 29% for honey bees) into context. We compared the proportion of

genes that were differentially expressed across embryonic and larval development in both

species, given that the molecular mechanisms of development are thought to be highly

conserved[44]. We identified 6089 and 6225 developmental DEGs in ants and honey bees,

respectively, including 2544 shared DEGs, representing 42% (2544/6089) and 41% (2544/6255)

of the total developmental DEGs in each species (Supplementary Fig. 5).

To identify which of the thousands of abdominal DEGs found in each species are

particularly important for queen abdominal expression (and presumably function), we performed

gene co-expression analysis, separately for each species. We focused on modules specifically

associated with queens because the majority of shared DEGs were queen-upregulated. We

identified a module of genes specifically associated with queen abdominal expression in each

species (N = 1006 genes in the module for ants, N = 1174 genes for honey bees). We identified

hub genes in each module (N = 92 genes in ants, N = 94 genes in honey bees), genes which are

centrally connected in networks and strongly associated with queen abdominal expression[45].

Many annotated hub genes are inferred to have functions associated with reproduction and

maternal effects (Supplementary Tables 8-9), including genes with known roles in caste

determination such as vitellogenin (*Vg* receptor was identified in each species)[20] and *vasa*[46],

while others are important maternal proteins such as *Smaug*[47] and *ovo*[48]. Furthermore, genes for

which *Drosophila melanogaster* orthologs are known to function in oogenesis (based on FlyBase

Gene Ontology[49]) were more highly connected within the queen abdominal modules than genes

not associated with oogenesis (Supplementary Fig. 6) for honey bees (Wilcoxon test; N = 649; P

< 0.001), though not for ants (N = 542; P = 0.114). Finally, we identified 181 genes which were

present in the queen abdominal module of both species. These genes tended to be queen-biased

(78.5% [142/181] upregulated in queens of both species) and were more centrally-located within

modules than genes found in only one species-specific module (Fig. 2c-d).

Caste bias is in part derived from ancestral sex bias

Given that our co-expression analysis indicated that many important queen-upregulated genes

are associated with oogenesis and overall female reproduction, we reasoned that caste-biased

expression would be linked to sex-biased expression (i.e. expression differences between

reproductive females and males). Indeed, there was a positive correlation between gene-wise $\log_2$

fold change between queen and worker abdomens and gene-wise $\log_2$ fold-change between

queen and male abdomens in both honey bees and pharaoh ants (Fig. 3a-b). Additionally, sex

bias itself was correlated between species (Fig. 3c). The correlation of caste bias and sex bias

was not restricted to the abdomen, as there were similar highly significant effects when

comparing transcriptomic profiles in head and thoracic tissues, albeit with weaker effect sizes

(Supplementary Fig. 7).

Given the association between shared caste bias and sex bias within pharaoh ants and

honey bees, we hypothesized that these shared caste-biased genes were derived from conserved

pathways that also underlie sexual dimorphism for reproductive physiology in distant relatives.

To test this hypothesis, we estimated sex-biased expression of orthologs in the fruit fly

Drosophila melanogaster using available data from male and mated female whole bodies[44].

Shared queen-biased abdominal DEGs tended to be upregulated in females in *D. melanogaster*

(Fig. 3d; one-sided Binomial Test for likelihood of shared queen-biased DEGs having $\log_2$ fold-

change > 0; P < 0.001; N = 566 shared queen DEGs), while shared worker-biased abdominal

DEGs tended to be upregulated in males (P < 0.001; N = 160 shared worker DEGs), indicative of

shared queen (social insects) and female (fly) down-regulation. Though we detected few shared

caste-associated DEGs in the head and thorax (N = 38 and N = 64, respectively), these DEGs

showed the same pattern, where orthologs of queen-biased DEGs were significantly more

female-biased in *D. melanogaster* than orthologs of worker-biased DEGs (Supplementary Fig.

8).

Expression plasticity across development, caste, and tissue is correlated between species

While we have emphasized the conservation of abdominal differential expression between

queens and workers in pharaoh ants and honey bees, differential expression based on either

reproductive caste or worker division of labor was largely not shared between species (Fig. 1).

Furthermore, genes were often differentially expressed across many stages and tissues,

sometimes in opposite directions (Supplementary Fig. 9; e.g., upregulated in queen heads but

downregulated in queen abdomens). To quantify the degree to which genes exhibited biased

expression according to reproductive caste across all developmental stages and tissues, we

calculated gene-wise "overall caste bias" in each species, where we defined overall caste bias as

the Euclidean distance of $\log_2$ fold-change across all queen/worker comparisons[49]. Similarly, we

defined "overall behavior bias" as the Euclidean distance of $\log_2$ fold-change across all

nurse/forager comparisons, separately for each species.

Across 1:1 orthologs, overall caste bias measured in ants was correlated to overall caste

bias measured in honey bees (Supplementary Fig. 10a; rho = 0.454, P < 0.001), and overall

behavior bias was similarly correlated between species (Supplementary Fig. 10b; rho = 0.221, P

< 0.001). Within species, overall caste and behavior bias were also correlated to each other

(Supplementary Fig. 11; ants: rho = 0.549, P < 0.001; honey bees: rho = 0.642, P < 0.001). This

indicates that plasticity in gene expression is correlated across contexts (caste versus behavior)

and species. GO terms associated with high overall caste bias were largely linked to metabolism,

while those associated with high overall behavior bias were largely linked to developmental

processes (Supplementary Table 10).

Characteristics of genes associated with caste and behavior

We compared overall caste bias and overall behavior bias to gene age, evolutionary rate, network

connectivity, and tissue-specificity to understand the general features of genes commonly

associated with caste (queen versus worker) or behavior (nursing versus foraging). Genes with

younger estimated evolutionary ages tended to exhibit higher overall caste bias (Fig. 4a-b) and

behavior bias (Supplementary Fig. 12a-b) compared in particular to ancient genes (Gamma

GLM; ant caste bias: $\chi^2 = 900.19$, honey bee caste bias: $\chi^2 = 1412.80$, ant behavior bias: $\chi^2 = $

316.36, honey bee behavior bias: $\chi^2 = 877.43$; $P < 0.001$ for all cases; N = 10520 in ant, N =

10011 in honey bees). Genes that were loosely connected (representing peripheral network

elements) in co-expression networks constructed across all samples tended to exhibit more caste

and behavior bias in comparison to highly connected genes (Fig. 4c-d; Supplementary Fig. 12c-

d). Similarly, genes with high tissue-specificity across 12 honey bee tissues tended to exhibit

higher values of caste and behavior bias in honey bees compared to more pleiotropic,

ubiquitously expressed genes (Supplementary Fig. 13), where tissue specificity was calculated

using available data[32]. Finally, genes that were rapidly evolving (as estimated by dN/dS) tended

to exhibit higher levels of caste and behavior bias (Fig. 4e-f; Supplementary Fig. 12e-f).

Importantly, while expression is correlated to overall caste and behavior bias, these results

remain highly significant when expression level is controlled for in partial correlation analyses

(Supplementary Table 11).

Discussion

Caste-based division of labor within social insect colonies is hypothesized to be derived from

conserved pathways regulating reproduction[11,13,16,17]. In this study, we identified a large set

(~1500) of genes with shared caste-biased abdominal expression in pharaoh ants and honey bees

(Fig. 1a), including many annotated genes with known roles in reproduction such as the

vitellogenin receptor[20] and *ovo*[48]. Our results are consistent with the notion that caste-biased

genes are derived from ancient plastically-expressed genes underlying female reproduction, as

genes upregulated in queen abdomens of both ants and honey bees tended to also be female-

biased in the distant insect relative *Drosophila melanogaster* (Fig. 3d). Previous studies had

failed to find large sets of genes repeatedly used for eusocial evolution[23–27], but no previous

comparative study investigated caste-biased expression in the abdomen.

The large overlap for abdominal caste-associated genes is notable because honey bees

and ants last shared a common ancestor approximately 160 million years ago[50], and this overlap

is nearly as much as we see for genes that were differentially expressed across developmental

stages (Supplementary Fig. 5). Shared developmental molecular mechanisms are presumably due

simply to shared ancestry and the deep conservation of developmental mechanisms[44,51]. The

similar level of overlap for caste-associated genes points to the large-scale recruitment of pre-

existing developmental and physiological machinery during the independent evolution of caste-

based division of labor in ant and honey bee lineages. Additionally, the association between sex-

bias in *D. melanogaster* and shared caste-bias in social insects extended to the head and thorax

(Supplementary Fig. 8), and caste- and sex-bias were correlated within species (Supplementary

Fig. 7). This indicates that although the strongest signature of overlap in caste-biased expression

occurred in the abdomen, the association of caste- and sex-bias is not simply driven by the

presence of ovaries but rather due to shared female reproductive physiology that is largely

conserved across insects.

While reproductive caste in complex eusocial societies such as ants and honey bees is

typically fixed in adulthood, the tasks performed by workers (specifically, nursing versus

foraging) change over the course of the worker's adult lifetime[18,33]. This plastic behavioral

change is known to be accompanied by a wide range of physiological changes and is regulated at

least in part by conserved physiological pathways, for example, those involving insulin

signaling, juvenile hormone, and vitellogenin[18,21]. However, we identified few genes that were

commonly differentially expressed between nurses and foragers in honey bees and pharaoh ants

(Fig. 1b), and the proportion of shared genes was much lower in comparison to genes underlying

abdominal differences between queens and workers. This could reflect a combination of different

roles of nurses and foragers between lineages[7] as well as differences in the precise molecular

relationships between these conserved pathways[20,52,53]. Nonetheless, we did identify a number of

enriched Gene Ontology categories associated with development and metabolism in each species

(Supplementary Tables 6-7), which is consistent with the notion that the transition from nurse to

forager is essentially a developmental process, and that common molecular pathways may

provide the raw genetic material for social evolution[14,26,27].

Conserved factors or pathways clearly play important roles in aspects of caste

development and function as well as the transition from nursing to foraging, but our results and

other studies indicate that the majority of the full transcriptomic architecture associated with

caste and age polyethism is not shared between species[26,29,33,36,37]. This lineage-specific

architecture is comprised of large groups of both orthologous genes with different expression

patterns and taxonomically-restricted genes (Fig. 1a-b). In contrast to the low amount of context-

specific overlap in differential expression, the overall degree of caste-associated plastic

expression across stages and tissues (overall caste bias) was correlated between species

(Supplementary Fig. 10a-b), and expression plasticity between queens and workers was

correlated to expression plasticity between nurses and foragers (Supplementary Fig. 11). Genes

with high levels of caste or behavior bias tended to exhibit a suite of network and evolutionary

features including being loosely connected in regulatory networks, evolutionarily young, and

rapidly evolving (Fig. 4; Supplementary Fig. 12) as well as displaying tissue-specific expression

profiles (Supplementary Fig. 10) in comparison to more ubiquitously-expressed genes.

These network and evolutionary characteristics have commonly been implicated for

genes underlying eusocial evolution[36,54–56], particularly in association with the worker caste[28–33].

While factors such as evolutionary age and rate to some degree cannot be reliably disentangled[57],

these characteristics together reflect relaxed selection on genes' coding sequences and expression

profile. This may indicate that caste-bias evolves from pre-existing expression plasticity[54]. This

could occur when genes that were previously tightly regulated in another context acquire biased

expression[58], which is possibly reflected in our results by the association between tissue

specificity and caste/behavior bias in honey bees (Supplementary Fig. 10). Alternatively, caste-

biased expression could evolve neutrally, in which genes with loosely regulated expression

patterns acquire caste-biased expression randomly, through neutral or slightly deleterious

fixations in regulatory sequences[56,59]. Our results are consistent with both mechanisms for the

evolution of caste bias: a large part of abdominal caste-bias seems to evolve through the novel

regulation of genes with sex-specific expression plasticity, while the bulk of the genes

underlying caste-based division of labor exhibit characteristics generally reflective of genes

which are weakly constrained.

Conclusions

Our study shows that the recruitment of a large core of conserved reproductive-associated genes,

which can be described as a "reproductive groundplan", is fundamental to the convergent

evolution of caste-based division of labor in ants and honey bees. However, our study also

reveals that the bulk of the full genetic architecture underlying the expression of social insect

caste-based division of labor varies between lineages. This is reflected by the general biology of

social insects, in that independently evolved societies share reproductive division of labor, the

main defining feature of eusociality, but also display a wide diversity of lineage-specific

adaptations[7]. Future studies including more species will be necessary to determine the generality

of the patterns (e.g., the precise numbers of shared and lineage-specific genes) we found. It is

likely that a relatively small number of core conserved genes exist as upstream hubs in

regulatory networks, and layered downstream of this core is a myriad of taxonomically-restricted

genes as well as conserved genes with lineage-specific expression patterns[6,32,33,60]. This is

consistent with models for the evolution of hierarchical developmental gene regulatory networks,

whereby a relatively small number of highly conserved genes act upstream to initiate gene

cascades (e.g., to set up body-patterning), while batteries of downstream genes are evolutionarily

labile and largely responsible for lineage-specific features[61]. Recent studies have made progress

elucidating the function of several core genes and pathways for caste[19,21,22,62]. Large-scale

transcriptomic studies such as ours serve a complimentary, indispensable role of identifying the

full suite of genes underlying caste-based division of labor in multiple independent lineages.

Methods

We collected parallel time series RNA-seq data of caste development in the pharaoh ant

Monomorium pharaonis and the honey bee *Apis mellifera*, including seven developmental stages

(egg, five larval stages, one pupal stage) plus each of three adult body segments (head, thorax,

abdomen) in both species (Supplementary Table 1). We collected three biological replicates of

each specific sample type. Each biological replicate contained a pool of individuals (N = 10 for

ants, N = 5 for honey bees) from the same colony, such that each biological replicate corresponds

to a colony. The only exception to this was mature honey bee queens, which were sampled from

separate unrelated colonies.

Differential Expression Analysis

To identify caste-associated differentially-expressed genes (DEGs), we performed differential

expression analysis between queens and workers at each developmental stage and tissue,

separately for each species. We removed lowly-expressed genes that did not meet one of two

criteria: 1) counts per million (CPM) greater than one in at least half the samples, or 2) CPM > 1

in all samples of a given tissue/stage/caste combination (to ensure tissue-specific genes were

retained). We removed 2350 lowly-expressed genes in ants, leaving 10804 genes for further

analysis, and we removed 2036 genes in bees, leaving 11775 genes for further analysis. We

constructed GLM-like models including replicate and caste and identified genes associated with

caste at each stage or tissue using EdgeR[63]. Similarly, to identify behavioral DEGs we performed

differential expression analyses between nurses and foragers for each tissue. To identify

developmental DEGs in each species, we constructed models with all larval and egg samples and

identified genes differentially expressed between any developmental stage, controlling for

overall caste differences. To estimate gene-wise sex-bias of *D. melanogaster* orthologs, we

downloaded available whole body RNA-seq data[44], consisting of one 5-day old and one 30-day

old fly of each sex, and performed differential expression analysis as above.

Coexpression Network Analysis

We performed plaid clustering, a non-deterministic biclustering algorithm[64]. Biclustering seeks

to identify groups of genes that are co-expressed across a specific subset of samples[65]. For each

species, we constructed networks with all samples to ensure proper resolution of gene-gene co-

expression relationships. We identified genes that were consistently associated with a queen-

abdomen specific bicluster across 1000 iterations, which we term queen abdominal modules (see

Supplemental Methods). We conservatively identified module hub genes as genes with intra-

module connectivity in at least the 90th percentile and abdominal $\log_2$ fold-change values greater

than 2 (representing a 4-fold increase in expression in queen relative to worker abdomens). We

calculated the connectivity of each gene as the weighted sum of the Pearson correlation of

expression between the given gene and all other genes, where we raised each correlation to the

6th power, the default value for weighted gene co-expression analysis[66]. Intra-module

connectivity (used in Fig. 2) represents the connectivity of genes within the queen abdominal

module to other genes within the module, while total network connectivity (used in Fig. 4)

represents the connectivity of genes across the entire transcriptome, after filtering out lowly-

expressed genes.

Estimation of tissue specificity

We downloaded available RNA-sequencing data on twelve tissues in *A. mellifera* worker nurses

and foragers[32]. To classify genes by their tissue specificity, we calculated τ, a commonly used

metric of expression specificity[67]. τ ranges from 0 to 1, where 0 indicates that genes are

ubiquitously expressed and 1 indicates that genes are exclusively expressed in one tissue.

Estimation of evolutionary rate

We estimated evolutionary rate using dN/dS, the ratio of non-synonymous to synonymous

nucleotide changes. We estimated pairwise dN/dS between each focal species and a second

closely related species with an available genome (*A. mellifera*: *A. cerana*; *M. pharaonis*: *S.*

invicta). For each 1:1 ortholog pair, we selected the longest transcript associated with the gene

for each pair of species. We aligned orthologous protein sequences using ClustalW[68], derived

nucleotide alignments from protein alignments using pal2nal[69], and estimated pairwise dN/dS of

nucleotide alignments using PAML, package codeml[70].

Partial correlation analysis

We performed partial Spearman correlations between overall bias and evolutionary/network

characteristics, controlling for the effect of expression.

Code availability

All scripts required to perform all analyses and generate figures and tables are available on

Github: https://github.com/warnerm/devnetwork.

Data availability

All data and scripts required to generate figures, tables, and perform statistical analyses are

available on Github: https://github.com/warnerm/devnetwork. Raw reads are deposited at DNA

Data Bank of Japan.

Acknowledgements

We would like to thank the following: Luigi Pontieri for photographs of pharaoh ants, Alex Wild

for photographs of honey bees, Chao Tong for compiling hymenopteran genomes for use in

phylostratigraphy and for comments on the manuscript, Rohini Singh for comments on the

manuscript, and Junhyong Kim, Mia Levine, and Justin Walsh for helpful comments and

discussion. This work was funded by the National Science Foundation (grant number IOS-

1452520 to TAL) and United States Department of Agriculture National Institute of Food and

Agriculture (grant number 2014-67013-21725 to TAL), and subsidy funding from Okinawa

Institute of Technology to ASM. The National Science Foundation also funded MRW (DGE-

1321851).

References

1. Stern, D. L. The genetic causes of convergent evolution. *Nat. Rev. Genet.* **14**, 751–764

(2013).

2. Conte, G. L., Arnegard, M. E., Peichel, C. L. & Schluter, D. The probability of genetic

parallelism and convergence in natural populations. *Proc. Biol. Sci.* **279**, 5039–5047 (2012).

3. Zhao, L. & Begun, D. J. Genomics of parallel adaptation at two timescales in *Drosophila.*

PLoS Genet. **13**, e1007016 (2017).

4. Shubin, N., Tabin, C. & Carroll, S. Deep homology and the origins of evolutionary

novelty. *Nature* **457**, 818–823 (2009).

5. Wilson, G. A. *et al.* Orphans as taxonomically restricted and ecologically important

genes. *Microbiology* **151**, 2499–2501 (2005).

6. Johnson, B. R. Taxonomically restricted genes are fundamental to biology and evolution.

Front. Genet. **9**, 407 (2018).

7. Wilson, E. O. *The Insect Societies.* (Harvard University Press, 1971).

8. Wheeler, D. E. Developmental and physiological determinants of caste in social

Hymenoptera: evolutionary implications. *Am. Nat.* **128**, 13–34 (1986).

9. Linksvayer, T. A. *et al.* Larval and nurse worker control of developmental plasticity and

the evolution of honey bee queen–worker dimorphism. *J. Evol. Biol.* **24**, 1939–1948 (2011).

10. Nijhout, H. F. Development and evolution of adaptive polyphenisms. *Evol. Dev.* **5**, 9–18

(2003).

11. West-Eberhard, M. J. Flexible strategy and social evolution. In *Animal Societies:*

Theories and Facts (eds Ito, Y., Brown, J. L., Kikkawa, J.), 35-51 (Scientific Societies

Press, 1987).

12. Linksvayer, T. A. & Wade, M. J. The evolutionary origin and elaboration of sociality in

the aculeate Hymenoptera: maternal effects, sib-social effects, and heterochrony. *Q. Rev.*

Biol. **80**, 317–336 (2005).

13. Amdam, G. V., Csondes, A., Fondrk, M. K. & Page, R. E., Jr. Complex social behaviour

derived from maternal reproductive traits. *Nature* **439**, 76–78 (2006).

14. Toth, A. L. & Robinson, G. E. Evo-devo and the evolution of social behavior. *Trends*

Genet. **23**, 334–341 (2007).

15. Toth, A. L. *et al.* Brain transcriptomic analysis in paper wasps identifies genes

associated with behaviour across social insect lineages. *Proc. Biol. Sci.* **277**, 2139–2148

(2010).

16. West-Eberhard, M. J. Wasp societies as microcosms for the study of development and

evolution. In *Natural history and evolution of paper wasps* (eds Turillazzi, S., West-

Eberhard, M. J.), 290-317 (Oxford University Press, 1996).

17. Amdam, G. V., Norberg, K., Fondrk, M. K. & Page, R. E., Jr. Reproductive ground plan

may mediate colony-level selection effects on individual foraging behavior in honey bees.

Proc. Natl. Acad. Sci. U. S. A. **101**, 11350–11355 (2004).

18. Robinson, G. E. Regulation of honey bee age polyethism by juvenile hormone. *Behav.*

Ecol. Sociobiol. **20**, 329–338 (1987).

19. Mutti, N. S. *et al.* IRS and TOR nutrient-signaling pathways act via juvenile hormone to

influence honey bee caste fate. *J. Exp. Biol.* **214**, 3977–3984 (2011).

20. Libbrecht, R. *et al.* Interplay between insulin signaling, juvenile hormone, and

vitellogenin regulates maternal effects on polyphenism in ants. *Proc. Natl. Acad. Sci. U. S.*

A. **110**, 11050–11055 (2013).

21. Gospocic, J. *et al.* The neuropeptide corazonin controls social behavior and caste

identity in ants. *Cell* **170**, 748–759.e12 (2017).

22. Chandra, V. *et al.* Social regulation of insulin signaling and the evolution of eusociality

in ants. *Science* **361**, 398–402 (2018).

23. Woodard, S. H. *et al.* Genes involved in convergent evolution of eusociality in bees.

Proc. Natl. Acad. Sci. U. S. A. **108**, 7472–7477 (2011).

24. Toth, A. L. *et al.* Shared genes related to aggression, rather than chemical

communication, are associated with reproductive dominance in paper wasps (*Polistes*

metricus). *BMC Genomics* **15**, 75 (2014).

25. Woodard, S. H., Bloch, G. M., Band, M. R. & Robinson, G. E. Molecular heterochrony

and the evolution of sociality in bumblebees (*Bombus terrestris*). *Proc. Biol. Sci.* **281**,

20132419 (2014).

26. Berens, A. J., Hunt, J. H. & Toth, A. L. Comparative transcriptomics of convergent

evolution: different genes but conserved pathways underlie caste phenotypes across lineages

of eusocial insects. *Mol. Biol. Evol.* **32**, 690–703 (2015).

27. Kapheim, K. M. *et al.* Genomic signatures of evolutionary transitions from solitary to

group living. *Science* **348**, 1139–1143 (2015).

28. Johnson, B. R. & Tsutsui, N. D. Taxonomically restricted genes are associated with the

evolution of sociality in the honey bee. *BMC Genomics* **12**, 164 (2011).

29. Ferreira, P. G. *et al.* Transcriptome analyses of primitively eusocial wasps reveal novel

insights into the evolution of sociality and the origin of alternative phenotypes. *Genome*

Biol. **14**, R20 (2013).

30. Feldmeyer, B., Elsner, D. & Foitzik, S. Gene expression patterns associated with caste

and reproductive status in ants: worker-specific genes are more derived than queen-specific

ones. *Mol. Ecol.* **23**, 151–161 (2014).

31. Harpur, B. A. *et al.* Population genomics of the honey bee reveals strong signatures of

positive selection on worker traits. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 2614–2619 (2014).

32. Jasper, W. C. *et al.* Large-scale coding sequence change underlies the evolution of

postdevelopmental novelty in honey bees. *Mol. Biol. Evol.* **32**, 334–346 (2015).

33. Mikheyev, A. S. & Linksvayer, T. A. Genes associated with ant social behavior show

distinct transcriptional and evolutionary patterns. *Elife* **4**, e04775 (2015).

34. Warner, M. R., Mikheyev, A. S. & Linksvayer, T. A. Genomic signature of kin selection

in an ant with obligately sterile workers. *Mol. Biol. Evol.* **34**, 1780–1787 (2017).

35. Patalano, S. *et al.* Molecular signatures of plastic phenotypes in two eusocial insect

species with simple societies. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 13970–13975 (2015).

36. Morandin, C. *et al.* Comparative transcriptomics reveals the conserved building blocks

involved in parallel evolution of diverse phenotypic traits in ants. *Genome Biol.* **17**, 43

(2016).

37. Qiu, B. *et al.* Towards reconstructing the ancestral brain gene-network regulating caste

differentiation in ants. *Nat Ecol Evol* **2**, 1782–1791 (2018).

38. Ometto, L., Shoemaker, D., Ross, K. G. & Keller, L. Evolution of gene expression in

fire ants: the effects of developmental stage, caste, and species. *Mol. Biol. Evol.* **28**, 1381–

1392 (2011).

39. Morandin, C. *et al.* Caste-biases in gene expression are specific to developmental stage

in the ant *Formica exsecta*. *J. Evol. Biol.* **28**, 1705–1718 (2015).

40. Lucas, E. R., Romiguier, J. & Keller, L. Gene expression is more strongly influenced by

age than caste in the ant *Lasius niger*. *Mol. Ecol.* **26**, 5058–5073 (2017).

41. Jones, B. M., Kingwell, C. J., Wcislo, W. T. & Robinson, G. E. Caste-biased gene

expression in a facultatively eusocial bee suggests a role for genetic accommodation in the

evolution of eusociality. *Proc. Biol. Sci.* **284**, (2017).

42. Branstetter, M. G. *et al.* Phylogenomic insights into the evolution of stinging wasps and

the origins of ants and bees. *Curr. Biol.* **27**, 1019–1025 (2017).

43. Winston, M. L. *The Biology of the Honey Bee*. (Harvard University Press, 1991).

44. Gerstein, M. B. *et al.* Comparative analysis of the transcriptome across distant species.

Nature **512**, 445–448 (2014).

45. Zhang, B. & Horvath, S. A general framework for weighted gene co-expression network

analysis. *Stat. Appl. Genet. Mol. Biol.* **4**, Article17 (2005).

46. Khila, A. & Abouheif, E. Evaluating the role of reproductive constraints in ant social

evolution. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **365**, 617–630 (2010).

47. Benoit, B. *et al.* An essential role for the RNA-binding protein *Smaug* during the

Drosophila maternal-to-zygotic transition. *Development* **136**, 923–932 (2009).

48. Mével-Ninio, M., Terracol, R., Salles, C., Vincent, A. & Payre, F. *ovo*, a *Drosophila*

gene required for ovarian development, is specifically expressed in the germline and shares

most of its coding sequences with *shavenbaby*, a gene involved in embryo patterning. *Mech.*

Dev. **49**, 83–95 (1995).

49. Gramates, L. S. *et al.* FlyBase at 25: looking to the future. *Nucleic Acids Res.* gkw1016

(2016).

50. Peters, R. S. *et al.* Evolutionary history of the Hymenoptera. *Curr. Biol.* **27**, 1013–1018

(2017).

51. Carroll, S. B. Evo-devo and an expanding evolutionary synthesis: a genetic theory of

morphological evolution. *Cell* **134**, 25–36 (2008).

52. Robinson, G. E. & Vargo, E. L. Juvenile hormone in adult eusocial Hymenoptera:

gonadotropin and behavioral pacemaker. *Arch. Insect Biochem. Physiol.* **35**, 559–583
(1997).

53. Guidugli, K. R. *et al. Vitellogenin* regulates hormonal dynamics in the worker caste of a
eusocial insect. *FEBS Lett.* **579**, 4961–4965 (2005).

54. Hunt, B. G. *et al.* Relaxed selection is a precursor to the evolution of phenotypic
plasticity. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 15936–15941 (2011).

55. Morandin, C., Mikheyev, A. S. & Pedersen, J. S. Evolutionary constraints shape
caste☐specific gene expression across 15 ant species. (2017).

56. Schrader, L., Helanterä, H. & Oettler, J. Accelerated evolution of developmentally
biased genes in the tetraphenic ant *Cardiocondyla obscurior*. *Mol. Biol. Evol.* **34**, 535–544
(2017).

57. Moyers, B. A. & Zhang, J. Phylostratigraphic bias creates spurious patterns of genome
evolution. *Mol. Biol. Evol.* **32**, 258–267 (2015).

58. Meisel, R. P. Towards a more nuanced understanding of the relationship between sex-
biased gene expression and rates of protein-coding sequence evolution. *Mol. Biol. Evol.* **28**,
1893–1900 (2011).

59. Helanterä, H. & Uller, T. Neutral and adaptive explanations for an association between
caste-biased gene expression and rate of sequence evolution. *Front. Genet.* **5**, 297 (2014).

60. Trible, W. & Kronauer, D. J. C. Caste development and evolution in ants: it's all about
size. *J. Exp. Biol.* **220**, 53–62 (2017).

61. Erwin, D. H. & Davidson, E. H. The evolution of hierarchical gene regulatory networks.
Nat. Rev. Genet. **10**, 141–148 (2009).

62. Simola, D. F. *et al.* Epigenetic (re)programming of caste-specific behavior in the ant

Camponotus floridanus. Science **351**, aac6633 (2016).

63. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for

differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140

(2010).

64. Lazzeroni, L. & Owen, A. Plaid models for gene expression data. *Stat. Sin.* **12**, 61–86

(2002).

65. Tanay, A., Sharan, R. & Shamir, R. Discovering statistically significant biclusters in

gene expression data. *Bioinformatics* **18 Suppl 1**, S136–44 (2002).

66. Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation network

analysis. *BMC Bioinformatics* **9**, 559 (2008).

67. Yanai, I. *et al.* Genome-wide midrange transcription profiles reveal expression level

relationships in human tissue specification. *Bioinformatics* **21**, 650–659 (2005).

68. Larkin, M. A. *et al.* Clustal W and Clustal X version 2.0. *Bioinformatics* **23**, 2947–2948

(2007).

69. Suyama, M., Torrents, D. & Bork, P. PAL2NAL: robust conversion of protein sequence

alignments into the corresponding codon alignments. *Nucleic Acids Res.* **34**, W609–12

(2006).

70. Yang, Z. PAML 4: phylogenetic analysis by maximum likelihood. *Mol. Biol. Evol.* **24**,

1586–1591 (2007).

Fig. 1. Number of differentially expressed genes (FDR < 0.1) between a) queens and workers and b) nurses and foragers at each developmental stage or tissue in ants (left) and honey bees (right). "Head", "thorax", and "abdomen" refer to body segments of adults, while "pupa" and "larva" refer to whole bodies. "No ortholog" refers to genes for which no 1:1 ortholog exists (either due to apparent duplication or complete lack or orthology), "not shared caste/task bias" refers to genes for which 1:1 orthologs can be identified but are only differentially expressed in one species, and "shared caste/task" bias refers to genes for which 1:1 orthologs are differentially expressed in both species. Insets show the proportion of each category of gene out of all differentially expressed genes at that stage or tissue. c) Proportion of abdominal DEGs by estimated evolutionary age (shading). "Shared queen/worker" indicates genes upregulated in queen or workers of both species.

*: the category "larva" represents differential expression across larvae of all stages for which caste can be identified (second to fifth larval stage). Source data are provided as a Source Data File, "Supplementary_Fig1a.txt", "Supplementary_Fig1b.txt", "Supplementary_Fig1c.txt".

Fig. 2. Abdominal caste bias (log$_2$ fold change queen versus worker) is correlated with
connectivity within the queen-abdomen module in a) ants (rho = 0.536, P < 0.001) and b) honey
bees (rho = 0.617, P < 0.001). Genes upregulated in queens are in red, while genes upregulated
in workers are in blue. Connectivity is proportional to the most highly connected gene in the
module. Connectivity within the queen abdominal module is higher for genes found in the
module for both species (shared) versus genes found in the module for only one species (not
shared) in c) ants and d) honey bees. Middle line represents median values, outer edges of
boxplot represent upper and lower quartiles, and whiskers represent a deviation of
1.5*(interquartile range) from the upper and lower quartiles. Source data are provided as a
Source Data File, "Supplementary_Fig2.txt", ***P < 0.001 (Wilcoxon test)
628
629

Fig. 3. Caste bias is linked to sex bias. Abdominal caste bias (queen vs worker log$_2$ fold-change) is correlated to abdominal sex bias (queen vs male log$_2$ fold-change) in a) *M. pharaonis* (rho = 0.715, P < 0.001) and b) *A. mellifera* (rho = 0.774, P < 0.001) and abdominal sex bias is correlated between the two species (rho = 0.280, P < 0.001) (c). Red indicates shared queen-biased abdominal DEGs, while blue indicates shared worker-biased abdominal DEGs. Grey indicates genes that did not exhibit shared expression patterns or were not differentially expressed. Lines in a-c indicate the trendline of a linear model. d) Shared queen-biased abdominal DEGs tend to be female-biased in *D. melanogaster* while shared worker-biased abdominal DEGs tend to be male-biased in *D. melanogaster* (likely reflecting down-regulation in females). Middle line represents median values, outer edges of boxplot represent upper and lower quartiles, and whiskers represent a deviation of 1.5*(interquartile range) from the upper and lower quartiles. Source data are provided as a Source Data File, "Supplementary_Fig3a.txt", "Supplementary_Fig3b.txt", "Supplementary_Fig3c.txt", "Supplementary_Fig3d.txt",.

666
Fig. 4. Genes that exhibit more caste bias across tissues and developmental stages have younger
estimated evolutionary ages (a,b) and tend to be loosely connected (c,d; ant: rho = -0.159, P <
0.001; honey bee: rho = -0.090, P < 0.001) and rapidly evolving (e,f; ant: rho = 0.157, P < 0.001;
honey bee: rho = 0.240, P < 0.001). "Overall caste bias" combines queen/worker $\log_2$ fold-
change values across all development stages and adult body segments. Connectivity is calculated
using all samples and genes and scaled proportionally to the highest value. In a and b, middle
line represents median values, outer edges of boxplot represent upper and lower quartiles, and
whiskers represent a deviation of 1.5*(interquartile range) from the upper and lower quartiles.
Source data are provided as a Source Data File, "Supplementary_Fig4.txt",
676