Spreading Depression and Seizure Unification Experimentally Observed in Epilepsy

Authors: Fatemeh Bahari[1,2]*, Paddy Ssentongo[1,2], Jiayang Liu[1,2], John Kimbugwe[1,2], Steven J. Schiff[1,2,3,4,5], Bruce J. Gluckman[1,2,3,4]*

Affiliations:

[1]Center for Neural Engineering, Pennsylvania State University, University Park, PA 16802, USA.

[2]Department of Engineering Science and Mechanics, Pennsylvania State University, University Park, PA 16802, USA.

[3]Department of Neurosurgery, Pennsylvania State College of Medicine, Hershey, PA 16802, USA.

[4]Department of Bioengineering, Pennsylvania State University, University Park, PA 16802, USA.

[5]Department of Physics, Pennsylvania State University, University Park, PA 16802, USA.

*Corresponding authors email addresses: Fatemeh Bahari, fzb116@psu.edu, Bruce J. Gluckman BruceGluckman@psu.edu

Number of figures and tables: 4

Acknowledgements:

We thank Ali Nabi, Balaji Shanmugasundaram, and Myles W. Billard for assistance in development of the acquisition hardware, and Carlos M. Curay for animal care. This work was supported by National Institute of Health grant R01EB019804, a Multidisciplinary grant from Citizens United for Research in Epilepsy (CURE), Pennsylvania State Institute for the Neuroscience from Pennsylvania Department of Health Tobacco Funds, and a doctoral Academic Computing Fellowship from Pennsylvania State University to F.B.

Author contributions:

F.B., S.J.S, B.J.G designed the research; P.S., J.K., F.B., and B.J.G. performed surgeries and animal care; J.L. and B.J.G. designed electronics and hardware; F.B. and B.J.G. analyzed data; F.B., S.J.S., and B.J.G. wrote the manuscript.

Recurrent spontaneous seizures have been considered the sine qua non of epilepsy. Epileptic

seizures were first associated with intense electrical activity measured from the brain in 1933 [1].

Spreading depression of brain activity was discovered by Leão in 1944 [2], and has since been shown

associated with migraine, stroke and traumatic brain injury [3,4]. Although for many years spreading

depression was thought to have only rare genetic links to epilepsy [5], recently it has been proposed

that seizure-triggered spreading depression could underlie sudden unexplained death in epilepsy

(SUDEP) [6]. The physiological underpinning of spreading depression is an electrical spreading

depolarization (SD) of brain cells. The depolarization is associated with a large, prolonged

negative shift in tissue potential that cannot be well detected with standard human

electroencephalographic (EEG) technology. Nevertheless, in recent years, computational models

of the dynamics of neuron biophysics have shown a linkage between seizures and SD [7], and have

predicted the possibility of dynamic trajectories linking the two phenomena [8]. We developed a

recording system for chronic, long-term measurements that provides a view of the complex

interactions between spontaneous epileptic seizures and spreading depolarizations in the intact

brain [9]. We here experimentally demonstrate these theoretically predicted trajectories in multiple

animal models of epilepsy, showing that spontaneous single seizures are frequently associated with

SD, and that SD may mediate between seizures within clusters. We therefore propose that the ictus

of epilepsy may be a combination of seizures and SD.

Main

20 We found, from thousands of days of long-term continuous DC-sensitive chronic recordings,

21 frequent instances of seizure associated SD in two very different chronic animal models of

22 epilepsy: the rat tetanus toxin (TeTX) model of temporal lobe epilepsy (TLE) [10,11] (Fig. 1A) and

23 the murine model of post-cerebral malaria (post-CM) epilepsy [9] (Fig. 1B). Shown in Fig. 1A-B are

typical examples of SD following seizures, with the seizures evident from band-pass filtered signals (1-55 Hz) in the upper panels, and the SD from the low-pass filtered signals (< 1 Hz) in the bottom panels. SD either overlapped with (Fig. 1A) or followed (Fig. 1B) the seizure. In both models, the amplitude, duration and propagation between electrodes of the slow shift in tissue electrical potential are consistent with changes in extracellular potential observed during experimentally induced SD in vitro, as shown in Fig. 1C as well as other reports of acute SD [12–16]. During the large negative deflection of the tissue potential in the very low frequency range, we observed marked suppression of EEG activity (Fig. 1A). This is consistent with the spreading depression subtype of SD. The later reappearance of seizure-like spiking activity correlates with the return of the very low-frequency potential to near its baseline value. This pattern is consistent with recovery dynamics from acutely-induced SD termed spreading convulsions [4,17].

Although seizure-related SDs often overlapped with demonstrable suppression of EEG activity, we also found post-ictal generalized EEG suppression (PGES) that occurred without concurrent SD (Fig. S2).

The large slow shifts in the tissue potential associated with SD reside in the very low frequency (1-100 mHz) range. Unless a wide digitization range and non-polarizing electrodes with sufficient surface area are used, within these ranges measurements are dominated by electrode-tissue electrochemical polarization driven by the input current of the amplifier. To eliminate the electrochemical-based signal components and to restrict the signal to what could be acquired with a standard digitizer, in most acquisition systems the measurements are high-pass filtered (e.g. > 1 Hz) and therefore slow potential shifts associated with SD are removed [18,19].

To simultaneously measure SD and field potentials we used custom-made micro reaction chamber (µRC) electrodes with electrodeposited iridium oxide films (EIROF) [20] with a direct current (DC)-

sensitive acquisition system [9]. The fidelity of this recording combination is demonstrated *in vitro* in Fig. 2. With an applied sinusoidal electric field, we observed only minor signal attenuation with 50 µm (diameter) µRC EIROF (Fig. 2B-C red trace) electrodes at frequencies relevant to SD (0.01 Hz). This is in contrast to 17 µm Nichrome electrodes (Fig. 2B-C black trace), which are included to illustrate that electrodes with poorly matched impedance – to the input impedance of the amplifier – will inherently high-pass filter the tissue potentials.

The measurements in the included animals from both models were sufficiently stable during the entire recording periods – typically more than 2 continuous weeks – to capture similar SD events as presented in Fig. 1. We operationally define "sufficient stability" as having baseline (low-frequency) fluctuations with amplitudes substantially less than 10 mV over hour-long periods. Shown in Fig. S3 are the maximal fluctuations per hour of the unfiltered left hippocampal electrode potential over 8 consecutive days of recording in one epileptic mouse first without seizures (Fig. S3 upper panel) and then with seizure-associated SDs (Fig. S3 bottom panel). For every hour during the seizure-free period (Fig. S3 upper panel), the hippocampal measurements fluctuate smoothly with a maximal range (< 6 mV) much smaller than the $\gg 10$ mV low-frequency potential differences associated with SD episodes. However, the within-hour fluctuations are much larger for the hours with seizures-associated SDs (magenta arrows in Fig. S3 bottom panel). The negative deflection in tissue potential during SD leads to distinct large variations in the recording electrode's potential. These DC fluctuations associated with SDs are fully separable from steady state potentials and mechanically or electronically induced artifacts.

The two animal models of epilepsy utilized in this study are fundamentally different. The TeTX model of temporal lobe epilepsy is induced with intrahippocampal injection of tetanus toxin which creates a focal lesion with modulated neuronal excitability, and produces secondary generalized

seizures. In this model seizures emanate from sites throughout ventral hippocampi and generalize to other hippocampal and cortical regions. SDs appeared to originate from variable sites throughout the dorsal-ventral hippocampi (Fig. S4A-C) with occasional episodes of SD invading the cortex (data not shown). SDs often occurred during or at the end of well-spaced seizures (Fig. S4B-C). In seizure clusters SD dynamics appeared to connect the individual seizures (Fig. 3A, and Fig. S4A cyan trace).

Across 124 cumulative days of recordings from 5 epileptic rats, 425 of 1256 seizures had co-occurring SD events (estimated rate = 33% ± 1%, see methods for the propagated error) (Fig. 3B). The murine model of post-CM epilepsy [9] mimics human conditions of post-infection acquired epilepsy with long epileptogenesis [21] and spontaneous unprovoked seizures. In this model the brain undergoes widespread damage, and the seizure origins and evolution patterns vary. We observed episodes of SD occurring during or after the seizures in one or both hippocampi (Fig. S5A-C) and/or cortex (Fig. S5D) in 171 of 442 total seizures (estimated rate = 38% ± 2%) (Fig. 3C). Collectively seizures were pooled from 21 epileptic mice with 1139 cumulative days of recordings. The SD pattern was uncorrelated with seizure origin or evolution (Fig. S6A-C), and SD was observed in all four mouse-parasite strain combinations studied (Fig. S6D).

In both animal models and all instances of SD the amplitude of the negative DC deflection, duration of the deflection, propagation, and existence of a quiescence period reflective of neuronal depolarization were consistent with SD characteristics, as was the frequent observation of spiking activity at SD offset [17]. The SD events in the TeTX rats were often concurrent with depression of EEG activity. Therefore we mark these SDs as spreading depression subtype of the spreading depolarization phenomenon.

In post-CM epileptic mice, the apparent propagation speed of SD from left to right hippocampus was approximately 6 mm/min. However, within hippocampi in the TeTX rats, with 3-4 hippocampal recording electrodes, the propagation speed of SD measured in the midst of prolonged and spatially extended seizures appears to be faster (6 to 40 mm/min). We hypothesize that, in these cases, the tissue is primed to support faster propagation perhaps analogous to the way that seizure propagation speeds can be parametrically modulated via neuronal excitability [22,23].

SD events are shown to generate long bouts of acidosis [24,25]. It is therefore important to note that our observations are not artifacts of the pH dynamics. Although the µRC EIROF electrodes are pH sensitive ([26], and as described in Methods), the expected change from SD acidosis yields positive deflections in the electrode potential, and therefore would *cancel* some of the actual negative shift in tissue potential observed during SD.

In order to confirm the linkage between SD and seizures, we scored data from age-matched control animals who had viable DC recordings for appearance of SD-like events. In rats controls included animals that did not receive TeTX injections (N = 2, 154 cumulative recording days), and rats with TeTX injection that did not develop epilepsy (N = 1, 80 recording days). In mice, control data included uninfected animals with no seizures (N = 17, 820 cumulative recording days), and animals rescued from CM that did not develop seizures (N = 6 mice, 282 cumulative recording days). No SD-like events were observed in any of these groups.

From a rest or steady state, a neuronal system exhibits excitability when in response to small perturbations it takes a dynamical trajectory determined by the presence of bifurcations in its underlying mathematical structure. Such trajectories have been extensively characterized for single neuron spiking [27,28], and with the incorporation of fast sodium and slower potassium dynamics for complex burst and seizure firing dynamics [29,30]. The dynamics characteristic of experimental

hypoxic seizure patterns can be described by utilizing oxygen as an experimental and theoretical bifurcation parameter [31]. Combination of oxygen and potassium as state variables, as well as ion mass and charge conservation and energy balance, uncovers the unification between steady-state, spikes, seizures, and spreading depolarization [7].The subdivision of the dynamics within the oxygen-potassium state space is illustrated in Fig. 4A (adapted from [7]). Cellular volume fraction as another bifurcation parameter further separates seizure from spreading depolarization dynamics during trajectories that visit them in sequence [8].

If one classifies brain state in our measurements into the discrete states: steady state (SS), seizure (SZ) and SD, then the time series of discrete states can be represented by continuous trajectories through the oxygen-potassium state space illustrated in Fig. 4A&C. Two of the most common state series we observe are represented by the trajectories illustrated in Fig. 4A (SS→SZ→SS in cyan, and SS→SZ→SD→SS in green). Even the transition dynamics observed in Fig. 4B, in which a seizure is followed by repeated SD events, is naturally described by the looping magenta trajectory in Fig. 4C.

Because the state space of the system includes un-represented slow variables such as extracellular volume and vascular tone, trajectories can cross in this two-dimensional projection. Validation of the linkage between such computational modeling and *in-vivo* observed dynamics can be done through casting the measurement time series into a discrete-state transition model, as illustrated in Fig. 4D, and then statistically correlating the transition probabilities ($\alpha_{i,j}$) as a function of the modeled state variables, such as tissue oxygen pressure and extracellular potassium.

Our recording system offers, for the first time, the opportunity to investigate the theoretically predicted complex interaction between seizures and spreading depolarizations in an intact brain. Within two fundamentally different animal models of chronic epilepsy, we observed direct

transitions and dynamical interplay between SDs and spontaneous seizures. Moreover, in the epileptic rats we often observe that seizures that occur in clusters are connected by SD events (Fig. 3A, Fig. S4A). Our findings suggest that beyond EEG, currently unobserved parameters such as extracellular volume, potassium, and tissue oxygenation are important in orchestrating the dynamics observed in epilepsy. Furthermore we propose an expanded definition for the event or ictus in epilepsy that encompasses dynamical trajectories that combine both seizures and SD, and better account for the electrical and behavioral manifestations during epilepsy.

Methods

All animal work was approved by and performed under administration of Institutional Animal

Care and Use Committee at the Pennsylvania State University.

Post-Cerebral Malaria Epilepsy Model:

Infection and Treatment:

Donor animals were infected from frozen stocks of Plasmodium berghei ANKA (PbANKA) or

Plasmodium berghei NK65 (Pb-NK65) parasite, and blood was drawn on day 7 for inoculation

into homologous experimental animals. Swiss Webster (SW), C57BL/6 (Charles Rivers

Laboratory) and CBA/CaJ (CBA, Jackson Laboratory) male mice were infected by intraperitoneal

injection of infected red blood cells from the donor animal. Age-matched control mice were

inoculated with red blood cells from un-infected donor animals, and otherwise received identical

drug and monitoring treatment.

Animals were treated by anti-malarial medication, Artesunate (64 mg/kg dosage), for seven days

starting from day 5 (C57BL/6-PbANKA or CBA-PbANKA), day 6 (SW-PbANKA) and day 7

(SW-PbNK65) post-infection. We based our treatment plan on observations in human cases where

epilepsy is a sequela of well-developed CM rather than cases with early detection and treatment.

Animal Surgery and Care:

Animals were implanted with electrodes to monitor brain activity at least five days post-treatment

following procedures described in [9]. Briefly, four stainless steel screws (#000 self-tapping, Morris

Co.) were placed over frontal (AP +1.5, ML ±2.5 mm) and S1 (AP -2.3, ML ±3.5 mm) cortices

for measurements of electrocorticogram (ECoG). Two custom-made 50 μm (diameter) ultra-low

impedance micro-reaction chamber (μRC)[20] electrodes were further enhanced by electrodeposition

of iridium oxide films (EIROF)[20] to provide DC stability and high charge passing capacity. The

electrodes were then implanted bilaterally in dorsal hippocampi (AP -2.3, ML ±2.0, DV -1.5 mm)

to provide hippocampal depth recordings. All coordinates were Bregma referenced.

Muscle activity was monitored via a 3 mm long 50 μm gold-plated 316L stainless steel wire placed into a bluntly dissected opening within the nuchal muscles and further secured with polyglycolic acid absorbable sutures through the subcutaneous tissue. A fraction of animals were implanted with a lead in the precordium to monitor electrocardiogram (ECG) activity. Electrodes were then secured in place and electrically isolated via a 3D printed multi-groove headmount embedded in dental cement.

At the completion of the surgery, animals were housed individually in custom-made plexiglass cages each containing two enclosures with dimensions 6"W×12"D×12"H with free access to food and water, and, following recovery, were cabled for continuous video and EEG monitoring. Implanted mice were cabled to the acquisition board which hangs down through a 1.5 inch hole in the ceiling of each enclosure. The board is connected to a custom-made commutator to allow for postural flexibility and free movement of the animals. A 12 hour light-dark cycle was maintained with the lights on between 6 am and 6 pm. During the dark period infrared LED arrays were used to illuminate the cage for continuous 24-hour video monitoring. Synchronized video (at 3 frames per second) and EEG recordings began once the animals were successfully cabled.

Rat Tetanus Toxin Model of Temporal Lobe Epilepsy:

Animal Surgery and Care:

Male and female Long-Evans rats, weighing 250-350 grams, were implanted with recording electrodes and received tetanus injections during the same surgical procedure following methods described previously [10]. In short, to induce epilepsy, 10-13 nano-grams of tetanus toxin (Santa Cruz Biotechnology, CAS 676570-37-9) dissolved in 1.3 microliters PBS mixed with 2% BSA were injected into the left ventral hippocampus (AP -5.15, ML +5.35, DV -7.65 mm) through a 30-gauge flexible cannula over 15 minutes.

Recording electrodes included stainless steel screws for measurements of electrocorticogram (ECoG) at coordinates of (AP +1.5, ML ±4 mm) and (AP -2, ML ±3 mm), and custom-made 50 μm (diameter) ultra-low impedance μRC, EIROF electrodes implanted in dorsal and ventral hippocampus (AP -2.5, ML ±2.0, DV -3.2 mm), (AP -3.9, ML ±2.2, DV -3.1 mm), (AP -5.15, ML -5.35, DV -7 mm), and (AP -6.0, ML 5.0, DV -5.5 mm) to provide differential measurements of the hippocampal local field potentials (LFP). For each hippocampal site, a bundle of two electrodes with ends 125-250 μ dorsally apart was used. All coordinates were Bregma referenced.

All rats received a lead in the precordium to monitor Electrocardiogram (ECG) activity. Electrodes were secured in place and electrically isolated via dental cement. The leads were then connected to the acquisition amplifier and all was encapsulated within a 3D printed headmount. At the completion of the surgery animals were returned to their individual homecages – standard autoclave-ready rat cages – with free access to food and water and maintained at a 12 hour light-dark cycle with lights on between 6 am and 6 pm.

Seven days post-recovery, rats were connected to a commutator at the top of the cage via a low-weight USB cable. A mini-computer (Raspberry Pi foundation, 3 model B) attached to the cage cover acquires data continuously to a network attached storage (NAS). A separate single-board computer (Raspberry Pi foundation, 3 model B) with a low-light level-compatible camera system similarly spools continuous, time-synchronized video data to the same NAS.

Animals were housed individually in custom-made plexiglass cages

Data Collection

For mice and rats all biopotentials were acquired at 24 bit resolution and 1 kHz sampling frequency (per channel) via our custom-made data acquisition system.

For mice acquisition system was designed to provide 8 channels of high quality biopotential recordings. The 24-bit digitization provides a dynamic range of 4.5 V with sub-microvolt

divisions. Therefore it accommodates the amplitude range and resolution to simultaneously resolve large shifts in tissue potentials 10-30 mV) associated with SD and normal activity in the range of few millivolts associated with field potentials. This feature complemented with DC-stable μRC electrodes will eliminate the need for analog high-pass filtering prior to digitization.

For rats, we extended the acquisition system to provide 16 channels and a 3-axis accelerometer and to fit within a 3D-printed head mounted box (amplifier size: 1"W×1"D×0.25"H).

The raw data contain the information in all frequencies, including bio-potentials from neural activity, spreading depolarization, and electrochemical changes.

Data Analysis

All recorded data were inspected via in-house written Labview (National Instruments) and MATLAB (MathWorks Inc.) programs that allow for simultaneous re-referencing, filtering, spectral analysis and annotation. All data were originally referenced to a reference screw electrode. For seizure and state of vigilance scoring, the raw ECoG and hippocampal depth recordings were band-pass filtered at 1-55 Hz and 1-125 Hz; respectively, to highlight field potentials and seizure dynamics. Electromyogram (EMG) and ECG data were band-pass filtered at 1-125 Hz to extract muscle activity and cardiac dynamics. 3-axis head acceleration was band-pass filtered at 1-100 Hz.

Seizure Detection

For mice seizures were detected manually, from band-pass filtered EEG, using custom-written in-house software within Labview environment. Spontaneous seizure activity of more than 10 seconds was identified and scored for origin and evolution pattern according to previously described criteria in Ssentongo et al. 2017. Inclusion criteria for putative seizure incidents were at least one low-noise hippocampal depth recording and at least one low-noise ECoG recording. Epilepsy criteria was defined as observation of at least 2 seizures at or after day 3 post-implant, and one seizure more than 26 days post-infection. All epileptic animals with at least one ECoG

and one hippocampal depth electrode with stable DC recordings throughout their lifetime were selected for further analysis.

For rats, seizures were detected automatically, from bandpass filtered EEG and head acceleration measurements, using in-house custom-written routines in Labview software and verified visually to determine precise onset and end times. Inclusion criteria for each hour of data analyzed were availability of at least one low-noise hippocampal LFP, at least one low-noise ECoG and at least one low-noise axis of head acceleration measurements. Spontaneous seizure activity of more than 10 seconds was marked as a seizure.

Spreading Depression Detection

A custom written MATLAB (Mathworks Inc.) script was used to find bouts of spreading depolarization. SD is characterized by significant negative DC changes in the signal. In order to detect such shifts, the raw EEG was low-pass filtered with a cutoff frequency of 2 Hz. We then calculated the first order time-derivative of the low-pass filtered signal and marked any instance of threshold (= 4 × standard deviation from the mean) crossing as an indicator of a SD bout. Time-series of detected SD events were visually inspected and verified to achieve temporal accuracy in SD end times which were identified as when DC potential recovered to the pre-seizure baseline.

DC Artifacts:

The changes in DC potential might not all be due to physiological phenomena. Transient changes in connectivity can cause transient, sharp shifts in electrode potential. These events are often caused by mechanical factors that temporarily affect the amplifier-headmount connection such as animal handling during cleaning and measurements of parasitemia levels. During these artifacts, the DC potential rapidly rises and then recovers to its baseline value through a short recharge period.

Our SD detection algorithm did not account for such DC shifts. Thus, time-series of detected SD events were visually inspected and verified to account for false detections. These transient incidents are distinguishable from baseline DC potential fluctuations because of their relatively larger amplitude and from SD-like events due to their relatively shorter duration.

Included Data

In both animal models of epilepsy, animals were included for further analysis if they had at least one ECoG and one hippocampal depth recordings with sufficient stability. As described previously, "sufficient stability" is defined as having baseline (low-frequency) fluctuations with amplitudes substantially less than 10 mV over hour-long periods. Therefore, all included animals had stable DC recordings throughout their lifetime.

For both models, we included epileptic animals (ones that experienced spontaneous, recurrent seizures according to the criteria previously described) and control animals. Controls included both animals that received neurological insult (CM or TeTX) and did not develop epilepsy and ones that did not receive CM or TeTX.

Post-Cerebral Malaria Epilepsy Model:

Overall data were collected from 21 post-CM epileptic mice with 1139 days of continuous recordings. Control data included uninfected control animals with no seizures (17 mice with over 820 cumulative recording days), and animals rescued from CM that did not develop seizures (6 mice with 282 cumulative recording days).

Rat Tetanus Toxin Model of Temporal Lobe Epilepsy:

Overall data were collected from 5 TeTX epileptic rats with 124 days continuous recordings. Control data included 2 rats with 74 recording days each that did not receive TeTX injections and one rat that received TeTX injection but did not develop epilepsy with 80 days of recording.

Collectively, the two models account for 2593 days of full-time continuous measurements.

Hardware Validation:

The large slow shifts in the tissue potential associated with SD reside in the very low frequency (1-100 mHz) ranges. Therefore, in order to capture these events the measurements need to be free of the electrode-tissue electrochemical polarization driven by the input current of the amplifier. This is in addition to the digitization range and resolution required to simultaneously record SD and field potentials.

For evaluation of electrode polarization under DC potential changes we prepared a conductive medium from 0.09% NaCl solution (10 times weaker than standard physiological saline). The solution was then poured into the experiment chamber which is built from plexiglass (blue area in Fig. 2A). The two conductive sections (area = 2×4 cm^2 and 2×0.5 cm^2) are separated from one another with molding clay (green piece) and the only electrical connectivity is from a glass pipette filled with 0.09% saline.

Within this setup, a spatially uniform, AC electric field was introduced by passing current (Analog Stimulus Isolator, AM Systems Model 2200) between a pair of large platinum plates (0.7×1.7 cm^2) embedded on either end of the chamber relatively far from the electrodes (distance between plates = 6 cm). Test electrodes were embedded inside the chamber at known distances from one another. Measurements were referenced to a stainless steel screw electrode placed in the chamber floor (one of the red circles in Fig. 2A).

The electrodeposited iridium oxide films (EIROF) are known to be pH sensitive[26]. Physiological brain pH variations occur in the time-scale of seconds to minutes. Therefore, our measurements in the very low frequency ranges are potentially combinations of changes induced by electrochemical reactions (pH) and electrical fields induced by depolarization of neural populations (SD).

We characterized the pH response of the custom-made EIROF deposited 50 μm micro-reaction chamber (μRC) electrodes in the physiologically relevant pH range of 6-8. The electrodes were

placed in the Britton-Robinson buffer solution while drop-wise addition of 0.2 M solutions of NaOH and H_2SO_4 set the appropriate pH range (measured via a calibrated pH-meter, Anaheim Scientific P771). The test electrodes were all referenced to a pH-insensitive Ag/AgCl pellet electrode (AM Systems, Catalog #550008). The pH-insensitivity of the Ag/AgCl pellet was confirmed via measurements against a double junction Ag/AgCl reference electrode (Beckman Coulter, 3.5 M KCl, Item #A57189). We found the pH sensitivity of our custom-made EIROF deposited μRC electrodes to be 70 mV/decade at body temperature of 37°C (data not shown).

Measurement of Spreading Depolarization in Rat Brain Slice:

In order to further validate our in vivo potential measurement of SD, we acutely induced cortical SD and measured tissue potential with our recording system designed and utilized for chronic measurements in epileptic rats (as previously described in Methods). The slice preparation and induction of cortical SD were in the context of experiments detailed in [32]. We assembled an array of μRC EIROF electrodes according to methods described in [23]. The assembly was then advanced into the slice while electrode potentials were continuously recorded. Shown in Fig. S1A is the position of electrodes with respect to the cortex. The colored crosses correspond to the colored traces of electrode potential time-series in Fig. S1C. The large negative deflection in each electrode potential was concurrent with the time SD reached the electrode as shown in Fig. S1B for the three of the recording electrodes (blue, green, black crosses in Fig. S1B).

Statistical Analysis:

The fraction of seizure-associated spreading depolarization events, 33% and 38% for TeTX and post-CM model respectively, is the count of seizures with SD from all observed seizures. These counts are estimates following the binomial count statistics. Therefore, the propagated error for these estimates is $\sqrt{f(1-f)/N}$, where N is the sample size and f is the estimated

1   $\frac{seizure-associated\ SD}{all\ seizures}$. The propagated estimation error was 0.02 and 0.01 for TeTX and post-CM

2   model respectively.

References

1. Berger, H. Über das elektrenkephalogramm des menschen. *Arch. Psychiatr. Nervenkr.* **98,** 231–254 (1933).

2. Leao, A. A. P. SPREADING DEPRESSION OF ACTIVITY IN THE CEREBRAL CORTEX. *J. Neurophysiol.* **7,** 359 LP-390 (1944).

3. Lauritzen, M. Pathophysiology of the migraine aura: The spreading depression theory. *Brain* **117,** 199–210 (1994).

4. Lauritzen, M. *et al.* Clinical relevance of cortical spreading depression in neurological disorders: Migraine, malignant stroke, subarachnoid and intracranial hemorrhage, and traumatic brain injury. *Journal of Cerebral Blood Flow and Metabolism* **31,** 17–35 (2011).

5. Rogawski, M. Migraine and epilepsy—shared mechanisms within the family of episodic disorders. *Jasper's Basic Mech. Epilepsies* (2012). doi:NBK98193 [bookaccession]

6. Aiba, I. & Noebels, J. L. Spreading depolarization in the brainstem mediates sudden cardiorespiratory arrest in mouse SUDEP models. *Sci. Transl. Med.* **7,** 282ra46-282ra46 (2015).

7. Wei, Y., Ullah, G. & Schiff, S. J. Unification of Neuronal Spikes, Seizures, and Spreading Depression. *J. Neurosci.* **34,** 11733–11743 (2014).

8. Ullah, G., Wei, Y., Dahlem, M. A., Wechselberger, M. & Schiff, S. J. The Role of Cell Volume in the Dynamics of Seizure, Spreading Depression, and Anoxic Depolarization. *PLoS Comput. Biol.* **11,** (2015).

9. Ssentongo, P. *et al.* A Murine Model to Study Epilepsy and SUDEP Induced by Malaria Infection. *Sci. Rep.* **7,** 43652 (2017).

10. Sedigh-Sarvestani, M. *et al.* Rapid eye movement sleep and hippocampal theta oscillations precede seizure onset in the tetanus toxin model of temporal lobe epilepsy. *J. Neurosci.* **34,** 1105–14 (2014).

11. Jefferys, J. G. R., Borck, C. & Mellanby, J. Chronic focal epilepsy induced by intracerebral tetanus toxin. *Ital. J. Neurol. Sci.* (1995). doi:10.1007/BF02229071

12. Canals, S. Longitudinal Depolarization Gradients Along the Somatodendritic Axis of CA1 Pyramidal Cells: A Novel Feature of Spreading Depression. *J. Neurophysiol.* **94,** 943–951 (2005).

13. Kunkler, P. E. & Kraig, R. P. Hippocampal spreading depression bilaterally activates the caudal trigeminal nucleus in rodents. *Hippocampus* **13,** 835–844 (2003).

14. Olsson, T. *et al.* Cell swelling, seizures and spreading depression: An impedance study. *Neuroscience* **140,** 505–515 (2006).

15. Martens-Mantai, T., Speckmann, E. J. & Gorji, A. Propagation of cortical spreading depression into the hippocampus: The role of the entorhinal cortex. *Synapse* **68,** 574–584 (2014).

16. Cain, S. M. *et al.* In vivo imaging reveals that pregabalin inhibits cortical spreading depression and propagation to subcortical brain structures. *Proc. Natl. Acad. Sci.* (2017). doi:10.1073/pnas.1614447114

17. Van Harreveld, A. & Stamm, J. S. Spreading cortical convulsions and depressions. *J. Neurophysiol.* **16,** 352–66 (1953).

18. Hartings, J. A. *et al.* Spreading Depolarizations and Late Secondary Insults after Traumatic Brain Injury. *J. Neurotrauma* (2009). doi:10.1089/neu.2009.0961

19. Herreras, O. Local Field Potentials: Myths and Misunderstandings. *Front. Neural Circuits* (2016). doi:10.3389/fncir.2016.00101

20. Shanmugasundaram, B. & Gluckman, B. J. Micro-reaction chamber microelectrodes especially for neural and biointerfaces. (2017).

21. Bahari, F., Ssentongo, P., Schiff, S. J. & Gluckman, B. J. A Brain-Heart Biomarker for Epileptogenesis. *J. Neurosci.* (2018).

22. Richardson, K. A., Schiff, S. J. & Gluckman, B. J. Control of Traveling Waves in the Mammalian Cortex. *Phys. Rev. Lett.* **94,** 028103 (2005).

23. Pinto, D. J. & Ermentrout, G. B. Spatially Structured Activity in Synaptically Coupled Neuronal Networks: I. Traveling Fronts and Pulses. *SIAM J. Appl. Math.* (2001). doi:10.1137/S0036139900346453

24. Tschirgi, R. D., Inanaga, K., Taylor, J. L., Walker, R. M. & Sonnenschein, R. R. Changes in Cortical pH and Blood Flow Accompanying Spreading Cortical Depression and Convulsion. *Am. J. Physiol. Content* **190,** 557–562 (1957).

25. Somjen, G. G. Acidification of interstitial fluid in hippocampal formation caused by seizures and by spreading depression. *Brain Res.* **311,** 186–188 (1984).

26. Papeschi, G., Bordi, S., Beni, C. & Ventura, L. Use of an iridium electrode for direct measurement of pI of proteins after isoelectric focusing in polyacrylamide gel. *Biochim. Biophys. Acta - Protein Struct.* **453,** 192–199 (1976).

27. Izhikevich, E. M. Neural Excitability, Spiking and Bursting. *Int. J. Bifurc. Chaos* **10,** 1171–1266 (2000).

28. Rinzel, J. & Ermentrout, G. B. Analysis of neural excitability and oscillations. *Methods in neuronal modeling* (1989).

29. Cressman, J. R., Ullah, G., Ziburkus, J., Schiff, S. J. & Barreto, E. The influence of sodium and potassium dynamics on excitability, seizures, and the stability of persistent states: I. Single neuron dynamics. *J. Comput. Neurosci.* **26,** 159–170 (2009).

30. Barreto, E. & Cressman, J. R. Ion concentration dynamics as a mechanism for neuronal bursting. *J. Biol. Phys.* (2011). doi:10.1007/s10867-010-9212-6

31. Wei, Y., Ullah, G., Ingram, J. & Schiff, S. J. Oxygen and seizure dynamics: II. Computational modeling. *J. Neurophysiol.* **112,** 213–223 (2014).

32. Whalen, A. J. *et al.* Control of Spreading Depression with Electrical Fields. *Sci. Rep.* (2018). doi:10.1038/s41598-018-26986-1

Figure 1. Examples of SD from chronic Recordings. (A) Spontaneous seizure with SD activity in the

hippocampus of a rat under the TeTX model of temporal lobe epilepsy. During the large negative deflection

in DC potential associated with SD (bottom traces), band-pass filtered field potentials (top traces) are

suppressed. Filter settings for traces shown: Seizure traces band-pass 1-50 Hz; SD traces low-pass filter

below 2 Hz; EPL, ECoG posterior left; EAL, ECoG anterior left; EPR, ECoG posterior right; EAR, ECoG

anterior right; HPL, hippocampal posterior left; HAL, hippocampal anterior left; HAR hippocampal

anterior right; HPR, hippocampal posterior right; HVR, hippocampal ventral right. **(B)** Example observed

from a mouse model of post-cerebral malaria epilepsy. SD waves are observed in the measured DC potential

(bottom traces) from hippocampal leads following a seizure (top traces). Seizure onset and offset are

marked by the red and green dashed lines, respectively. The gold line marks the onset of the first seizure-

associated SD. Filter settings for traces shown: Seizure traces band-pass 1-50 Hz; SD traces low-pass filter

below 2 Hz. D_L, depth hippocampus left and D_R, depth hippocampus right, E_{LA}, ECoG left anterior, E_{RA},

ECoG right anterior and E_{LP}, ECoG left posterior, E_{RP}, ECoG right posterior. **(C)** Shape of experimentally

1   induced hippocampal SD (Top; adapted with permission from Canals et al., 2006) and (Bottom, in house

2   slice electrophysiology recordings with the DC-sensitive acquisition system, for more details see Fig. S1).

Figure 2. Hardware validation for stable DC and AC measurements. (A) Schematic of the

experimental chamber for in vitro hardware validation. The chamber is built from plexiglass and

contains 0.09% saline solution (blue area). The two sides of the chamber are electrically isolated from one

another via clay (green area) and the only connection is through a pipette (black lines) that forms a salt

bridge. The test electrodes are placed on either side of the chamber in the conductive solution. Two screw

electrodes (indicated by the red circles) are used as reference and ground. An electric field is imposed by a

set of platinum electrodes (indicated by the gray boxes) on the two sides of the chamber. Test electrodes,

reference and ground screws are connected to the acquisition system. The sinusoidal electric field with

different frequencies is applied by driving current from an isolated current supply (Analog Stimulus

Isolator, AM Systems Model 2200) through platinum plates at either end of the chamber away from the

testing electrodes. Testing electrodes are matched pairs of 50 um (in diameter) micro-reaction chamber

EIROF, 1 mm x 2.5 mm silver/silver chloride pellets (AM Systems, Catalog #550010), and 17 um

Nichrome wires at variable known distances apart. **(B) In vitro measurements of phase and amplitude**

of the electrode potentials across applied near-DC sinusoid electric field. The large Ag/AgCl pellet

(blue trace) phase and amplitude follow the sinusoid AC field (gray trace) at all frequencies shown here (0.01, 0.02, 0.04 Hz). While the EIROF electrodes (red trace) show some level of degradation both in phase and amplitude for these very low frequencies, they still sufficiently track the field. However the Nichrome electrodes (black trace) show a large lag and distorted signal. **(C) Low frequency sensitivity of the recording system.** The EIROF can sense and track the field across a wide range of frequencies with only minor attenuation at very low frequencies, in contrast to the much smaller and higher impedance Nichrome electrode that inherently acts as a high-pass filter for the tissue potentials and shows substantial phase and amplitude distortion.

Figure 3. SD is associated with a substantial percentage of seizures and mediates seizure clusters in

rat TeTX model of temporal lobe epilepsy. We identified seizure clusters as seizure events occurring

within at least 10 minutes of each other. **(A)** Two seizures occur within 5 minutes with SDs occurring during

and prior to seizures; the posterior and anterior left hippocampal leads capture SDs immediately prior to

and after the second seizure (HPL and HAL traces). **(B, C)** Cumulative count of seizure-associated SDs as

a function of seizures rank-ordered by time of occurrence in each epileptic model. Seizure-associated SDs

occurred in 33% of the collected seizures in epileptic rats **(B)** and 38% of the collected seizures in epileptic

mice **(C)**. Estimated rates have propagated errors less than 2% (see methods).

(Top traces A) Bandpass filtered field potentials. (Bottom traces A) Raw (unfiltered) hippocampal and

cortical measurements. Filter settings for traces shown: Seizure traces band-pass 1-50 Hz; SD traces no

low-pass filter. EPL; ECoG posterior left, EAL; anterior left, HPL; hippocampal posterior left LFP, HAL;

hippocampal anterior left LFP, HAR; hippocampal anterior right LFP, HPR; hippocampal posterior right

LFP, HVR; hippocampal ventral right LFP (contralateral side to the tetanus toxin lesion), EPR; ECoG posterior right, EAR; ECoG anterior right.

Figure 4. Bifurcations of the neuronal transmembrane dynamics. Based on computational models of the neuronal dynamics, the system (brain) can transition between steady state (SS), spreading depolarization (SD), and seizures (SZ). **(A, C)** Two-parameter bifurcation based on extracellular oxygen and potassium concentrations; adapted from (Wei et al. 2014). Experimental measurements are described by state-space trajectories such as SS→SZ→SS (cyan arrow in A), SS→SZ→SD→SS (green arrow in **A** and traces in Fig. 1A, Fig. S1B-C), and SS→SZ→SD→SS→SD→SS (magenta arrow in **C**, and cyclic SD in traces in **B**). These transitions can be cast into a discrete state transition model **(D)** to further calculate the transition probabilities based on variables such as extracellular oxygen, potassium concentrations, and cellular volume fraction (Ullah et al., 2015).

(Top traces B) Bandpass filtered field potentials. (Bottom traces B) Raw (unfiltered) hippocampal and cortical measurements. Filter settings for traces shown: Seizure traces band-pass 1-50 Hz; SD traces no low-pass filter. EPL; ECoG posterior left, EAL; anterior left, HPL; hippocampal posterior left LFP, HAL;

1   hippocampal anterior left LFP, HAR; hippocampal anterior right LFP, HPR; hippocampal posterior right

2   LFP, EPR; ECoG posterior right, EAR; ECoG anterior right.

1   Extended Data

Supplementary Figure 1: Measurement of induced SD in coronal slice of rat brain. Measurement of tissue electrical potential was performed with our recording system designed for chronic recordings from TeTX epileptic rats, in the context of slice electrophysiology experiments described in (Whalen et al., 2018). **(A)** Micro-reaction chamber EIROF electrodes were placed in the experimental chamber such that their tips touched the cortex (array was designed as in (Pinto and Ermentrout, 2001)). The spreading depolarization was induced using pipette injection of KCl. Cortical spreading depolarization was tracked using intrinsic optical imaging (IOS) as described in (Whalen et al. 2018) **(B)** and measurement of electrical tissue potential via the electrodes that were in tissue (each colored cross in A, B corresponds to the same color time-series in C) **(C)**. The frames captured in **(B)** are normalized to the background (first 100 frames prior to injection) and are 18 seconds apart in time. The color indicates intensity differences from the background. Each frame corresponds to a numbered gray dashed line in **(C)** and the colored crosses correspond to the potential traces.

Supplementary Figure 2: Post-ictal generalized suppression occurs in the absence of spreading

depolarization. (A) Power in all hippocampal and cortical channels is normalized to the baseline power

calculated from one hour of recordings. Normalized power distribution rises at the beginning of the seizure

and decreases much (20 dB) below the baseline values (30 seconds of baseline is highlighted via the gray

box) immediately after the seizure; similar to postictal suppression descriptions. Duration between five

seconds prior to the seizure and 15 seconds after the seizure is highlighted via the red box. Seizure onset

and offset are marked by the red and green dashed lines with triangle markers. **(B)** Band-pass filtered EEG

prior to the seizure indicates no large amplitude abnormal/epileptic like activity (gray panel). A short period

of postictal generalized EEG suppression (PGES) is apparent from the flat EEG immediately after the

seizure ends (after the green dashed line in red panel). This period lasts for approximately 10 seconds after

which the EEG recovers to baseline activity. **(C)** Seizures are concurrent with small shifts in tissue potential

as indicated in the raw EEG (EEG-DC) and previous reports (Mayanagi and Walker, 1975; Mader et al.,

2005). The tissue potential returns to baseline by the end of the seizure, and there is no large negative shift

during the postictal suppression of EEG activity. This is consistent with there NOT being SD concurrent

with the PGES. Red and green dashed lines mark the onset and offset of the seizure, respectively.

Filter settings for traces shown: EEG traces were band-pass filtered (1-50 Hz) prior to calculation of power

(A) and for presentation (B); DC traces in (C) were not filtered. EAR; ECoG anterior right, HAR;

hippocampal anterior right LFP, HPR; hippocampal posterior right LFP, HVR; hippocampal ventral right

LFP (contralateral side to the tetanus toxin lesion), EPL; ECoG posterior left, HPL; hippocampal posterior

left LFP, HAL; hippocampal anterior left LFP, THAL; ventrolateral thalamic anterior left, EAL; ECoG

anterior left.

Supplementary Figure 3: Stability of *in vivo* DC measurements over periods of days to weeks. In each

hour long block of data, the unfiltered potential range (max-min) for each hippocampal depth electrode (D_L

and D_R) was calculated. The range of the referential D_L measurements over 8 consecutive days (top trace)

remains stable with small fluctuations between 0.56–6.23 mV during a seizure free period of 8 days.

However the SD episodes (magenta arrows) associated with seizures in 3 different days during an 8-day

period (bottom trace) cause large fluctuations in the DC potential. Traces are from one epileptic animal.

Supplementary Figure 4: Types of spreading depolarization in the rat TeTX model of temporal lobe epilepsy. Negative DC deflections were observed across dorsal-ventral hippocampi in the electrodes recording local field potentials (LFP). Spreading depolarization either occurred prior to and during seizures in seizure clusters suggesting that SD can mediate the seizures **(A)** and/or occur following seizures **(B, C)**.

Red and green dashed lines mark the onset and offset of the seizure. The onset of the first measured SD in each panel is marked by a gold dashed line.

(Top traces) Bandpass filtered field potentials. (Bottom traces) Raw (unfiltered) hippocampal and cortical measurements. Filter settings for traces shown: Seizure traces band-pass 1-50 Hz; SD traces low-pass filter below 2 Hz. EPL; ECoG posterior left, EAL; anterior left, HPL; hippocampal posterior left LFP, HAL; hippocampal anterior left LFP, HAR; hippocampal anterior right LFP, HPR; hippocampal posterior right LFP, HVR; hippocampal ventral right LFP (contralateral side to the tetanus toxin lesion), EPR; ECoG posterior right, EAR; ECoG anterior right.

Seizures in (A.a, B.a, C.a) correspond to SD events in (A.b, B.b, C.b) respectively.

Supplementary Figure 5: Types of spreading depression in the post-CM epileptic mice. Negative DC deflections were observed mostly in hippocampal depth electrodes. Incidents of spreading depolarization were measured unilaterally **(B.b)** and bilaterally **(C.b)** in hippocampus often following or during a secondarily generalized seizure. At times the spreading depression occurred during a seizure and was concurrent with EEG suppression which was followed by a short run of high-amplitude spikes **(A.b)**. Red and green dashed lines mark the onset and offset of the seizure. **(D.b)** An example of cortical SD (in E_{LP})

starting during a secondarily generalized seizure with cortical origin (E_{LP}). The onset of the first measured SD in each panel is marked by a gold dashed line.

(Top traces) Bandpass filtered field potentials. A small focal hippocampal seizure is marked by the green box. (Bottom traces) Low-pass filtered hippocampal and cortical measurements. Filter settings for traces shown: Seizure traces band-pass 1-50 Hz; SD traces low-pass filter below 2 Hz. D_L, depth hippocampus left and D_R, depth hippocampus right, E_{LA}, ECoG left anterior, E_{RA}, ECoG right anterior and E_{LP}, ECoG left posterior, E_{RP}, ECoG right posterior.

Seizures in (A.a, B.a, C.a, D.a) correspond to SD events in (A.b, B.b, C.b, D.b) respectively.

Supplementary Figure 6: Origin and pattern of evolution in fraction of seizures followed by

spreading depolarization in post-CM mice. Out of all epileptic animals, we included the ones with

chronic, stable DC measurements. **(A)** Distribution of fraction of seizures in all cohorts of the model; yellow

lines represent seizures from animals with viable DC recordings selected for further analysis. **(B)**

Distribution of fraction of seizures in epileptic animals with viable DC recordings. **(C)** Fraction of seizures

followed by spreading depolarization events out of the total seizures pooled from animals with stable DC

recordings. The SD rate here is conditioned on the mouse/parasite combinations, seizure origin, and seizure

evolution. Note that a small number of cortical origin seizures are accompanied by SD events. **(D)**

Cumulative count of seizure-associated SDs as a function of seizures rank-ordered by time of occurrence

for each epileptic cohort.

References for the Extended Data

Mader EC, Fisch BJ, Carey ME, Villemarette-Pittman NR (2005) Ictal onset slow potential shifts recorded with hippocampal depth electrodes. Neurol Clin Neurophysiol.

Mayanagi Y, Walker AE (1975) DC potentials of temporal lobe seizures in the monkey. J Neurol 209:199–215 Available at: http://link.springer.com/10.1007/BF00312542.

Pinto DJ, Ermentrout GB (2001) Spatially Structured Activity in Synaptically Coupled Neuronal Networks: I. Traveling Fronts and Pulses. SIAM J Appl Math.

Whalen AJ, Xiao Y, Kadji H, Dahlem MA, Gluckman BJ, Schiff SJ (2018) Control of Spreading Depression with Electrical Fields. Sci Rep.