Patterns of fish utilisation in a tropical Indo-Pacific mangrove-coral seascape, New

Caledonia

Fish utilisation of a mangrove forest in New Caledonia

Authors: Dubuc, A[1,2,]*, Waltham, N.[1,2], Baker, R.[3], Marchand, C.[4], Sheaves M.[1,2]

[1] College of Science and Engineering, James Cook University, Townsville, Queensland, Australia

[2] TropWATER (Centre for Tropical Water & Aquatic Ecosystem Research), James Cook University,

Townsville, Queensland, Australia

[3] Department of Marine Sciences, University of South Alabama, Dauphin Island Sea Lab, Alabama,

United Sates of America

[4] ISEA (Institute of Exact and Applied Sciences), University of New Caledonia, Noumea, New

Caledonia

* Corresponding author

E-mail: alexia.dubuc1@my.jcu.edu.au (AD)

14   Abstract (300 words max)

Mangrove forests are important habitats for fish. However, their utilisation by fish,

and the specific values they confer, are still not fully understood. This study

describes how fish use mangrove forests in an Indo-Pacific mangrove-coral reef

seascape. Sampling was conducted using underwater video cameras (UVCs) to

describe spatial and temporal variations in fish assemblages across a small-scale (~

2.5 km^2) system, and over the tidal and lunar cycle. UVCs were deployed in the two

main component habitats of mangrove forests: at the mangrove forest edge, and

inside the forest (5 m from the forest edge), to establish patterns of utilisation of fish

across the tidal and lunar cycle. Proximity to coral reefs had a strong influence on

the mangrove fish community, as most fish recorded were reef-associated. Juveniles

of 12 reef species were observed, including two species classified as vulnerable on the IUCN list, and one endemic species. Fish assemblages on the mangrove edge differed significantly from those inside the forest. Most fish utilised the forest edge, with few species making regular use of in-forest habitats, supporting the contention that most fish species remain on the edge and potentially retreat into the forest for opportunistic feeding, or when threatened by larger predators. Species-specific patterns of utilisation varied across the tidal and lunar cycle. Small differences in depth profiles and substrate across the small-scale system had a significant effect on fish assemblages, highlighting the importance of accounting for spatial heterogeneity in these factors. These data provide important information for managers to implement adequate conservation strategies that include broader interconnected habitat mosaics.

Introduction

Mangrove systems are part of a mosaic of productive coastal habitats [1] that provide a variety of services to fish and human populations [2, 3]. Mangrove forests are a fundamental component habitat of mangrove systems [4], and confer many of the attributes that make them highly valuable fish habitats [5-9]. However, studies have shown varying degree of mangrove forest utilisation, with for instance a higher contribution of reef fish species to mangrove fish assemblages in the Caribbean compared to several places in the Indo-Pacific [10-21]. These observations suggest that not all mangrove forests provide equivalent values to fish. Moreover, recent work in mesotidal Australia suggests that few fish penetrate beyond the forest boundary [15], suggesting that the use of mangrove forests is spatially heterogeneous. This new evidence raises the question relating to the specific ways

in which mangrove forests are utilised by fish. More studies are needed to

characterise fish assemblages in mangrove forests with different settings (coastal,

estuarine, island, embayment), different tidal ranges (micro-, meso- or macrotidal),

proximity of other high value habitats such as seagrass beds and coral reefs, or

climatic zones [21-24]. A better understanding of how mangrove forest utilisation

varies spatially and temporally would provide new insights to help explaining the

contrasting results found in the literature.

In many parts of the Indo-Pacific, the tidal range is greater than in the

Caribbean, where mangrove forests are usually continually available to fish [3].

Intertidal mangrove forests are challenging environments, most notably because

they are only available to most aquatic organisms while they are flooded at high tide

[1, 24, 25]. The intermittent availability of mangrove forests may explain the low use

by fish in the Indo-Pacific [23]. Indeed, tidal variation (extent, duration and frequency

of flooding) generates a range of constraints for fish utilising mangrove forests. Most

evident is the decrease in water depth and eventual drainage of the forest as the tide

ebbs, forcing fish out of intertidal mangrove forest zones. Several studies have

indeed demonstrated that fish undertake regular migrations in tidally driven

mangrove systems, with different patterns of mangrove use according to fish

species, lunar cycle (neap vs spring tide) and tidal phase (flooding vs ebbing) [15,

26-29]. Migration of fish in response to tidal movements results in substantial

connectivity between the three major tropical coastal habitats: coral reefs, seagrass

beds and mangrove forests [25, 30], giving rise to the idea that mangrove forests are

part of a wider interconnected habitat mosaic [1]. Therefore, investigating tidal and

spatial variations in fish assemblages in mangrove forests is a crucial step towards

fully appreciating the value and functioning of the whole tropical coastal ecosystem.

The difficulty of sampling these habitats goes a long way towards explaining

the paucity of information available on fish assemblages inside mangrove forests

[20, 31]. The use of conventional techniques such as underwater visual censuses or

netting techniques is restricted across much of the Australasian region where

saltwater crocodiles (*Crocodylus porosus*) are common, and where dense mangrove

forests reduce the efficiency of most net-based approaches [15]. Recently,

underwater video has been successfully applied to study in-forest fish assemblages

[15, 32, 33], most notably because it overcomes a lot of sampling issues,

substantially reduces field labour intensity, and allows for high-temporal and -spatial

resolution data collection simultaneously in different habitats, such as the edge and

the inside of a mangrove forest [34].

In this study we deployed underwater cameras on the edge and inside a

mangrove forest [15, 26] coupled with high frequency depth loggers to record spatio-

temporal variations in fish assemblages in a microtidal Indo-Pacific mangrove-coral

reef seascape. We identified fish species that use the mangrove forest, and used an

array of exploratory data analyses and modelling techniques to describe how fish

utilisation changes between the forest edge and in-forest habitats, and how fish

assemblages vary across the tidal cycle.

Materials and methods

Study site

Our study focused on a relatively pristine mangrove forest in Bouraké, South

Province of New Caledonia (21° 56.971S, 165° 59.481E; Fig 1). New Caledonia is

an archipelago located in the South West Pacific, 1,500 km east of Australia. New

Caledonia has the largest lagoon in the world, partly registered on the UNESCO

World Heritage list. New Caledonia experiences a semi-arid to tropical climate with

annual total rainfall of 1,000 mm, and a mixed semi-diurnal microtidal regime

(maximum 1.8 m tidal range). Bouraké receives little freshwater inflow with no

defined drainage.

The area comprises approximately 2.5 km² of mangrove forest dominated by

Rhizophora stylosa on the seaward edge and *Avicennia marina* on the landward

margin, with a large semi-enclosed central lagoon (1.2 km long, 60 m wide, 1-2 m

depth). A channel (20-70 m wide, 2-6 m depth, 700 m long) connects the main

lagoon to the coastal waters of Pritzbuer Bay (~ 20 km^2). The channel comprises two

sheltered inlets (approximately 0.01 km² each), and a shallow (1-2 m depth) coral

reef platform that extends from the middle of the channel to the edge of the

mangrove forest. Corals could be seen right on the edge of the forest in some

places. We chose a study location where coral reefs occur in close proximity to

mangrove forests (Fig 1), effectively a seascape comparable to the Caribbean

coastline [35], so we could relate our findings to this other ecoregion [36] where the

tidal range is smaller.

Fig 1. Map and picture of the study system in Bouraké, South Province, New
Caledonia. The nine study sites in the mangrove channel sampled from the 21
February to 1 March 2017 are represented by their respective number. Light grey
areas represent mangrove forest, dark grey areas represent mainland, and white
areas represent water.

Data collection

Fish using the mangrove forest were examined on an inland/offshore gradient

along the channel (Fig 1). To assess differences in fish assemblage composition

between edge and inside the forest, 4 paired sampling were conducted (sites 1 to 8).

Each paired sampling consisted of two sites within 5-7 m distance; the even site

number of the paired sampling was located on the mangrove forest edge (defined as

the boundary between mangrove prop-roots and bare substrate), and the odd site

number located about 5 m inside the forest. Site 9 (considered an edge site in the

analyses) was located on the reef platform of the innermost bay, at the edge of

scattered mangrove trees slightly away from the main forest. The substrate at sites 1

and 9 consisted of dead corals, small live coral boulders and sand, while on other

edge sites it comprised mainly dead corals and small and larger live coral boulders.

The substrate was homogeneous and consisted of silt material at in-forest sites.

Fish assemblages were sampled using underwater video cameras (UVCs;

Model ATC9K Oregon Scientific) to investigate tidal variations in fish assemblages

simultaneously on the edge and inside the forest. Unbaited UVCs mounted on stable

bases were deployed at each site during neap (21 to 23 February 2017) and spring

tides (28 February to 1 March 2017). A sampling day consisted of cameras first

deployed at sites early in the morning (first light), continuing until the battery was

discharged, and, with a replacement battery, again deployed mid-afternoon at all

sites until the battery was discharged (recording lasted between 2h and 2.5h). Four

sampling days were completed (two during neap tides and two during spring tides).

Cameras were positioned around 7 cm above the substrate, facing towards the

channel. A marker mounted on a flexible rod (3 mm diameter, 0.5 m long) was

placed 0.5 m in front of the camera lens as a visibility indicator to ensure a minimum

visibility of 0.5 m was achieved in all videos. Visibility was very good and consistent

during the sampling period, and fish could be identified confidently up to approximately 2 m from the UVCs in all videos. As depth is one of the main limiting factors to mangrove accessibility, tidal variations in water depth (cm) were measured every 15 minutes at each site with depth loggers (In-Situ Inc. Rugged Troll 100 model). James Cook University issued a permit for a limited impact research to deploy underwater cameras in New Caledonia (no endangered or protected species were involved as no collection of any specimen was conducted). The study area does not benefit from any special protection, therefore, access and activities are not restricted, and no specific permit was required to sample.

Data extraction from videos

While UVCs allow large amounts of data to be gathered quickly in the field, considerable time is required to process these videos. Therefore, we subsampled the acquired video footage. From the two neap tide sampling days, one day was randomly selected and videos at all sites were processed for that day. For the remaining sampling day, all videos were processed from five sites; being the reef platform (site 9) and two pairs of in-forest and forest edge sites (sites 5-8). These sites were selected so one replicate for a site located on the reef platform, and two replicates of paired sites not located on the reef-platform were available. Considering this selection, the sites were then randomly chosen. The same selection was applied to the two sampling days conducted during spring tides.

Once sediments had settled after camera deployment (typically 2-3 min), videos were viewed using VLC media player (VideoLAN, 2001) and subdivided in 5-min intervals to follow tidal variations in fish assemblages. The occurrence of each fish taxon in each 5-min interval was recorded. Only presence/absence data were

recorded to avoid biases induced by count data when using UVCs [15]. Fish were

identified to the lowest possible taxonomic level. Features useful in discriminating

species within some genera or families such as *Plectorhinchus* spp.*,* Mugilidae spp.,

or Gobiidae spp. were difficult to distinguish in videos, therefore these taxa were

identified to genus or family level only. When possible, juvenile fish were identified

based on colour patterns and body shape. Any activity such as feeding, hiding,

cruising or escaping was also noted. All fish identifications were validated by two

additional experts. For each 5-min interval video processed, the information

concerning the date of sampling, site, time of day, habitat (edge vs in-forest), lunar

phase (neap vs spring), tide direction (flooding vs ebbing), and corresponding water

depth was recorded (Appendix S1).

Data analyses

An index depending on observation per unit effort, similar to the catch per unit

effort index (CPUE) when dealing with fishing techniques, was developed to

calculate frequencies of occurrence of taxa from the video data. The frequency of

occurrence of each taxon was calculated per site (the total number of 5-min intervals

in which a taxon was observed at a site was divided by the total number of 5-min

intervals recorded at this site). Only taxa with a frequency of occurrence ≥ 0.05 at

one or more sites were retained for analyses (referred to as "common taxa"). Taxa

with a frequency of occurrence < 0.05 (referred to as "rare taxa") were excluded from

analyses.

Non-metric multidimensional scaling (nMDS) was used as an exploratory

analysis to assess differences in fish assemblages among sites during spring and

neap tides. The frequency of occurrence of each common taxon was calculated per

site per lunar phase. Data were square root transformed (SQRT) to decrease the impact of extreme values, and an nMDS analysis based on Bray-Curtis dissimilarities, the most appropriate distance measure when using abundance data [37], was conducted. Clusters within the nMDS were determined by conducting an overlay cluster analysis at 40% and 45% similarity on the dissimilarity matrix of all frequencies of occurrence. A two-way analysis of similarity (ANOSIM) was used to test whether there were significant differences in fish assemblages between sites and lunar phase. Pearson correlations exceeding $R > 0.7$ between the ordination and taxa were used to fit vectors on the nMDS plot. All analyses were performed using PRIMER 6 software [38]. Additionally, frequencies of occurrence of each common taxon at in-forest and edge sites were calculated and plotted using horizontal bar plots to further investigate differences in fish assemblage composition between the two habitats.

To investigate the factors impacting fish presence/absence, a General Linear Mixed Model (GLMM) was conducted using the package "glmm" in R [39]. The GLMM was conducted on all the 5-min intervals recorded (Appendix S1) with presence/absence of any common taxa for each 5-min interval (1 if any common taxa were observed in the 5-min interval, or 0 if no common taxa were observed) as the response variable, "Depth", "Habitat" (edge vs in-forest), "Lunar phase" (neap vs spring), and "Time of day" (morning vs afternoon) as the fixed factors, and "Site", "Date", "Tide direction" (flooding vs ebbing) and a nested effect of "Site" within "Habitat" as the random factors, using a Bernoulli distribution and a logit link function.

Cumulative depth frequency curves were plotted for each site to highlight differences in temporal dynamics. To further understand how fish utilisation varies across depth, variations of SQRT frequencies of occurrence across depth, over

flooding and ebbing tide on the edge and in-forest were assessed using a General

Additive Mixed Model (GAMM). Each 5-min interval was allocated to a class of water

depth of 10 cm (from 10-20 cm to 120-130 cm) according to the water depth value

recorded, and the SQRT frequencies of occurrence of each common taxon was

calculated per class of water depth during flooding and ebbing tide per habitat (the

total number of 5-min intervals in which a taxon was observed at a class of depth

during flooding and ebbing tide per habitat was divided by the total number of 5-min

intervals recorded for this same sample unit). Frequencies of occurrence were SQRT

to reduce the impact of extreme values. To avoid false absence recordings, taxa

never recorded in the habitat of interest were not considered (i.e. if a taxon was

never recorded in-forest during the study it was not included in the in-forest

analysis). To run the GAMM, SQRT frequencies of occurrence were used as the

response variable, "Depth" as a smooth term, and "Habitat" and "Tide direction" as

parametric terms using a Gaussian distribution and an identity link function. "Habitat"

was included in the model to avoid any nesting issue. The model was built using the

package "mgcv" in R [40]. Patterns of variations of SQRT frequencies of occurrence

were then investigated graphically using boxplots to examine the variations of

average SQRT frequencies of occurrence among taxa at each depth interval, and a

LOESS curve was fitted to the data to analyse the general pattern of habitat use.

Patterns of mangrove habitat use for each taxon were then plotted using a LOESS

curve and individually assessed graphically to examine similarities and classify

patterns of fish occurrence across depth. Taxa were grouped in similar patterns if

their maximum average occurrence was observed at a similar tidal stage. Three

equivalent tidal stages were defined for this purpose: Low tide (between 10-20 and

40-50 cm); Mid tide (between 50-60 and 80-90 cm); High tide (between 90-100 and

120-130 cm).

Results

Fish composition

Fifty-six video deployments were analysed (totalling more than 118h of video).

Seventy-two taxa from 29 families were recorded, with 36 common taxa (frequency

of occurrence ≥ 0.05 on at least one site) retained for further statistical analyses

(Table 1). Most species recorded were marine and reef-associated [41] including fish

of families Scaridae, Chaetodontidae, Pomacanthidae, Siganidae, Acanthuridae,

Lutjanidae, or Labridae.

Table 1. Summary of all the families and taxa identified at Bouraké, New
Caledonia.

Family	Taxon	Family	Taxon
Acanthuridae	*Acanthurus auranticavus*[*][1]	Haemulidae	*Plectorhinchus lineatus*
	Acanthurus grammoptilus[2]		*Plectorhinchus spp.*[2]
	Acanthurus sp. cf blochii		*Pomadasys argenteus*[2]
	Ctenochaetus sp.	Hemiramphidae	*Hyporhamphus sp.*
	Zebrasoma velifer	Labridae	*Choerodon graphicus*
Apogonidae	*Fibramia lateralis*[*][3]		*Labridae spp.*
	Ostorhinchus septemstriatus	Lethrinidae	*Lethrinus harak*[*][2]
Belonidae	*Belonidae spp.*		*Lethrinus lentjan*[1]
Blenniidae	*Blenniidae spp.*		*Lethrinus obsoletus*
Carangidae	*Caranx ignobilis*	Lutjanidae	*Lutjanus argentimaculatus*[*][3]
	Caranx papuensis[1]		*Lutjanus fulviflamma*[4]
	Caranx sp.		*Lutjanus fulvus*[1]
Chaetodontidae	*Chaetodon auriga*[*][2]		*Lutjanus russellii*[2]
	Chaetodon bennetti[2]	Monodactylidae	*Monodactylus argenteus*[*][1]
	Chaetodon ephippium	Mugilidae	*Mugilidae spp.*[*][3]
	Chaetodon flavirostris	Mullidae	*Mulloidichthys flavolineatus*[2]
	Chaetodon lineolatus[2]		*Parupeneus ciliatus*
	Chaetodon lunula[2]		*Parupeneus indicus*[4]
	Chaetodon melannotus		*Upeneus tragula*[1]
	Chaetodon speculum	Pomacanthidae	*Pomacanthus sexstriatus*[2]
	Chaetodon vagabundus[2]		

	Heniochus acuminatus [1]	Pomacentridae	**_Neopomacentrus spp._** [2]
Clupeidae	**_Clupeidae spp._** [2]	Scaridae	**_Bolbometopon muricatum_**
Diodontidae	*Diodon hystrix*		**_Scarus sp. cf ghobban_** [1]
Ephippidae	*Platax pinnatus*	Scatophagidae	*Scatophagus argus*
Fistulariidae	*Fistularia spp.*	Serranidae	*Epinephelus caeruleopunctatus*
Gerreidae	*Gerres filamentosus*		*Epinephelus lanceolatus*
	_Gerres oyena_* [4]		*Epinephelus malabaricus*
Gobiidae	*Amblygobius linki*		*Epinephelus sp.*
	Amblygobius nocturnus	Siganidae	**_Siganus canaliculatus_** [1]
	Amoya gracilis [3]		**_Siganus lineatus_*** [2]
	Asterropteryx sp. cf striata [4]		*Siganus punctatus*
	Cryptocentrus leptocephalus [2]	Sparidae	**_Acanthopagrus sp. cf akazakii_** [2]
	Eviota sp.	Sphyraenidae	*Sphyraena barracuda*
	Exyrias puntang	Tetraodontidae	*Arothron hispidus*
	_Gobiidae spp._* [3]		
	Gobiidae spp.2		
	Redigobius balteatus [3]		

Taxa highlighted in bold represent the common taxa (frequency of occurrence ≥ 0.05 on at least one site) that were kept for statistical analyses. Taxa highlighted with a * represent the 10 most common taxa. Taxa or families underlined mean that juveniles were potentially observed for that taxon or for at least one taxon within the family that could not be identified. The superscript number corresponds to the pattern of mangrove habitat utilisation across depth followed by the taxon as described in Fig 6: 1) High-tide users; 2) Mid-tide users; 3) Low-tide users; 4) Generalist users.

Fish composition varied significantly among sites (ANOSIM: R = 0.793, p < 0.001), with distinct assemblages generating three and four different clusters at 40 % and 45 % similarity respectively on the nMDS plot (Fig 2). At 40 % similarity, the 1st cluster comprised all the samples conducted in-forest. The samples were characterised by a lower taxonomic richness (23 common taxa; Fig 3) dominated by *Fibramia lateralis* and all the taxa belonging to the Gobiidae family (except *Cryptocentrus leptocephalus* and *Asterropteryx* spp.), that were the only taxa recorded almost exclusively at in-forest sites (Fig 3). The 2nd cluster comprised all the samples conducted on the edge but site 7 at spring tide. The samples were characterised by a higher taxonomic richness (34 common taxa; Fig 3), among which 10 taxa, mostly reef-associated, significantly contributed to the fish

assemblage composition at edge sites (Fig 2). Site 7 at spring tide was an outlier

and made the 3rd cluster driven by the abnormally high occurrence of

Neopomacentrus spp. (Fig 2). Interestingly at 45 % similarity, another cluster was

generated, separating deep edge and shallow edge sites (Fig 4, Fig 2). Three

species of snappers, *Lutjanus fulviflamma, Lutjanus argentimaculatus and Lutjanus*

russellii were the only three species not showing apparent preference for edge or in-

forest sites as they were almost evenly recorded on the two habitats (Fig 3), and

therefore did not significantly characterised any of the two habitats (Fig 2). Replicate

samples plotted close to each other and were grouped in the same clusters (Fig 2).

Lunar phase did not significantly influence fish assemblages (ANOSIM: R = 0.2, p >

0.2).

Fig 2. nMDS analysis performed on square root transformed frequencies of
occurrence for each taxon per site per tide. Edge sites are represented by
squares and in-forest sites by circles. Sites sampled at neap tide are coloured in
deep pink, and sites sampled at spring tide in navy blue. Solid green and dotted blue
ellipses represent overlay clusters determined at 45 and 40 % similarity respectively.
Vectors represent taxa with a Pearson correlation with the ordination R > 0.7.

Fig 3. Proportion of time spent by each of the common taxa on the edge
versus inside the forest. edge=green and in-forest=brown. Proportions range from
0 to 1, 1 corresponding to a taxon exclusively recorded on the edge or in-forest and
0.5 corresponding to a taxon recorded on the edge as frequently as in-forest.

"Habitat" (GLMM: z = -2.637; p < 0.005), "Lunar phase" (GLMM: z = -9.406; p

< 0.001) and "Depth" (GLMM: z = -2.118; p < 0.05) significantly influenced the

presence/absence of taxa. However, "Time of day" (GLMM: z = -1.519; p > 0.1), date

of sampling (GLMM: z = 1.555; p > 0.05), "Tide direction" (GLMM: z = 0.991; p > 0.1)

and sites within a same habitat (GLMM: z = 1.394; p > 0.05) did not significantly

influence presence/absence of taxa. Further data exploration following the GLMM

results showed that at edge sites there was a higher proportion of 5-min intervals in

which a taxon was observed compare to in-forest sites. Similarly, during neap tides,

there was a higher proportion of 5-min intervals in which a taxon was observed

compare to spring tides (Appendix S2).

Tidal variations in fish assemblages

Average depth was substantially shallower at in-forest than edge sites (neap

tides (mean ± SE): 34 ± 0.57 and 55 ± 0.66 cm respectively; spring tides: 48 ± 1.11

and 71 ± 1.23 cm respectively), as was maximum depth (95 cm and 133 cm

respectively; Fig 4). Moreover, in-forest sites were exposed (i.e. not flooded) for 4-5

h per day during neap tides, and 2-3 h per day during spring tides, while sites on the

edge were always submerged. Sites could be classified into three groups according

to depth profiles (Fig 4): deep edge sites (sites 3, 5, 7; maximum depth 133 cm);

shallow edge sites (sites 1 and 9; maximum depth: 107 cm); in-forest sites (sites 2,

4, 6, 8; maximum depth: 95 cm; Fig 4).

Fig 4. Site-specific cumulative depth frequencies. Each colour represents a
paired edge and in-forest site, and edge sites are represented by dashed lines and
in-forest sites by solid lines. The dashed dark line indicates the mean cumulative
depth frequencies across all edge sites. The solid black line indicates the mean
cumulative depth frequencies across all in-forest sites. An example is provided to
help interpret the figure (for 25 % of the recorded time depth was on average equal
or below 37 cm at edge sites).

As the GLMM showed that depth had a significant effect on presence/absence

of taxa, a GAMM was used to further explore the response of frequencies of

occurrence of fish across increasing and decreasing depth (equivalent to flooding

and ebbing tide) and by habitat. SQRT frequencies of occurrence of fish significantly

varied across depth (GAMM: $F = 6.756$; $p < 0.001$; Appendix S3), and significantly

differed between "Habitat" (GAMM: $F = 39.792$; $p < 0.001$) and "Tide direction"

(GAMM: $F = 9.056$; $p < 0.005$). Magnitude of variations in SQRT frequencies of

occurrence across depth was higher at edge than in-forest sites (Figs 5a and 5b,

Appendix S3). However, the patterns were similar between the two habitats, with

overall frequencies of occurrence highest at mid-tide, especially mid-ebb tide, and

lowest at extreme depth values (low or high tide; Figs 5a and 5b, Appendix S3).

Fig 5. Boxplots of average square root transformed frequency of occurrence of
common taxa across depth on a) edge sites; b) in-forest sites. The blue line is
the LOESS curve representing the general pattern of habitat use for all common taxa
considered. Shaded area around the LOESS curve represents the 95% confidence
interval. On Fig 5b, interval 90-80 cm has been removed as no data were recorded.

Similarities in mangrove forest utilisation among common taxa clearly

determined 4 main patterns of utilisation: 1) taxa with higher frequencies of

occurrence at high tide (90-130 cm; High-tide users); 2) taxa with higher frequencies

of occurrence at mid-tide (50-90 cm; Mid-tide users); 3) taxa with higher frequencies

at low tide (10-50 cm; Low-tide users); 4) taxa with similar frequencies of occurrence

across depth (Generalist users); Fig 6, Appendix S4).

Fig 6. Patterns of mangrove habitat utilisation across the depth profile. The
curves represent the LOESS curves constructed with the square root transformed
frequencies of occurrence of fish across depth. Common taxa followed four main
patterns of mangrove habitat utilisation across depth: 1) taxa using mangrove
habitats at high tide (High-tide users); 2) taxa using mangrove habitats at mid-tide
(Mid-tide users); 3) taxa accessing and leaving mangrove habitats at low tide (Low-
tide users); 4) taxa without any apparent preferences for depth (Generalist users).
Table 1 identifies the taxa allocated to each category.

358 Discussion

Understanding the spatial and temporal variations in the use of mangrove

habitats by fish is important when considering conservation and resource

management to protect mangrove ecosystems from human and natural

disturbances. This study highlights that the edge and inside of mangrove forests, the

two major component habitats of mangrove forests, featured distinct taxonomic

diversity and fish assemblage composition. Fish assemblages varied significantly

across the tidal cycle, with species-specific patterns of mangrove habitat utilisation.

Spatial differences in water depth among sites within a same habitat also seemed to

influence fish assemblages across this mangrove/coral system. While only a small

portion of the species observed on nearby coral reefs were recorded in Bourake, we

found that this mangrove forest does have a role in supporting reef fish species,

emphasising the importance of Indo-Pacific mangroves as valuable fish habitats.

The setting of this tropical mangrove/coral system influenced the nature of the

fish assemblages recorded. At least 72 taxa made use of this relatively small

mangrove/coral system, with most species classified as tropical marine and reef-

associated [41]. Most taxa recorded have not been identified as mangrove-

associated in previous studies in coastal mangroves in the west Pacific [15, 16, 42],

suggesting that their presence is linked to the proximity of coral reefs, further

supporting the contention that mangrove-coral habitats are interconnected.

Conversely, many fish families important in other Indo-Pacific mangrove systems

remote from coral reefs, such as Leiognatidae, Ambassidae, Sillaginidae,

Terapontidae, or Toxotidae [15, 16, 28, 43] were not recorded in Bouraké. Most

individuals observed were at a sub-adult stage, however, juveniles where

occasionally recorded for several taxa. Juveniles of at least 12 reef fish species were

commonly recorded (*Lutjanus fulviflamma*, *Lutjanus argentimaculatus, Lutjanus*

russellii, *Lethrinus* spp. (2 species), *Bolbometopon muriculatum*, *Siganus lineatus*,

Caranx sp., *Epinephelus caeruleopunctatus*, *Scarus* sp., *Acanthurus* sp.,

Neopomacentrus sp.). Additionally, relatively small individuals of *Epinephelus*

lanceolatus and *Acanthopagrus akazakii* were observed. Thus, the fish community

using this system consisted of a substantial number of juvenile reef species,

including juveniles of two species classified as vulnerable on the IUCN list (*E.*

lanceolatus and *B. muricatum*), and one endemic species (*A. akazakii*) [44]. These

findings highlight that near-coral mangrove habitats in the Central Indo-Pacific, such

as Bouraké, have a role in providing habitats for juvenile reef fish in parallel to the

situation in the Tropical Atlantic [12, 45, 46].

While early studies concluded that high connectivity between coral reefs and

mangroves had little influence on mangrove fish assemblages [18, 28, 47], recent

evidence suggest that in many instances there is a high occurrence of reef-

associated fish in mangroves adjacent to reefs [10, 19, 20, 48]. While supporting this

idea, the current study emphasises that the utilisation and value of mangrove forests

vary locally and cannot be generalised from one system to another [10, 15, 21].

This study highlighted clear spatial variations in fish assemblages across the

two different habitats mangrove edge and mangrove in-forest. Indeed, fish

assemblages were distinctly different between the mangrove edge and just a few

meters inside the mangrove forest. Most fish were recorded cruising on the edge of

the mangrove forest, while sightings inside the mangrove forest were sparser. Two

main hypotheses, namely increased food supply and providing shelter, have been

suggested to explain why fish use mangrove forests. However, neither of these two

hypotheses were confirmed by the current study as few foraging activities were

recorded and few individuals were observed actively sheltering among mangrove

prop-roots. In fact, few species made regular use of the mangrove forest, supporting

the idea that most fish species simply remain on the edge and potentially retreat into

the forest for opportunistic feeding, or to escape presence of larger predators [7].

This result aligns with the observations in estuarine mangrove forests of northern

Australia where few species made regular use of the in-forest habitat [15]. These two

habitats (edge and in-forest) probably confer different values to fish, however, fish

could benefit from most attributes that physically attract them in mangrove systems

[8] by using the mangrove fringe without venturing into the forest. This result

supports the idea that high tidal range leading to forest drainage limits the use of

mangrove forests in the Indo-Pacific compare to the Caribbean. Accessing the forest

could be disadvantageous because of increased risk of becoming trapped after the

tide falls, but could also be linked to adverse water quality such as low dissolved

oxygen that develops at low tide [49].

Fish assemblages exhibited small-scale spatial (dozens of meters)

heterogeneity, particularly along the forest edge compared to in-forest sites. There

was a clear distinction in terms of fish assemblages in the nMDS plot between sites

1 and 9, and sites 3, 5 and 7. This pattern could be explained by water depth profile

and substrate differences, with sites 1 and 9 featuring dead corals, small live coral

boulders and sand, and shallow depth, while other edge sites also had dead corals,

along with small and larger live coral boulders but lacked sand, and experienced

deeper depth. Conversely, all the in-forest sites were quite similar in terms of fish

assemblages, suggesting that they provide a homogeneous habitat with similar

substrate and depth profile throughout the system. Johnston and Sheaves (2007)

[50] also identified species-specific responses to different small-scale habitats

according to their depth and substrate composition. The importance of accounting for

spatial heterogeneity of fish assemblages when characterising the habitat value of a

system, or when using fish assemblages as a bio-indicator of ecological change or

ecosystem health [51], was highlighted by Becker et al. (2012) [52] who observed

the influence of small spatial scale changes in water depth and substrate

composition on fish assemblages at seagrass beds in South Africa.

Fish assemblages varied temporally across the tidal cycle. Tide-induced depth

variations have been linked to changes in fish assemblages [26, 28, 52, 53]. This

result was corroborated here as fish assemblages varied across depth, with more

fish observed during mid-tides, especially at mid- ebbing tides, and most species

generally avoiding extreme shallow or deep water. In fact, fish displayed species-

specific responses to depth with four main patterns identified: 1) taxa using

mangrove habitats at high tide (High-tide users); 2) taxa using mangrove habitats at

mid-tide (Mid-tide users); 3) taxa accessing and leaving mangrove habitats at low

tide (Low-tide users); 4) taxa without any apparent preferences for depth (Generalist

users). Patterns 3 and 4 mainly comprised taxa that frequently used mangrove

habitats such as *Fibramia lateralis*, *Lutjanus argentimaculatus*, *Siganus lineatus*,

Gerres oyena, or taxa belonging to the Gobiidae family [15], while the other two

profiles comprised mainly marine and reef-associated species. In essence, rather

than accessing mangrove habitats as soon as they become available, many species

seem to use mangrove habitats only for a restricted period of time. Other studies that

looked at variations in fish assemblages across the tidal cycle also reported species-

specific responses to the depth profile and highlighted that species using mangrove

habitats extensively were accessing them at a shallower depth than other less

frequently observed species [26, 52-56]. Factors driving these tidal migrations are

not fully understood, and the fact that species do not enter mangrove habitats as

soon as they become available may suggest that these patterns could be the result

of behavioural adaptations to avoid adverse water conditions such as low dissolved

oxygen that can occur early or late in the tide [49]. Species using extensively

mangrove habitats could be adapted to tolerate lower depth and adverse dissolved

oxygen conditions compare to species that would occasionally use mangrove

habitats when they are more suitable. More studies are needed to link tidal fish

migrations with dissolved oxygen conditions in mangrove habitats because dissolved

oxygen is likely to be a critical environmental factor determining the value of these

habitats.

Lunar phase was another influential factor responsible for temporal variations

in mangrove habitats utilisation by fish. More fish were detected during neap tide

than spring tide, however, taxonomic richness and fish assemblage composition

were similar. These data oppose previous studies that observed more fish at spring

tide than neap tide [57-59]. These authors suggest that spring tides result in more

habitats available and for longer duration, attracting more fish. We firstly thought this

was an artefact of the methodology, with fish disappearing from the field of view as

water became too deep. However, we compared fish occurrence within the same

depth intervals between neap and spring tides, and fish presence was still

substantially lower during spring tides, which suggests that there may be another

explanation. One explanation could be that at spring tides fish can access more

intertidal habitats, reducing the probability of encounter with the UVCs. We also

observed very strong currents in the channel and along the mangrove edge during

spring tides that could reduce the time fish can benefit from using mangrove habitats

as the energy needed to remain on the mangrove edge may be too high.

483 Conclusion

The results here provide further support that within a mangrove forest, the

inside and edge of the forest are two distinct habitats characterised by different fish

assemblages. The study mangrove forest plays a role in maintaining a substantial

number of fish species. However, the habitats use was species-specific, suggesting

that utilisation and value need to be considered species-by-species if we want to fully

understand the role mangrove systems play in maintaining fish communities. The

high spatial and temporal heterogeneity of fish assemblages complicates the

characterisation of mangrove forests value and utilisation, suggesting that results

from one location are unlikely to be applicable to other systems more broadly. This is

an important conclusion for managers when considering to adapt conservation

strategies from other locations, to local-specific habitat mosaics.

495 Acknowledgements

496 We thank Dr. Adrien Jacotot, Dr. Thanh Nho Nguyen and Mrs Clara Hass for their

497 assistance in the field. We greatly appreciated suggestions from members of the

498 Science for Integrated Coastal Ecosystem Management group to improve the

499 manuscript.

500 References

1.   Sheaves M. Nature and consequences of biological connectivity in mangrove
systems. Mar Ecol Prog Ser. 2005;302:293-305.
2.   Hogarth PJ. The Biology of Mangroves: Oxford University Press; 1999.
3.   Nagelkerken I, Blaber SJM, Bouillon S, Green P, Haywood M, Kirton LG, et al.
The habitat function of mangroves for terrestrial and marine fauna: A review. Aquat
Bot. 2008;89(2):155-85.
4.   Vance DJ, Haywood MDE, Heales DS, Kenyon RA, Loneragan NR, Pendrey
RC. How far do prawns and fish move into mangroves? Distribution of juvenile
banana prawns *Penaeus merguiensis* and fish in a tropical mangrove forest in
northern Australia. Mar Ecol Prog Ser. 1996;131:115-24.
5.   Nagelkerken I, Faunce CH. Colonisation of artificial mangroves by reef fishes
in a marine seascape. Estuar Coast Shelf Sci. 2007;75(3):417-22.
6.   Bell JD, Pollard DA, Burchmore JJ, Pease BC, Middleton MJ. Structure of a
fish community in a temperate tidal mangrove creek in Botany Bay, New South
Wales. Mar Freshw Res. 1984;35(1):33-46.

7.   Laegdsgaard P, Johnson C. Why do juvenile fish utilise mangrove habitats? J
Exp Mar Biol Ecol. 2001;257(2):229-53.
8.   Nagelkerken I, Faunce CH. What makes mangroves attractive to fish? Use of
artificial units to test the influence of water depth, cross-shelf location, and presence
of root structure. Estuar Coast Shelf Sci. 2008;79(3):559-65.
9.   Beck MW, Heck KL, Able KW, Childers DL, Eggleston DB, Gillanders BM, et
al. The identification, conservation, and management of estuarine and marine
nurseries for fish and invertebrates. Bioscience. 2001;51(8):633-41.
10.   Dorenbosch M, Verberk W, Nagelkerken I, van der Velde G. Influence of
habitat configuration on connectivity between fish assemblages of Caribbean
seagrass beds, mangroves and coral reefs. Mar Ecol Prog Ser. 2007;334:103-16.
11.   Dorenbosch M, Grol MGG, Christianen MJA, Nagelkerken I, van der Velde G.
Indo-Pacific seagrass beds and mangroves contribute to fish density coral and
diversity on adjacent reefs. Mar Ecol Prog Ser. 2005;302:63-76.
12.   Mumby PJ, Edwards AJ, Arias-Gonzalez JE, Lindeman KC, Blackwell PG,
Gall A, et al. Mangroves enhance the biomass of coral reef fish communities in the
Caribbean. Nature. 2004;427(6974):533-6.
13.   Nagelkerken I, van der Velde G, Gorissen MW, Meijer GJ, van't Hof T, den
Hartog C. Importance of mangroves, seagrass beds and the shallow coral reef as a
nursery for important coral reef fishes, using a visual census technique. Estuar Coast
Shelf Sci. 2000;51(1):31-44.
14.   Lal PN. Coastal fisheries and the management of mangrove resources in Fiji.
SPC Fisheries Newsletter. 1984;31(15-23):15-23.
15.   Sheaves M, Johnston R, Baker R. Use of mangroves by fish: new insights
from in-forest videos. Mar Ecol Prog Ser. 2016;549:167-82.
16.   Robertson AI, Duke NC. Mangrove fish-communities in tropical Queensland,
Australia: Spatial and temporal patterns in densities, biomass and community
structure. Mar Biol. 1990;104(3):369-79.
17.   Paillon C, Wantiez L, Kulbicki M, Labonne M, Vigliola L. Extent of mangrove
nursery habitats determines the geographic distribution of a coral reef fish in a
South-Pacific archipelago. PLoS One. 2014;9(8):e105158.
18.   Thollot P. Importance des mangroves pour la faune ichtyologique des récifs
coralliens de Nouvelle-Calédonie. Cybium. 1992;16(4):331-44.
19.   Barnes L, Bellwood DR, Sheaves M, Tanner JK. The use of clear-water non-
estuarine mangroves by reef fishes on the Great Barrier Reef. Mar Biol.
2012;159(1):211-20.
20.   Unsworth RKF, Garrard SL, Salinas De León P, Cullen LC, Smith DJ, Sloman
KA, et al. Structuring of Indo-Pacific fish assemblages along the mangrove–seagrass
continuum. Aquat Biol. 2009;5(1):85-95.
21.   Unsworth RKF, De Leon PS, Garrard SL, Jompa J, Smith DJ. High
connectivity of Indo-Pacific seagrass fish assemblages with mangrove and coral reef
habitats. Mar Ecol Prog Ser. 2008;353:213-24.
22.   Sheaves M, Johnston R. Influence of marine and freshwater connectivity on
the dynamics of subtropical estuarine wetland fish metapopulations. Mar Ecol Prog
Ser. 2008;357:225-43.
23.   Igulu MM, Nagelkerken I, Dorenbosch M, Grol MGG, Harborne AR. Mangrove
habitat use by juvenile reef fish: meta-analysis reveals that tidal regime matters more
than biogeographic region. PloS one. 2014;9(12):e114715.

24.   Baker R, Sheaves M, Johnston R. Geographic variation in mangrove flooding
and accessibility for fishes and nektonic crustaceans. Hydrobiologia. 2015;762(1):1-
14.
25.   Unsworth RKF, Bell JJ, Smith DJ. Tidal fish connectivity of reef and sea grass
habitats in the Indo-Pacific. J Mar Biol Assoc UK. 2007;87(05).
26.   Ellis WL, Bell SS. Tidal influence on a fringing mangrove intertidal fish
community as observed by in situ video recording: implications for studies of tidally
migrating nekton. Mar Ecol Prog Ser. 2008;370:207-19.
27.   Meynecke J-O, Poole GC, Werry J, Lee SY. Use of PIT tag and underwater
video recording in assessing estuarine fish movement in a high intertidal mangrove
and salt marsh creek. Estuar Coast Shelf Sci. 2008;79(1):168-78.
28.   Laroche J, Baran E, Rasoanandrasana NB. Temporal patterns in a fish
assemblage of a semiarid mangrove zone in Madagascar. J Fish Biol. 1997;51(1):3-
20.
29.   Krumme U. Patterns in tidal migration of fish in a Brazilian mangrove channel
as revealed by a split-beam echosounder. Fish Res. 2004;70(1):1-15.
30.   Krumme U. Diel and Tidal Movements by Fish and Decapods Linking Tropical
Coastal Ecosystems. In: Nagelkerken I, editor. Ecological Connectivity among
Tropical Coastal Ecosystems. Dordrecht; 2009. p. 271-324.
31.   Rozas LP, Minello TJ. Estimating densities of small fishes and decapod
crustaceans in shallow estuarine habitats: A review of sampling design with focus on
gear selection. Estuaries. 1997;20(1):199-213.
32.   Reis JA, Giarrizzo T, Barros F. Tidal migration and cross-habitat movements
of fish assemblage within a mangrove ecotone. Mar Biol. 2016;163(5):13.
33.   Dunbar K, Baker R, Sheaves M. Effects of forest width on fish use of fringing
mangroves in a highly urbanised tropical estuary. Mar Freshw Res. 2017;68(9):1764-
70.
34.   Kimball ME, Able KW. Tidal Migrations of Intertidal Salt Marsh Creek Nekton
Examined with Underwater Video. Northeast Nat. 2012;19(3):475-86.
35.   Harborne AR, Mumby PJ, Micheli F, Perry CT, Dahlgren CP, Holmes KE, et
al. The Functional Value of Caribbean Coral Reef, Seagrass and Mangrove Habitats
to Ecosystem Processes. Adv Mar Biol. 2006;50:57-189.
36.   Spalding MD, Fox HE, Allen GR, Davidson N, Ferdaña ZA, Finlayson M, et al.
Marine Ecoregions of the World: A Bioregionalization of Coastal and Shelf Areas.
Bioscience. 2007;57(7):573-83.
37.   Bray JR, Curtis JT. An Ordination of the Upland Forest Communities of
Southern Wisconsin. Ecol Monogr. 1957;27(4):325-49.
38.   Clarke K, Gorley RN. Primer v6: User Manual/Tutorial. Plymouth Marine
Laboratory: Plymouth, UK; 2006.
39.   Knudson C, Geyer CJ, Benson S. Generalized Linear Mixed Models via
Monte Carlo Likelihood Approximation. R Package Version. 1.3.0. 2018.
40.   Wood SN. Generalized Additive Models: An Introduction With R: Chapman
and Hall/CRC; 2006. 391 p.
41.   Froese R, Pauly D. Fishbase 2017 [cited 2017 10 April]. Available from:
www.fishbase.org.
42.   Sheaves M, Johnston R, Abrantes K. Fish fauna of dry tropical and
subtropical estuarine floodplain wetlands. Mar Freshw Res. 2007;58(10):931-43.
43.   Thollot P. Les poissons de mangrove de Nouvelle-Calédonie: caractérisation
du peuplement et relations avec les autres communautés ichtyologiques du lagon.
ORSTOM. 1989;52:1-63.

44.   Iwatsuki Y, Kimura S, Yoshino T. A new sparid, *Acanthopagrus akazakii*, from
New Caledonia with notes on nominal species of *Acanthopagrus*. Ichthyol Res.
2006;53(4):406-14.
45.   Sedberry GR, Carter J. The Fish Community of a Shallow Tropical Lagoon in
Belize, Central America. Estuaries. 1993;16(2):198-215.
46.   Nagelkerken I, Roberts C, van der Velde G, Dorenbosch M, van Riel MC,
Cocheret de la Morinière E, et al. How important are mangroves and seagrass beds
for coral-reef fish? The nursery hypothesis tested on an island scale. Mar Ecol Prog
Ser. 2002;244:299-305.
47.   Quinn NJ, Kojis BJ. Does the presence of coral reefs in proximity to a tropical
estuary affect the estuarine fish assemblage? Proc 5th Int Coral Reef Congr.
1985;5:445-50.
48.   Olds AD, Albert S, Maxwell PS, Pitt KA, Connolly RM. Mangrove-reef
connectivity promotes the effectiveness of marine reserves across the western
Pacific. Global Ecol Biogeogr. 2013;22(9):1040-9.
49.   Mattone C, Sheaves M. Patterns, drivers and implications of dissolved oxygen
dynamics in tropical mangrove forests. Estuar Coast Shelf Sci. 2017;197:205-13.
50.   Johnston R, Sheaves M. Small fish and crustaceans demonstrate a
preference for particular small-scale habitats when mangrove forests are not
accessible. J Exp Mar Biol Ecol. 2007;353(2):164-79.
51.   Whitfield A, Elliott M. Fishes as indicators of environmental and ecological
changes within estuaries: A review of progress and some suggestions for the future.
J Fish Biol. 2002;61:229-50.
52.   Becker A, Coppinger C, Whitfield AK. Influence of tides on assemblages and
behaviour of fishes associated with shallow seagrass edges and bare sand. Mar
Ecol Prog Ser. 2012;456:187-99.
53.   Harborne AR, Talwar B, Brooks EJ. The conservation implications of spatial
and temporal variability in the diurnal use of Bahamian tidal mangrove creeks by
transient predatory fishes. Aquat Conserv: Mar Freshwat Ecosyst. 2016;26(1):202-
11.
54.   Reis-Filho JA, Giarrizzo T, Barros F. Tidal migration and cross-habitat
movements of fish assemblage within a mangrove ecotone. Mar Biol.
2016;163(5):111.
55.   Castellanos-Galindo GA, Krumme U, Willis TJ. Tidal influences on fish
distributions on tropical eastern Pacific rocky shores (Colombia). Mar Ecol Prog Ser.
2010;416:241-54.
56.   Bretsch K, Allen DM. Tidal migrations of nekton in salt marsh intertidal creeks.
Estuar Coasts. 2006;29(3):474-86.
57.   Hampel H, Cattrijsse A, Vincx M. Tidal, diel and semi-lunar changes in the
faunal assemblage of an intertidal salt marsh creek. Estuar Coast Shelf Sci.
2003;56(3-4):795-805.
58.   Krumme U, Saint-Paul U, Rosenthal H. Tidal and diel changes in the structure
of a nekton assemblage in small intertidal creeks in northern Brazil. Aquat Living
Resour. 2004;17:215-29.
59.   Ramos JAA, Barletta M, Dantas DV, Lima ARA, Costa MF. Influence of moon
phase on fish assemblages in estuarine mangrove tidal creeks. J Fish Biol.
2011;78(1):344-54.
661

Supporting information

S1 Appendix. Presence/absence of all common taxa for each 5-minute interval
recorded during the study.

S2 Appendix. Percentage of 5-minutes intervals with no common taxa
observed at a) neap tide vs spring tide and b) edge vs in-forest habitats.

S3 Appendix. General Additive Mixed Model for a) edge sites and b) in-forest
sites.

S4 Appendix. Similarities in species-specific patterns of mangrove utilisation
across depth: a) High-tide users; b) Mid-tide users; c) Mid-tide users (continue)
d) Low-tide users; e) Generalist users.

Transform: Square root
Resemblance: S17 Bray Curtis similarity

Neap edge
Neap in-forest
Spring edge
Spring in-forest

Similarity
——— 45
········· 40

Cryptocentrus leptocephalus

Gobiidae spp.
Fibramia lateralis

Lethrinus harak

Mulloidichthys flavolineatus

Parupeneus indicus

Scarus sp

Gerres oyena

Chaetodon bennetti

Monodactylus argenteus

Chaetodon vagabundus

Neopomacentrus spp.

2D Stress: 0.1