Patterns of fish utilisation in a tropical Indo-Pacific mangrove-coral seascape, New

Caledonia

Fish utilisation of a mangrove forest in New Caledonia

Authors : Dubuc, A[1,2],*, Waltham, N.[1,2], Baker, R.[3], Marchand, C.[4], Sheaves M. [1,2]

[1] College of Science and Engineering, James Cook University, Townsville, Queensland, Australia

[2] TropWATER (Centre for Tropical Water & Aquatic Ecosystem Research), James Cook University,

Townsville, Queensland, Australia

[3] Department of Marine Sciences, Dauphin Island Sea Lab, University of South Alabama, Dauphin

Island Sea Lab, Alabama, United Sates of America

[4] IMPMC, Institut de Recherche pour le Développement (IRD), UPMC, CNRS, MNHN, Nouméa, New

Caledonia, France

* Corresponding author

E-mail: alexia.dubuc1@my.jcu.edu.au (AD)

Abstract (300 words max)

Mangrove forests are important habitats for fish. However, their utilisation by fish, and

the specific values they confer, are still not fully understood. This study details how

fish use mangrove forests in an Indo-Pacific mangrove-coral reef seascape. Sampling

was conducted using underwater video cameras (UVCs) to describe spatial and

temporal variations in fish assemblages across a small-scale (~ 2.5 km^2) system, and

over the tidal and lunar cycle. UVCs were deployed in the two main component

habitats of mangrove forests: at the mangrove forest edge, and inside the forest (5 m

from the forest edge). The patterns of utilisation of fish were established across the

tidal and lunar cycle. Proximity to coral reefs had a strong influence on the mangrove

fish community, as most fish recorded were reef-associated. Juveniles of 12 reef

species were observed, including two species classified as vulnerable on the IUCN list and one endemic species. Fish assemblages on the mangrove edge differed substantially from those inside the forest. Most fish utilised the forest edge with few species making regular use of in- forest habitats, supporting the contention that most fish species remain on the edge and potentially retreat into the forest for opportunistic feeding, or when threatened by larger predators. Species-specific patterns of utilisation varied across the tidal and lunar cycle. Small differences in depth profiles and substrate across the small-scale system had a substantial effect on fish assemblages, highlighting the importance of accounting for spatial heterogeneity in these factors. These data provide important information for managers to implement adequate conservation strategies that includes broader interconnected habitat mosaics.

Introduction

Mangrove forests are a fundamental component habitat of mangrove systems [1], and confer many of the attributes that attract juvenile fish [2, 3]. However, recent work in mesotidal Australia suggests that few fish there penetrate beyond the forest boundary [4]. This brings into question the specific ways in which mangrove forests are utilised and indicates the need for studies to characterise fish assemblages at locations with different tidal ranges and contexts, and to investigate how widespread such patterns of utilisation of mangrove forests are.

At a broader scale, studies have found that some mangrove forests, especially in the Indo-Pacific, seem to be used less intensively as nursery or feeding grounds by reef fish [4-6] than in the Caribbean [7-9]. These observations suggest that not all mangrove forests provide equivalent functions and services. This dissimilarity has

been attributed to differences in the settings in which mangrove forests occur (coastal,

estuarine, island, embayment), different tidal ranges (micro-, meso- or macrotidal),

proximity of other high value habitats such as seagrass beds and coral reefs, or

climatic zones [10-13]. Comparing fish assemblages between the Indo-Pacific and the

Caribbean in mangrove habitats with similar micro-tidal settings would provide crucial

information on how those factors influence the value and functioning of mangrove

habitats.

Intertidal mangrove forests are challenging environments, most notably

because they are only available to most aquatic organisms while they are flooded at

high tide [13-15]. In many parts of the Indo-Pacific, the tidal range is greater than in

the Caribbean, where mangrove forests are usually continually available to fish [2].

The intermittent availability of mangrove forests may explain the low use by fish in the

Indo-Pacific [11]. Tidal variation (extent, duration and frequency of flooding) generates

a range of constraints for fish utilising mangrove forests. Most obvious is the decrease

in water depth and eventual drainage of the forest as the tide ebbs. Several studies

have demonstrated that fish undertake regular migrations in tidally driven mangrove

systems, with different patterns of mangrove use according to fish species, lunar cycle

(neap vs spring tide) and tidal phase (flooding vs ebbing) [4, 16-19]. Migration of fish

in response to tidal movements results in substantial connectivity between the three

major tropical coastal habitats: coral reefs, seagrass beds and mangrove forests [14,

20], giving rise to the idea that mangrove forests are part of a wider interconnected

habitat mosaic [15]. Therefore, investigating tidal and spatial variations in fish

assemblages in mangrove forests is a crucial step towards fully appreciating the value

and functioning of the whole tropical coastal ecosystem.

The difficulty of sampling these habitats goes a long way to explaining the

paucity of information available on fish assemblages inside mangrove forests [21, 22].

The use of conventional techniques such as underwater visual censuses or netting

techniques is restricted across much of the Australasian region where saltwater

crocodiles (*Crocodylus porosus*) are common, and where dense mangrove forests

reduce the efficiency of most net-based approaches [4]. Recently, underwater video

has been successfully applied to study in-forest fish assemblages [4, 23], most notably

because it overcomes a lot of sampling issues, substantially reduces field labour

intensity, and allows for high -temporal and -spatial resolution data collection

simultaneously in different habitats, such as the edge and the inside of a mangrove

forest [24].

We used underwater cameras on the edge and inside a mangrove forest [4,

16], coupled with high frequency depth loggers, to examine how fish use mangrove

forests in a micro-tidal Indo-Pacific mangrove-coral reef seascape. In particularly we

assessed the fish species that used the mangrove forest, how their utilisation changed

between the forest edge and in-forest habitats, and how fish assemblages changed

across the tidal cycle.

Materials and methods

Study site

Our study focused on a relatively pristine mangrove forest in Bouraké, South

Province of New Caledonia (Fig 1). New Caledonia experiences a semi-arid to tropical

climate with annual total rainfall of 1,000 mm, and a mixed semi-diurnal mircrotidal

regime (maximum 1.8 m tidal range). Bouraké receives little freshwater inflow with no

defined drainage.

The area comprises approximately 2.5 km² of mangrove forest dominated by

Rhizophora stylosa on the seaward edge and *Avicennia officinalis* on the landward

margin, with a large semi-enclosed central lagoon (1.2 km long, 60 m wide, 1-2 m

depth). A channel (20-70 m wide, 2-6 m depth, 700 m long) connects the main lagoon

to the coastal waters of Pritzbuer Bay (~ 20 km²). The channel comprises two

sheltered inlets (approximately 0.01 km² each), and a shallow (1-2 m depth) coral reef

platform that extends from the middle of the channel to the edge of the mangrove

forest. Coral reefs occur up to a first few meters into the forest in some places (Fig 1).

We chose a mangrove-coral reef seascape, a setting comparable to the one in which

mangrove forests occur in the Caribbean, so we could relate our findings to this other

ecoregion [25] where many studies have been conducted.

Fig 1. Map and picture of the study system in Bouraké, South Province, New
Caledonia. The nine study sites in the mangrove channel sampled from the 21
February to 1 March 2017 are represented by their respective number. Light grey
areas represent mangrove forest, dark grey areas represent mainland, and white
areas represent water.

Data collection

The fish community using the mangrove forest was examined at nine sites on

an inland/offshore gradient along the channel (Fig 1). To assess differences in fish

assemblage composition between edge and inside the forest, sites 1 to 8 comprised

4 paired sites where each odd site number was located on the mangrove forest edge,

and even site numbers were located approximately 5 m inside the mangrove forest.

Site 9 (considered an edge site in the analyses) was located on the reef platform of

the innermost bay, at the edge of scattered mangrove trees slightly away from the

main forest. The substrate at sites 1 and 9 consisted of dead corals, small live coral

boulders and sand, while on other edge sites it comprised mainly dead corals and

small and larger live coral boulders. The substrate was homogeneous and consisted

of silt material at in-forest sites.

Fish assemblages were sampled using underwater video cameras (UVCs;

Model ATC9K Oregon Scientific) to investigate tidal variations in fish assemblages

simultaneously on the edge and inside the forest. Unbaited UVCs mounted on stable

bases were deployed at each site during neap (21 to 23 February 2017) and spring

tides (28 February to 1 March 2017). A sampling day consisted of cameras first

deployed at sites early in the morning (first light), continuing until the battery was fully

discharged, and again deployed mid-afternoon at all sites until the battery was again

fully discharged (recording lasted between 2h and 2.5h). A total of four sampling days

were completed (two during neap tides and two during spring tides). Cameras were

positioned around 7 cm above the substrate, facing towards the channel. A marker

mounted on a flexible rod (3 mm diameter, 0.5 m long) was placed 0.5 m in front of

the camera lens as a visibility indicator to ensure a minimum visibility of 0.5 m was

achieved in all videos. Visibility was very good during the sampling period, and fish

could be identified confidently up to approximately 2 m from the UVCs in all videos.

As depth is one of the main limiting factors to mangrove accessibility, tidal variations

in water depth (cm) were measured every 15 minutes at each site with depth loggers

(In-Situ Inc. Rugged Troll 100 model).

Data analyses

While UVCs allow large amounts of data to be gathered quickly in the field,

considerable time is required to process these videos. Therefore, we subsampled the

acquired video footage. From the two neap tide sampling days, one day was randomly

selected and videos at all sites were processed for that day. For the remaining

sampling day, all videos were processed from five sites; being the reef platform (site

9) and two pairs of in-forest and forest edge sites (sites 5-8). The same selection was

applied to the two sampling days conducted during spring tides.

Once sediments had settled after camera deployment (typically 2-3 min), videos

were viewed using VLC media player (VideoLAN, 2001) and subdivided in 5-min

intervals to follow tidal variations in fish assemblages. The occurrence of each fish

taxon in each 5-min interval was recorded. Only presence/absence data were

recorded to avoid biases induced by count data when using UVCs [4]. Fish were

identified to the lowest possible taxonomic level. Features useful in discriminating

species within some genera or families such as *Plectorhinchus* spp.*,* Mugilidae spp.,

or Gobiidae spp. were difficult to distinguish in videos, therefore these taxa were

identified to genus or family level only. Where possible, juvenile fish were identified

based on colour patterns and body shape. Any activity such as feeding, hiding,

cruising or escaping was also noted. All fish identifications were validated by two

additional experts.

The frequency of occurrence of each taxon was calculated per site (the total

number of 5-min intervals in which a taxon was observed at a site was divided by the

total number of 5-min intervals recorded at this site). Only taxa with a frequency of

occurrence ≥ 0.05 at one or more sites were retained for analyses (referred to as

"common taxa"). Taxa with a frequency of occurrence < 0.05 (referred to as "rare taxa")

were excluded from analyses.

Frequency of occurrence of each common taxon in-forest and on the edge was

calculated and plotted using horizontal bar plots to investigate differences graphically.

The same methodology was followed to investigate differences in frequency of

occurrence of fish between neap and spring tides. As fewer fish were observed during

spring tides, a logistic regression was performed in R using the presence of any

common taxa for each 5-min interval as the dependent variable (1 if any common taxa

were observed in the 5-min interval, or 0 if no common taxa were observed), and

"Lunar phase" (neap vs spring) as the independent variable, to determine whether

"Lunar phase" had a significant effect on the presence/absence of taxa.

Depth cumulative time curves were plotted for each site to highlight differences

in temporal dynamics. Patterns of mangrove habitat use over flooding and ebbing tide

on the edge and in-forest were investigated graphically for each common taxon. Each

5-min interval was allocated to a class of water depth of 10 cm (from 10-20 cm to 120-

130 cm) according to the water depth value recorded. The mean frequency of

occurrence of each common taxon was then calculated per class of water depth during

flooding and ebbing tide. Boxplots were used to represent the average square root

transformed frequency of occurrence for each common taxon at each depth interval,

and a LOESS curve was used to analyse the general pattern of habitat use. To avoid

false absence recordings, only taxa recorded in the habitat of interest were

considered. Frequencies of occurrence were square root transformed to reduce the

impact of extreme values.

Non-metric multidimensional scaling (nMDS) was used to compare fish

assemblages among sites during spring and neap tides. The frequency of occurrence

of each common taxon was calculated per site per lunar phase. Data were square root

transformed to decrease the impact of extreme values, and an nMDS analysis based

on Bray-Curtis dissimilarities, the most appropriate distance measure when using

abundance data [26], was performed using the package "vegan" in R [27]. Groupings

within the nMDS were determined using similarity profile tests (SIMPROF) based on

hierarchical clustering at a significance level of 95 %.

Results

200 　　　Fifty-six video deployments were processed (totalling more than 118h of video).

201 Seventy-two taxa from 29 families were recorded, with 36 common taxa (frequency of

202 occurrence ≥ 0.05) retained for further statistical analyses (Table 1). Most species

203 recorded were marine and reef-associated [28] including fish of families Scaridae,

204 Chaetodontidae, Pomacanthidae, Siganidae, Acanthuridae, Lutjanidae, or Labridae.

205 Table 1. Summary of all the families and taxa identified at Bouraké, New
206 Caledonia.

Family	Taxon
Acanthuridae	*Acanthurus auranticavus** [1]
	Acanthurus grammoptilus [2]
	Acanthurus sp. cf blochii
	Ctenochaetus sp.
	Zebrasoma velifer
Apogonidae	*Fibramia lateralis** [4]
	Ostorhinchus septemstriatus
Belonidae	*Belonidae spp.*
Blenniidae	*Blenniidae spp.*
Carangidae	*Caranx ignobilis*
	Caranx papuensis [1]
	Caranx sp.
Chaetodontidae	*Chaetodon auriga** [3]
	Chaetodon bennetti [2]
	Chaetodon ephippium
	Chaetodon flavirostris
	Chaetodon lineolatus [2]
	Chaetodon lunula [2]
	Chaetodon melannotus
	Chaetodon speculum
	Chaetodon vagabundus [2]
	Heniochus acuminatus [1]

Table 1 (continued)

Family	Taxon
Haemulidae	*Plectorhinchus lineatus*
	Plectorhinchus spp. [2]
	Pomadasys argenteus [2]
Hemiramphidae	*Hyporhamphus sp.*
Labridae	*Choerodon graphicus*
	Labridae spp.
Lethrinidae	*Lethrinus harak** [1]
	Lethrinus lentjan [1]
	Lethrinus obsoletus
Lutjanidae	*Lutjanus argentimaculatus** [4]
	Lutjanus fulviflamma [3]
	Lutjanus fulvus [3]
	Lutjanus russellii [4]
Monodactylidae	*Monodactylus argenteus** [1]
Mugilidae	*Mugilidae spp.** [4]
Mullidae	*Mulloidichthys flavolineatus* [2]
	Parupeneus ciliatus
	Parupeneus indicus [3]
	Upeneus tragula [1]
Pomacanthidae	*Pomacanthus sexstriatus* [2]
Pomacentridae	*Neopomacentrus spp.* [2]

Clupeidae	*Clupeidae spp.* [2]	Scaridae	<u>*Bolbometopon muricatum*</u>
Diodontidae	*Diodon hystrix*		<u>***Scarus sp. cf ghobban***</u> [1]
Ephippidae	*Platax pinnatus*	Scatophagidae	*Scatophagus argus*
Fistulariidae	*Fistularia spp.*	Serranidae	<u>*Epinephelus caeruleopunctatus*</u>
Gerreidae	*Gerres filamentosus*		<u>*Epinephelus lanceolatus*</u>
	Gerres oyena* [3]		*Epinephelus malabaricus*
Gobiidae	*Amblygobius linki*		*Epinephelus sp.*
	Amblygobius nocturnus	Siganidae	<u>***Siganus canaliculatus***</u> [1]
	Amoya gracilis [4]		<u>***Siganus lineatus****</u> [3]
	Asterropteryx sp. cf striata [3]		*Siganus punctatus*
	Cryptocentrus leptocephalus [4]	Sparidae	<u>***Acanthopagrus sp. cf akazakii***</u> [2]
	Eviota sp.	Sphyraenidae	*Sphyraena barracuda*
	Exyrias puntang	Tetraodontidae	*Arothron hispidus*
	Gobiidae spp.* [4]		
	Gobiidae spp.2		
	Redigobius balteatus [4]		

Taxa highlighted in bold represent the taxa with frequency of occurrence ≥ 0.05 on at least 1 site that were kept for statistical analyses. Taxa highlighted with a * represent the 10 most common taxa. Taxa or families underlined mean that juveniles were potentially observed for that taxon or for at least one taxon within the family that could not be identified. The superscript number corresponds to the pattern of mangrove habitat utilisation across depth followed by the taxon (Fig 5).

Taxonomic richness was lower at in-forest sites compared to edge sites (21 and 34 common taxa recorded respectively; Fig 2a). *Fibramia lateralis,* and all the common taxa belonging to the Gobiidae family, except *Cryptocentrus leptocephalus and Asterropteryx* spp., were frequently observed at in-forest sites compared to edge sites (Fig 2a). *Lutjanus fulviflamma* was the only common taxon recorded evenly in-forest and edge sites. All the other common taxa were more frequently observed at edge sites (Fig 2a).

Fig 2. Proportion of time spent by each of the most common taxa a) on the edge versus inside the forest, and b) at neap tide versus spring tide. On Fig 2a: edge=green and in-forest=brown, and on Fig 2b: neap=deep pink and spring=blue navy Proportions range from 0 to 1, 1 corresponding to a taxon exclusively recorded a) on the edge or in-forest or b) at neap tide or spring tide, and 0.5 corresponding to a taxon recorded a) on the edge as frequently as in-forest or b) at neap tides as frequently as at spring tides.

Lunar phase had a significant impact on presence of common taxa and

frequencies of occurrence. Twenty-five of the 36 common taxa had higher frequency

of occurrence during neap compared to spring tides (Fig 2b). Moreover, the logistic

regression highlighted a significant relationship between lunar phase and fish

presence ($p < 0.001$), with fish observed less frequently during spring than neap tides.

Average depth was substantially shallower at in-forest than edge sites (neap

tides (mean ± SE): 34 ± 0.57 and 55 ± 0.66 cm respectively; spring tides: 48 ± 1.11

and 71 ± 1.23 cm respectively) as was maximum depth (95 cm and 133 cm

respectively; Fig 3). Moreover, in-forest sites were exposed (i.e. not flooded) for 4-5 h

per day during neap tides, and 2-3 h per day during spring tides, while sites on the

edge were always submerged. Sites could be classified into three groups according

to depth profiles (Fig 3): deep edge sites (sites 3, 5, 7; maximum depth 133 cm);

shallow edge sites (sites 1 and 9; maximum depth: 107 cm); in-forest sites (sites 2, 4,

6, 8; maximum depth: 95 cm; Fig 3).

Fig 3. Site-specific cumulative time curves for depth. Each colour represents a
paired edge and in-forest site, and edge sites are represented by dashed lines and
in-forest sites by solid lines. The dashed dark line indicates the mean cumulative
time for depth across all edge sites. The solid black line indicates the mean
cumulative time for depth across all in-forest sites.

Frequencies of occurrence of fish varied across the tidally driven depth profile

(Figs 4a and 4b). Patterns of fish occurrence were quite similar between edge and in-

forest sites with overall frequencies of occurrence highest at mid-tide, especially mid-

ebb tide (Figs 4a and 4b), and lowest at extreme depth values (low or high tide).

Fig 4. Boxplots of average square root transformed frequency of occurrence
for each common taxon across depth on a) edge sites; b) in-forest sites. The
blue line is the LOESS curve representing the general pattern of habitat use for all
common taxa considered. Shaded area around the LOESS curve represents the
95% confidence interval.

We identified 4 main taxa-specific patterns of utilisation across the depth profile:

1) taxa using mangrove habitats at high tide (High tide users); 2) taxa using mangrove

habitats at mid-tide (flood or ebb; Mid-tide users); 3) taxa without any apparent

preferences for depth (Generalist users); 4) taxa accessing and leaving mangrove

habitats at low depth (Shallow water users) (Fig 5).

Fig 5. Taxa-specific patterns of mangrove habitat utilisation across the depth
profile. The curves represent the LOESS curves constructed with the square root
transformed (SQRT) frequency of occurrence of fish across depth. Common taxa
followed four main patterns of mangrove habitat utilisation across depth: 1) taxa
using mangrove habitats at high tide (High tide users); 2) taxa using mangrove
habitats at mid-tide (flood or ebb; Mid-tide users); 3) taxa without any apparent
preferences for depth (Generalist users); 4) taxa accessing and leaving mangrove
habitats at low depth (Shallow water users).

Fish composition varied greatly among sites, with distinct assemblages

generating four different clusters on the nMDS plot (Fig 6). The 1st cluster comprised

all the samples conducted in-forest, the 2nd cluster comprised 5 of the 6 samples

conducted at shallow edge sites (Fig 3), the 3rd cluster comprised all the samples

conducted at site 5, and the 4th cluster comprised 3 of the 4 samples conducted at site

7 and 1 sample conducted at site 3. The samples conducted on the first day (X) and

the ones conducted on the following day (Xrep) during the same lunar phase plotted

close to each other on the nMDS plot and were for most of them grouped in the same

clusters. Three sites were outliers (1, 3 and 7) during spring tides and did not group

into any cluster. This could be explained by the presence of sedentary species

(*Cryptocentrus leptocephalus* at site 1, and *Neopomacentrus sp.* at site 7), thus if the

camera was placed in front of their home, the same individual was most of the time

recorded across the entire video driving the fish assemblage composition

(*Neopomacentrus* spp. and). Lunar phase did not influence fish assemblages given

that the same sites during spring and neap tides grouped together in the same

respective clusters.

Fig 6. nMDS analysis performed on square root transformed frequency of
occurrence for each taxon per site per tide. Edge sites are represented by
squares and in-forest sites by circles. Sites sampled at neap tide are coloured in
deep pink, and sites sampled at spring tide in navy blue. Dotted line ellipses
represent clusters determined using similarity profile tests (SIMPROF) based on
hierarchical clustering at a significance level of 95 %.

Discussion

Mangrove edge and mangrove forest, the two major component habitats of the

Bouraké mangrove system, featured distinct taxonomic diversity and fish assemblage

composition. Fish assemblages varied substantially across the tidal cycle, with

species-specific patterns of mangrove habitat utilisation. Spatial differences in water

depth among sites within a same habitat also seemed to influence fish assemblages

across this mangrove/coral system.

The setting of this tropical mangrove/coral system influenced the nature of the

fish assemblages recorded. At least 72 taxa made use of this relatively small

mangrove/coral system, with most species classified as tropical marine and reef-

associated [28]. Most taxa recorded have not been identified as mangrove-associated

in previous studies in coastal estuarine mangroves in the west Pacific [4, 29, 30],

suggesting that their presence is linked to the proximity of coral reefs, further

supporting the contention that mangrove-coral habitats are interconnected.

Conversely, many families important in other Indo-Pacific mangrove systems remote

from coral reefs, such as Leiognatidae, Ambassidae, Sillaginidae, Terapontidae, or

Toxotidae [4, 18, 29, 31] were not recorded in Bouraké. Juveniles of at least 12 reef

fish species were recorded (*Lutjanus fulviflamma*, *Lutjanus argentimaculatus,*

Lutjanus russellii, *Lethrinus* spp. (2 species), *Bolbometopon muriculatum*, *Siganus*

lineatus, *Caranx* sp., *Epinephelus caeruleopunctatus*, *Scarus* sp., *Acanthurus* sp.,

Neopomacentrus sp.). Additionally, relatively small individuals of *Epinephelus*

lanceolatus and *Acanthopagrus akazakii* were observed. Therefore, juveniles of two

species classified as vulnerable on the IUCN list (*E. lanceolatus* and *B. muricatum*),

and one endemic species (*A. akazakii*) [32] used this system. These findings highlight

that near-coral mangrove habitats in the Central Indo-Pacific, such as Bouraké,

function as important habitats for juvenile reef fish in parallel to the situation in the

Tropical Atlantic [8, 33, 34].

While early studies concluded that high connectivity between coral reefs and

mangroves had little influence on mangrove fish assemblage [5, 18, 35], recent

evidence suggest that in many instances there are high occurrence of reef-associated

fish in mangroves adjacent to reefs [6, 7, 22, 36]. While supporting this idea, the

current study emphasises that the utilisation and value of mangrove systems vary

locally and cannot be generalised without understanding of location-specific factors [4,

7, 12].

Fish assemblages varied temporally across the tidal cycle. Tide-induced depth

variation has been linked to changes in fish assemblages [16, 18, 37]. This result was

corroborated here as fish assemblages varied across depths, with more fish observed

during mid-tides, and most species generally avoiding extreme shallow or deep water.

In fact, fish seem to display species-specific responses to depth with four main

patterns identified: taxa using mangrove habitats at high tide (High tide users); 2) taxa

using mangrove habitats at mid-tide (flood or ebb; mid-tide users); 3) taxa without any

apparent preferences for depth (Generalist users); 4) taxa accessing and leaving

mangrove habitats at low depth (Shallow water users). The first two patterns mainly comprised taxa known to use mangrove habitats extensively such as *Fibramia lateralis*, *Lutjanus argentimaculatus*, *Siganus lineatus*, *Gerres oyena*, or species belonging to the Gobiidae family [4], while the other two profiles comprised mainly marine and reef-associated species. In essence, rather than accessing mangrove habitats as soon as they become available, many species seem to use mangrove habitats only for a restricted period of time later in the tide. Other studies that looked at variations in fish assemblages across the tidal cycle also reported species-specific responses to the depth profile and highlighted that species using mangrove habitats extensively were accessing them at shallower depth than other less frequently observed species [16, 37-41]. These patterns could be the result of behavioural adaptations to avoid adverse DO conditions that can occur early or late in the tide [42]. Species using extensively mangrove habitats must be adapted to tolerate lower depth and DO conditions compare to species occasionally using mangrove habitats when they are more suitable. More studies are needed to link tidal fish migrations with DO conditions in mangrove habitats because DO is likely be a critical environmental factor determining the value of these habitats.

Lunar phase was another influential factor, largely responsible for temporal variations in mangrove utilisation by fish. More fish were detected during neap tide than spring tide, however, taxonomic richness and fish assemblage composition were similar. These data contradict previous studies that observed more fish at spring tide than neap tide [43-45]. These authors suggest that spring tides result in more habitats available and for longer duration, attracting more fish. We firstly thought that this result was an artefact of the methodology, with fish disappearing from the field of view as water became too deep. However, we compared fish abundance within the same

depth intervals between neap and spring tides, and fish abundance was still

significantly lower during spring tides, which suggests that there may be another

explanation. One explanation could be that at spring tides fish can access more

intertidal habitats, reducing the probability of encounter with the UVCs. We also

observed very strong currents in the channel and along the mangrove edge that could

prevent fish from benefiting from using mangrove habitats as the energy needed to

remain on the mangrove edge may be too high.

This study highlighted clear spatial variations in fish assemblages across the

different mangrove habitats. Indeed, fish assemblages were distinctly different

between the mangrove edge and just a few meters inside the mangrove forest. Most

fish were recorded cruising on the edge of the mangrove forest, while sightings inside

the mangrove forest were sparser. The two hypotheses that are usually made to

explain why fish use mangrove forests relate to foraging and finding shelter, however,

neither were confirmed by the current study as few foraging activities were observed

and few individuals were observed actively sheltering among mangrove prop-roots. In

fact, few species made regular use of the mangrove forest, supporting the idea that

most fish species simply remain on the edge and potentially retreat into the forest for

opportunistic feeding, or when threatened by larger predators [46]. This result aligns

with the observations in estuarine mangrove forests of northern Australia where few

species make regular use of the in-forest habitat [4]. These two habitats (edge and in-

forest) probably confer different values to fish, however, fish could benefit from most

attributes that physically attract them in mangrove systems [47] by using the mangrove

fringe, without venturing into the forest. Accessing the forest could be

disadvantageous because of adverse dissolved oxygen [42] or because of increased

risk of stranding.

Fish assemblages exhibited small-scale spatial (ten or so meters)

heterogeneity, particularly along the forest edge compared to in-forest sites. There

was a clear distinction in terms of fish assemblages in the nMDS plot between sites 1

and 9, and sites 3, 5 and 7. This pattern could be explained by water depth profile

differences, or because of differences in substrate, with sites 1 and 9 featuring dead

corals, small live coral boulders and sand, and shallow depth, while other edge sites

also had dead corals, along with small and larger live coral boulders but lacked sand,

and deeper depth. Conversely, all the in-forest sites were quite similar in terms of fish

assemblages, suggesting that they provide a homogeneous habitat with similar

substrate and depth profile throughout the system. Johnston and Sheaves (2007) [48]

also identified species-specific responses to different small-scale habitats according

to their depth and substrate composition. The importance of accounting for spatial

heterogeneity of fish assemblages was highlighted by Becker et al. (2012) [37] who

observed the influence of small spatial scale changes in water depth and substrate

composition on fish assemblages at seagrass beds in South Africa. It is important to

consider small-scale spatial heterogeneity when characterising the habitat value of a

system to fish, or when using fish assemblages as a bio-indicator of ecological change

or ecosystem health [49].

Conclusion

The settings in which mangrove systems occur is important in determining the

values they provide to fish. Within a mangrove system, mangrove forest and mangrove

edge proved to be two distinct habitats from the point of view of fish. The values

reflected in their patterns of utilisation were species-specific, suggesting that utilisation

and value need to be considered species-by-species if we are to fully understand the

role mangrove systems play in maintaining fish communities. The high spatial and

temporal heterogeneity of fish assemblages complicates the characterisation of

mangrove systems value and utilisation, suggesting that simplistic models are unlikely

to be suitable and that care needs to be exercised when transferring understanding to

a new location. These data provide important information for managers to implement

adequate conservation strategies that includes location-specific knowledge and

integration to broader interconnected habitat mosaics.

Acknowledgements

420 We thank Dr. Adrien Jacotot, Dr. Thanh Nho Nguyen and Mrs Clara Hass for their

421 assistance in the field. We greatly appreciated suggestions from members of the

422 Science for Integrated Coastal Ecosystem Management group to improve the

423 manuscript.

References

1.   Vance DJ, Haywood MDE, Heales DS, Kenyon RA, Loneragan NR, Pendrey
RC. How far do prawns and fish move into mangroves? Distribution of juvenile banana
prawns *Penaeus merguiensis* and fish in a tropical mangrove forest in northern
Australia. Mar Ecol Prog Ser. 1996;131:115-24.
2.   Nagelkerken I, Blaber SJM, Bouillon S, Green P, Haywood M, Kirton LG, et al.
The habitat function of mangroves for terrestrial and marine fauna: A review. Aquat
Bot. 2008;89(2):155-85.
3.   Beck MW, Heck KL, Able KW, Childers DL, Eggleston DB, Gillanders BM, et al.
The identification, conservation, and management of estuarine and marine nurseries
for fish and invertebrates. Bioscience. 2001;51(8):633-41.
4.   Sheaves M, Johnston R, Baker R. Use of mangroves by fish: new insights from
in-forest videos. Mar Ecol Prog Ser. 2016;549:167-82.
5.   Thollot P. Importance des mangroves pour la faune ichtyologique des récifs
coralliens de Nouvelle-Calédonie. Cybium. 1992;16(4):331-44.
6.   Barnes L, Bellwood DR, Sheaves M, Tanner JK. The use of clear-water non-
estuarine mangroves by reef fishes on the Great Barrier Reef. Mar Biol.
2012;159(1):211-20.
7.   Dorenbosch M, Verberk W, Nagelkerken I, van der Velde G. Influence of habitat
configuration on connectivity between fish assemblages of Caribbean seagrass beds,
mangroves and coral reefs. Mar Ecol Prog Ser. 2007;334:103-16.

8.   Mumby PJ, Edwards AJ, Arias-Gonzalez JE, Lindeman KC, Blackwell PG, Gall
A, et al. Mangroves enhance the biomass of coral reef fish communities in the
Caribbean. Nature. 2004;427(6974):533-6.
9.   Nagelkerken I, van der Velde G, Gorissen MW, Meijer GJ, van't Hof T, den
Hartog C. Importance of mangroves, seagrass beds and the shallow coral reef as a
nursery for important coral reef fishes, using a visual census technique. Estuar Coast
Shelf Sci. 2000;51(1):31-44.
10.   Sheaves M, Johnston R. Influence of marine and freshwater connectivity on the
dynamics of subtropical estuarine wetland fish metapopulations. Mar Ecol Prog Ser.
2008;357:225-43.
11.   Igulu MM, Nagelkerken I, Dorenbosch M, Grol MGG, Harborne AR. Mangrove
habitat use by juvenile reef fish: meta-analysis reveals that tidal regime matters more
than biogeographic region. PloS one. 2014;9(12):e114715.
12.   Unsworth RKF, De Leon PS, Garrard SL, Jompa J, Smith DJ. High connectivity
of Indo-Pacific seagrass fish assemblages with mangrove and coral reef habitats. Mar
Ecol Prog Ser. 2008;353:213-24.
13.   Baker R, Sheaves M, Johnston R. Geographic variation in mangrove flooding
and accessibility for fishes and nektonic crustaceans. Hydrobiologia. 2015;762(1):1-
14.
14.   Unsworth RKF, Bell JJ, Smith DJ. Tidal fish connectivity of reef and sea grass
habitats in the Indo-Pacific. J Mar Biol Assoc UK. 2007;87(05).
15.   Sheaves M. Nature and consequences of biological connectivity in mangrove
systems. Mar Ecol Prog Ser. 2005;302:293-305.
16.   Ellis WL, Bell SS. Tidal influence on a fringing mangrove intertidal fish
community as observed by in situ video recording: implications for studies of tidally
migrating nekton. Mar Ecol Prog Ser. 2008;370:207-19.
17.   Meynecke J-O, Poole GC, Werry J, Lee SY. Use of PIT tag and underwater
video recording in assessing estuarine fish movement in a high intertidal mangrove
and salt marsh creek. Estuar Coast Shelf Sci. 2008;79(1):168-78.
18.   Laroche J, Baran E, Rasoanandrasana NB. Temporal patterns in a fish
assemblage of a semiarid mangrove zone in Madagascar. J Fish Biol. 1997;51(1):3-
20.
19.   Krumme U. Patterns in tidal migration of fish in a Brazilian mangrove channel
as revealed by a split-beam echosounder. Fish Res. 2004;70(1):1-15.
20.   Krumme U. Diel and Tidal Movements by Fish and Decapods Linking Tropical
Coastal Ecosystems. In: Nagelkerken I, editor. Ecological Connectivity among
Tropical Coastal Ecosystems: SPRINGER; 2009. p. 271-324.
21.   Rozas LP, Minello TJ. Estimating densities of small fishes and decapod
crustaceans in shallow estuarine habitats: A review of sampling design with focus on
gear selection. Estuaries. 1997;20(1):199-213.
22.   Unsworth RKF, Garrard SL, Salinas De León P, Cullen LC, Smith DJ, Sloman
KA, et al. Structuring of Indo-Pacific fish assemblages along the mangrove–seagrass
continuum. Aquat Biol. 2009;5(1):85-95.
23.   Reis JA, Giarrizzo T, Barros F. Tidal migration and cross-habitat movements of
fish assemblage within a mangrove ecotone. Mar Biol. 2016;163(5):13.
24.   Kimball ME, Able KW. Tidal Migrations of Intertidal Salt Marsh Creek Nekton
Examined with Underwater Video. Northeast Nat. 2012;19(3):475-86.
25.   Spalding MD, Fox HE, Allen GR, Davidson N, Ferdaña ZA, Finlayson M, et al.
Marine Ecoregions of the World: A Bioregionalization of Coastal and Shelf Areas.
Bioscience. 2007;57(7):573-83.

26.   Bray JR, Curtis JT. An Ordination of the Upland Forest Communities of
Southern Wisconsin. Ecol Monogr. 1957;27(4):325-49.
27.   Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin P, O'Hara RB, et al.
Vegan: Community Ecology Package. R Package Version. 2.0-10. 2013.
28.   Froese R, Pauly D. Fishbase 2017 [cited 2017 10 April]. Available from:
www.fishbase.org.
29.   Robertson AI, Duke NC. Mangrove fish-communities in tropical Queensland,
Australia: Spatial and temporal patterns in densities, biomass and community
structure. Mar Biol. 1990;104(3):369-79.
30.   Sheaves M, Johnston R, Abrantes K. Fish fauna of dry tropical and subtropical
estuarine floodplain wetlands. Mar Freshw Res. 2007;58(10):931-43.
31.   Thollot P. Les poissons de mangrove de Nouvelle-Calédonie: caractérisation
du peuplement et relations avec les autres communautés ichtyologiques du lagon.
ORSTOM. 1989;52:1-63.
32.   Iwatsuki Y, Kimura S, Yoshino T. A new sparid, *Acanthopagrus akazakii*, from
New Caledonia with notes on nominal species of *Acanthopagrus*. Ichthyol Res.
2006;53(4):406-14.
33.   Sedberry GR, Carter J. The Fish Community of a Shallow Tropical Lagoon in
Belize, Central America. Estuaries. 1993;16(2):198-215.
34.   Nagelkerken I, Roberts C, van der Velde G, Dorenbosch M, van Riel MC,
Cocheret de la Morinière E, et al. How important are mangroves and seagrass beds
for coral-reef fish? The nursery hypothesis tested on an island scale. Mar Ecol Prog
Ser. 2002;244:299-305.
35.   Quinn NJ, Kojis BJ. Does the presence of coral reefs in proximity to a tropical
estuary affect the estuarine fish assemblage? Proc 5th Int Coral Reef Congr.
1985;5:445-50.
36.   Olds AD, Albert S, Maxwell PS, Pitt KA, Connolly RM. Mangrove-reef
connectivity promotes the effectiveness of marine reserves across the western Pacific.
Global Ecol Biogeogr. 2013;22(9):1040-9.
37.   Becker A, Coppinger C, Whitfield AK. Influence of tides on assemblages and
behaviour of fishes associated with shallow seagrass edges and bare sand. Mar Ecol
Prog Ser. 2012;456:187-99.
38.   Harborne AR, Talwar B, Brooks EJ. The conservation implications of spatial
and temporal variability in the diurnal use of Bahamian tidal mangrove creeks by
transient predatory fishes. Aquat Conserv: Mar Freshwat Ecosyst. 2016;26(1):202-11.
39.   Reis-Filho JA, Giarrizzo T, Barros F. Tidal migration and cross-habitat
movements of fish assemblage within a mangrove ecotone. Mar Biol.
2016;163(5):111.
40.   Castellanos-Galindo GA, Krumme U, Willis TJ. Tidal influences on fish
distributions on tropical eastern Pacific rocky shores (Colombia). Mar Ecol Prog Ser.
2010;416:241-54.
41.   Bretsch K, Allen DM. Tidal migrations of nekton in salt marsh intertidal creeks.
Estuar Coasts. 2006;29(3):474-86.
42.   Mattone C, Sheaves M. Patterns, drivers and implications of dissolved oxygen
dynamics in tropical mangrove forests. Estuar Coast Shelf Sci. 2017;197:205-13.
43.   Hampel H, Cattrijsse A, Vincx M. Tidal, diel and semi-lunar changes in the
faunal assemblage of an intertidal salt marsh creek. Estuar Coast Shelf Sci.
2003;56(3-4):795-805.

44.   Krumme U, Saint-Paul U, Rosenthal H. Tidal and diel changes in the structure
of a nekton assemblage in small intertidal creeks in northern Brazil. Aquat Living
Resour. 2004;17:215-29.
45.   Ramos JAA, Barletta M, Dantas DV, Lima ARA, Costa MF. Influence of moon
phase on fish assemblages in estuarine mangrove tidal creeks. J Fish Biol.
2011;78(1):344-54.
46.   Laegdsgaard P, Johnson C. Why do juvenile fish utilise mangrove habitats? J
Exp Mar Biol Ecol. 2001;257(2):229-53.
47.   Nagelkerken I, Faunce CH. What makes mangroves attractive to fish? Use of
artificial units to test the influence of water depth, cross-shelf location, and presence
of root structure. Estuar Coast Shelf Sci. 2008;79(3):559-65.
48.   Johnston R, Sheaves M. Small fish and crustaceans demonstrate a preference
for particular small-scale habitats when mangrove forests are not accessible. J Exp
Mar Biol Ecol. 2007;353(2):164-79.
49.   Whitfield A, Elliott M. Fishes as indicators of environmental and ecological
changes within estuaries: A review of progress and some suggestions for the future.
J Fish Biol. 2002;61:229-50.

Supporting information

**S1 Appendix. Presence/absence of all common taxa for each 5-minute interval
recorded during the study.**

Fig 1

Fig 2

Fig 3

Fig 4

Fig 5

Fig 6