Title: Crop residues in wheat-oilseed rape rotation system: a pivotal, shifting platform for

microbial meetings

Authors: Lydie Kerdraon [1], Marie-Hélène Balesdent [1], Matthieu Barret [2], Valérie Laval [1] *,

Frédéric Suffert [1] *

* contributed equally

Address: [1] UMR BIOGER, INRA, AgroParisTech, Université Paris-Saclay, 78850 Thiverval-

Grignon, France; [2] UMR IRHS, INRA, Agrocampus Ouest, Université d'Angers, 49071

Beaucouzé, France

* frederic.suffert@inra.fr; tel +33 (0)1 30 8154 35; ORCID 0000-0001-6969-3878

* valerie.laval@inra.fr; tel +33 (0)1 30 81 45 16; ORCID 0000-0002-3667-3878

Abstract

Crop residues are a crucial ecological niche with a major biological impact on agricultural

ecosystems. In this study we used a combined diachronic and synchronic field experiment

based on wheat-oilseed rape rotations to test the hypothesis that plant is a structuring factor of

microbial communities in crop residues, and that this effect decreases over time with their

likely progressive degradation and colonization by other microorganisms. We characterized an

entire fungal and bacterial community associated with 150 wheat and oilseed rape residue

samples at a plurennial scale by metabarcoding. The impact of plant species on the residue

microbiota decreased over time and our data revealed turnover, with the replacement of

oligotrophs, often plant-specific genera (such as pathogens) by copiotrophs, belonging to

more generalist genera. Within a single cropping season, the plant-specific genera and species

were gradually replaced by taxa that are likely to originate from the soil. These changes

occurred more rapidly for bacteria than for fungi, known to degrade complex compounds. Overall, our findings suggest that crop residues constitute a key fully-fledged microbial ecosystem. Taking into account this ecosystem, that has been neglected for too long, is essential, not only to improve the quantitative management of residues, the presence of which can be detrimental to crop health, but also to identify groups of beneficial micro-organisms. Our findings are of particular importance, because the wheat-oilseed rape rotation, in which no-till practices are frequent, is particularly widespread in the European arable cropping systems.

Keywords

community succession, microbial diversity, oilseed rape, residue microbiota, wheat

Background

Crop residues are an essential living element of agricultural soils. Smil [1] stressed that they "should be seen not as wastes but as providers of essential environmental services, assuring the perpetuation of productive agrosystems". When left in the field in the period between two successive crops, rather than being buried immediately, crop residues contribute to the formation of soil organic carbon, improve soil structure, prevent erosion, filter and retain water, reduce evaporation from the soil surface, and increase the diversity and activity of micro-organisms in the ground [2]. No-till practices are becoming increasingly widespread, as they take advantage of these attributes [3]. However, such practices are often considered likely to increase the risk of disease epidemics [4–6]. Indeed, several leaf-, stem-, head-, and fruit-infecting micro-organisms, classified as "residue-borne" or "stubble-borne" pathogens, are dependent on host residues for survival during the period between successive crops and for the production of inoculum for their next attack [7, 8]. The epidemiological contribution of residues as an effective source of inoculum is well-established but difficult to quantify [e.g. 9] and generalise, because the nature of survival structures depends on the biology of the species. The situation is rendered even more complex by the presence of several species reported to act as crop pathogens in plants as endophytes, without symptom development in the plant, and in the soil and plant residues as saprophytes. Taking into account the inoculum from stubble-borne pathogens and possible competition with other micro-organisms, it appears likely that the expression of a disease is the consequence of an imbalance between a potentially pathogenic species and the rest of the microbial community, rather than the consequence of the mere presence of this species [10].

Residues constitute a crucial ecological niche, not only for pathogenic species, but also for non-pathogenic and beneficial species. Residues can be viewed as both a fully-fledged matrix and a transient compartment, because they originate from the plant (temporal link), are

in close contact with the soil (spatial link) and degrade over the following cropping season, at

rates depending on the plant species, the cropping practices used [11], and the year (climate

effect). It remains unknown whether the succession of microbial communities in residues is

driven primarily by plant tissue degradation or edaphic factors [12]. Many studies have

investigated the structure of the microbial communities present during the life cycle of the

plant [e.g. 13–15], but few have investigated the microbiota associated with plant residues.

Several ecological studies have investigated the impact of the residue compartment on the

structure of soil microbial communities [2, 16–19], but not the impact of the soil compartment

on structure of the residue communities. The detritusphere, defined as the part of the soil

attached to residues [12, 20, 21], is the most extensive and broad hotspot of microbial life in

the soil [22]. The residue compartment and the detritusphere are located in close physical

proximity but are considered by microbiologists to be separate trophic and functional niches

[23]. A description of the residue communities and the specific changes in these communities

over time might, therefore, help agronomists to understand the impact of cropping practices

on crop productivity. Fungi and bacteria play important roles in the degradation of plant

tissues in debris (cellulose, hemicellulose, lignin), but the interactions between them within

the microbial community remain unclear, due to the lack of information about their origins

(air-borne, soil-borne or plant-borne), their individual functions and the drivers of community

structure in residues.

Crop rotation induces changes in the composition of the soil microbial community and

usually reduces pathogen pressure [e.g. 18]. For instance, wheat yields benefit from "break

crops" such as oilseed rape or other non-host crops to break the life-cycle of wheat-specific

pathogens [24]. We focused here on the wheat-oilseed rape rotation, one of the most widely

used cropping systems in Europe. In 2017, the areas under bread wheat and oilseed rape in

France were 5.0 million ha and 1.4 million ha [25], respectively. As oilseed rape usually

recurs every three years in the rotation and is used almost systematically either directly before

or directly after wheat, we estimate that this classical rotation is used on almost 4.2 million ha

every year. Half the area occupied by these two crops is now grown without tillage, with at

least some of the residues of the preceding crop left on the soil [26]. The issue addressed here

is thus directly relevant to more than 2 million ha, or about one tenth of the total arable area in

France.

In this study, we deliberately focused on crop residues as a neglected, transient, but

fully-fledged half-plant/half-soil compartment without describing the soil microbial

communities, considering that it has been already performed in several studies [e.g. 27, 28].

We tested the specific hypothesis that plant is a structuring factor of bacterial and fungal

communities in residues, and that this effect decreases over time, as contact with the soil

induce progressive colonization of residues by other microorganisms. Over the last few years,

high-throughput metabarcoding has become an indispensable tool for studying the ecology of

such complex microbial communities [29], partly due to the difficulties in isolating fungal and

bacterial species and growing them in axenic conditions. We used this approach to describe

and compare changes in the microbial community of wheat and oilseed-rape residues left on

the soil surface of three cultivated fields during two cropping seasons. We investigated

whether the three main determinants of the diversity of fungal and bacterial communities -

plant species, cropping season, and cropping system (monoculture vs. rotation, focusing on

wheat residues) - affected the microbiota of crop residues.

Methods

Experimental design

Field plots and rotations – An extensive field experiment based on a wheat (W)-oilseed

rape (O) rotation cropping system was carried out during the cropping seasons of 2015-2016

and 2016-2017 at the Grignon experimental station (Yvelines, France; 48°51'N, 1°58'E). This area is characterised by an oceanic climate (temperate, with no dry season and a warm summer). A combined diachronic and synchronic strategy [30] was used to investigate the dynamics of the residue microbial communities both over a two-year period on the same plot and along a chronosequence substituting spatial differences (three plots) for time differences. A first monoculture plot (WWW) was sown with the winter wheat cultivar Soissons. This plot had been cropped with wheat since 2007 and was used in previous epidemiological studies focusing on the impact of wheat debris on the development of Septoria tritici blotch [e.g. 31, 32, 33]. Two other plots were cropped with oilseed rape cv. Alpaga and wheat cv. Soissons in rotation (OWO, adjacent to the WWW plot, and WOW, located 400 m away; Fig. 1). The size of the three plots was identical (20 m × 100 m). The OWO and WWW plots are characterized by a silty clay loam soil and plot WOW is characterized by a silty loam soil. Soil texture of the three plots is presented in Additional Table S1. The three plots were not tilled during the two cropping seasons. The wheat and oilseed rape residues were left on the soil surface after harvest. Soil was superficially disturbed to a depth of 10 cm with a disc harrow 6 weeks later (late September), leaving a large portion of residue on the surface. Crops were managed in a conventional way following local practices (nitrogen fertilization, insecticide and herbicide treatments). No fungicide was sprayed on the leaves during the study.

Residue sampling - Wheat and oilseed rape residues from the previous crop were collected over the two cropping seasons. The changes in the microbial communities during residue degradation were described on the basis of four sampling periods each year (October, December, February, and May) Sampling dates are presented in Additional Table S2. A supplementary sample was taken in July 2016, and *a posteriori* in July 2017, to characterise the plant microbiota before the residues came into contact with the soil. For each sampling period, residues samples were collected at soil surface from five points in each plot, 20 m

apart, along a linear transect (Fig. 1). Each sample was composed of twelve pieces of wheat

residue or four pieces of oilseed rape residue. The five sampling points were located at the

same place in the plots during the two years of the experiment.

DNA extraction - Residues were cut to take off remaining roots, rinsed with water to

remove the soil and air-dried in laboratory conditions. They were then cut into small pieces,

pooled in a 50 mL bowl and crushed with a Retsch™ Mixer Mill MM 400 for 60 seconds at

30 Hz in liquid nitrogen, in a zirconium oxide blender. The crushed powder was stored in

50 mL Falcon tubes at -80°C until DNA extraction. We transferred 40 mg of crushed residues

to a 2.0 mL Eppendorf tube, which was stored to -80°C. Total environmental DNA (eDNA)

was extracted according to the TriZol® Reagent protocol (Invitrogen, according to the

manufacturer's instructions). Two independent extractions were performed per sample, giving

a total of 300 eDNA samples. The two extractions were considered as technical replicates.

PCR and Illumina sequencing

Fungal and bacterial community profiles were estimated by amplifying ITS1 and the

v4 region of the 16S rRNA gene, respectively. Amplifications were performed with the

ITS1F/ITS2 [34] and 515f/806r [35] primers. All PCRs were run in a reaction volume of 50

μL, with 1x Qiagen Type-it Multiplex PCR Master Mix (Type-it® Microsatellite PCR kit Cat

No./ID: 206243), 0.2 μM of each primer, 1x Q-solution® and 1 μL DNA (approximately 100

ng). The PCR mixture was heated at 95°C for 5 minutes and then subjected to 35 cycles of

amplification [95°C (1 min), 60°C (1 min 30 s), 72°C (1 min)] and a final extension step at

72°C (10 min). PCR products were purified with Agencourt® AMPure® XP (Agencourt

Bioscience Corp., Beverly, MA). A second round of amplification was performed with 5 μL of

purified amplicons and primers containing the Illumina adapters and indexes. PCR mixtures

were heated at 94°C for 1 min, and then subjected to 12 cycles of amplification [94°C (1

min), 55°C (1 min), 68°C (1 min)] and a final extension step at 68°C (10 min). PCR products

were purified with Agencourt® AMPure® XP and quantified with Invitrogen QuantIT™

PicoGreen®. Purified amplicons were pooled in equimolar concentrations in five independent

batches, and the final concentration of each batch was determined with the qPCR NGS library

quantification kit (Agilent). The five independent batches were sequenced in five independent

runs with MiSeq reagent kit v3 (300bp PE).

Sequence processing

Fastq files were processed with DADA2 v1.6.0 [36], using the parameters described in

the workflow for "Big Data: Paired-end" [37]. The only modification made relative to this

protocol was a change in the truncLen argument according to the quality of the sequencing

run. Each run was analysed separately. Taxonomic affiliations for amplicon sequence variants

(ASV) generated with DADA2 were assigned with a naive Bayesian classifier on the RDP

trainset 14 for bacteria [38] and the UNITE 7.1 database for fungi [39].

Only ASV detected in both technical replicates were conserved to ensure robustness

[40] and were then added together. ASV classified as "Cyanobacteria/Chloroplast", or not

classified at the phylum level, were discarded from the datasets. This resulted in suppression

of 1.2% of reads for fungi (4.2% of unclassified ASV), and 1.5% of reads for bacteria (4.9%

of unclassified ASV and 1.3% of ASV affiliated to Cyanobacteria/Chloroplast). The

remaining ASV were normalised according to the proportion of reads within each sample

[41].

Microbial community analyses

Microbial community profiles were obtained for 100 wheat residue samples and 50

oilseed rape residue samples. The alpha-diversity was estimated for each sample by

calculating the Shannon index [42], completed by Faith's phylogenetic diversity (PD; [43]; [44]) for bacterial communities. The compositional dissimilarity between samples (beta-diversity) was estimated by the Bray-Curtis dissimilarity index.

Factors taken into account in the microbial community analyses were plant (wheat; oilseed rape), cropping system (monoculture; rotation), cropping season (2015-2016; 2016-2017), sampling period (July; October; December; February; May), and sampling plot (WWW; WOW; OWO; Fig. 1). We used a model to test these effects on each aforementioned index in a general way, and then conducted post-hoc contrasts to characterize the differences. A complete model combining all the factors could not have been used because the experimental design did not include an oilseed rape monoculture field plot. Of note oilseed rape monoculture is considered as an agronomic nonsense. Thus, a first model including the plant effect but not the cropping system effect (plant * cropping season * sampling period * sampling plot) was applied using the data set from plots in rotation only (WOW and OWO). The effect of the cropping system (monoculture; rotation) was estimated separately using a second model (cropping system * cropping season * sampling period) applied on the data set from wheat residues only; the sampling plot factor was not included in this second model as it would have been confounding with the cropping system factor.

Shannon index was calculated for both bacterial and fungal communities with the ggpubr package in R [45] and Faith's phylogenetic diversity was calculated for bacterial communities with the picante package [46]. The effect of each factor on the Shannon index was assessed with two complementary ANOVA. A Kruskal-Wallis test also performed to assess significant differences in microbial diversity with time for each cropping season and sampling plot. Wilcoxon pairwise tests were also performed to compare the effects of sampling periods. Divergences were considered significant if $p < 0.05$.

The effect of each factor on the Bray-Curtis dissimilarity index was assessed with two complementary PERMANOVA using the adonis2 function of the vegan R package (version 2.4-4 [47] with "margin" option, used to determine the effect of each term in the model, including all other variables, to avoid sequential effects. They were visualized by multidimensional scaling (MDS) on the Bray-Curtis dissimilarity index with the phyloseq package in R (version 1.22.3 [48] and completed for bacterial communities by incorporating phylogenetic distances using the UniFrac distance matrix. After the aggregation of ASV for each sampling condition "sampling period/cropping year * crop within a rotation", the betapart R package [49] was used to determine whether temporal changes in community composition were due to turnover (i.e. replacement of ASV between two sampling periods) or nestedness (gain or loss of ASV between two sampling periods). The effect of the plant on the microbial communities associated with residues during degradation was also assessed with PERMANOVA on each sampling period, for each cropping season.

The genus composition of fungal and bacterial communities was analysed with a cladogram based on genus names. Only genera observed in three biological samples harvested on the same plot were incorporated into the cladogram. A cladogram representing the number of ASV for each genus, read percentage, occurrence and distribution for each sample, was constructed with the Interactive Tree Of Life (iTOL [50]) online tool for phylogenetic trees.

To illustrate taxonomic changes over time, especially between plant-derived communities and communities involved later in the colonization of the residues, we focused on seasonal shifts (increase, decrease or stability) in the relative abundance of a selection of some fungal and bacterial genera and tested their statistical significance (Wilcoxon tests between sampling periods).

Results

The bacterial and fungal communities associated with wheat (W) and oilseed rape (O) crop residues were characterised on three plots: the wheat monoculture (WWW), and two oilseed rape-wheat rotation plots (WOW and OWO) (Fig. 1). We assessed the composition of these microbial communities four times per year, during two consecutive cropping seasons (in October, December, February, and May). An additional time point (in July) was also included for identification of the micro-organisms present on the plant before contact with the soil. The analysis of raw sequence datasets for the 150 samples of wheat and oilseed rape residues collected over the two cropping seasons resulted in the grouping of 14,287,970 bacterial and 9,898,487 fungal reads into 2,726 bacterial and 1,189 fungal amplicon sequence variants (ASV). ASV not detected in both technical replicates (5.4% of bacterial reads and 1.5% of fungal reads) were removed from the datasets. Total number of reads remaining after ASV filtering is presented in Additional Table S3.

Alpha diversity of microbial communities

Diversity dynamics, assessed by calculating the Shannon index, differed between the two cropping seasons and between fungi and bacteria. The diversity was significantly impacted by most of factors, including cropping system (monoculture; rotation) for wheat residues (Table 1; Fig. 2). Oilseed rape residues supported less fungal diversity and as much bacterial diversity than wheat residues in rotation. In addition, the diversity was significantly higher in wheat grown in monoculture than in wheat grown in rotation for both bacteria and fungi.

Fungal diversity increased over time in 2015-2016, whereas the differences between the samples in 2016-2017 did not reflect a gradual increase as the minimum was reached in December. Bacterial diversity followed a quite similar trend with however a significant decreased from February to May, more pronounced for wheat residues than for oilseed rape

260 residues during the first cropping season; this trend was also illustrated by the Faith's

261 phylogenetic diversity (Additional Figure S1). The climatic conditions during residue

262 degradation (Additional Table S4) or differences in initial diversity on the plant before harvest

263 may explain the less marked trends observed between the two cropping seasons.

Comparison of microbial communities associated with residues (beta diversity)

We analysed the effects of plant, cropping system, cropping season, and sampling

period on communities using the Bray-Curtis dissimilarity index and PERMANOVA. There

was remarkably little heterogeneity between the five samples collected in the same sampling

plots (Fig. 3) and the number of biological samples was, therefore, sufficient to assess

differences due to the variables of interest (i.e. plant cropping season, sampling period, and

cropping system). This result was confirmed by the structure of bacterial communities

visualized by incorporating phylogenetic distances using the UniFrac distance matrix

(Additional Fig. S2).

The structure of bacterial and fungal communities is influenced by plant species and

cropping system – Oilseed rape and wheat residues presented different sets of ASV, for both

bacterial and fungal communities (Fig. 3). Plant species was the main factor explaining

differences between the communities, accounting for 38.4% for fungi and 26.6% of the

variance for bacteria, as established with PERMANOVA (Table 2). For wheat, the cropping

system (rotation; monoculture) accounted for 10.5% of the variance for fungal community

composition and 6.6% of the variance for bacterial community composition.

The percentage of variance explained by the plant decreased over time for fungal

community structure (e.g. from 75% in October 2016 to 40% in May), while for bacteria the

decrease of the percentage of variance explained by the plant was less pronounced (from 65%

to 50%). The percentages of variance associated with the plant for each date are presented in Additional Table S5.

Community structures change over time – Cropping season was the main temporal factor underlying changes in community structure, accounting for 11.8% of the variance for bacteria when considering only the plots in rotation (or 24.8% when considering only the wheat residues without sampling plot effect), and 13.1% of the variance for fungi when considering only the plots in rotation (or 29.2% when considering the wheat residues without sampling plot effect; Table 2). Sampling period had also a significant impact on community composition, accounting for 14.9% of the variance for bacteria when considering only the plots in rotation (or 20.4% when considering only the wheat residues without sampling plot effect), and 6.8% of the variance for fungi when considering only the plots in rotation (or 12.3% when considering the wheat residues without sampling plot effect). Theoretically, changes in ASV composition result from turnover (replacement of ASV between two sampling periods) and nestedness (gain or loss of ASV between two sampling periods [49]). We found that the dissimilarity between sampling periods was smaller for bacterial than for fungal ASV structure. By decomposing the dissimilarity between sampling periods, we found that, for fungi, 94% (± 5%) of dissimilarity was explained by turnover for oilseed rape, 89% (±15%) for wheat in monoculture, and 80% (±13%) for wheat in rotation. For bacteria, 69% (±17%) of dissimilarity was explained by turnover for oilseed rape, 61% (±19%) for wheat in monoculture, and 80% (±16%) for wheat in rotation. Decomposition of dissimilarity between sampling periods is presented in Additional Table S6.

Changes in communities, by genus

We characterised potential taxonomic differences in communities over time by analysing wheat and oilseed rape residues separately. ASV were aggregated together at genus

level, resulting in 84 fungal and 184 bacterial genera for wheat, and 63 fungal and 186 bacterial genera for oilseed rape. For the sake of clarity, most 60 genera of fungi and bacteria were presented in Fig. 4 and Fig. 5, respectively. All the detected genera and their evolution over time were presented in Additional Fig. S3, S4, S5 and S6. For both plant species, we identified genera that disappeared or displayed a significant decrease in relative abundance over time. The seasonal shifts of some genera and their significance are presented in Additional Fig. S6. Among these genera, some are known to be associated with plants, such as *Alternaria*, *Acremonium* [14, 51, 52], *Cryptococcus* [53], *Sarocladium* [54] and *Cladosporium* [13, 51–54].

Some of the fungal species detected on wheat, such as *Oculimacula yallundae* (all ASV of *Oculimacula* genera), *Zymoseptoria tritici* and *Pyrenophora tritici-repentis,* (Fig. 4, Additional Fig. S3) are known to be pathogenic. Some of the species detected on oilseed rape, such as *Verticillium* spp., *Leptosphaeria maculans* (= *Plenodomus maculans*) and *Leptosphaeria biglobosa* (= *Plenodomus biglobosa*), are also known to be pathogenic. Strikingly, *L. maculans* and *L. biglobosa* predominated over the other taxa. *Verticillium longisporum*, *V. dahlia* and *V. albo-atrum* were mostly detected during the second sampling year (Fig. 4, Additional Fig. S4). As samples were collected in two different fields, it was not possible to determine whether the occurrence of *Verticillium* spp., a soil-borne pathogen complex causing *Verticillium* wilt [55], was affected more by year or by the soil contamination. *Acremonium*, *Clonostachys* and *Alternaria* genera, which have also been described as associated with plants [56], were detected in the early sampling periods. Their relative abundances decreased over time (Additional Fig. S7). Most of the genera that were not present at early sampling points and with relative abundances increasing over time (e.g. *Coprinellus*, *Psathyrella*, *Torula*, *Tetracladium*, and *Exophiala*) were common to wheat and

oilseed rape residues (Fig. 4). These genera can thus be considered as probably derived

primarily from the surrounding soil.

For bacteria, the difference in the genera detected between the two plants species was

less marked than for fungi, as 146 genera were common to wheat and oilseed rape residues

(Fig. 5). These 146 genera corresponded to the 98.7% most prevalent reads for wheat and

97.5% most prevalent genus reads for oilseed rape. *Proteobacteria* was the predominant

phylum the first year. The most prevalent proteobacterial subgroup was *Alphaproteobacteria*,

with a high prevalence of *Rhizobiales* and *Sphingomonadales*. *Rhizobium* and *Neorhizobium*,

two major genera from *Rhizobiales*, decreased in abundance between October and May in

both wheat and oilseed rape (Additional Fig. S7). *Sphingomonadales* genera were much more

abundant on wheat than on oilseed rape, especially *Sphingomonas* (Fig. 5). *Bacteroidetes*

genera, including *Pedobacter* in particular, were frequently detected and their prevalences

tended to be stable for oilseed rape residues, and to decrease for wheat residues (Additional

Fig. S7). In parallel, an increase in *Actinobacteria*, particularly *Nocarioides*, was observed.

Major differences between July and October were observed for oilseed rape, consistent with

the beta-diversity analysis, in which the percentage dissimilarity between July and October

was high, due to both species extinction and turnover. *Gammaproteobacteria* were highly

abundant on oilseed rape in July. Their frequency then decreased rapidly from October to

May, due largely to the decrease in *Pseudomonas* (Fig. 5, Additional Fig. S7). In parallel, we

observed an increase in the levels of *Alphaproteobacteria*, especially *Rhizobium* and

Sphingomonas, between July and October. A small decrease in levels of

Gammaproteobacteria was observed between July and October for wheat in rotation, whereas

the percentage of reads associated with this class increased between July and December for

wheat in monoculture, due largely to the decrease in *Pantoea* and *Enterobacteria* (Fig. 5,

Additional Fig. S7). The abundance of *Bacteroidetes*, especially *Pedobacter* and *Flavobacterium*, also increased between July and October.

Discussion

Most studies on crop residues have focused on their impact on soil microbial communities [16], and the rare studies investigating the impact of soil on residue communities focused exclusively on bacteria [27, 28] or fungi [57]. Most of these studies were conducted on residues from a single year. Bastian et al. [12] established an extensive description of the species present in the soil, detritusphere and wheat residues, using sterilised residues and soil in a microcosm. In this study, we showed, under natural conditions, that three main factors (plant species, cropping season, rotation) simultaneously influence the composition of both fungal and bacterial communities present on residues. This study is the first to investigate the total fungal and bacterial communities associated with wheat and oilseed rape residues by a metabarcoding approach over two consecutive years. The very low variability of the communities for the five replicates is remarkable and shows that our strategy would be appropriate for comparing the effects of different treatments on microbial communities.

Crop residues should be viewed as a shifting platform for microbial meeting strongly affected by plant species

Oilseed rape and wheat residues contained different sets of micro-organisms before soil contact and during the firsts sampling periods after harvest. Similar results were previously obtained for the bacterial communities of buried crop residues [28]. Consistent with the findings of this previous study, the divergence between wheat and oilseed rape bacterial communities was probably due to differences in the chemical compounds present in the plants. The rapid change in the community observed at early stages of residue degradation

for oilseed rape may be explained by the modification of simple compounds (sugars, starch,

etc.), whereas wheat is composed of more complex compounds (lignin) and is, therefore,

broken down less quickly, resulting in a slower change in the microbial community [28].

Overall, the change in bacterial community composition highlights turnover between

copiotrophs and oligotrophs. Although copiotrophy and oligotrophy are physiological traits,

several attempts have been made to classify microorganisms as oligotrophs and copiotrophs

based on phylogeny [58]. According to this generalization, bacterial and fungal taxa whose

relative abundances are significantly decreased during succession belong mainly to

copiotroph. These taxa include for instance *Alternaria*, *Cladosporium*, *Massilia* and

Pseudomonas (Additional file: Fig. S5). In contrast, the relative abundances of oligotrophic

taxa such as *Coprinellus* or *Nocardiodes* increased during residues degradation, which could

be indicative of the superior abilities of these micro-organisms to degrade complex polymers.

The initial fungal communities were structured mostly by the presence of species

originating from the plant, several of which were highly specialised on the host plant. These

species were gradually replaced by more generalist species, which colonised the residues of

both plants. Most of these generalists, such as *Exophiala*, *Coprinellus* and *Torula,* are known

to be soil-born [59, 60], or involved in degradation, such as *Coprionopsis* [61]. The host-

specific fungi identified in our study included a large number of ascomycetes known to be

foliar pathogens (*O. yallundae, Fusarium* sp. and *Gibberella* sp*., Z. tritici, P. tritici-repentis,*

Parastagonospora nodorum, Monographella nivalis, L. biglobosa and *L. maculans*). The

lifestyles of some pathogens are well-documented, as for *Z. tritici, P. tritici-repentis* and *L.*

maculans. The decrease with time in levels of *Z. tritici* and other pathogens in wheat residues

contrasts with the persistence of *L. maculans* and *L. biglobosa* in oilseed rape residues. These

three pathogens are all known to reproduce sexually on the residues of their host plant [31,

62], but the life cycle of *L. maculans* is characterised by systemic host colonisation through

intracellular growth in xylem vessels [63], whereas the development of *Z. tritici* is localised

and exclusively extracellular [64]. Oilseed rape residues thus provide *L. maculans* with

greater protection than is provided to *Z. tritici* by wheat residues. This likely explains

differences in the persistence of the two pathogens and in the temporal dynamics of ascospore

release: over up to two years for *L. maculans* [65, 66] but only a few months for *Z. tritici* [31,

67]. The predominance of *L. maculans* on oilseed rape residues was not surprising given that

the oilseed rape cultivar Alpaga is known to be susceptible to *L. maculans*, but the high

abundance of *L. biglobosa* was much more remarkable. One surprising finding of our study

was the constant association of *L. maculans* with *L. biglobosa* on residues. Indeed, *L.*

biglobosa is known to be more associated with upper-stem lesions [68], and its presence in

large amounts on residues has never before been reported.

Our findings are consistent with current epidemiological knowledge of emblematic

wheat and oilseed rape diseases, but they highlight our lack of knowledge concerning the

lifestyles of many other fungal pathogens present on residues. A key point to be taken into

account is that the trophic status of many species known to be principally pathogenic or non-

pathogenic is not definitive [69]. For instance, *Alternaria infectoria* is sometimes described as

a pathogen of wheat [13, 70], sometimes as an endophyte [71], and has even been tested as a

potential biocontrol agent against *Fusarium pseudograminearum* on wheat [72]. Crop

residues, half-plant/half-soil, should be the focus of future studies aiming to disentangle the

succession of microbial species with different lifestyles and to characterise their relative

impacts on the development of currently minor, but potentially threatening diseases.

The residue microbiota should be analysed in a dynamic manner, both within and

between years

The results of our study highlight the importance of conducting multi-year studies

focusing on ecological dynamics both within and between years in natural conditions. Year

had a strong effect on both bacterial and fungal communities. Fluctuations of climatic

conditions (temperature, rainfall, wind) have a major impact on pathogenesis (disease triangle

concept [73]) and on the saprophytic survival of plant pathogens during interepidemic periods

[74]. The two years of our study were marked by similar means of 10-day mean temperatures,

but large differences in rainfall: mean 10-day cumulative rainfall in the first year was almost

twice that in the second (Additional Table S5). The colonisation of residues by late colonisers

may be affected by such climatic differences: in wheat, most prevalent degrading fungi (like

Coprinellus, *Psathyrella*, *Coprinopsis*) were almost absent in the second year of the study.

There was also considerable dissimilarity between the bacterial communities associated with

each of the two years. For example, genus *Enterobacter*, which was highly abundant in the

second year, was barely detectable in the first year.

Crop rotation has little impact on residue microbial communities

Oilseed rape is never grown in monoculture, so the effect of crop rotation was assessed

only for wheat. The effect of rotation on residue microbial communities was much smaller

than the effect of year (cropping season). It was more marked for fungi, for which diversity

was greater in monoculture than in rotation. The use of a rotation may prevent the most

strongly specialised species, in this case fungi, from becoming established, regardless of their

pathogenicity. This finding is consistent with the greater development of some diseases in

monoculture conditions, which promote the maintenance of pathogens through the local

presence of primary inoculum. For instance, the presence of *P. tritici-repentis*, agent of tan

spot disease, in the wheat monoculture plot and its absence from wheat-oilseed rape plots is

consistent with epidemiological knowledge indicating that this disease can be controlled by

leaving a sufficient interval between consecutive wheat crops in the same field [75].

Lesson to be learned from the residue microbial communities for the sustainable

management of debris-borne diseases: a delicate balance between pathogenic and

beneficial micro-organisms

The maintenance of crop residues at the surface of the cultivated soil increases the

microbial diversity of the soil and, in some ways, helps to maintain good functional

homeostasis [76]. However, conservation practices tend to increase the risk of foliar diseases

[4–6]. Most disease management strategies focus on epidemic periods, during which the

pathogen and its host are in direct contact. Interepidemic periods are also crucial for pathogen

development, although during these periods the primary inoculum is not directly in contact

with the new crop whilst not present in the field. Indeed, by carrying the sexual reproduction

of several fungal pathogens, residues contribute to the generation and transmission of new

virulent isolates potentially overcoming resistance genes, during monocyclic epidemics, as

described for oilseed rape canker caused by *L. maculans* [77], but also polycyclic epidemics,

as described for Septoria tritici blotch caused by *Z. tritici* [78].

However, the results of our study suggest that residues should not only be considered as

a substrate for pathogens and a potential source of inoculum. Indeed, we detected several

fungi identified as beneficial or even biocontrol agents in previous studies, such as

Clonostachys rosea, *Aureobasidium pullulans*, *Chaetomium globosum* and *Cryptococcus* spp.

C. rosea, which was detected in both oilseed rape and wheat residues, has been reported to

limit the sexual and asexual reproduction of *Didymella rabiei* on chickpea residues by

mycoparasitism [79]. It has also been reported to be effective against *Fusarium culmorum* on

wheat plants, through antibiosis during the epidemic period [80], and on wheat residues,

through antagonism during the interepidemic period [81]. *Cladosporium* sp., which were abundant in our study, have also been reported to inhibit the development of *P. tritici-repentis* on wheat plants [82] and of *Fusarium* sp. on wheat residues [81]. The presence of these fungal species on wheat and oilseed rape residues is of potential interest for future analyses of interactions. Due to the use of a low-resolution marker for bacterial characterisation, we were unable to identify similarly the bacteria potentially interacting with pathogenic fungi. For instance, the presence of *Pseudomonas* spp. suggests possible interactions both with other microbial species and with the host plant [83], but the nature of the potential interactions is indeterminate: species of the *Pseudomonas fluorescens* group are known to be beneficial to plants, whereas *Pseudomonas syringae* and *Pseudomonas aeruginosa* are known to be pathogens of plants and even humans.

Although our study reveals the presence of genera or species reported in the literature as biocontrol agents, it has not yet shown any interaction between them and the pathogens. This experimental study (sampling effort, residue treatments, etc.) was not designed to characterize such interactions. A strategy involving the inference of microbial interaction networks from metabarcoding datasets might help to identify the species beneficial against pathogens, through competition, antagonism or parasitism. This however requires a more analytical, comparative experimental approach, that goes beyond the only description of shifts in natural communities composition: for example, using different "treatments" in a broad sense (e.g. artificial inoculation with a species or a group of species, change of biotic or abiotic environmental conditions, etc.) in order to modify interaction networks and so highlight the impact of some groups of micro-organisms on the whole community or a given species.

Conclusion

This study shows that crop residues, which can be seen as half-plant/half-soil transient compartment, constitute a pivotal fully-fledged microbial ecosystem that has received much less attention than the phyllosphere and rhizosphere to date. This study therefore fills a gap in knowledge of the communities present on crop residues under natural conditions. It confirms that the microbiote of crop residues should be taken into account in the management of residue-borne diseases. Taking into account this ecosystem is essential, not only to improve the quantitative management of crop residues, but also to identify groups of beneficial micro-organisms naturally present. The beneficial elements of the microbial community should be preserved, or even selected, characterised and used as biological control agents against the pathogens that complete their life cycle on the residues. These results are particularly important in that wheat-oilseed rape rotations are among the most widespread arable cropping systems in France and Europe.

Acknowledgements

This study was performed in collaboration with the GeT core facility, Toulouse, France (http://get.genotoul.fr) and was supported by *France Génomique National Infrastructure*, funded as part of "*Investissement d'avenir*" program managed by *Agence Nationale pour la Recherche* (contract ANR-10-INBS-09). We thank Martial Briand (INRA, UMR IRHS) and Dr. Gautier Richard (INRA, UMR IGEPP) for assistance with bioinformatic analyses, and Dr. Thierry Rouxel (INRA, UMR BIOGER) for improving and clarifying this manuscript. We thank Julie Sappa for her help correcting our English. We finally thank the reviewers for their insightful comments on the paper, as these comments led us to improve the correctness of several analyses.

Funding

529 This study was supported by a grant from the European Union Horizon Framework 2020

530 Program (EMPHASIS Project, Grant Agreement no. 634179) covering the 2015-2019 period.

Availability of data and materials

533 The raw sequencing data is available from the European Nucleotide Archive (ENA) under the

534 study accession PRJEB27255 (Sample SAMEA4723701 to SAMEA4724326). We provide

535 the command-line script for data analysis and all necessary input files as Additional File 2.

Authors' contributions

538 LK, FS, VL, MHB, MB conceived the study, participated in its design, and wrote the

539 manuscript. LK conducted the experiments and analysed the data. FS and VL supervised the

540 project. All authors read and approved the final manuscript.

Ethics approval and consent to participate

543 Not applicable

Consent for publication

546 Not applicable

Competing interests

549 The authors declare that they have no competing interests.

References

1. Smil, V. (1999). Crop residues: agriculture's largest harvest crop residues incorporate

more than half of the world's agricultural phytomass. BioScience *49*, 299–308.

2. Govaerts, B., Mezzalama, M., Unno, Y., Sayre, K.D., Luna-Guido, M., Vanherck, K.,

Dendooven, L., and Deckers, J. (2007). Influence of tillage, residue management, and

crop rotation on soil microbial biomass and catabolic diversity. Appl. Soil Ecol. *37*, 18–

30.

3. Derpsch, R., Friedrich, T., Kassam, A., and Hongwen, L. (2010). Current status of

adoption of no-till farming in the world and some of its main benefits. Biol Eng *3*, 25.

4. Bailey, K.L. (1996). Diseases under conservation tillage systems. Can. J. Plant Sci. *76*,

635–639.

5. Bailey, K.., and Lazarovits, G. (2003). Suppressing soil-borne diseases with residue

management and organic amendments. Soil Tillage Res. *72*, 169–180.

6. Bockus, W.W., and Shroyer, J.P. (1998). The impact of reduced tillage on soilborne

plant pathogens. Annu. Rev. Phytopathol. *36*, 485–500.

7. Bockus, W.W. (1998). Control strategies for stubble-borne pathogens of wheat. Can. J.

Plant Pathol. *20*, 371–375.

8. Cook, R.J., Boosalis, M.G., and Doupnik, B. (1978). Influence of crop residues on plant

diseases. Crop Residue Manag. Syst. 147–163.

9. Morais, D., Sache, I., Suffert, F., and Laval, V. (2016). Is the onset of Septoria tritici

blotch epidemics related to the local pool of ascospores? Plant Pathol. *65*, 250–260.

10. Vayssier-Taussat, M., Albina, E., Citti, C., Cosson, J.-F., Jacques, M.-A., Lebrun, M.-

H., Le Loir, Y., Ogliastro, M., Petit, M.-A., Roumagnac, P., et al. (2014). Shifting the

paradigm from pathogens to pathobiome: new concepts in the light of meta-omics.

Front. Cell. Infect. Microbiol. *4*.

11. Hadas, A., Kautsky, L., Goek, M., and Erman Kara, E. (2004). Rates of decomposition

of plant residues and available nitrogen in soil, related to residue composition through

simulation of carbon and nitrogen turnover. Soil Biol. Biochem. *36*, 255–266.

12.   Bastian, F., Bouziri, L., Nicolardot, B., and Ranjard, L. (2009). Impact of wheat straw

decomposition on successional patterns of soil microbial community structure. Soil

Biol. Biochem. *41*, 262–275.

13.   Hertz, M., Jensen, I.R., Jensen, L.Ø., Thomsen, S.N., Winde, J., Dueholm, M.S.,

Sørensen, L.H., Wollenberg, R.D., Sørensen, H.O., Sondergaard, T.E., et al. (2016). The

fungal community changes over time in developing wheat heads. Int. J. Food Microbiol.

222, 30–39.

14.   Qin, Y., Fu, Y., Dong, C., Jia, N., and Liu, H. (2016). Shifts of microbial communities

of wheat (*Triticum aestivum L.*) cultivation in a closed artificial ecosystem. Appl.

Microbiol. Biotechnol. *100*, 4085–4095.

15.   Vacher, C., Hampe, A., Porté, A.J., Sauer, U., Compant, S., and Morris, C.E. (2016).

The phyllosphere: microbial jungle at the plant–climate interface. Annu. Rev. Ecol.

Evol. Syst. *47*, 1–24.

16.   Bending, G.D., Turner, M.K., and Jones, J.E. (2002). Interactions between crop residue

and soil organic matter quality and the functional diversity of soil microbial

communities. Soil Biol. Biochem. *34*, 1073–1082.

17.   Degrune, F., Theodorakopoulos, N., Dufrêne, M., Colinet, G., Bodson, B., Hiel, M.-P.,

Taminiau, B., Nezer, C., Daube, G., and Vandenbol, M. (2016). No favorable effect of

reduced tillage on microbial community diversity in a silty loam soil (Belgium). Agric.

Ecosyst. Environ. *224*, 12–21.

18.   Sommermann, L., Geistlinger, J., Wibberg, D., Deubel, A., Zwanzig, J., Babin, D.,

Schlüter, A., and Schellenberg, I. (2018). Fungal community profiles in agricultural

soils of a long-term field trial under different tillage, fertilization and crop rotation

conditions analyzed by high-throughput ITS-amplicon sequencing. PLoS ONE *13*,

e0195345.

19. Zelenev, V.V., van Bruggen, A.H.C., and Semenov, A.M. (2005). Short-term wavelike
dynamics of bacterial populations in response to nutrient input from fresh plant
residues. Microb. Ecol. *49*, 83–93.

20. Poll, C., Brune, T., Begerow, D., and Kandeler, E. (2010). Small-scale diversity and
succession of fungi in the detritusphere of rye residues. Microb. Ecol. *59*, 130–140.

21. Poll, C., Marhan, S., Ingwersen, J., and Kandeler, E. (2008). Dynamics of litter carbon
turnover and microbial abundance in a rye detritusphere. Soil Biol. Biochem. *40*, 1306–
1321.

22. Kuzyakov, Y., and Blagodatskaya, E. (2015). Microbial hotspots and hot moments in
soil: Concept & review. Soil Biol. Biochem. *83*, 184–199.

23. Nicolardot, B., Bouziri, L., Bastian, F., and Ranjard, L. (2007). A microcosm
experiment to evaluate the influence of location and quality of plant residues on residue
decomposition and genetic structure of soil microbial communities. Soil Biol. Biochem.
39, 1631–1644.

24. Hilton, S., Bennett, A.J., Chandler, D., Mills, P., and Bending, G.D. (2018). Preceding
crop and seasonal effects influence fungal, bacterial and nematode diversity in wheat
and oilseed rape rhizosphere and soil. Appl. Soil Ecol.

25. AGRESTE, 2018. Données de production Grandes Cultures (national + régions) au
01/02/2018. Excel file,
http://agreste.agriculture.gouv.fr/IMG/xls/2018_017donneesdeptgrandescultures.xls

26. AGRESTE, 2008. Un tiers des grandes cultures semées en 2006 sans retournement des
sols. Dans le sillon du non-labour. Agreste Primeur, n°207,
http://agreste.agriculture.gouv.fr/IMG/pdf/primeur207.pdf

27. Pascault, N., Nicolardot, B., Bastian, F., Thiébeau, P., Ranjard, L., and Maron, P.-A.

(2010a). *In situ* dynamics and spatial heterogeneity of soil bacterial communities under

different crop residue management. Microb. Ecol. *60*, 291–303.

28. Pascault, N., Cécillon, L., Mathieu, O., Hénault, C., Sarr, A., Lévêque, J., Farcy, P.,

Ranjard, L., and Maron, P.-A. (2010b). *In situ* dynamics of microbial communities

during decomposition of wheat, rape, and alfalfa residues. Microb. Ecol. *60*, 816–828.

29. Hanson, B.M., and Weinstock, G.M. (2016). The importance of the microbiome in

epidemiologic research. Ann. Epidemiol. *26*, 301–305.

30. Costa Junior, C., Corbeels, M., Bernoux, M., Píccolo, M.C., Siqueira Neto, M., Feigl,

B.J., Cerri, C.E.P., Cerri, C.C., Scopel, E., and Lal, R. (2013). Assessing soil carbon

storage rates under no-tillage: Comparing the synchronic and diachronic approaches.

Soil Tillage Res. *134*, 207–212.

31. Suffert, F., and Sache, I. (2011). Relative importance of different types of inoculum to

the establishment of *Mycosphaerella graminicola* in wheat crops in north-west Europe:

Wheat debris as local source of *M. graminicola* inoculum. Plant Pathol. *60*, 878–889.

32. Morais, D., Laval, V., Sache, I., and Suffert, F. (2015). Comparative pathogenicity of

sexual and asexual spores of *Zymoseptoria tritici* (Septoria tritici blotch) on wheat

leaves. Plant Pathol. *64*, 1429–1439.

33. Suffert, F., Goyeau, H., Sache, I., Carpentier, F., Gélisse, S., Morais, D., Delestre G.

(2017) Epidemiological trade-off between intra- and interannual scales in the evolution

of aggressiveness in a local plant pathogen population. Evol. Appl. *11*, 768–780.

34. Buée, M., Reich, M., Murat, C., Morin, E., Nilsson, R.H., Uroz, S., and Martin, F.

(2009). 454 Pyrosequencing analyses of forest soils reveal an unexpectedly high fungal

diversity: Research. New Phytol. *184*, 449–456.

35.   Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-Lyons, D., Lozupone, C.A.,

Turnbaugh, P.J., Fierer, N., and Knight, R. (2011). Global patterns of 16S rRNA

diversity at a depth of millions of sequences per sample. Proc. Natl. Acad. Sci. USA

108, 4516–4522.

36.   Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A., and Holmes,

S.P. (2016). DADA2: High-resolution sample inference from Illumina amplicon data.

Nat. Methods *13*, 581–583.

37.   DADA2   Pipeline   Tutorial   (1.8).   Available   at:

https://benjjneb.github.io/dada2/tutorial.html. (Accessed: 26th June 2018)

38.   Cole, J.R., Wang, Q., Cardenas, E., Fish, J., Chai, B., Farris, R.J., Kulam-Syed-

Mohideen, A.S., McGarrell, D.M., Marsh, T., Garrity, G.M., et al. (2009). The

Ribosomal Database Project: improved alignments and new tools for rRNA analysis.

Nucleic Acids Res. *37*, D141–D145.

39.   Abarenkov, K., Nilsson, R.H., Larsson, K.-H., Alexander, I.J., Eberhardt, U., Erland, S.,

Høiland, K., Kjøller, R., Larsson, E., Pennanen, T., et al. (2010). The UNITE database

for molecular identification of fungi – recent updates and future perspectives. New

Phytol. *186*, 281–285.

40.   Ficetola, G.F., Pansu, J., Bonin, A., Coissac, E., Giguet-Covex, C., De Barba, M.,

Gielly, L., Lopes, C.M., Boyer, F., Pompanon, F., et al. (2015). Replication levels, false

presences and the estimation of the presence/absence from eDNA metabarcoding data.

Mol. Ecol. Resour. *15*, 543–556.

41.   McMurdie, P.J., and Holmes, S. (2014). Waste not, want not: why rarefying

microbiome data is inadmissible. PLoS Comput. Biol. *10*, e1003531.

42.   Bo-Ra, K., Shin, J., Guevarra, R.B., Lee, J.H., Kim, D.W., Seol, K.H., Lee, J.H., Kim,

H.B., and Isaacson, R.E. (2017). Deciphering diversity indices for a better

understanding of microbial communities. J. Microbiol. Biotechnol. *27*, 2089–2093.

43.   Tucker, C.M., Cadotte, M.W., Carvalho, S.B., Davies, T.J., Ferrier, S., Fritz, S.A.,

Grenyer, E., Helmus, M.R., Jin, L.S., Mooers, A.O., et al. (2017) A guide to

phylogenetic metrics for conservation, community ecology and macroecology. Biol.

Rev. *92*, 698–715.

44.   Faith, D.P. (1992) Conservation evaluation and phylogenetic diversity. . Biol.

Conserv. *61*, 1–10.

45.   Kassambara, A. (2017). ggpubr: "ggplot2" Based Publication Ready Plots. R package

version 0.1.5.999.

46.   Kembel, S.W, Cowan, P.D., Helmus, M.R., Cornwell, W.K., Morlon, H., Ackerly,

D.D., and Webb, C.O. (2010) Picante: R tools for integrating phylogenies and ecology.

Bioinformatics *26*, 1463–1464.

47.   Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D.,

Minchin, P.R., O'Hara, R.B., Simpson, G.L., Solymos, P., et al. (2017). vegan:

Community Ecology Package. R package version 2.4-4. https://CRAN.R-

project.org/package=vegan.

48.   McMurdie, P.J., and Holmes, S. (2013). phyloseq: An R package for reproducible

interactive analysis and graphics of microbiome census data. PLoS ONE *8*, e61217.

49.   Baselga, A., and Orme, C.D.L. (2012). betapart: an R package for the study of beta

diversity. Methods Ecol. Evol. *3*, 808–812.

50.   Letunic, I., and Bork, P. (2016). Interactive tree of life (iTOL) v3: an online tool for the

display and annotation of phylogenetic and other trees. Nucleic Acids Res. *44*, W242–

W245.

51.   Bateman, G.L., and Kwaśna, H. (1999). Effects of number of winter wheat crops grown successively on fungal communities on wheat roots. Appl. Soil Ecol. *13*, 271–282.

52.   Vujanovic, V., Mavragani, D., and Hamel, C. (2012). Fungal communities associated with durum wheat production system: A characterization by growth stage, plant organ and preceding crop. Crop Prot. *37*, 26–34.

53.   Larran, S., Perelló, A., Simón, M.R., and Moreno, V. (2002). Isolation and analysis of endophytic microorganisms in wheat (*Triticum aestivum* L.) leaves. World J. Microbiol. Biotechnol. *18*, 683–686.

54.   Grudzinska-Sterno, M., Yuen, J., Stenlid, J., and Djurle, A. (2016). Fungal communities in organically grown winter wheat affected by plant organ and development stage. Eur. J. Plant Pathol. *146*, 401–417.

55.   Zhou, L., Hu, Q., Johansson, A., and Dixelius, C. (2006). *Verticillium longisporum* and *V. dahliae*: infection and disease in *Brassica napus*. Plant Pathol. *55*, 137–144.

56.   Zhang, Q., Zhang, J., Yang, L., Zhang, L., Jiang, D., Chen, W., and Li, G. (2014). Diversity and biocontrol potential of endophytic fungi in *Brassica napus*. Biol. Control *72*, 98–108.

57.   Naseri, B., Davidson, J.A., and Scott, E.S. (2008). Survival of *Leptosphaeria maculans* and associated mycobiota on oilseed rape stubble buried in soil. Plant Pathol. *57*, 280–289.

58.   Ho, A., Di Lonardo, D.P., and Bodelier, P.L.E. (2017). Revisiting life strategy concepts in environmental microbial ecology. FEMS Microbiol Ecol. *93*, doi:10.1093/femsec/fix006.

59.   Gorfer, M., Blumhoff, M., Klaubauf, S., Urban, A., Inselsbacher, E., Bandian, D., Mitter, B., Sessitsch, A., Wanek, W., and Strauss, J. (2011). Community profiling and

gene expression of fungal assimilatory nitrate reductases in agricultural soil. ISME J. *5*,

1771–1783.

60.   Swer, H., Dkhar, M.S., and Kayang, H. (2011). Fungal population and diversity in

organically amended agricultural soils of Meghalaya, India. J. Org. Syst. *6*, 3–12.

61.   Turbe-Doan, A., Arfi, Y., Record, E., Estrada-Alvarado, I., and Levasseur, A. (2013).

Heterologous production of cellobiose dehydrogenases from the basidiomycete

Coprinopsis cinerea and the ascomycete *Podospora anserina* and their effect on

saccharification of wheat straw. Appl. Microbiol. Biotechnol. *97*, 4873–4885.

62.   Rouxel, T., and Balesdent, M.H. (2005). The stem canker (blackleg) fungus,

Leptosphaeria maculans, enters the genomic era. Mol. Plant Pathol. *6*, 225–241.

63.   Hammond, K.E., Lewis, B.G., and Musa, T.M. (1985). A systemic pathway in the

infection of oilseed rape plants by *Leptosphaeria maculans*. Plant Pathol. *34*, 557–565.

64.   Ponomarenko A., S.B. Goodwin, and G.H.J. Kema. 2011. Septoria tritici blotch (STB)

of wheat. *Plant Health Instructor*. DOI:10.1094/PHI-I-2011-0407-01

65.   Hershman, D.E. (1995). Etiology of canola blackleg in Kentucky and seasonal

discharge patterns of *Leptosphaeria maculans* ascospores from infested canola stubble.

Plant Dis. *v. 79*, 1225–1229.

66.   Petrie, G.A. (Agriculture and A.-F.C. (1995). Patterns of ascospore discharge by

Leptosphaeria maculans (blackleg) from 9- to 13-month-old naturally-infected

rapeseed/canola stubble from 1977 to 1993 in Saskatchewan. Can. Plant Dis. Surv. Can.

67.   Eyal, Z., Scharen, A.L., Prescott, J.M., and Ginkel, M.V. (1987). The septoria diseases

of wheat : concepts and methods of disease management (CIMMYT).

68.   Fitt, B.D.L., Brun, H., Barbetti, M.J., and Rimmer, S.R. (2006). World-wide importance

of phoma stem canker (*Leptosphaeria maculans* and *L. biglobosa*) on oilseed rape

(*Brassica napus*). Eur. J. Plant Pathol. *114*, 3–15.

69. Fesel, P.H., and Zuccaro, A. (2016). Dissecting endophytic lifestyle along the

parasitism/mutualism continuum in *Arabidopsis*. Curr. Opin. Microbiol. *32*, 103–112.

70. Nicolaisen, M., Justesen, A.F., Knorr, K., Wang, J., and Pinnschmidt, H.O. (2014).

Fungal communities in wheat grain show significant co-existence patterns among

species. Fungal Ecol. *11*, 145–153.

71. Larran, S., Perelló, A., Simón, M.R., and Moreno, V. (2007). The endophytic fungi

from wheat (*Triticum aestivum* L.). World J. Microbiol. Biotechnol. *23*, 565–572.

72. Lakhesar, D.P.S., Backhouse, D., and Kristiansen, P. (2010). Nutritional constraints on

displacement of *Fusarium pseudograminearum* from cereal straw by antagonists. Biol.

Control *55*, 241–247.

73. Brader, G., Compant, S., Vescio, K., Mitter, B., Trognitz, F., Ma, L.-J., and Sessitsch,

A. (2017). Ecology and genomic insights into plant-pathogenic and plant-nonpathogenic

endophytes. Annu. Rev. Phytopathol. *55*, 61–83.

74. Leplat, J., Friberg, H., Abid, M., and Steinberg, C. (2013). Survival of *Fusarium*

graminearum, the causal agent of *Fusarium* head blight. A review. Agron. Sustain. Dev.

33, 97–111.

75. Bockus, W.W., and Claassen, M.M. (1992). Effects of crop rotation and residue

management practices on severity of tan spot of winter wheat. Plant Dis. *76*, 633–636.

76. Van Bruggen, A.H.C., and Semenov, A.M. (2000). In search of biological indicators for

soil health and disease suppression. Appl. Soil Ecol. *15*, 13–24.

77. Daverdin, G., Rouxel, T., Gout, L., Aubertot, J.-N., Fudal, I., Meyer, M., Parlange, F.,

Carpezat, J., and Balesdent, M.-H. (2012). Genome structure and reproductive

behaviour influence the evolutionary potential of a fungal phytopathogen. PLoS Pathog.

8, e1003020.

78. Kema, G.H.J., Mirzadi Gohari, A., Aouini, L., Gibriel, H.A.Y., Ware, S.B., van den

Bosch, F., Manning-Smith, R., Alonso-Chavez, V., Helps, J., Ben M'Barek, S., et al.

(2018). Stress and sexual reproduction affect the dynamics of the wheat pathogen

effector AvrStb6 and strobilurin resistance. Nat. Genet. *50*, 375–380.

79. Dugan, F.M., Lupien, S.L., Hernandez-Bello, M., Peever, T.L., and Chen, W. (2005).

Fungi resident in chickpea debris and their suppression of growth and reproduction of

Didymella rabiei under laboratory conditions. J. Phytopathol. *153*, 431–439.

80. Hue, A.G., Voldeng, H.D., Savard, M.E., Fedak, G., Tian, X., and Hsiang, T. (2009).

Biological control of fusarium head blight of wheat with *Clonostachys rosea* strain

ACM941. Can. J. Plant Pathol. *31*, 169–179.

81. Luongo, L., Galli, M., Corazza, L., Meekes, E., Haas, L.D., Van Der Plas, C.L., and

Köhl, J. (2005). Potential of fungal antagonists for biocontrol of *Fusarium spp.* in wheat

and maize through competition in crop debris. Biocontrol Sci. Technol. *15*, 229–242.

82. Li, B.C., and Sutton, J.C. (1995). Evaluation of leaf-associated microorganisms for

biocontrol of tan spot in wheat foliage. Fitopatol. Bras. *20*, 545–552.

83. Silby, M.W., Winstanley, C., Godfrey, S.A.C., Levy, S.B., and Jackson, R.W. (2011).

Pseudomonas genomes: diverse and adaptable. FEMS Microbiol. Rev. *35*, 652–680.

Figure and table captions

Figure 1 - Experimental layout of the experiment. **(A)** Plots (WWW, WOW and OWO) used

during the two cropping seasons of the experiment at the INRA Grignon experimental station

(Yvelines, France). WWW: plot cropped with winter wheat since 2007. WOW and OWO:

plots cropped with a wheat-oilseed rape rotation since 2014. Wheat straw and oilseed rape

debris were chopped at harvest and a large portion of residue was then left on the surface. The

dashed line indicates the sampling transect. **(B)** Oilseed rape residues in a plot cropped with wheat (OWO or WOW). **(C)** Wheat residues in a plot cropped with oilseed rape (WOW or OWO). **(D)** Wheat residues in the wheat monoculture crop (WWW).

Figure 2 - Fungal **(A)** and bacterial **(B)** diversity in plants (July) and residues (October; December; February; May), as assessed with the Shannon index, according to sampling period, the crop within a rotation (oilseed rape in WOW and OWO; wheat in WWW; wheat in WOW and OWO) and the cropping season (2015-2016; 2016-2017). Each box represents the distribution of Shannon index for five sampling points. Kruskal-Wallis tests were performed for each "crop within a rotation * cropping season" combination (*p*-values are given under each graph). Wilcoxon tests between sampling periods were performed when the Kruskal-Wallis test revealed significant differences. Samples not sharing letters are significantly different.

Figure 3 - Structure of the fungal **(A)** and bacterial **(B)** communities present in oilseed rape and wheat residues, according to compositional dissimilarity (Bray-Curtis dissimilarity index), after multidimensional scaling (MDS). The two MDS were performed on the overall dataset and faceted according to the sampling period. Each point represents one sample corresponding to a cropping season (shape: 2015-2016; 2016-2017; 2017-2018) and crop within a rotation (colour: oilseed rape in rotation, i.e. in WOW and OWO; wheat monoculture, i.e. in WWW; wheat in rotation, i.e. in WOW and OWO).

Figure 4 – Distribution of the 60 most prevalent fungal genera detected in wheat residues in the five samples for each sampling date. Unclassified genera were removed from the visualisation.

Figure 5 – Distribution of the 60 most prevalent bacterial genera detected in wheat residues in

the five samples for each sampling date. Unclassified genera were removed from the

visualisation.

Table 1 - Results of the ANOVA performed to assess the effects of plant (wheat; oilseed rape),

cropping season (2015-2016; 2016-2017), sampling period (July, October; December;

February; May) and sampling plot on the Shannon index of the fungal and bacterial

communities present in oilseed rape and wheat residues from the plots in rotation (OWO;

WOW). The effect of the cropping system (monoculture; rotation) was estimated separately

with a second ANOVA performed on the wheat residue samples dataset only (wheat in

monoculture, i.e. in WWW; wheat in rotation, i.e. in WOW and in OWO).

Table 2 - Results of the PERMANOVA performed to assess the effects of plant (wheat;

oilseed rape), cropping season (2015-2016; 2016-2017), sampling period (July; October;

December; February; May) and sampling plot on the Bray-Curtis dissimilarity index of the

fungal and bacterial communities present in oilseed rape and wheat residues from the plots in

rotation (OWO; WOW). The effect of the cropping system (monoculture; rotation) was

estimated separately with a second PERMANOVA performed on the wheat residue samples

data set only (wheat in monoculture, i.e. in WWW; wheat in rotation, i.e. in WOW and in

OWO). PERMANOVAs were performed using the adonis2 function with "margin" option.

Additional files

List of supplementary figures:

Figure S1 - Bacterial diversity in plants (July) and residues (October, December, February, May), as assessed with the Faith's Phylogenetic Diversity index (PD).

Figure S2 - Structure of the bacterial communities present in oilseed rape and wheat residues visualized by incorporating phylogenetic distances using the UniFrac distance matrix.

Figure S3 - Distribution of the most prevalent fungal genera detected in wheat residues.

Figure S4 - Distribution of the most prevalent fungal genera detected in oilseed rape residues.

Figure S5 - Distribution of the most prevalent bacterial genera detected in wheat residues.

Figure S6 - Distribution of the most prevalent bacterial genera detected in oilseed rape residues.

Figure S7 - Seasonal shift in the relative abundance of a selection of fungal and bacterial genera present on wheat and oilseed rape residues.

List of supplementary tables:

Table S1 - Soil texture of the three plots.

Table S2 - Sampling dates of wheat and oilseed rape plants (July) and residues (October, December, February and May) for each cropping season.

Table S3 - Total number of reads and percentage (in brackets) remaining after ASV filtering.

Table S4 - Summary of meteorological data (temperature, rainfall) for the INRA Grignon experimental station (Yvelines, France).

Table S5 - Plant effect (wheat vs. oilseed rape) on community dispersion.

Table S6 - Decomposition of dissimilarity due to temporal changes in fungal (F) and bacterial (B) community.

WOW		
	Crop residues	**Crop**
2015-2016	**W**heat	Oilseed Rape (Cv. Alpaga)
2016-2017	**O**ilseed Rape	Wheat (Cv. Soissons)
July 2017	**W**heat	

WWW		
	Crop residues	**Crop**
2015-2016	**W**heat	Wheat (Cv. Soissons)
2016-2017	**W**heat	Wheat (Cv. Soissons)
July 2017	**W**heat	

OWO		
	Crop residues	**Crop**
2015-2016	**O**ilseed Rape	Wheat (Cv. Soissons)
2016-2017	**W**heat	Oilseed Rape (Cv. Alpaga)
July 2017	**O**ilseed Rape	

Reads number

512   8192   131072

J : July
O : October   F : February
D : December   M : May

Reads number

1024 16384 262144

J : July
O : October   F : February
D : December   M : May

Table 1 - Results of the ANOVA performed to assess the effects of plant (wheat; oilseed rape), cropping season (2015-2016; 2016-2017), sampling period (July, October; December; February; May) and sampling plot on the Shannon index of the fungal and bacterial communities present in oilseed rape and wheat residues from the plots in rotation (OWO; WOW). The effect of the cropping system (monoculture; rotation) was estimated separately with a second ANOVA performed on the wheat residue samples dataset only (wheat in monoculture, i.e. in WWW; wheat in rotation, i.e. in WOW and in OWO).

		Fungi		Bacteria	
	Tested factors	F value	p-value	F value	p-value
Plots in rotation	Plant	22.42	< 0.001	0.33	0.567
	Cropping season	14.19	< 0.001	35.28	< 0.001
	Sampling period	14.25	< 0.001	37.73	< 0.001
	Sampling plot	31.06	< 0.001	61.57	< 0.001
Wheat residues	Cropping system	46.42	< 0.001	23.66	< 0.001
	Cropping season	51.43	< 0.001	33.34	< 0.001
	Sampling period	6.72	< 0.001	13.89	< 0.001

Table 2 - Results of the PERMANOVA performed to assess the effects of plant (wheat; oilseed rape), cropping season (2015-2016; 2016-2017), sampling period (July; October; December; February; May) and sampling plot on the Bray-Curtis dissimilarity index of the fungal and bacterial communities present in oilseed rape and wheat residues from the plots in rotation (OWO; WOW). The effect of the cropping system (monoculture; rotation) was estimated separately with a second PERMANOVA performed on the wheat residue samples data set only (wheat in monoculture, i.e. in WWW; wheat in rotation, i.e. in WOW and in OWO). PERMANOVAs were performed using the adonis2 function with "margin" option.

Data set	Factors	Fungi		Bacteria	
		R^2	*p*-value	R^2	*p*-value
Plots in rotation	Plant	0.384	< 0.001	0.266	< 0.001
	Cropping season	0.131	< 0.001	0.118	< 0.001
	Sampling period	0.068	< 0.001	0.149	< 0.001
	Sampling plot	0.100	< 0.001	0.053	< 0.001
Wheat residues	Cropping system	0.105	< 0.001	0.066	< 0.001
	Cropping season	0.292	< 0.001	0.248	< 0.001
	Sampling period	0.123	< 0.001	0.204	< 0.001

Figure S1 - Bacterial diversity in plants (July) and residues (October, December, February, May), as assessed with the Faith's Phylogenetic Diversity index (PD), according to sampling period, the crop within a rotation (oilseed rape in OWO or WOW, wheat in WWW, wheat in WOW or OWO) and the cropping season (2015-2016, 2016-2017). Each box represents the distribution of PD for five sampling points. Kruskal-Wallis tests were performed for each "crop within a rotation * cropping season" combination (p-values are given under each graph). Wilcoxon tests between sampling periods were performed when the Kruskal-Wallis test revealed significant differences. Samples not sharing letters are significantly different.

Figure S2 - Structure of the bacterial communities present in oilseed rape and wheat residues visualized by incorporating phylogenetic distances using the UniFrac distance matrix. MDS were performed on the overall dataset and faceted according to the sampling period. Each point represents one sample corresponding to a cropping season (shape: 2015-2016; 2016-2017; 2017-2018) and crop within a rotation (colour: oilseed rape in rotation, i.e. in WOW and OWO; wheat monoculture, i.e. in WWW; wheat in rotation, i.e. in WOW and OWO).

Figure S3 - Distribution of the most prevalent fungal genera detected in wheat residues. (**A**) Cladogram of the most prevalent genera. Genera were filtered according to their occurrence (at least three times in the five sampling points for each "crop within a rotation * cropping season * sampling period" combination). Unclassified genera were removed from the tree. (**B**) Number of ASV of each genus. (**C**) Occurrence of each ASV in the 100 samples of wheat residues. (**D**) Percentage of reads for each genus. (**E**) Distribution of each genus in the five samples per date (increasing number of reads shown on a scale running from yellow to red).

Figure S4 - Distribution of the most prevalent fungal genera detected in oilseed rape residues. (**A**) Cladogram of the most prevalent genera. Genera were filtered according to their occurrence (at least three times in the five sampling points for each "crop within a rotation * cropping season * sampling period") combination. Unclassified genera were removed from the tree. (**B**) Number of ASV for each genus. (**C**) Occurrence of each ASV in the 50 samples of oilseed rape residues. (**D**) Percentage of reads for each genus. (**E**) Distribution of each genus in the five samples per date (increasing number of reads shown on a scale from yellow to red).

Figure S5 - Distribution of the most prevalent bacterial genera detected in wheat residues. **(A)** Cladogram of the most prevalent genera. Genera were filtered according to their occurrence (at least three times in five sampling points for each "crop within rotation * cropping season * sampling period" combination). Unclassified genera were removed from the tree. **(B)** Number of ASV for each genus. **(C)** Occurrence of each ASV in the 100 samples of wheat residues. **(D)** Number of reads for each genus. **(E)** Distribution of each genus in the five samples per date (increasing numbers of reads on a scale running from yellow to red). Due to the number of genera, the plot is separated in [1] proteo- and [2] other bacteria.

A — Taxon
B — Number of haplotypes
C — Number of reads
D — Occurrence
E — Wheat (Monoculture) / Wheat (Rotation)

Taxon	Number of haplotypes	Occurrence
Neochlamydia	8	11
Fimbriimonas	4	17
Deinococcus	3	28
Gemmatimonas	3	6
Nitrolancea	4	59
Litorilinea	4	12
Tepidisphaera	7	37
Pirellula	3	10
Gemmata	5	7
Opitutus	18	20
Cerasicoccus	3	6
Luteolibacter	18	76
Verrucomicrobium	13	71
Roseimicrobium	5	9
Prosthecobacter	18	33
Carnobacterium	1	5
Lactobacillus	1	4
Enterococcus	2	7
Lysinibacillus	1	8
Exiguobacterium	1	10
Staphylococcus	2	15
Bacillus	3	12
Fluviicola	18	57
Epilithonimonas	8	82
Chryseobacterium	13	97
Moheibacter	1	22
Flavobacterium	151	100
Spirosoma	14	95
Hymenobacter	72	99
Fibrella	3	66
Adhaeribacter	9	41
Larkinella	5	32
Rhodocytophaga	5	9
Cytophaga	10	28
Dyadobacter	49	100
Rudanella	2	4
Rubrivirga	1	13
Mucilaginibacter	49	97
Pedobacter	79	100
Sphingobacterium	22	49
Arcticibacter	4	10
Ferruginibacter	24	88
Flavisolibacter	19	79
Taibaiella	25	88
Flavitalea	5	51
Parasegetibacter	6	29
Chitinophaga	11	49
Segetibacter	9	53
Terrimonas	20	83
Gaiella	4	14
Aquihabitans	3	12
Ferrimicrobium	1	7
Patulibacter	2	51
Solirubrobacter	9	38
Conexibacter	11	32
Streptomyces	6	47
Sanguibacter	2	79
Promicromonospora	3	32
Lentzea	2	8
Nakamurella	5	48
Mycobacterium	6	80
Cellulomonas	5	49
Arthrobacter	4	56
Williamsia	2	50
Rhodococcus	5	91
Kineococcus	3	90
Kineosporia	7	84
Pseudokineococcus	1	25
Intrasporangium	1	10
Phycicoccus	1	49
Aquipuribacter	2	15
Nocardioides	39	97
Aeromicrobium	8	98
Marmoricola	4	67
Blastococcus	3	32
Geodermatophilus	1	100
Modestobacter	2	18
Actinoplanes	7	78
Micromonospora	6	24
Verrucosispora	1	15
Phytohabitans	1	13
Catenuloplanes	1	44
Rathayibacter	1	82
Frondihabitans	2	99
Herbiconiux	1	26
Clavibacter	1	84
Cryobacterium	1	39
Microbacterium	1	10
Agromyces	3	6
Leucobacter	1	12

Number of reads scale: 5000000

Wheat (Monoculture): 2015 - 2016 (Oct., Dec., Feb., May), 2016 - 2017 (Jul., Oct., Dec., Feb., May), 2017 (Jul.)

Wheat (Rotation): 2015 - 2016 (Oct., Dec., Feb., May), 2016 - 2017 (Jul., Oct., Dec., Feb., May), 2017 (Jul.)

Color scale: 1 — 348166

[2]

Figure S6 - Distribution of the most prevalent bacterial genera detected in oilseed rape residues. **(A)** Cladogram of most prevalent genera. Genera were filtered according to their occurrence (at least three times in five sampling points for each "crop within rotation * cropping season * sampling period" combination). Unclassified genera were removed from the tree. **(B)** Number of ASV for each genus. **(C)** Occurrence of each ASV in the 49 samples of oilseed rape residues. **(D)** Number of reads for each genus. (E) Distribution of each genus in the five samples per date (increasing number of reads shown on a scale running from yellow to red). Due to the number of genera, the plot is separated in [1] proteo- and [2] other bacteria.

	A	B	C	D	E
	Taxon	Haplotypes number	Reads percent	Occurrence	

Column E header: 2015 - 2016 (Oct. Dec. Feb. May), 2016 - 2017 (Jul. Oct. Dec. Feb. May), 2017 (Jul.)

Taxa (top to bottom):
Parachlamydia 11, 10
Fimbriimonas 4, 16
Truepera 3, 8
Gemmatimonas 3, 10
Ignavibacterium 2, 6
Nitrolancea 4, 34
Litorilinea 4, 20
Lactobacillus 1, 3
Lysinibacillus 1, 5
Bacillus 3, 8
Opitutus 18, 29
Cerasicoccus 3, 3
Luteolibacter 18, 41
Verrucomicrobium 13, 29
Prosthecobacter 18, 36
Tepidisphaera 7, 7
Rubinisphaera 1, 9
Pirellula 3, 15
Gemmata 5, 11
Zavarzinella 5, 6
Fluviicola 18, 39
Crocinitomix 1, 5
Epilithonimonas 8, 21
Chryseobacterium 13, 47
Moheibacter 1, 34
Flavobacterium 151, 50
Spirosoma 14, 6
Hymenobacter 72, 26
Fibrella 3, 5
Adhaeribacter 9, 16
Larkinella 5, 16
Rhodocytophaga 5, 12
Cytophaga 10, 26
Dyadobacter 49, 50
Pedobacter 79, 50
Solitalea 3, 10
Sphingobacterium 22, 49
Arcticibacter 4, 26
Mucilaginibacter 49, 24
Olivibacter 4, 34
Sediminibacterium 3, 12
Ferruginibacter 24, 26
Flavihumibacter 1, 8
Flavisolibacter 19, 27
Taibaiella 25, 45
Flavitalea 5, 35
Niastella 3, 6
Chitinophaga 11, 19
Hydrotalea 1, 10
Segetibacter 9, 13
Terrimonas 20, 40
Solirubrobacter 9, 16
Conexibacter 11, 25
Aquihabitans 3, 24
Ferrimicrobium 1, 9
Streptomyces 6, 25
Sanguibacter 2, 47
Promicromonospora 3, 37
Nakamurella 5, 5
Mycobacterium 6, 20
Nocardiopsis 2, 10
Cellulomonas 5, 22
Phycicoccus 1, 22
Williamsia 2, 15
Rhodococcus 5, 34
Amycolatopsis 1, 9
Lentzea 2, 12
Kineococcus 3, 14
Kineosporia 7, 11
Nesterenkonia 1, 6
Arthrobacter 4, 32
Blastococcus 3, 8
Geodermatophilus 1, 19
Actinoplanes 7, 28
Micromonospora 6, 16
Catenuloplanes 1, 19
Nocardioides 39, 49
Aeromicrobium 8, 50
Marmoricola 4, 14
Frondihabitans 2, 18
Herbiconiux 1, 11
Clavibacter 1, 7
Cryobacterium 1, 22
Lysinimonas 1, 3
Microbacterium 1, 12
Leifsonia 1, 19
Leucobacter 1, 7
Agromyces 3, 28

Scale bar: 5000000 (Reads percent)
Heatmap color scale: 1 to 386282

Figure S7 - Seasonal shift in the relative abundance of a selection of fungal **(A)** and bacterial **(B)** genera present on wheat and oilseed rape residues according to the system (wheat monoculture **[1]**, wheat in rotation **[2]**, oilseed rape in rotation **[3]**) and the cropping season (2015-2016, 2016-2017). Due to the plant impact (wheat and oilseed rape) in the fungal community, the fungal genera used here as examples are different for the two plants, unlike the case of the bacterial community. Each box represents the distribution of genera relative abundance for the five sampling points. Samples not sharing letters are significantly different (Wilcoxon tests between sampling periods).

A.1.

A.2.

A.3.

B.3.

Table S1 - Soil texture of the three plots (WWW, OWO and WOW).

	WWW, OWO	WOW
Clay (%)	27.4	18.2
Silt (%)	53.2	61.2
Sand (%)	18.8	20.4

Table S2 - Sampling dates of wheat and oilseed rape plants (July) and residues (October, December, February, and May) for each cropping season.

Cropping season	July	October	December	February	May
2015-2016	/	23.10.2015	04.12.2015	26.02.2016	19.05.2016
2016-2017	11.07.2016	17.10.2016	05.12.2016	06.02.2016	15.05.2017
2017-2018	07.07.2017	/	/	/	/

Table S3 - Total number of reads and percentage (in brackets) remaining after ASV filtering.

	After DADA2	After replicate suppression	After taxon suppression
Bacterial reads	14,287,970	13,509,461 (94.6%)	13,228,976 (92.6%)
Fungal reads	9,898,487	9,753,628 (98.5%)	9,628,995 (97.3%)
Bacterial haplotypes	19,235	2,905	2,726
Fungal haplotypes	3,587	1,241	1,189

Table S4 - Summary of meteorological data (temperature, rainfall) for the INRA Grignon experimental station (Yvelines, France), obtained from the CLIMATIK INRA database (https://intranet.inra.fr/climatik_v2/) from July 1st to May 31st of the following year, for the cropping seasons 2015-2016 and 2016-2017.

	10-day mean temperature (°C)		10-day cumulative rainfall (mm)	
	2015-2016	2016-2017	2015-2016	2016-2017
Mean	11.2	10.8	22.6	12.3
Minimum	2.0	0.9	0	0
Maximum	21.8	21.4	131	55

Table S5 - Plant effect (wheat vs. oilseed rape) on community dispersion. This effect was tested by applying the adonis function of the vegan R package to the Bray-Curtis dissimilarity index (PERMANOVA). *P*-values (not shown) were all < 0.02.

	Fungi				Bacteria			
	All	2015-2016	2016-2017	2017	All	2015-2016	2016-2017	2017
July	0.372	/	0.611	0.755	0.423	/	0.540	0.696
October	0.495	0.612	0.755	/	0.367	0.520	0.659	/
December	0.486	0.688	0.691	/	0.370	0.573	0.641	/
February	0.429	0.541	0.651	/	0.409	0.643	0.611	/
May	0.273	0.337	0.401	/	0.315	0.435	0.508	/

Table S6 - Decomposition of dissimilarity due to temporal changes in fungal (F) and bacterial (B) community composition. Total dissimilarity is broken down into turnover (replacement of ASV) and nestedness (gain or loss of ASV).

Crop within a rotation	Season	Sampling period compared	Total dissimilarity		Turnover		Nestedness	
			F	B	F	B	F	B
Oilseed rape	2015-2016	Oct. - Dec.	0.622	0.318	0.618	0.219	0.005	0.099
Oilseed rape	2015-2016	Dec. - Feb.	0.650	0.321	0.577	0.290	0.073	0.031
Oilseed rape	2015-2016	Feb. - May	0.591	0.390	0.565	0.202	0.027	0.188
Oilseed rape	2016-2017	Jul. - Oct.	0.652	0.554	0.648	0.250	0.004	0.304
Oilseed rape	2016-2017	Oct. - Dec.	0.620	0.353	0.549	0.217	0.071	0.136
Oilseed rape	2016-2017	Dec. - Feb.	0.585	0.353	0.516	0.276	0.068	0.077
Oilseed rape	2016-2017	Feb. - May	0.529	0.384	0.529	0.342	0.000	0.042
Wheat in monoculture	2015-2016	Oct. - Dec.	0.427	0.330	0.425	0.142	0.002	0.188
Wheat in monoculture	2015-2016	Dec. - Feb.	0.444	0.294	0.416	0.190	0.028	0.104
Wheat in monoculture	2015-2016	Feb. - May	0.444	0.458	0.424	0.255	0.020	0.203
Wheat in monoculture	2016-2017	Jul. - Oct.	0.438	0.346	0.424	0.300	0.014	0.046
Wheat in monoculture	2016-2017	Oct. - Dec.	0.463	0.330	0.257	0.113	0.207	0.217
Wheat in monoculture	2016-2017	Dec. - Feb.	0.386	0.248	0.311	0.200	0.075	0.048
Wheat in monoculture	2016-2017	Feb. - May	0.344	0.332	0.341	0.213	0.004	0.120
Wheat in rotation	2015-2016	Oct. - Dec.	0.425	0.317	0.409	0.157	0.016	0.160
Wheat in rotation	2015-2016	Dec. - Feb.	0.472	0.266	0.370	0.185	0.102	0.081
Wheat in rotation	2015-2016	Feb. - May	0.505	0.347	0.432	0.311	0.073	0.035
Wheat in rotation	2016-2017	Jul. - Oct.	0.498	0.313	0.427	0.272	0.071	0.041
Wheat in rotation	2016-2017	Oct. - Dec.	0.541	0.287	0.292	0.214	0.249	0.073
Wheat in rotation	2016-2017	Dec. - Feb.	0.350	0.284	0.292	0.284	0.059	0.000
Wheat in rotation	2016-2017	Feb. - May	0.424	0.334	0.329	0.287	0.095	0.047
Mean			0.496	0.341	0.436	0.234	0.060	0.107