Preclinical Modeling of Exposure to a Global Marine Bio-Contaminant:

Effects of In Utero Domoic Acid Exposure on Neonatal Behavior and Infant Memory

Kimberly S. Grant[1,3], Brenda Crouthamel[1,3], Caroline Kenney[3], Noelle McKain[1,3], Rebekah Petroff[1], Sara Shum[4], Jing Jing[4], Nina Isoherranen[2,4] and Thomas M. Burbacher[1,2,3]

[1]Department of Environmental and Occupational Health Sciences, University of Washington, Seattle, Washington, USA, [2]Center on Human Development and Disability, University of Washington, Seattle, Washington, USA, [3]Washington National Primate Research Center, Seattle, Washington, USA, [4]Department of Pharmaceutics, University of Washington, Seattle, Washington, USA

Running Title: Domoic Acid and Infant Development

Corresponding author: Kimberly S. Grant

Abstract

Domoic Acid (DA) is a naturally-occurring marine neurotoxin that is increasingly recognized as an important public health issue. Prenatal DA exposure occurs through the maternal consumption of contaminated shellfish/finfish. To better understand the fetal risks associated with DA, we initiated a longitudinal, preclinical study focused on the reproductive and developmental effects of chronic, low-dose oral DA exposure. To this end, 32 adult female Macaca fascicularis monkeys were orally dosed with 0, 0.075 or 0.15 mg/kg/day DA on a daily basis throughout breeding and pregnancy. The doses included the proposed human Tolerable Daily Intake (TDI) (0.075 mg/kg/day) for DA. Adult females were bred to nonexposed males. To evaluate behavioral development during early infancy, offspring were administered a neonatal exam modeled after the human Neonatal Behavior Assessment Scale and a series of visual recognition memory problems. Results indicated that DA does not impact early survival reflexes or responsivity to the environment during the neonatal period. Findings from recognition memory tests showed that exposed and control infants demonstrated robust novelty scores when test problems were relatively easy to solve. However, infants in the 0.15 mg/kg DA group failed to demonstrate a novelty response when problems became more challenging. This study is the first to evaluate the offspring neurodevelopmental effects of prenatal DA exposure in a translational primate model and results indicate subtle but significant consequences on emerging memory processes.

Keywords: domoic acid, prenatal exposure, infant, memory, macaque

1.0 Introduction

Domoic Acid (DA) is a naturally-occurring marine biotoxin that is usually produced by microalgae from the genus *Pseudo-nitzschia*. Mammalian exposures, including human, occur through the consumption of contaminated shellfish and finfish and elevated levels of DA have been documented in ocean waters and coastal regions around the world (Lefebvre and Robertson, 2010, https://www.whoi.edu/science/B/redtide/HABdistribution/HABmap.html.) Based upon an acute poisoning episode involving contaminated mussels on Prince Edward (PEI), Canada in 1987, human symptomology is known to include gastrointestinal distress, confusion, transient and permanent memory loss, coma and death (Perl et al., 1990 a,b). Indeed, the clinical label for human DA poisoning, "Amnesic Shellfish Poisoning", reflects its distinct action on the nervous system that selectively targets cognition. Imaging studies of PEI survivors using Positron Emission Tomography indicated reductions in cerebral glucose metabolism in the hippocampus and the amygdala (Teitelbaum et al, 1990a,b). While the episode of human poisoning in Canada provided data related to the acute effects of DA exposure at very high levels (the estimated exposures ranged from 60 to 290 mg DA or about 1 to 5 mg/kg for a 60 kg person (Perl 1990b, Jefferey et al., 2004), the more subtle effects of sub-toxic, oral DA exposure are largely unknown. Of specific concern are the potential impacts of chronic, low-dose exposure in populations where high shellfish consumption is the dietary norm (Grattan et al., 2016a). Recent results from an environmental epidemiology study focused on chronic, low dose DA exposure reported that the high consumption of razor clams (>15 per week) in adult members of three Native American tribes living on the Western coast of the U.S. was associated with significant memory decrements when compared to a reference group of non-consumers or low-consumers (Grattan et al., 2016b). Deficits in memory were significant enough to impact daily living skills (Grattan et al., 2018).

To date, the developmental neurotoxicity of DA, the focus of this study, has been primarily explored in preclinical rodent models and adverse effects are known to be dose and route dependent (for review see Grant et al., 2010). While the literature base is relatively small, a theme of heightened fetal sensitivity characterized by hippocampal injury and neurological effects can be extracted from the literature (for reviews see Costa et al., 2010; Lefebvre and Robertson, 2010; Doucette et al., 2016). Domoic acid readily crosses the placenta and enters the fetal brain but does not produce structural malformations in exposed pups (Maucher Fuquay et al., 2012). Despite cessation of DA exposure at birth, progressive damage to regions of the hippocampus have been documented over the first 30 days of postnatal life (Dakshinamurti et al., 1993). Results from a number of studies suggest that a single in-utero exposure can cause

long-term changes in behavioral development, primarily targeting cognition, social behavior and emotionality and experimental results are frequently accompanied by gender-related differences in performance (Levin et al., 2005, Doucette et al., 2007; Adams et al., 2008; Burt et al., 2008; Tanemura et al., 2009, Ryan et al., 2011; Gill et al., 2012; Marriott et al., 2012, Zuloaga et al., 2016, Shiotani et al., 2017). Early postnatal exposure is associated with a time-dependent profile of neurotoxicity that is characterized by hyperactivity, stereotypic scratching, convulsions and death (Xi et al, 1997). These effects occurred at DA doses 40 times lower than what is required to produce similar clinical symptomology in adult rodents. Reductions in prepulse inhibition (PPI), the normal suppression of the startle response produced by a preceding, non-startling cue (Graham, 1975), have been observed in DA-exposed animals after developmental exposure (Marriot et al, 2012, Adams, 2008). This phenomenon is frequently disrupted in humans with various psychiatric/neurological disorders such as schizophrenia (Takahashi et al., 2011, Haß et al., 2017). To date, there have been no published reports on the effects of prenatal or postnatal DA exposure in human or nonhuman primate infants.

As described above, there is robust evidence from rodent studies that DA can pose significant risks to the fetus but the real-world danger of prenatal exposure in humans has not been explored. Defining these risks to protect human health will depend upon our understanding of this complex chemical when ingested orally at low doses on a chronic basis. This is particularly important because the harmful algal blooms that produce DA are becoming more toxic and widespread as physical processes in ocean waters adapt to warming temperatures (McKibben et al., 2016; Zhu et al., 2017). The research reported in this manuscript is the first to investigate the developmental toxicity of chronic, low-level, oral DA exposure in a nonhuman primate model. The macaque monkey model was selected for this study based on key maternal reproductive and infant developmental similarities shared with humans (Grant and Rice, 2008). The results presented in this paper focus on neonatal behavior and infant visual recognition memory. The dosing levels selected for this investigation were strongly focused on environmental relevance and bracket the proposed human Tolerable Daily Intake (TDI) for DA (Marien, 1996, Costa et al., 2010, Wekell et al., 2004). Results from this investigation provide meaningful translational data on the fetal risks of maternal oral DA exposure during pregnancy in a closely-related species.

2.0 Materials and Methods

2.1 Chemicals and Reagents

DA was purchased from BioVectra (Charlottetown, PE, Canada). Certified calibration solution for DA was purchased from National Research Council Canada (Ottawa, ON, Canada). Optima grade water, methanol, acetonitrile, and formic acid used for bioanalytical assays were purchased from Fisher Scientific (Pittsburgh, PA).

2.2 Animal Selection and Study Design

Detailed descriptions of the animals and study design have been reported in Burbacher et al., 2018. Briefly, 32 adult female *Macaca fascicularis* were assigned to either a control group (N=10), a 0.075 mg/kg/day DA group (N=11) or a 0.15 mg/kg/day DA group (N=11) to establish groups with similar average ages and weights. The average age of females was ~ 7 years of age and the average weight was ~3.5 Kg. All females were nulliparous. Three adult males were used for breeding; males were between 7.5 and 8.5 years of age and weighed between 5.4 and 7.9 kg. The study design was based on past studies from our laboratory of in utero exposure to other developmental neurotoxicants such as methylmercury and methanol (Burbacher et al., 1988, 2004) and included a prebreeding, breeding, pregnancy and delivery period. Protocols were implemented to observe signs of DA toxicity (Clinical Observations), detect menstrual bleeding, evaluate general health status (weights, body index) and collect blood for DA analysis without sedation. The adult females had unrestricted access to water and were fed Lab Diet High Protein Monkey Diet biscuits twice a day, once approximately 2 hours before the daily DA dose and once approximately 5 hours after dosing. Animals were housed in individual communicating cages that were equipped with grooming bars that allowed adjacent females contact 24 hours/day. Frozen treats, fresh fruit, music, animated movies and toys were provided to the females on a routine basis. The adult males were housed individually in a separate room away from the females and provided the same suite of enrichments. All animal procedures strictly followed the Animal Welfare Act and the Guide for Care and Use of Laboratory Animals of the National Research Council. Study protocols were approved by the University of Washington Institutional Animal Care and Use Committee to meet the highest standards of ethical conduct and compassionate use of animals in research.

2.3 Maternal DA Exposure

Females were dosed daily with DA solution or sweetened water prior to breeding and throughout pregnancy (Jing et al., 2018). Dosing was done in a blinded manner and all dosing

solutions were quality controlled and analyzed for DA. To conduct voluntary oral dosing in this study, females were trained with positive reinforcement to drink sweetened water from a syringe offered by the tester. The study protocol required that females continue to be trained until they were reliably drinking from a syringe with no visible spillage. To insure all testers were blind to the dosing history of the animals, solutions containing both DA and 5% sugar water or just 5% sugar water alone were made weekly and stored at 4°C with only animal identifiers on the vials. Each morning, the tester would vortex the vial for approximately 5-10 seconds, and draw 1 ml of solution into a syringe. Dosing occurred at approximately the same time every morning, seven days a week. Dosing was calculated on a mg/kg basis based on weekly weights prior to pregnancy. Males were not exposed to DA. Detailed descriptions related to maternal DA exposure and plasma DA concentrations can be found in Burbacher et al., 2018. Briefly, the plasma DA concentrations 5 hours after dosing were significantly higher for the 0.15 mg/kg/day DA exposure group when compared to the 0.075 mg/kg/day DA exposure group both before and during pregnancy (3.42 ng/ml vs 1.31 ng/ml, before pregnancy; 2.93 ng/ml vs 0.93 ng/ml, during pregnancy).

2.4 Offspring

Twenty-eight infants were delivered over a 1 year period, 9 in the control group, 9 in the 0.075 mg/kg/day DA group and 10 in the 0.15 mg/kg/day DA group. Infants were nursery-reared at the Infant Primate Research Laboratory at the University of Washington. This laboratory is equipped to provide 24/7 husbandry and veterinary care for infant monkeys and has set the contemporary standards for the nursery rearing of nonhuman primates (Sackett et al., 2006). Briefly, DA-exposed and control infants were bottle-fed and housed in a temperature-controlled nursery with human incubators until they were able regulate their body temperatures and self-feed. At about 3 weeks of age, animals were transferred to the main rearing quarters and housed in single cages equipped with hanging surrogates and cloth diapers. To facilitate normal social development and future reproductive competence, subjects were placed in stable, mixed-sex groups five days/week in playrooms equipped with a ramp, shelves, swinging chains, toys and a viewing window for tester observations. There was a high rate of C-section deliveries in all of the DA exposure groups. Offspring characteristics such as birth weight, crown-rump length and head circumference were similar across the 3 experimental groups (Burbacher et al., 2018). One female in the 0.15 mg/kg/day DA exposure group was vaginally delivered pre-term (<150 days) and had the lowest birthweight at 240

grams. The remaining infants exhibited birthweights and anthropometric measures consistent with normal, full-term newborns for this species.

2.5 Maternal and Infant Plasma DA Concentrations: Maternal plasma DA concentrations at delivery were significantly higher than newborn plasma DA concentrations for both DA exposure groups (2.23 ng/ml vs 1.26 ng/ml for the 0.15 mg/kg/day DA exposure group and 1.64 ng/ml vs 0.44 ng/ml for the 0.075 mg/kg/day DA exposure group). Newborn plasma DA concentrations were significantly higher in the 0.15 mg/kg/day DA exposure group compare to the 0.075 mg/kg/day DA exposure group (1.26 ng/ml vs 0.44 ng/ml) (for details, see Burbacher et al., 2018). Plasma DA concentrations decreased after birth and were below the limit of quantification (0.31 ng/ml) by 36 hours of life.

2.6 Offspring Assessments

All testers who participated in the evaluation of infants were blind to the exposure history of the animals. Behavioral testers were trained by experienced research personnel and then tested for reliability before collecting data.

Neonatal Assessment of Behavior: Our neonatal assessment is adapted from the Brazelton Neonatal Behavior Assessment Scale (NBAS) used with human infants (Brazelton and Nugent, 2011). The assessment is composed of 5 reflex items and 6 behavioral responses as well as an overall evaluation of the behavioral state of the infant during testing. Tests began with removing the infant from their home-cage and swaddling the infant in a cloth diaper, leaving the upper and lower limbs exposed. Initially, the grasping response of the hands and clasping of the feet were evaluated by placing the index finger against the palm of the hand and sole of the feet. Resistance of the arms and legs to extension was then evaluated by gently but firmly pulling the limbs away from the body. To elicit the rooting and snout reflex, the infant was touched on the corners of the mouth and then on the nose, respectively. Sucking was evaluated by inserting a nipple in the mouth and evaluating the strength of the sucking response. Next, a metal lid was dropped onto a counter to elicit an auditory startle response. The tester then made a lip-smacking noise at the back of each ear to elicit the auditory orienting response. A small colorful toy was held in front of the infant and slowly moved from side to side to examine visual orientation and following. The infant was held above the counter and then quickly moved in the direction of the counter to elicit placing. The arms and legs of the infant were then placed around a rolled-up diaper and strength of grasp evaluated. On the

last test item, the infant was placed in a supine position to elicit the righting response. Finally, the tester scored the general behavioral state and temperament of the infant during the evaluation. Offspring were tested every other day during the first 3 weeks of life.
.

Assessment of Object and Social Visual Recognition Memory: A series of recognition memory problems adapted from the Fagan Test of Infant Intelligence (FTII) and the work of Jocelyn Bachevalier at Emory University (Zeamer et al., 2015) were used to evaluate early cognitive processing. Using the paired-comparison paradigm (Fantz, 1956, 1964), this test makes use of the infant's propensity to prefer novel over familiar visual stimuli and is considered a measure of emerging information processing skills (for review see Burbacher and Grant, 2012). Trials were presented in a preferential looking apparatus where age-graded digitized images were presented on two computer monitors (see Figure 1) at 200 and 210 days of postconceptual age (roughly equivalent to postnatal ages 35 and 45).

For each trial, the examiner held the swaddled infant in front of the images on the computer monitors and recorded visual fixations to the test stimuli. Looking duration and

frequency data were obtained from foot pedals depressed by the tester when the infant actively looked at the right or left stimulus. A physical barrier prevented the examiner from seeing the test stimuli. On immediate (no delay) recognition memory problems, the infant was presented with 2 identical visual stimuli until a predetermined amount of looking time had accrued (familiarization time varied from 20 to 30 seconds). When the familiarization period was completed, a 2-part test trial was presented where the previously-viewed stimulus was paired with a novel image. Each part of the test trial (A and B) was 5 seconds in length (10 seconds total) and triggered by the first directed look of the infant to one of the two test stimuli. The position of the familiar and novel stimuli was reversed between Part A and B of the test trial to control for side preferences. Twelve problems using simple colored images of everyday objects were presented during the initial testing session, which included delays (0, 10, 30 seconds) in between the familiarization period and the test trials. Four memory problems using social stimuli (macaque faces) were presented during the second testing session. At the end of each session, the examiner provided a rating of the subjects overall behavior. Ratings ranged from 1 (asleep or drowsy) to 5 (extremely agitated, untestable). Prior to testing, infants were given a human visual acuity screening test to ensure that each infant had a minimum of 20/800 vision (Teller and Boothe, 1979, Teller, 1997).

3.0 Statistics and Results

3.1 Neonatal Assessment of Behavior: The 5 reflex items, 6 behavioral-response items and the overall evaluation of attention, irritability and consolability during the assessment were used to develop 4 independent factors (Behavioral State, Reflexes, Muscle Tone and Responsivity (see Table 1) in a manner similar to that used in previous studies (Jacobson et al., 1984, Burbacher et al., 1999, Sagiv et al., 2009). Behavioral State was scored from 1 (attentive, cooperative and easy to console) to 4 (inattentive, fussy, and difficult to console). Reflex, Muscle Tone and Responsivity items were scored from weak (1) to strong (4) based on the strength of the response. The overall mean of the items included in the factor was calculated for each test. For the analysis, data were averaged over 2-day blocks beginning on day 2 of age. If infants were tested more than once within a 2-day block, the average score over the 2 days was used in the analysis.

Table 1. Distribution of Test Items for Neonatal Behavioral Scale

Factors	Factor Test Items

Behavioral state	Attention	Irritability	Consolability		
Reflexes	Rooting	Righting	Snout	Placing	Sucking
Muscle tone	Resistance	Grasping	Clasping		
Responsivity	Startle	Orient[1]	Follow[2]		

[1] Orient to Auditory and Visual Stimuli

[2] Follow Visual Stimuli

A summary of the results for the 4 factors from the neonatal assessment is shown in Figure 2. Repeated measures ANOVA models were used to compare the scores for the different DA exposure groups over time, as well as to examine whether or not sex of the offspring or delivery type (C-section or natural delivery) affected the outcomes. Results of the ANOVAs did not indicate a DA exposure effect on any of the 4 factors ($p > 0.05$, all tests). Scores for 3 of the 4 factors changed with age (Reflexes, Muscle Tone and Responsivity) ($p < 0.02$, all test). Sex of the offspring and delivery type did not affect the outcomes directly and these factors did not interact with DA exposure to affect the outcomes ($p > 0.05$, all tests).

Figure 2. Results of Neonatal Assessment during First 3 Weeks of Postnatal Life

3.2 Assessment of Object and Social Visual Recognition Memory: Percent looking time to the novel stimulus during test trials (novelty score) is consider a proxy for recognition memory as some aspects of the familiar stimulus must be encoded in memory for the novelty response to occur. For the initial test session that included delays, novelty scores were calculated for each problem and a mean novelty score was then calculated for the 3 problems within each of the 3 delay periods. For the second test session using social stimuli, novelty scores were calculated for each of the 4 problems and a mean novelty score was then calculated across the problems. Data from infants with ratings of 1 (asleep, drowsy) or 5 (untestable) were not used in the analysis. The novelty preference score is the primary outcome for this test but additional variables can be extracted from the test sessions to better understand the patterns of looking to the stimuli presented during the familiarization and test periods. These variables include mean test fixation (average duration of fixations directed to test stimuli), shift rate (# of times fixations shift from one target to the other) and mean exposure time (time required to reach familiarization criteria). These variables are particularly valuable in cases where a group novelty preference is not displayed.

For the initial test session, the mean novelty scores obtained for each infant for the 3 delay periods were used to test the performance of the DA exposure groups against chance looking (50%) in a single sample t-test. The same process was used to test the performance of the DA exposure groups against chance for the second test session using social stimuli. The results of the initial test session indicated that each of the 3 DA exposure groups exhibited novelty scores significantly above chance for all of the delay periods (p<0.05, all tests). The novelty scores for the control and 0.075 mg/kg/day DA exposure group on the second test using social stimuli were also significantly above change (p<0.05). However, scores for animals in the 0.15 mg/kg/day DA exposure group were not significantly above chance (p=0.29) on these test problems, indicating random looking behavior.

Analysis of variance models were used to test for differences in the performance of the DA exposure groups as well as to examine whether or not sex of the offspring or delivery type (C-section or natural delivery) affected the outcomes. Results did not indicate significant differences in the novelty scores for the 3 DA exposure groups for either test session ($p>0.05$) (see Table 2). Sex of the offspring and delivery type did not affect the outcomes directly and these factors did not interact with DA exposure to affect the outcomes ($p> 0.05$, all tests).

Table 2: Mean ± SE (Range) Percent Looking to Novel Stimuli[1]

DA Exposure Group	Delay Test			Social Stimuli Test
	0 Second	10 Second	30 Second	
Controls (N=9)	60.1±4.0	75.4±3.5	68.8±5.5	62.4±4.0[2]
	(31.7-74.2)	(59.2-95.0)	(47.8-95.0)	(44.4-73.3)
0.075 mg/kg/day (N=9)	60.7±4.4	66.6±4.3	70.0±4.9	63.3±3.5
	(42.2-85.5)	(54.1-85.4)	(51.2-96.5)	(39.1-72.4)
0.15 mg/kg/day (N=10)	74.7±4.5	67.8±4.8	67.8±6.1	**53.9±3.5***
	(51.4-94.9)	(34.0-85.4)	(44.0-100.0)	(34.7-73.3)

[1] Values are presented as means ± SE percent time looking at novel stimuli and range of novelty scores.

[2] N=8 (1 infant tested at wrong age, dropped from analysis)

*No significant novelty preference, $p<.05$

Discussion

This study is the first to investigate the effects of prenatal DA exposure on infant neurodevelopment in a preclinical nonhuman primate model. Neonatal behavioral examinations in the nursery indicated that there were no significant differences between control and DA-exposed infants on Brazelton-derived factors of behavioral state, reflexes, muscle tone and responsivity. Findings from the recognition memory assessment at approximately 35 and 45 days postnatal showed that controls and DA-exposed infants demonstrated significant novelty preferences (memory scores) when test problems were composed of simple, highly-discriminable images of everyday objects. It was only when complex social stimuli with decreased discriminability were used that infants in the 0.15 mg/kg DA group failed to provide

evidence of memory. In studies with human infants, problems using social stimuli are considered more difficult (i.e. require greater study time to elicit a comparable novelty response) than problems using images of abstract patterns (Rose, 1980). The sensitivity of social stimuli to individual differences in information processing is widely recognized and the reason that this class of stimuli is employed in human studies with both clinical and research cohorts, including infants exposed to chemicals in-utero.

The novelty test paradigm was selected for use in this study because demonstration of a novelty response requires the ability to discriminate, remember and categorize information. Infant macaque monkeys demonstrate a significant novelty preference by 4-6 weeks of postnatal age and the strength of this response increases over the first several months of life (Gunderson and Sackett, 1984, Bachevalier et al., 1993, Zeamer et al., 2010). Longitudinal work with nonhuman primates has provided insights into the neural substrates of this fundamental memory system (Bachevalier, 2015). While visual recognition memory is dependent on hippocampal functioning, there is mounting evidence that novelty preferences during the first few months after birth are also supported by the perirhinal cortex (e.g. Zeamer & Bachevalier, 2013). A diminished capacity to solve recognition memory problems has been observed in studies of prenatal neurotoxicant exposure with both human and macaque monkey infants (for review see Burbacher and Grant, 2012). In human infants, reduced novelty preference scores have been associated with prenatal exposure to environmental contaminants such as phthalates (Ipapo et al., 2017), polychlorinated biphenyls (Boucher et al., 2014), methylmercury (Oken et al., 2005), and chlordecone (Dallaire et al., 2012) as well as recreational drugs such as cocaine (Singer et al., 2005, Chiriboga et al., 2007) and alcohol (Jacobson et al., 1985, 2002). Comparative studies with macaque monkeys have demonstrated the sensitivity of this cognitive test paradigm to naturally-occurring perinatal risk factors (Gunderson et al., 1987, 1989) and developmental exposure to neurotoxicants such as methylmercury (Gunderson et al., 1986, 1988), alcohol (Clarren et al., 1992) and methanol (Burbacher et al., 1999).

One mechanism that has been proposed to account for individual differences in performance on infancy tests of visual recognition memory is speed of processing. The results of this study are consistent with this theoretical approach. When test images were simple representations of objects, DA-exposed infants demonstrated significant novelty responses even when 10 and 30 second delay periods were introduced. This suggests that the time available during the familiarization period on these problems was adequate to fully process and

encode the test stimuli. When more complex, less-discriminable test stimuli were used and processing abilities were challenged during the familiarization period, exposed infants were unable to demonstrate recognition. One underlying reason for this performance deficit may be diminished processing speed. In humans, individual differences on cognitive tests are due to a number of variables, but speed of information processing, along with attention and working memory, is considered central to performance (Kail, 1994, Fagan, 2000, McFarland, 2017). In a longitudinal birth cohort study conducted in New York, preterm human infants exhibited significantly reduced novelty preferences relative to their full-term counterparts (Rose et al., 2001) but deficits in performance were eliminated when familiarization periods were lengthened to provide more time to process test stimuli (Rose et al., 1980). When children in this cohort were reevaluated at 11 years of age, structural equation modeling identified speed of processing and attention as attributes of cognition that were associated with long-term impairment and essential to complex mental operations (Rose et al., 2011). Reductions in the processing speed of visual information have also been identified in infants and children after prenatal exposure to neurotoxicants such as alcohol (Burden et al., 2005) and dichlorodiphenyltrichloroethane (DDT) (Gaspar et al., 2015).

Given that DA has demonstrated latent neurotoxicity in other studies, it is reasonable to hypothesize that the effects documented in this study may not fully represent the long-term consequences of prenatal exposure. After early-life chemical or drug exposure, adverse consequences on health and behavior may be transient or attenuated by neural processes that compensate for structural brain damage (Rice and Barone, 2000). Later in life and particularly during the aging process, chemically-induced changes in the brain may be expressed when neural reserves are less able to mask preexisting injuries. Research with humans and multiple animal species has linked DA exposure with the latent onset of neurological effects. This was first identified in a human case of acute DA poisoning where, after a relatively symptom-free year, the subject developed temporal lobe epilepsy and died (Cendes, 1995). At autopsy, this patient showed serious bilateral hippocampal sclerosis. Adult rats exposed to DA also exhibit changing neurological symptomology and after recovery from acute toxicity, enter a period of clinical silence before the onset of progressive, recurrent seizures (Muha and Ramsdell, 2011, Tiedeken and Ramsdell, 2013). In young California sea lions receiving veterinary treatment for DA toxicosis, complete recovery from acute poisoning can be followed by a progressive and frequently lethal epilepsy-like syndrome that is not clinically manifest until months after cessation of exposure (Ramsdell and Gulland, 2014). The authors suggest that epileptic

disease is a delayed manifestation of DA poisoning in affected animals. Long-term follow-up of the animals in the present study will provide a unique opportunity to study latent neurotoxicity following prenatal exposure in a highly translational primate model. This is particularly important because general observations of the infants in this cohort provided anecdotal and video evidence of intentional tremors in several DA-exposed infants.

The results described in this manuscript are the first to characterize the effects of prenatal DA exposure on neonatal behavior and visual recognition memory in infant macaque monkeys. While the long-term consequences of early changes in neurodevelopment are not known, it has been cogently argued that in humans, small and even subtle impacts of developmental neurotoxicants, when amortized across populations and generations, can have influential and significant impacts on society (Bellinger, 2012). The animals in this birth cohort are now being evaluated on a suite of developmental exams to study the effects of DA on learning and memory, social behavior, temperament and EEG activity during late infancy and pre-adolescence. Results from studies such as these will be critical to defining the neurodevelopmental risks posed by fetal DA exposure at levels close to the human TDI and will identify key domains of behavior and health that are vulnerable to long-term disruption.

References

Adams, A.L., Doucette, T.A., Ryan, C.L., 2008. Altered pre-pulse inhibition in adult rats treated neonatally with domoic acid. Amino Acids 35, 157–160. https://doi.org/10.1007/s00726-007-0603-3

Bachevalier, J., Brickson, M., Hagger, C., 1993. Limbic-dependent recognition memory in monkeys develops early in infancy. Neuroreport., 4(1):77-80.

Bachevalier, J., 2015. Commentary: The development of hippocampal-dependent memory functions: Theoretical comments on Jabès and Nelson review (2015). Int. J. Behav. Dev. https://doi.org/10.1177/0165025415573644

Bellinger, D.C., 2012. A strategy for comparing the contributions of environmental chemicals and other risk factors to neurodevelopment of children. Environ. Health Perspect. 120, 501–507. https://doi.org/10.1289/ehp.1104170

Boucher, O., Muckle, G., Jacobson, J.L., Carter, R.C., Kaplan-Estrin, M., Ayotte, P., Dewailly, É., Jacobson, S.W., 2014. Domain-specific effects of prenatal exposure to PCBs, mercury, and lead on infant cognition: Results from the environmental contaminants and child development study in nunavik. Environ. Health Perspect. 122, 310–316. https://doi.org/10.1289/ehp.1206323

Brazelton, T.B., Nugent, J.K., 2011. Neonatal behavioral assessment scale, Clinics in developmental medicine; no. 88. MacKeith Press, 4th ed.

Burbacher, T.M., Mohamed, M.K., Mottett, N.K., 1988. Methylmercury effects on reproduction and offspring size at birth. Repro.Tox., 1(4), 267–78.

Burbacher, T., Shen, D., Grant, K., Sheppard, L., Damian, D., Ellis, S., Liberato, N., 1999. Reproductive and offspring developmental effects following maternal inhalation exposure to methanol in nonhuman primates. Research Report (Health Effects Institute), (89), i–ii, 1-117; discussion 119-133.

Burbacher, T.M., Grant, K.S., Shen, D.D., Sheppard, L., Damian, D., Ellis, S., Liberato, N., 2004. Chronic maternal methanol inhalation in nonhuman primates (Macaca fascicularis): Reproductive performance and birth outcome. Neurotoxicol. Teratol. 26, 639–650. https://doi.org/10.1016/j.ntt.2004.06.001

Burbacher, T.M., Grant, K.S., 2012. Measuring infant memory: Utility of the visual paired-comparison test paradigm for studies in developmental neurotoxicology. Neurotoxicol. Teratol. https://doi.org/10.1016/j.ntt.2012.06.003

Burbacher, T., Grant, K., Petroff, R., Crouthamel, B., Stanley, C., McKain, N., Shum, S., Jing, J., Isoherranen, N. preprint. Effects of Chronic, Oral Domoic Acid Exposure on Maternal Reproduction and Infant Birth Characteristics in a Preclinical Primate Model. Neurotoxicology and Teratology. Biorxiv. https://doi.org/10.1101/440354

Burden, M.J., Jacobson, S.W., Jacobson, J.L., 2005. Relation of prenatal alcohol exposure to cognitive processing speed and efficiency in childhood. Alcohol. Clin. Exp. Res. 29, 1473–1483. https://doi.org/10.1097/01.alc.0000175036.34076.a0

Burt, M.A., Ryan, C.L., Doucette, T.A., 2008. Altered responses to novelty and drug reinforcement in adult rats treated neonatally with domoic acid. Physiol. Behav. 93, 327–336. https://doi.org/10.1016/j.physbeh.2007.09.003

Cendes, F., Andermann, F., Carpenter, S., Zatorre, R.J., Cashman, N., 1995. Temporal lobe epilepsy caused by domoic acid intoxication: Evidence for glutamate receptor–mediated excitotoxicity in humans. Ann. Neurol. 37, 123–126. https://doi.org/10.1002/ana.410370125

Chiriboga, C.A., Kuhn, L., Wasserman, G.A., 2007. Prenatal cocaine exposures and dose-related cocaine effects on infant tone and behavior. Neurotoxicol. Teratol. 29, 323–30. https://doi.org/10.1016/j.ntt.2006.12.002

Clarren, S.K., Astley, S.J., Gunderson, V.M., Spellman, D., 1992. Cognitive and behavioral deficits in nonhuman primates associated with very early embryonic binge exposures to ethanol. J. Pediatr. 121, 789–796. https://doi.org/10.1016/S0022-3476(05)81917-1

Costa, L.G., Giordano, G., Faustman, E.M., 2010. Domoic acid as a developmental neurotoxin. Neurotoxicology 31, 409–423. https://doi.org/10.1016/j.neuro.2010.05.003

Dakshinamurti, K., Sharma, S.K., Sundaram, M., Watanabe, T., 1993. Hippocampal changes in developing postnatal mice following intrauterine exposure to domoic acid. J. Neurosci. 13, 4486–4495.

Dallaire, R., Muckle, G., Rouget, F., Kadhel, P., Bataille, H., Guldner, L., Seurin, S., Chajès, V., Monfort, C., Boucher, O., Pierre Thomé, J., Jacobson, S.W., Multigner, L., Cordier, S., 2012. Cognitive, visual, and motor development of 7-month-old Guadeloupean infants exposed to chlordecone. Environ. Res. 118, 79–85. https://doi.org/10.1016/J.ENVRES.2012.07.006

Doucette, T.A., Ryan, C.L., Tasker, R.A., 2007. Gender-based changes in cognition and emotionality in a new rat model of epilepsy. Amino Acids 32, 317–322. https://doi.org/10.1007/s00726-006-0418-7

Doucette, T.A., Tasker, R.A., 2016. Perinatal domoic acid as a neuroteratogen, in: Current Topics in Behavioral Neurosciences. Springer, Cham, pp. 87–110. https://doi.org/10.1007/7854_2015_417

Fagan III, J.F., 2000. A Theory of Intelligence: Implications for Society. Psychol. Public Policy, Law 6, 168–179. https://doi.org/10.1037/1076-8971.6.1.168

Fantz, R.L., 1956. A Method for Studying Early Visual Development. Percept. Mot. Skills 6, 13–15. https://doi.org/10.2466/PMS.6..13-15

Fantz, R.L., 1964. Visual experience in infants: Decreased attention to familiar patterns relative to novel ones. Science (80). 146, 668–670. https://doi.org/10.1126/science.146.3644.668

Gaspar, F.W., Harley, K.G., Kogut, K., Chevrier, J., Mora, A.M., Sjödin, A., Eskenazi, B., 2015. Prenatal DDT and DDE exposure and child IQ in the CHAMACOS cohort. Environ. Int. 85, 206–212. https://doi.org/10.1016/j.envint.2015.09.004

Gill, D.A., Perry, M.A., McGuire, E.P., Pérez-Gómez, A., Tasker, R.A., 2012. Low-dose neonatal domoic acid causes persistent changes in behavioural and molecular indicators of stress response in rats. Behav. Brain Res. 230, 409–417. https://doi.org/10.1016/j.bbr.2012.02.036

Graham, F.K., 1975. The more or less startling effects of weak prestimulation. Psychophysiology 12, 238–248. https://doi.org/10.1111/j.1469-8986.1975.tb01284.x

Grant, K.S., Burbacher, T.M., Faustman, E.M., Grattan, L.M., 2010. Domoic acid: Neurobehavioral consequences of exposure to a prevalent marine biotoxin. Neurotoxicol. Teratol. 32, 132–141. https://doi.org/10.1016/j.ntt.2009.09.005

Grant, K.S., Rice, D.C., 2008. Exposure to Environmental Chemicals and Developmental Risk, in: Burbacher, T.M., Sackett, G.P., Grant, K.S. (Eds.), Primate Models of Children's Health and Developmental Disabilities. Elsevier, pp. 377–419.

Grattan, L.M., Boushey, C.J., Tracy, K., Trainer, V.L., Roberts, S.M., Schluterman, N., Morris, J.G.J., 2016b. The association between razor clam consumption and memory in the CoASTAL cohort. Harmful Algae 57, 20–25. https://doi.org/10.1016/j.hal.2016.03.011

Grattan, L.M., Boushey, C.J., Liang, Y., Lefebvre, K.A., Castellon, L.J., Roberts, K.A., Toben, A.C., Morris, J.G.J., 2018. Repeated Dietary Exposure to Low Levels of Domoic Acid and Problems with Everyday Memory: Research to Public Health Outreach. Toxins (Basel). 10, 103. https://doi.org/10.3390/toxins10030103

Grattan, L.M., Holobaugh, S., Morris, J.G.J., 2016a. Harmful algal blooms and public health. Harmful Algae. https://doi.org/10.1016/j.hal.2016.05.003

Gunderson, V.M., Grant, K.S., Burbacher, T.M., Fagan, J.F., Mottet, N.K., 1986. The effect of low-level prenatal methylmercury exposure on visual recognition memory in infant crab-eating macaques. Child Dev. 57, 1076–1083. https://doi.org/10.1111/j.1467-8624.1986.tb00268.x

Gunderson, V.M., Grant-Webster, K.S., Sackett, G.P., 1989. Deficits in visual recognition in low birth weight infant pigtailed monkeys (Macaca nemestrina). Child Dev 60, 119–127.

Gunderson, V.M., Grant-Webster, K.S., 1987. Visual Recognition Memory in High- and Low-Risk Infant Pigtailed Macaques (Macaca Nemestrina). Dev. Psychol. 23, 1983–1987.

Gunderson, V.M., Grant-Webster, K.S., Burbacher, T.M., Mottet, N.K., 1988. Visual recognition memory deficits in methylmercury-exposed Macaca fascicularis infants. Neurotoxicol. Teratol. 10, 373–379. https://doi.org/10.1016/0892-0362(88)90041-4

Gunderson, V.M., Sackett, G.P., 1984. Development of pattern recognition in infant pigtailed macaques (Macaca nemestrina). Dev. Psychol. 20, 418–426. https://doi.org/10.1037/0012-1649.20.3.418

Haß, K., Bak, N., Szycik, G.R., Glenthøj, B.Y., Oranje, B., 2017. Deficient prepulse inhibition of the startle reflex in schizophrenia using a cross-modal paradigm. Biol. Psychol. 128, 112–116. https://doi.org/10.1016/j.biopsycho.2017.07.016

Ipapo, K.N., Factor-Litvak, P., Whyatt, R.M., Calafat, A.M., Diaz, D., Perera, F., Rauh, V., Herbstman, J.B., 2017. Maternal prenatal urinary phthalate metabolite concentrations and visual recognition memory among infants at 27 weeks. Environ. Res. 155, 7–14. https://doi.org/10.1016/j.envres.2017.01.019

Jacobson, J.L., Fein, G.G., Jacobson, S.W., Schwartz, P.M., 1984. Factors and clusters for the Brazelton Scale: An investigation of the dimensions of neonatal behavior, Dev. Psychol. 20, 339-353.

Jacobson, S.W., Fein, G.G., Jacobson, J.L., Schwartz, P.M., Dowler, J.K., 1985. The effect of intrauterine PCB exposure on visual recognition memory. Child Dev. 56, 853–860. https://doi.org/10.1111/j.1467-8624.1985.tb00158.x

Jacobson, S.W., 1998. Specificity of neurobehavioral outcomes associated with prenatal alcohol exposure. Alcohol Clin. Exp. Res., 22(2), 313-20.

Jeffery, B., Barlow, T., Moizer, K., Paul, S., Boyle, C., 2004. Amnesic shellfish poison. Food and Chem. Toxicol., 42(4), 545–557. http://doi.org/10.1016/j.fct.2003.11.010

Jing J, Petroff R, Shum S, Crouthamel B, Topletz AR, Grant KS, Burbacher TM, Isoherranen N., 2018. Toxicokinetics and Physiologically Based Pharmacokinetic Modeling of the Shellfish Toxin Domoic Acid in Nonhuman Primates. Drug Metab Dispos. Feb;46(2):155-165.

Kail, R., Salthouse, T.A., 1994. Processing speed as a mental capacity. Acta Psychol. (Amst). 86, 199–225.

Lefebvre, K.A., Robertson, A., 2010. Domoic acid and human exposure risks: A review. Toxicon 56, 218–230. https://doi.org/10.1016/j.toxicon.2009.05.034

Levin, E.D., Pizarro, K., Pang, W.G., Harrison, J., Ramsdell, J.S., 2005. Persisting behavioral consequences of prenatal domoic acid exposure in rats. Neurotoxicol. Teratol. 27, 719–725. https://doi.org/10.1016/j.ntt.2005.06.017

Mariën, K., 1996. Establishing tolerable Dungeness crab (Cancer magister) and razor clam (Siliqua patula) domoic acid contaminant levels. Environ. Health Perspect. 104, 1230–6. https://doi.org/10.1289/ehp.961041230

Marriott, A.L., Ryan, C.L., Doucette, T.A., 2012. Neonatal domoic acid treatment produces alterations to prepulse inhibition and latent inhibition in adult rats. Pharmacol. Biochem. Behav. 103, 338–344. https://doi.org/10.1016/j.pbb.2012.08.022

Maucher Fuquay, J., Muha, N., Wang, Z., Ramsdell, J.S., 2012. Elimination Kinetics of Domoic Acid from the Brain and Cerebrospinal Fluid of the Pregnant Rat. Chem. Res. Toxicol. 25, 2805–2809. https://doi.org/10.1021/tx300434s

McFarland, D.J., 2017. How neuroscience can inform the study of individual differences in cognitive abilities. Rev. Neurosci. 28, 343–362. https://doi.org/10.1515/revneuro-2016-0073

McKibben, S.M., Peterson, W., Wood, A.M., Trainer, V.L., Hunter, M., White, A.E., 2017. Climatic regulation of the neurotoxin domoic acid. PNAS 114, 239–244. https://doi.org/10.1073/pnas.1606798114

Muha, N., Ramsdell, J.S., 2011. Domoic acid induced seizures progress to a chronic state of epilepsy in rats. Toxicon 57, 168–171. https://doi.org/10.1016/j.toxicon.2010.07.018

Oken, E., Wright, R.O., Kleinman, K.P., Bellinger, D., Amarasiriwardena, C.J., Hu, H., Rich-Edwards, J.W., Gillman, M.W., 2005. Maternal fish consumption, hair mercury, and infant cognition in a U.S. cohort. Environ. Health Perspect. 113, 1376–1380. https://doi.org/10.1289/ehp.8041

Perl, T.M., Bedard, L., Kosatsky, T., Hockin, J.C., Todd, E.C., McNutt, L.A., Remis, R.S., 1990a. Amnesic shellfish poisoning: a new clinical syndrome due to domoic acid. Canada Dis. Wkly. Rep. 16 Suppl 1, 7–8.

Perl, T.M., Bedard, L., Kosatsky, T., Hockin, J.C., Todd, E.C.D., Remis, R.S., Bédard, L., 1990b. An Outbreak of Toxic Encephalopathy Caused by Eating Mussels Contaminated with Domoic Acid. N. Engl. J. Med. 322, 1775–1780. https://doi.org/10.1056/NEJM199006213222504

Ramsdell, J.S., Gulland, F.M., 2014. Domoic acid epileptic disease. Mar. Drugs 12, 1185–1207. https://doi.org/10.3390/md12031185

Rice, D., Barone, S., 2000. Critical periods of vulnerability for the developing nervous system: Evidence from humans and animal models. Environ. Health Perspect. 108, 511–533. https://doi.org/10.1289/ehp.00108s3511

Rose, S.A., 1980. Enhancing visual recognition memory in preterm infants. Dev. Psychol. 16:85-92.

Rose, S.A., Feldman, J.F., Jankowski, J.J., 2001. Attention and recognition memory in the 1st year of life: A longitudinal study of preterm and full-term infants. Dev. Psychol. 37, 135–151. https://doi.org/10.1037//0012-1649.37.1.135

Rose, S.A., Feldman, J.F., Jankowski, J.J., Van Rossem, R., 2011. Basic information processing abilities at 11years account for deficits in IQ associated with preterm birth. Intelligence 39, 198–209. https://doi.org/10.1016/j.intell.2011.03.003

Ryan, C.L., Robbins, M.A, Smith, M., Gallant, I.C., Adams-Marriott, A.L., Doucette, T.A., 2011. Altered social interaction in adult rats following neonatal treatment with domoic acid. Physiol. Behav. 102, 291–295. https://doi.org/10.1016/j.physbeh.2010.11.020

Sackett, G.P., Ruppentahal, G.C., Elias, K. (Eds.), 2006. Nursery Rearing of Nonhuman Primates in the 21st Century. Springer US, Boston, MA. https://doi.org/10.1007/978-0-387-25640-5

Sagiv, S.K., Nugent, J.K., Brazelton, T.B., Choi, A.L., Tolbert, P.E., Altshul, L.M., Korrick, S.A., 2008. Prenatal organochlorine exposure and measures of behavior in infancy using the Neonatal Behavioral Assessment Scale (NBAS). Environ. Health Perspect. 116(5):666-73. doi: 10.1289/ehp.10553

Shiotani, M., Cole, T.B., Hong, S., Park, J.J.Y., Griffith, W.C., Burbacher, T.M., Workman, T., Costa, L.G., Faustman, E.M., 2017. Neurobehavioral assessment of mice following repeated oral exposures to domoic acid during prenatal development. Neurotoxicol. Teratol. 64, 8–19. https://doi.org/10.1016/J.NTT.2017.09.002

Shum, S., Kirkwood, J.S., Jing, J., Petroff, R., Crouthamel, B., Grant, K.S., Burbacher, T.M., Nelson, W.L., Isoherranen, N., 2018. Validated HPLC-MS/MS Method To Quantify Low Levels of Domoic Acid in Plasma and Urine after Subacute Exposure. ACS Omega 3, 12079–12088. https://doi.org/10.1021/acsomega.8b02115

Singer, L.T., Eisengart, L.J., Minnes, S., Noland, J., Jey, A., Lane, C., Min, M.O., 2005. Prenatal cocaine exposure and infant cognition. Infant Behav. Dev. 28, 431–444. https://doi.org/10.1016/j.infbeh.2005.03.002

Takahashi, H., Hashimoto, R., Iwase, M., Ishii, R., Kamio, Y., Takeda, M., 2011. Prepulse inhibition of startle response: Recent advances in human studies of psychiatric disease. Clin. Psychopharmacol. Neurosci. https://doi.org/10.9758/cpn.2011.9.3.102

Tanemura, K., Igarashi, K., Matsugami, T.-R., Aisaki, K., Kitajima, S., Kanno, J., 2009. Intrauterine environment-genome interaction and Children's development (2): Brain structure impairment and behavioral disturbance induced in male mice offspring by a single intraperitoneal administration of domoic acid (DA) to their dams. J. Toxicol. Sci. 34, SP279-SP286. https://doi.org/10.2131/jts.34.SP279

Teitelbaum, J., Zatorre, R.J., Carpenter, S., Gendron, D., Cashman, N.R., 1990a. Neurological Sequelae of Domoic Acid Intoxication. Symp. Domoic Acid Toxic. 16, 9–12. https://doi.org/10.2174/13816128236661701241

Teitelbaum, J., Zatorre, R.J., Carpenter, S., Gendron, D., Evans, A., Gjedde, A., Cashman, N., 1990b. Neurologic Sequelae of Domoic Acid Intoxication due to the Ingestion of Contaminated Mussels. N. Engl. J. Med. 322, 1781–1787.

Teller, D., 1997. First glances: the vision of infants, the Friedenwald lecture. Invest. Ophthalmol. Vis. Sci. 38, 2183–2203.

Teller, D.Y., Boothe, R., 1979. Development of vision in infant primates. Trans Ophthalmol Soc U K 99, 333–337.

Tiedeken, J.A., Ramsdell, J.S., 2013. Persistent neurological damage associated with spontaneous recurrent seizures and atypical aggressive behavior of domoic acid epileptic disease. Toxicol. Sci. 133, 133–143. https://doi.org/10.1093/toxsci/kft037

Wekell, J.C., Jurst, J., Lefebvre, K. A., 2004. The origin of the regulatory limits for PSP and ASP toxins in shellfish. J. Shellfish Res. 23, 927–930.

Woods Hole Oceanographic Institution, n.d. The Harmful Algae Page: Distribution of HABs in the US. https://www.whoi.edu/science/B/redtide/HABdistribution/HABmap.html Accessed October 27, 2018.

Xi, D., Peng, Y.G., Ramsdell, J.S., 1997. Domoic acid is a potent neurotoxin to neonatal rats. Nat. Toxins 5, 74–79. https://doi.org/10.1002/(SICI)(1997)5:2<74::AID-NT4>3.0.CO;2-I

Zeamer, A., Heuer, E., Bachevalier, J., 2010. Developmental trajectory of object recognition memory in infant rhesus macaques with and without neonatal hippocampal lesions. J Neurosci. 30(27), 9157-65. doi: 10.1523/JNEUROSCI.0022-10.2010

Zeamer, A., Bachevalier, J., 2013. Long-term effects of neonatal hippocampal lesions on novelty preference in monkeys. Hippocampus, 23(9),745-50. doi: 10.1002/hipo.22139.

Zeamer, A., Richardson, R.L., Weiss, A.R., Bachevalier, J., 2015. The development of object recognition memory in rhesus macaques with neonatal lesions of the perirhinal cortex. Dev. Cogn. Neurosci. 11, 31–41. https://doi.org/10.1016/j.dcn.2014.07.002

Zhu, Z., Qu, P., Fu, F., Tennenbaum, N., Tatters, A.O., Hutchins, D.A., 2017. Understanding the blob bloom: Warming increases toxicity and abundance of the harmful bloom diatom Pseudo-nitzschia in California coastal waters. Harmful Algae 67, 36–43. https://doi.org/10.1016/j.hal.2017.06.004

Zuloaga, D.G.G., Lahvis, G.P.P., Mills, B., Pearce, H.L.L., Turner, J., Raber, J., 2016. Fetal domoic acid exposure affects lateral amygdala neurons, diminishes social investigation and alters sensory-motor gating. NeuroToxicology 53, 132–140. https://doi.org/10.1016/j.neuro.2016.01.007