A transgenic zebrafish line for *in vivo* visualisation of neutrophil myeloperoxidase

A transgenic zebrafish line expressing fluorescently-tagged human myeloperoxidase in neutrophils presents a useful tool for investigating neutrophil granule dynamics *in vivo*, without disrupting neutrophil migration to inflammatory stimuli.

Kyle D. Buchan[1], Nienke W.M. de Jong[2], Michiel van Gent[2], Julia Kolata[2], Simon J. Foster[3], Jos A.G. van Strijp[2], Stephen A. Renshaw[1]

1. The Bateson Centre and Department of Infection, Immunity and Cardiovascular Disease, University of Sheffield, Western Bank, Sheffield, S10 2TN, UK.

2. Department of Medical Microbiology, University Medical Center Utrecht, Utrecht University, Utrecht, The Netherlands.

3. Department of Molecular Biology and Biotechnology, University of Sheffield, Western Bank, Sheffield, S10 2TN, UK.

Abstract

The neutrophil enzyme myeloperoxidase (MPO) is a major enzyme utilised by neutrophils to generate reactive oxygen species (ROS), amplifying their capacity for destroying pathogens and regulating inflammation. Despite its roles in the innate immunity, the importance of MPO in preventing infection is unclear, as individuals with MPO deficiency are asymptomatic with the exception of an increased risk of candidiasis. Dysregulation of MPO activity is also linked with inflammatory conditions such as atherosclerosis, emphasising a need to understand the roles of the enzyme in greater detail. Consequently, new tools for investigating granular dynamics *in vivo* can provide useful insights into how MPO localises within neutrophils, aiding understanding of its role in preventing and exacerbating disease. The zebrafish is a powerful model for investigating the immune system *in vivo*, as it is genetically tractable, and optically transparent.

To visualise MPO activity within zebrafish neutrophils, we created a genetic construct that expresses human MPO as a fusion protein with a C-terminal fluorescent tag, driven by the neutrophil-specific promoter *lyz*. After introducing the construct into the zebrafish genome by Tol2 transgenesis, we established the *Tg(lyz:MPO-mEmerald,cmlc2:eGFP)sh496* line, and confirmed transgene expression in zebrafish neutrophils. We observed localisation of MPO-mEmerald within a subcellular location resembling neutrophil granules, mirroring MPO in human neutrophils. In Spotless (*mpx^NL144*) larvae - which express a non-functional zebrafish myeloperoxidase - the MPO-mEmerald transgene does not disrupt neutrophil migration to sites of infection or inflammation, suggesting that it is a suitable line for the study of neutrophil granule function. We also describe a method for genotyping

44   Spotless (*mpx^{NL144}*) fish, which will be useful in future studies concerning

45   myeloperoxidase in zebrafish.

46   We present a novel transgenic line that can be used to investigate neutrophil

47   granule dynamics *in vivo* without disrupting neutrophil behaviour, with potential

48   applications in studying processing and maturation of MPO during development.

Introduction

The enzyme Myeloperoxidase (MPO) enhances the microbicidal potential of neutrophils by converting hydrogen peroxide (H_2O_2) into the highly toxic antimicrobial compound hypochlorous acid (HOCl) (1), and by forming radicals by oxidating substrates including phenols, nitrate and tyrosine residues (2). MPO is located in the primary granules of neutrophils, which deliver MPO and other bactericidal compounds to invading pathogens by fusing with phagocytic vesicles, accelerating pathogen destruction. MPO is the most abundant protein in the primary granules of human neutrophils (3), and consequently neutrophils are able to produce high levels of HOCl to deliver a highly potent antimicrobial response that is capable of killing a broad variety of major pathogens (4–6). The important pathogen *Staphylococcus aureus* produces a specific virulence factor against MPO, highlighting its importance in resisting staphylococcal infection (7).

Beyond its role in bolstering the antimicrobial defence, MPO is also an important regulator of inflammation. The arrival of neutrophils at the wound site marks the initial steps of the anti-inflammatory response, as MPO is delivered to the wound site to consume H_2O_2 and reduce inflammatory signalling (8,9). There is also a link between aberrant MPO activity and inflammatory conditions: overactivity is associated with cardiovascular disease, multiple sclerosis and glomerulonephritis (10–12), while MPO deficiency has been implicated in pulmonary fibrosis and atherosclerosis (13,14), highlighting its critical role in immune homeostasis. MPO deficiency is a relatively common condition affecting 1 in every 2,000-4,000 people across Europe and North America (15), with no major health risks apart from a susceptibility to *Candida albicans* infections (16). This observation is in stark contrast

to people with chronic granulomatous disease (CGD), who lack a working enzyme

that is essential for the initiation of the oxidative defence known as NADPH-oxidase.

Those with CGD are unable to generate an adequate respiratory burst to destroy

microbes. Unlike MPO deficiency, those with CGD experience frequent life-

threatening infections from a wide range of pathogens (17), and consequently, the

role of MPO is largely unclear when observed in the context of other oxidative

enzymes and compounds. Further studies are required to understand the complex

roles of MPO in the immune system.

The zebrafish is a powerful model for studying physiology and pathology *in*

vivo and has been used to model many important conditions ranging from

neurodegenerative disorders such as Alzheimers (18), to cancers including

melanoma (19) and leukaemia (20). They are optically transparent, making them

amenable to imaging studies and produce high numbers of offspring, which permits

the application of high-throughput approaches. Another major advantage of the

zebrafish is their genetic tractability, facilitating the introduction of large genetic

constructs into the genome, often expressing fluorescent proteins driven by tissue-

specific promoters (21). Several studies have utilised these features to create

transgenic lines labelling macrophages (22) and neutrophils (23) to image the innate

immune response during infection (24) and inflammation (25).

MPO can be measured using a variety of cytochemical and cytometry-based

approaches (26), however there are relatively few tools that allow granular MPO to

be visualised *in vivo* and in real time. Mouse models that permit imaging of

neutrophil granules and MPO do exist (12,27), however murine MPO lacks several

transcription factor binding domains (28), and is expressed at 1/10 the level found in

human neutrophils (29), raising concerns over whether a murine model can fully

represent human MPO.

In this study, we have generated a transgenic zebrafish line expressing

fluorescently-labelled human MPO in zebrafish neutrophils, as a tool towards

investigating the roles of MPO during infection and inflammation. The MPO

transgene (*lyz:*MPO-mEmerald,*cmlc2*:eGFP) was successfully expressed in

zebrafish neutrophils and the resulting protein appears to be trafficked to the primary

granules, recapitulating expression of MPO in human neutrophils. Additionally, we

showed that the MPO-mEmerald enzyme does not disrupt neutrophil recruitment to

sites of injury and infection. In the future, *Tg(lyz:MPO-mEmerald,cmlc2:eGFP)sh496*

zebrafish may prove to be a useful tool for investigating MPO and imaging granular

dynamics *in vivo* and in real-time.

Methods

Zebrafish Husbandry

Zebrafish (*Danio rerio*) were raised and maintained under the Animals (Scientific Procedures) Act 1986 using standard protocols (30). Adult zebrafish were hosted in UK Home Office-approved aquaria at the Bateson Centre, University of Sheffield, and kept under a 14/10 light/dark regime at 28°C.

Cloning

The plasmid used for introducing the transgene into the zebrafish genome (pDestTol2CG2 *lyz*:MPO-mEmerald *cmlc2*:eGFP) was created by Gateway cloning (21). Briefly, the MPO-mEmerald gene was incorporated into an expression vector by first digesting the mEmerald-MPO-N-18 plasmid (Addgene plasmid #54186, Dr. Michael Davidson's lab), and ligating the MPO-mEmerald fusion protein gene into the multiple cloning site vector pME MCS. pME MCS MPO-mEmerald then became a middle-entry vector after BP reaction with the pDONR221 plasmid. The final construct was created by an LR reaction combining a 5' vector containing the *lyz* promoter, the middle entry vector pDONR221 MPO-mEmerald, a 3' vector containing a polyadenylation site, and the destination vector pDestTol2CG2.

Microinjection of *lyz*:MPO-mEmerald Construct DNA

Construct DNA of the donor plasmid pDestTol2CG2 *lyz*:MPO-mEmerald *cmlc2*:eGFP was injected into zebrafish embryos at the one-cell stage with 10ng/µl of Tol2 transposase RNA, according to published protocols (30).

Zebrafish Tailfin Transection

Tg(lyz:MPO-mEmerald,cmlc2:eGFP)sh496 zebrafish at 3 days post-fertilisation were anaesthetised by immersion in E3 supplemented with 4.2% Tricaine and complete transection of the tail was performed with a sterile scalpel. For imaging of larvae, a Nikon® custom-build wide-field microscope was used: Nikon® Ti-E with a CFI Plan Apochromat λ 10X, N.A.0.45 objective lens, a custom built 500 µm Piezo Z-stage (Mad City Labs, Madison, WI, USA) and using Intensilight fluorescent illumination with ET/sputtered series fluorescent filters 49002 and 49008 (Chroma, Bellow Falls, VT, USA) was used. Analysis was performed using Nikon's® NIS Elements software package.

Bacterial Culture Preparation

To prepare a liquid overnight culture of *S. aureus*, 5ml of BHI broth medium (Oxoid) was inoculated with a colony of *S. aureus* strain USA300, and incubated at 37°C overnight with shaking. To prepare *S. aureus* for injection, 50ml of BHI media was inoculated with 500µl of overnight culture, and incubated for roughly 2 hours at 37°C with shaking. The OD_{600} of each culture was measured and 40ml of the remaining culture harvested by centrifugation at 4,500g for 15 minutes at 4°C. The pellet was then resuspended in a volume of PBS appropriate to the bacterial dose required. Once the pellets were resuspended they were then kept on ice until required.

Spotless (*mpx^NL144*) Fish

The Spotless (*mpx^NL144*) mutant line contains a C to T mutation at nucleotide 1126 of the *mpx* RefSeq mRNA sequence (NM_212779), resulting in a premature stop codon (31). A detailed protocol of Sudan Black B staining can be found in

Supplementary File 1. After staining, fixed larvae were imaged using a Nikon®

Extended focus SMZ1500 stereomicroscope with a Prior Z-drive and transmitted and

reflected illumination. Images captured using a DS-Fi1 Nikon® colour camera. Data

analysis was performed using Nikon's® NIS Elements software package.

Microscopy of Neutrophil Granules

Microscopy of neutrophil granules in *Tg(lyz:MPO-mEmerald,cmlc2:eGFP)sh496*

larvae was performed using a Zeiss® Axiovert LSM 880 Airyscan confocal

microscope with 63x Plan Apochromat oil objective (NA 1.4). Cells were illuminated

with a 488 nm argon laser and/or a 561 nm diode laser. Images were processed

using the Zeiss® microscope software and analysed using Zen Black.

Statistics

All data were analysed (Prism 7.0, GraphPad Software, San Diego, CA, USA) using

a two-way ANOVA with Bonferroni post-test to adjust for multiple comparisons.

Results

Creation of a transgenic zebrafish expressing human myeloperoxidase

To create a transgenic zebrafish that expresses a fluorescently-tagged human myeloperoxidase (MPO), we created a genetic construct using Gateway cloning that contains the MPO gene with a C-terminal fusion of the fluorescent protein mEmerald, driven by the neutrophil-specific promoter *lyz* (Figure 1A). After the construct was successfully assembled, it was introduced into the zebrafish genome by Tol2-mediated transgenesis. Successful expression in zebrafish neutrophils was confirmed by inducing transgenesis in the transgenic red neutrophil line *Tg(lyz:nfsB-mCherry)sh260* (Figure 1CD). Injected larvae were then screened at 3 days post fertilisation (dpf) for mEmerald expression and colocalisation with mCherry expression. Figure 1CD shows double-transgenic neutrophils expressing both mEmerald and mCherry in the primary haematopoietic tissue of the zebrafish larvae, the caudal haematopoietic tissue (CHT) (indicated in Figure 1B) (32). This observation confirms that the construct is successfully expressed and suggests that it co-localises with zebrafish neutrophils. We also noted that in double-transgenic neutrophils, there appeared to be a differential subcellular localisation between mEmerald and mCherry signal, with mCherry localised to areas with no visible mEmerald signal (Figure 1D).

MPO-mEmerald is stably expressed in zebrafish neutrophils

To secure adult zebrafish with stable germline integrations of the *lyz:*MPO-mEmerald transgene, larvae that transiently expressed the transgene were identified, raised and outcrossed to determine whether the transgene was inherited by their offspring.

An adult that produced larvae with a cell population labelled with mEmerald was

identified and its progeny raised to produce fish stably expressing the MPO

transgene, with the designation *Tg(lyz:MPO-mEmerald,cmlc2:eGFP)sh496*. To verify

whether the *lyz*:MPO-mEmerald transgene was expressed in neutrophils of stably

transgenic fish, they were crossed to the red neutrophil reporter line *Tg(lyz:nfsB-*

mCherry)sh260, and screened for any co-expression of fluorescent proteins. Both

transgenes were expressed in neutrophils throughout the CHT (Figure 2),

demonstrating that *lyz*:MPO-mEmerald is expressed in zebrafish neutrophils in

stably transgenic larvae.

MPO-mEmerald is trafficked to a subcellular location

As MPO is located in the primary granules of neutrophils prior to delivery to the

phagosome (1), we wished to determine whether the *lyz*:MPO-mEmerald transgene

recapitulates MPO expression in human neutrophils. To investigate the intracellular

localisation of the MPO transgene, *Tg(lyz:MPO-mEmerald,cmlc2:eGFP)sh496* fish

were outcrossed to *Tg(lyz:nfsB-mCherry)sh260* fish, and at 3dpf the double-

transgenic larvae were imaged in high detail using an Airyscan confocal microscope.

Both transgenes are expressed in the same cells, with MPO-mEmerald localising

with a granular subcellular distribution (Figure 3), suggesting that the MPO-

mEmerald fusion protein is trafficked to and packaged within neutrophil granules.

Additionally, as observed in Figure 1D, complete labelling of a neutrophil with

mCherry is visible, while mEmerald is only observed on specific subcellular locations

(Figure 3C), suggesting that the subcellular location of mCherry and mEmerald

differs.

MPO-mEmerald does not disrupt neutrophil migration

In addition to its role in potentiating ROS generation in neutrophils, MPO also influences neutrophil migration to inflammatory stimuli (8). Accordingly, we sought to determine whether *Tg(lyz:MPO-mEmerald,cmlc2:eGFP)sh496* fish exhibit disrupted neutrophil migration to inflammatory and infectious stimuli. To answer these questions, *Tg(lyz:MPO-mEmerald,cmlc2:eGFP)sh496* fish were crossed to *Tg(lyz:nfsB-mCherry)sh260* and at 3dpf their larvae were separated into two groups: non-humanised (*lyz*:nfsB-mCherry only) and humanised (*lyz*:MPO-mEmerald; *lyz*:nfsB-mCherry) to determine how expression of MPO-mEmerald affects these responses.

To assess neutrophil migration to inflammatory stimuli, we used a tailfin-transection model that initiates neutrophil recruitment to a vertically transected tailfin injury in zebrafish larvae (25). Non-humanised and humanised larvae were injured and neutrophil recruitment to the site of injury was imaged at 3 and 6 hours post injury (hpi) (Figure 4A). Both groups exhibited comparable migration of neutrophils to the site of injury at 3 and 6hpi (Figure 4B), suggesting that *lyz*:MPO-mEmerald does not interfere with neutrophil recruitment to sites of injury.

To determine whether the neutrophil response to infection is affected by expression of *lyz*:MPO-mEmerald, we used an otic vesicle infection model to investigate neutrophil recruitment (33,34). After separating larvae into non-humanised and humanised groups, they were injected into the otic vesicle with either a PBS vehicle control or *S. aureus* USA300 at 3dpf. The larvae were then fixed in paraformaldehyde at 4 hours post infection (hpi) and stained with Sudan Black B to detect neutrophils. Injection of *S. aureus* USA300 induces robust recruitment of neutrophils to the otic vesicle, with comparable numbers recruited between non-

humanised and humanised larvae (Figure 4CD). This confirms that expression of the

lyz:MPO-mEmerald transgene does not interfere with neutrophil recruitment to sites

of infection.

Genotypic and functional identification of myeloperoxidase-null Spotless

(*mpx^{NL144}*) larvae

While the *lyz*:MPO-mEmerald transgene is expressed in zebrafish neutrophils in a

manner that recapitulates expression in human neutrophils, it was still unknown

whether MPO-mEmerald is expressed as a functional enzyme. To determine

whether MPO was functional, we sought to create a zebrafish that expresses only

human MPO by removing expression of the endogenous zebrafish myeloperoxidase

(*mpx*) from the *Tg(lyz:MPO-mEmerald,cmlc2:eGFP)sh496* line. This was achieved

using an existing zebrafish line known as Spotless (*mpx^{NL144}*), which possesses a

premature stop codon in the first exon of the *mpx* gene (31). Once acquired, we

aimed to cross the Spotless line to our *Tg(lyz:MPO-mEmerald,cmlc2:eGFP)sh496*

line to create a line that expresses only human MPO. Before we could create this

line, it was necessary to develop a genotyping protocol that could accurately identify

Spotless *mpx^{NL144}* fish.

The *mpx^{NL144}* allele can be identified by PCR amplification of the mutated

gene from genomic DNA, followed by restriction digest of the PCR product. The

restriction enzyme *Bts*CI recognises 5' GG ATG NN 3' sites in DNA, one of which is

present within the mutated *mpx^{NL144}* gene (GGA TGA) but not the wild-type *mpx^{wt}*

gene (GGA CGA), allowing the enzyme to determine the presence of a *mpx^{NL144}*

allele (Figure 5A). The PCR primers were successful in amplifying the region in the

mpx gene from *mpx^{wt}*, *mpx^{NL144}* and *mpx^{wt/NL144}* groups (Figure 5B), and once

digested with *Bts*CI produced different DNA fragments depending on the *mpx^{NL144}*

allele of the fish (Figure 5C), confirming *Bts*CI digestion as an efficient means of

identifying the *mpx^{NL144}* allele. The accuracy of the restriction digest was confirmed

further by sequencing the PCR products, confirming that the fish identified by

restriction digest each have the specific basepair in the expected position (Figure

5D). After adults were genotyped, their larvae were then assessed for functional

myeloperoxidase expression using the myeloperoxidase-dependent stain Sudan

Black B (8), which verified the genotyping results (Figure 5E).

MPO-mEmerald is non-functional in zebrafish neutrophils

To create zebrafish larvae expressing only human MPO, the *Tg(lyz:MPO-*

mEmerald,cmlc2:eGFP)sh496 line was crossed to the Spotless line to create

zebrafish that express the *lyz*:MPO-mEmerald transgene and do not produce

functional endogenous *mpx*. Once created, *Tg(lyz:MPO-*

mEmerald,cmlc2:eGFP)sh496; mpx^{NL144} larvae were stained with the

myeloperoxidase-dependent stain Sudan Black B to determine whether this

conferred staining; these larvae were compared against three sibling control groups

– *mpx^{NL144}*, *mpx$^{wt/NL144}$* and *Tg(lyz:MPO-mEmerald,cmlc2:eGFP)sh496; mpx$^{wt/NL144}$*.

All groups tested containing a functioning *mpx* allele (*mpx$^{wt/NL144}$*, *Tg(lyz:MPO-*

mEmerald,cmlc2:eGFP)sh496; mpx$^{wt/NL144}$) stained with Sudan Black B, indicating

that the stain identifies functional endogenous myeloperoxidase (Figure 6). As

expected, the negative control group did not stain (*mpx^{NL144}*), but surprisingly, neither

did the *Tg(lyz:MPO-mEmerald,cmlc2:eGFP)sh496; mpx^{NL144}* human MPO-only

larvae, indicating that the *lyz*:MPO-mEmerald transgene does not produce a

functional MPO enzyme (Figure 6).

Discussion

In this study, we created a transgenic line expressing a fluorescently-tagged human myeloperoxidase in zebrafish neutrophils. Expression in neutrophils was determined by observing expression of *lyz*:MPO-mEmerald in the fluorescent red neutrophil line, *Tg(lyz:nfsB-mCherry)sh260*, which expresses mCherry in the cytoplasm of zebrafish neutrophils. Both transgenes were expressed within the same cell population, confirming that the *lyz*:MPO-mEmerald transgene labels neutrophils (Figure 1, 2). However, as MPO localises with the primary granules of neutrophils, it was essential that the fluorescent signal observed in the *lyz*:MPO-mEmerald line should differ from the cytoplasmic signal observed in the *Tg(lyz:nfsB-mCherry)sh260* line. This was observed in several instances; in double transgenic neutrophils, distinct areas of the cell remain unlabelled with mEmerald (Figures 1, 3) suggesting that MPO is translated and trafficked to a subcellular location that is distinct from the cytoplasm. This observation is also evident in Airyscan confocal images (Figure 3C), where a large unlabelled area of a double-transgenic neutrophil is visible in the mEmerald channel. This is likely to be a region of the cell that is inaccessible to the primary granules, for example the nucleus, and could be verified using a fluorescent nuclear probe.

In addition to the *lyz*:MPO-mEmerald and *lyz*:nfsB-mCherry signals being distinct, double-transgenic neutrophils contain small intracellular foci of mEmerald signal (Figure 3), suggesting that MPO-mEmerald might be targeted to the primary granules. To confirm that primary granules are labelled, imaging experiments could determine whether these bodies fuse with maturing phagosomes during infection to potentiate the ROS burst. Additionally, primary granules may be distinguished using

peroxidase-sensitive stains such as TSA or O-dianisidine (35–37), and may be used

to confirm the observations shown here.

In addition to the role of MPO in antibacterial defence, it is also an important

enzyme regulating the migration of neutrophils to sites of infection and inflammation,

primarily by mediating H_2O_2 flux (8). Using a combination of approaches for studying

neutrophil migration, we found that expression of the *lyz*:MPO-mEmerald transgene

does not interfere with neutrophil recruitment to sites of infection and inflammation

(Figure 4). Currently, there are no existing tools for visualising neutrophil granules *in*

vivo, and measuring MPO is limited to cytochemical and cytometry-based

approaches (26). Therefore, the *Tg(lyz:MPO-mEmerald,cmlc2:eGFP)sh496*

transgenic line may be used to study granule dynamics *in vivo* without disrupting

neutrophil function.

In order to determine whether MPO-mEmerald is produced as a functional

enzyme, it was necessary to produce zebrafish that do not express endogenous

zebrafish myeloperoxidase. We describe here a genotyping protocol that can be

used to identify Spotless (*mpx^{NL144}*) fish, an *mpx*-null mutant line created in a

separate study (31). This was accomplished by amplifying a region present in the

first exon of the *mpx* gene, followed by restriction digest with *Bts*CI (Figure 5BCD);

this was then functionally verified using the myeloperoxidase-dependent stain Sudan

Black B (8) (Figure 5E). We believe this to be a useful and robust method for

identifying Spotless fish, and may be useful in future studies.

Once a robust method for identifying Spotless fish was established, the

Spotless line was crossed to the *Tg(lyz:MPO-mEmerald,cmlc2:eGFP)sh496* line to

generate a line that expresses human MPO, and does not express zebrafish Mpx

(known as *Tg(lyz:MPO-mEmerald,cmlc2:eGFP)sh496; mpx^{NL144}*). By comparing

staining with Sudan Black B with sibling controls, we found that MPO-mEmerald is

not produced as a functional enzyme, as *lyz*:MPO-mEmerald expression does not

complement staining in the *mpx*[NL144] background. It is unclear why the *Tg(lyz:MPO-*

mEmerald,cmlc2:eGFP)sh496 line does not produce functional MPO, however it is

important to note that MPO is a complex glycoprotein enzyme that undergoes

numerous tightly regulated post-translational modifications. Before mature MPO is

produced, the peptide associates with calreticulin and calnexin in the endoplasmic

reticulum before undergoing a series of proteolytic events leading to insertion of a

haem group and dimerisation of the enzyme, followed by glycosylation and ending

with granule targeting (38). The importance of each step in producing a functional

enzyme is unclear, however studies of myeloperoxidase-deficient individuals suggest

that targeting to the primary granules universally correlates with functional MPO

(15,39–41), and *in vitro* studies show that dimerisation is not required for enzyme

function (7,42,43). Additionally, the discrepancy is unlikely to lie with calnexin and

calreticulin, as they possess roughly 70% amino acid identity with the human

chaperones, and are important during development of the zebrafish lateral line (44).

Differences at any other stages may lead to incomplete MPO maturation and

function   in   the   zebrafish   and   consequently,   the   *Tg(lyz:MPO-*

mEmerald,cmlc2:eGFP)sh496 line may also be useful in investigating how MPO is

processed and targeted to the granules during development.

Conclusion

Conclusion

We have generated a transgenic zebrafish line expressing fluorescently labelled

human MPO within its neutrophils. The enzyme is non-functional and does not

interfere with neutrophil recruitment to sites of infection or inflammation, suggesting

that it may be used to study granule dynamics *in vivo* interfering with neutrophil

behaviour. Additionally, the *Tg(lyz:MPO-mEmerald,cmlc2:eGFP)sh496* line may be

used to investigate processing and targeting of MPO during development, which is

currently uncharacterised *in vivo*. Lastly, we provide a protocol for genotyping

endogenous myeloperoxidase-null Spotless (*mpx^{NL144}*) fish, which will prove useful in

future studies investigating myeloperoxidase in the zebrafish.

References

1. Klebanoff SJ, Kettle a. J, Rosen H, Winterbourn CC, Nauseef WM. Myeloperoxidase: a front-line defender against phagocytosed microorganisms. Vol. 93, Journal of Leukocyte Biology. 2012. p. 185–98.

2. Kettle AJ, Winterbourn CC. Myeloperoxidase: a key regulator of neutrophil oxidant production. Redox Rep. 1997 Feb 13;3(1):3–15.

3. Borregaard N, Cowland JB. Granules of the human neutrophilic polymorphonuclear leukocyte. Blood. 1997 May 15;89(10):3503–21.

4. Decleva E, Menegazzi R, Busetto S, Patriarca P, Dri P. Common methodology is inadequate for studies on the microbicidal activity of neutrophils. J Leukoc Biol. 2006;79(1):87–94.

5. Aratani Y, Kura F, Watanabe H, Akagawa H, Takano Y, Suzuki K, et al. Critical Role of Myeloperoxidase and Nicotinamide Adenine Dinucleotide Phosphate – Oxidase in High-Burden Systemic Infection of Mice with Candida albicans. J Infect Dis. 2002;(April):1833–7.

6. Hirche TO, Gaut JP, Heinecke JW, Belaaouaj A. Myeloperoxidase plays critical roles in killing Klebsiella pneumoniae and inactivating neutrophil elastase: effects on host defense. J Immunol. 2005 Feb 1;174(3):1557–65.

7. de Jong NWM, Ramyar KX, Guerra FE, Nijland R, Fevre C, Voyich JM, et al. Immune evasion by a staphylococcal inhibitor of myeloperoxidase. Proc Natl Acad Sci. 2017 Aug 29;114(35):9439–44.

8. Pase L, Layton JE, Wittmann C, Ellett F, Nowell CJ, Reyes-Aldasoro CC, et al. Neutrophil-delivered myeloperoxidase dampens the hydrogen peroxide burst after tissue wounding in zebrafish. Curr Biol. 2012;22(19):1818–24.

9. Schürmann N, Forrer P, Casse O, Li J, Felmy B, Burgener A, et al. Myeloperoxidase targets oxidative host attacks to Salmonella and prevents collateral tissue damage. Nat Microbiol. 2017 Jan 23;2(January):16268.

10. Kutter D, Devaquet P, Vanderstocken G, Paulus JM, Marchal V, Gothot A. Consequences of Total and Subtotal Myeloperoxidase Deficiency: Risk or Benefit ? Acta Haematol. 2000;104(1):10–5.

11. Yang JJ, Pendergraft WF, Alcorta DA, Nachman PH, Hogan SL, Thomas RP,

et al. Circumvention of normal constraints on granule protein gene expression

in peripheral blood neutrophils and monocytes of patients with antineutrophil

cytoplasmic autoantibody-associated glomerulonephritis. J Am Soc Nephrol.

2004 Aug 1;15(8):2103–14.

12. Chen JW, Breckwoldt MO, Aikawa E, Chiang G, Weissleder R.

Myeloperoxidase-targeted imaging of active inflammatory lesions in murine

experimental autoimmune encephalomyelitis. Brain. 2008;131(4):1123–33.

13. Brennan M-L, Anderson MM, Shih DM, Qu X-D, Wang X, Mehta AC, et al.

Increased atherosclerosis in myeloperoxidase-deficient mice. J Clin Invest.

2001 Feb 15;107(4):419–30.

14. Shvedova AA, Kapralov AA, Feng WH, Kisin ER, Murray AR, Mercer RR, et al.

Impaired clearance and enhanced pulmonary inflammatory/fibrotic response to

carbon nanotubes in myeloperoxidase-deficient mice. PLoS One. 2012 Mar

30;7(3):e30923.

15. DeLeo FR, Goedken M, McCormick SJ, Nauseef WM. A novel form of

hereditary myeloperoxidase deficiency linked to endoplasmic

reticulum/proteasome degradation. J Clin Invest. 1998 Jun 15;101(12):2900–9.

16. Lehrer RI, Cline MJ. Leukocyte myeloperoxidase deficiency and disseminated

candidiasis: the role of myeloperoxidase in resistance to Candida infection. J

Clin Invest. 1969;48(8):1478–88.

17. Assari T. Chronic Granulomatous Disease; fundamental stages in our

understanding of CGD. Med Immunol. 2006;5:4.

18. Paquet D, Bhat R, Sydow A, Mandelkow E-M, Berg S, Hellberg S, et al. A

zebrafish model of tauopathy allows in vivo imaging of neuronal cell death and

drug evaluation. J Clin Invest. 2009 May;119(5):1382–95.

19. Haldi M, Ton C, Seng WL, McGrath P. Human melanoma cells transplanted

into zebrafish proliferate, migrate, produce melanin, form masses and

stimulate angiogenesis in zebrafish. Angiogenesis. 2006;9(3):139–51.

20. Langenau DM, Traver D, Ferrando AA, Kutok JL, Aster JC, Kanki JP, et al.

Myc-induced T cell leukemia in transgenic zebrafish. Science. 2003 Feb

7;299(5608):887–90.

21. Kwan KM, Fujimoto E, Grabher C, Mangum BD, Hardy ME, Campbell DS, et

al. The Tol2kit: A multisite gateway-based construction Kit for Tol2 transposon

transgenesis constructs. Dev Dyn. 2007 Nov;236(11):3088–99.

22.   Ellett F, Pase L, Hayman JW, Andrianopoulos A, Lieschke GJ. mpeg1
promoter transgenes direct macrophage-lineage expression in zebrafish.
Blood. 2011 Jan 27;117(4):e49-56.

23.   Hall C, Flores MV, Storm T, Crosier K, Crosier P. The zebrafish lysozyme C
promoter drives myeloid-specific expression in transgenic fish. BMC Dev Biol.
2007 May 4;7(1):42.

24.   Elks PM, Brizee S, van der Vaart M, Walmsley SR, van Eeden FJ, Renshaw S
a, et al. Hypoxia inducible factor signaling modulates susceptibility to
mycobacterial infection via a nitric oxide dependent mechanism. PLoS Pathog.
2013 Jan;9(12):e1003789.

25.   Renshaw SA, Loynes CA, Trushell DMI, Elworthy S, Ingham PW, Whyte MKB.
A transgenic zebrafish model of neutrophilic inflammation. Blood. 2006 Dec
15;108(13):3976–8.

26.   Pulli B, Ali M, Forghani R, Schob S, Hsieh KLC, Wojtkiewicz G, et al.
Measuring myeloperoxidase activity in biological samples. PLoS One. 2013 Jul
5;8(7):e67976.

27.   Kikushima K, Kita S, Higuchi H. A non-invasive imaging for the in vivo tracking
of high-speed vesicle transport in mouse neutrophils. Sci Rep. 2013 Dec
31;3(1):1913.

28.   Nauseef WM. The proper study of mankind. J Clin Invest. 2001 Feb
3;107(4):401–3.

29.   Rausch PG, Moore TG. Granule enzymes of polymorphonuclear neutrophils: A
phylogenetic comparison. Blood. 1975 Dec;46(6):913–9.

30.   Nüsslein-Volhard C, Dahm R. Zebrafish: A Practical Approach. Practical.
Oxford; New York: Oxford University Press, c2002.; 2002.

31.   Elks PM, van der Vaart M, van Hensbergen V, Schutz E, Redd MJ, Murayama
E, et al. Mycobacteria counteract a TLR-mediated nitrosative defense
mechanism in a zebrafish infection model. PLoS One. 2014 Jun
26;9(6):e100928.

32.   Murayama E, Kissa K, Zapata A, Mordelet E, Briolat V, Lin H-F, et al. Tracing
Hematopoietic Precursor Migration to Successive Hematopoietic Organs

during Zebrafish Development. Immunity. 2006 Dec;25(6):963–75.

33.   Benard EL, van der Sar AM, Ellett F, Lieschke GJ, Spaink HP, Meijer AH.

Infection of zebrafish embryos with intracellular bacterial pathogens. J Vis Exp.

2012 Mar 15;(61):1–8.

34.   Deng Q, Sarris M, Bennin D a, Green JM, Herbomel P, Huttenlocher A.

Localized bacterial infection induces systemic activation of neutrophils through

Cxcr2 signaling in zebrafish. J Leukoc Biol. 2013 May;93(5):761–9.

35.   Gilbert CS, Parmley RT, Rice WG, Kinkade JM. Heterogeneity of peroxidase-

positive granules in normal human and Chediak- Higashi neutrophils. J

Histochem Cytochem. 1993;41(6):837–49.

36.   Lieschke GJ, Oates AC, Crowhurst MO, Ward AC, Layton JE. Morphologic

and functional characterization of granulocytes and macrophages in embryonic

and adult zebrafish. Blood. 2001 Nov 15;98(10):3087–96.

37.   Robertson AL, Holmes GR, Bojarczuk AN, Burgon J, Loynes CA, Chimen M,

et al. A Zebrafish Compound Screen Reveals Modulation of Neutrophil

Reverse Migration as an Anti-Inflammatory Mechanism. Sci Transl Med. 2014

Feb 26;6(225):225ra29-225ra29.

38.   Hansson M, Olsson I, Nauseef WM. Biosynthesis, processing, and sorting of

human myeloperoxidase. Arch Biochem Biophys. 2006 Jan 15;445(2):214–24.

39.   Nauseef WM. Lessons from MPO deficiency about functionally important

structural features. Jpn J Infect Dis. 2004;57(5):4–5.

40.   Nauseef WM, Cogley M, McCormick S. Effect of the R569W missense

mutation on the biosynthesis of myeloperoxidase. J Biol Chem.

1996;271(16):9546–9.

41.   Nauseef WM, McCormick S, Goedken M. Impact of missense mutations on

biosynthesis of myeloperoxidase. Redox Rep. 2000;5(4):197–206.

42.   Moguilevsky N, Garcia-Quintana L, Jacquet A, Tournay C, Fabry L, Piérard L,

et al. Structural and biological properties of human recombinant

myeloperoxidase produced by Chinese hamster ovary cell lines. Eur J

Biochem. 1991 May 8;197(3):605–14.

43.   Andrews PC, Krinsky NI. The reductive cleavage of myeloperoxidase in half,

producing enzymically active hemi-myeloperoxidase. J Biol Chem. 1981 May

10;256(9):4211–8.

44.   Hung I-C, Cherng B-W, Hsu W-M, Lee S-J. Calnexin is required for zebrafish

posterior lateral line development. Int J Dev Biol. 2013;57(5):427–38.

Acknowledgements

K.D.B performed all experiments with assistance from M.v.G, N.W.M.d.J, and J.K. S.A.R, J.A.G.v.S and S.J.F conceived the study and designed experiments. K.D.B and S.A.R wrote the manuscript with significant input from all authors. Thank you to Annemarie Meijer's group for providing the Spotless (mpx^{NL144}) line, to the Bateson Centre aquarium staff for all their help, and to the Wolfson Light Microscopy Facility.

Conflicts of Interest

The authors declare no conflicts of interest.

Funding Information

This work was supported in part by AMR cross-council funding from the MRC to the SHIELD consortium "Optimising Innate Host Defence to Combat Antimicrobial Resistance" MRNO2995X/1. Microscopy studies were supported by a Wellcome Trust grant to the Molecular Biology and Biotechnology/Biomedical Science Light Microscopy Facility (GR077544AIA).

Figure 1. Transient expression of the *lyz*:MPO-mEmerald transgene labels

zebrafish neutrophils

A) Schematic of the *lyz*:MPO-mEmerald *cmlc2*:eGFP construct, which includes the

neutrophil-specific promoter (*lyz*), the MPO gene with a C-terminal fluorescent tag

(MPO-mEmerald) and a green heart marker to aid optimisation of transgenesis

(*cmlc2:*eGFP). **B)** A zebrafish larva at 3 days post fertilisation (dpf), with the caudal

haematopoietic tissue (CHT) indicated by the red box. **C)** The CHT of a double-

transgenic *Transient lyz*:MPO-mEmerald,*cmlc2*:eGFP; *Tg(lyz:nfsB-mCherry)sh260*

larva with a population of neutrophils expressing both mEmerald and mCherry. The

white arrowhead indicates the neutrophil enlarged below. **D)** Enlarged view of a

neutrophil expressing mEmerald and mCherry.

A

B

Figure 2. Transgenic *Tg(lyz:MPO-mEmerald,cmlc2:eGFP)sh496* zebrafish

stably express the transgene in zebrafish neutrophils

A) A brightfield view of a double-transgenic *Tg(lyz:MPO-*

mEmerald,cmlc2:eGFP)sh496; *Tg(lyz:nfsB-mCherry)sh260* zebrafish larva at 3dpf.

The dashed white box indicates the enlarged region shown in B). **B)** mEmerald and

mCherry expression in the CHT of the larva shown in A).

Figure 3. The *lyz:*MPO-mEmerald transgene labels intracellular neutrophil granules

A) A 3dpf zebrafish larva, the field of view shown in B) is outlined by the red box. **B)** An Airyscanner confocal image of neutrophils within the CHT of a double-transgenic *Tg(lyz:MPO-mEmerald,cmlc2:eGFP)sh496; Tg(lyz:nfsB-mCherry)sh260* larva at 3dpf. **C)** An enlarged image of the neutrophil highlighted by the dashed white box in B). Scale bars **B)** 20μm and **C)** 5μm.

Figure 4. Transgene expression does not disrupt neutrophil recruitment to sites of injury or infection

A) Non-humanised (*lyz*:nfsB-mCherry only) and humanised (*lyz*:MPO-mEmerald*; lyz*:nfsB-mCherry) 3dpf larvae with tailfins transected to induce neutrophil recruitment; dashed outline represents the area in which neutrophils were counted. Scale bar = 250μm. **B)** Neutrophils present at the site of injury at 3 and 6 hours post injury (hpi); blue points denote the representative images in A). Error bars shown are mean ± SEM (n=45 over three independent experiments); groups were analysed using an ordinary two-way ANOVA and adjusted using Bonferroni's multiple comparisons test; ns, p>0.9999. **C)** Non-humanised and humanised larvae injected with either a PBS vehicle control or 1,400cfu *S. aureus* USA300 into the otic vesicle at 3dpf, then fixed in paraformaldehyde at 4 hours post infection (hpi) and stained with Sudan Black B to detect neutrophils; dashed white outline indicates the otic vesicle. **D)** Neutrophils present at the otic vesicle at 4hpi. Scale bars = 250μm. Error bars shown are mean ± SEM (n=25 over two independent experiments); groups were analysed using an ordinary two-way ANOVA and adjusted using Bonferroni's multiple comparisons test. ****, p<0.0001; ns, p>0.9999.

 Figure 5. Genotyping and verifying *mpx*-null zebrafish larvae

A) Diagram of a WT (*mpx^{wt}*) and mutated (*mpx^{NL144}*) gene, showing the *Bts*CI restriction site cutting only the mutated *mpx^{NL144}* gene. **B)** PCR amplification of the *mpx* gene from the genomic DNA of *mpx^{wt}*, *mpx^{wt/NL144}* and *mpx^{NL144}* fish – fragment 312bp; control DNA is a positive control from a separate genotyping experiment. Hyperladder 1kb. **C)** Diagnostic digest of the PCR product from *mpx^{wt}*, *mpx^{wt/NL144}* and *mpx^{NL144}* fish. Band sizes: *mpx^{wt}*- 312bp, *mpx^{NL144}*- 230bp, *mpx^{wt/NL144}*- 312bp and 230bp. Hyperladder 100bp plus. **D)** DNA sequencing of the PCR products to confirm the accuracy of the *Bts*CI digest. **E)** *mpx^{wt}*, *mpx^{NL144}* and *mpx^{wt/NL144}* larvae fixed at 4dpf and stained with Sudan Black B. Larvae with at least one functional *mpx* allele stained (57/58 *mpx^{wt}*, 20/20 *mpx^{wt/NL144}*) and larvae that do not produce Mpx did not stain (32/32 *mpx^{NL144}*). Inset shows an enlarged view of the region indicated by the dashed white box. Scale bar = 200µm.

mpx^{wt/NL144}

18/18

Tg(lyz:MPO-mEmerald,cmlc2:eGFP)sh496; mpx^{wt/NL144}

16/16

mpx^{NL144}

20/20

Tg(lyz:MPO-mEmerald,cmlc2:eGFP)sh496; mpx^{NL144}

22/22

 Figure 6. Larvae expressing only human MPO do not stain with

 myeloperoxidase-dependent Sudan Black B

Four groups of larvae were fixed at 4dpf and stained with Sudan Black B: $mpx^{wt/NL144}$,

$Tg(lyz:MPO\text{-}mEmerald,cmlc2:eGFP)sh496$; $mpx^{wt/NL144}$, mpx^{NL144} and $Tg(lyz:MPO\text{-}$

$mEmerald,cmlc2:eGFP)sh496$; mpx^{NL144}. $mpx^{wt/NL144}$ and $Tg(lyz:MPO\text{-}$

$mEmerald,cmlc2:eGFP)sh496$; $mpx^{wt/NL144}$ stained (18/18, 16/16 respectively);

mpx^{NL144} and $Tg(lyz:MPO\text{-}mEmerald,cmlc2:eGFP)sh496$; mpx^{NL144} did not stain

(20/20, 22/22 respectively). Dashed outline indicates the enlarged region shown

adjacent. Scale bar = 200μm.

lyz:MPO-mEmerald,*cmlc2*:eGFP   *lyz:nfsB-mCherry*

C 18:00:43.294	18:00:43.294	18:00:43.294
mEmerald 20μm	mCherry 20μm	Merge 20μm
D		
mEmerald 5μm	mCherry 5μm	Merge 5μm

lyz:MPO-mEmerald,cmlc2:eGFP   *lyz:nfsB-mCherry*

mCherry mEmerald Merge

20 μm 20 μm 20 μm

mCherry mEmerald Merge

5 μm 5 μm 5 μm

T   A   C   A   ✗

T   A   T   A   ✓
STOP

T G G A <u>C</u> G A

T G G A <u>T</u> G A

T G G A <u>c/T</u> G A

57/58

32/32

20/20