An infant mouse model of influenza virus transmission demonstrates the role of virus-specific shedding, humoral immunity, and sialidase expression by colonizing *Streptococcus pneumoniae*.

Mila Brum Ortigoza,[a] Simone Blaser,[b] M. Ammar Zafar,[c,*] Alexandria Hammond,[c] Jeffrey N. Weiser,[c,#]

[a] Department of Medicine, Division of Infectious Diseases, New York University School of Medicine, New York, New York, USA

[b] New York University School of Medicine, New York, New York, USA

[c] Department of Microbiology, New York University School of Medicine, New York, New York, USA

Running Title: Infant mouse model of influenza virus transmission

Address correspondence to Jeffrey N. Weiser, jeffrey.weiser@nyulangone.org

* Present address: Department of Microbiology and Immunology, Wake Forrest School of Medicine, Winston-Salem, NC

Word count: Abstract (368), Text (4692)

ABSTRACT

The pandemic potential of influenza A viruses (IAV) depends on the infectivity of

the host, transmissibility of the virus, and susceptibility of the recipient. While virus traits

supporting IAV transmission have been studied in detail using ferret and guinea pig

models, there is limited understanding of host traits determining transmissibility and

susceptibility because current animal models of transmission are not sufficiently

tractable. Although mice remain the primary model to study IAV immunity and

pathogenesis, the efficiency of IAV transmission in adult mice has been inconsistent.

Here we describe an infant mouse model which support efficient transmission of IAV.

We demonstrate that transmission in this model requires young age, close contact,

shedding of virus particles from the upper respiratory tract (URT) of infected pups, the

use of a transmissible virus strain, and a susceptible recipient. We characterize

shedding as a marker of infectiousness that predicts the efficiency of transmission

among different influenza virus strains. We also demonstrate that transmissibility and

susceptibility to IAV can be inhibited by humoral immunity via maternal-infant transfer of

IAV-specific immunoglobulins, and modifications to the URT milieu, via sialidase activity

of colonizing *Streptococcus pneumoniae* (Spn). Due to its simplicity and efficiency, this

model can be used to dissect the host's contribution to IAV transmission and explore

new methods to limit contagion.

IMPORTANCE

This study provides insight into the role of the virus strain, age, immunity, and

URT flora on IAV shedding and transmission efficiency. Using the infant mouse model,

we found that: (a) differences in viral shedding of various IAV strains is dependent on

specific hemagglutinin (HA) and/or neuraminidase (NA) proteins; (b) host age plays a

key role in the efficiency of IAV transmission; (c) levels of IAV-specific immunoglobulins

are necessary to limit infectiousness, transmission, and susceptibility to IAV; and (d)

expression of sialidases by colonizing Spn antagonize transmission by limiting the

acquisition of IAV in recipient hosts. Our findings highlight the need for strategies that

limit IAV shedding, and the importance of understanding the function of the URT

bacterial composition in IAV transmission. This work reinforces the significance of a

tractable animal model to study both viral and host traits affecting IAV contagion, and its

potential for optimizing vaccines and therapeutics that target disease spread.

INTRODUCTION

Influenza virus infections continue to cause 140,000-700,000 hospitalizations and

12,000-56,000 deaths in the United States annually (1). For the 2017-2018 season

alone, more than 900,000 people were hospitalized and 80,000 people died from

influenza (2). Despite the availability of vaccines which have been efficacious at

preventing hospitalizations, morbidity, and mortality, evidence that the inactivated

influenza virus (IIV) vaccine blocks virus acquisition, shedding, or transmission has

been limited in animal models (3-7). In addition, the low vaccination coverage (in the

population) and low vaccine effectiveness (due to viral antigenic drift) likely contributes

to the limited effects of the IIV vaccine (8, 9). Likewise, available therapeutics, primarily

neuraminidase inhibitors (NI), have been shown to be effective at reducing the duration

of illness if treatment is initiated within 24 hours of symptom onset (10-13). However, NI

treatment of index cases alone shows limited effectiveness reducing viral shedding or

transmission, possibly due to its short therapeutic window (10, 11, 14, 15). These

limitations of our current options to prevent disease spread highlight a critical aspect of

the IAV ecology that needs further study: contagion.

While IAV transmission has been studied in human, ferret, and guinea pig

models, there is a general lack of understanding about the host influence on viral

transmission, because none of these models are easily manipulated. Hence, scientific

progress to date has emphasized viral genetics, viral tropism, and environmental

impacts on transmission (16-19). While these factors contribute to knowledge about IAV

contagion, host characteristics that could affect transmissibility, including the highly

variable composition of the URT flora, remain largely unexplored.

This knowledge gap could be addressed with the use of mice, whose practical

features (small, inexpensive, inbred), expansive reagent repertoire, and availability of

genetically-modified hosts allows for studies of extraordinary intricacy providing a

significant research advantage. Since the 1930s, the mouse model has been essential

in understanding IAV immunity and pathogenesis, and early studies described its

usefulness in evaluating IAV transmission (20, 21). However, the use of mice for

studying IAV transmission has been largely disregarded due to marked differences

among studies and low transmission rates (22-24). Nevertheless, recent reports have

revived the potential of the murine species as an IAV transmission model (23-28).

Therefore, in this study, we sought to reevaluate the mouse as a tool to study the

biology of IAV contagion, particularly the contribution of host factors.

RESULTS

Infant mice support efficient influenza virus transmission.

Given the remarkable capacity of infant mice to support IAV transmission among

littermates (25), we sought to validate and optimize the infant mouse as a potential new

model to study IAV transmission. Restricted URT infection of infant C57BL/6J pups in a

litter (index) was performed with low volume intranasal (IN) inoculum (3µl) using IAV

strain A/X-31 (H3N2) (24, 29). Intralitter transmission was assessed in littermates

(contact/recipient) by measuring virus from retrograde tracheal lavages at 4-5 days

post-infection (p.i.) (Fig.1A). A/X-31 virus was selected to model transmission because

of its intermediate virulence in mice (30) and ability to replicate in the URT to high titers

with natural progression to the lungs, simulating key features of the infectious course in

humans. Furthermore, the 50% mouse infectious dose (MID_{50}) in this model is 4-5

plaque forming units (PFU), suggesting high susceptibility to A/X-31 infection.

Transmission efficiency was observed to be 100% when index and contact pups were

housed together at the time of IAV inoculation (Fig.1B). Transmission declined the

longer the index and recipient pups were housed apart prior to being in direct contact,

and was completely eliminated when housed together after 72 hours of separation

(Fig.1C and Fig. S1). This observation suggested that in this model, transmission from

index to recipient is most effective within the first 72-hour period of contact.

To determine the window of viral acquisition in recipient mice, an 8-day IAV

transmission experiment was performed (Fig. 1D). The observed growth of IAV in the

URT of recipient pups suggested that *de novo* virus acquisition occurred between 2-3

days after contact with the index. Hence, the infectious window for the index pups

corresponded with the timing of IAV acquisition in contact pups.

Given that pups gain weight as they grow, morbidity in this model was assessed

by observing a decrease in weight gain during the infectious period. Mild morbidity of

pups was observed in both the index and contact groups, with complete recovery from

IAV infection by 10 days p.i. (Fig. 1E).

Direct contact between pups is required for influenza virus transmission.

Because infant mice need their mother for survival during the first 21 days of life,

they cannot be separately housed, therefore this model cannot differentiate between

airborne versus droplet routes of transmission. To distinguish between direct and

indirect contact routes of transmission, the mother and housing contents were evaluated

as potential fomites. This was done by daily switching the mothers or the cages with

bedding between infected and uninfected litters, respectively (Fig. S2A-B). Inefficient or

no transmission was observed, suggesting that direct (close) contact between pups is

the main mode of transmission. Occasionally during a transmission experiment, the

mother in the cage became infected with IAV from close contact with her infected pups

(Fig. S2C). Although the acquisition of IAV in the mother was a rare event, we did not

observe a decline in transmission in contact pups when the mother did not become

infected, despite the mother being capable of transmitting IAV to her pups if she were to

be inoculated with IAV as the index case (Fig. 2D).

Role of shedding of influenza virus from the upper respiratory tract.

To determine the correlates of transmission, an assay was developed to quantify

infectious virus expelled from the nasal secretions of pups. This assay allowed us to

follow the journey of particle exit from index pups to acquisition by contact pups over the

course of the transmission period. Index pups in a litter were infected with A/X-31 and

cohoused with uninfected littermates for 10 days. The nares of each mouse was gently

dipped in viral media daily, and virus titers were assessed for each sample (Fig. 3A).

We observed that index pups, like in humans (31), began shedding virus from day 1 p.i.,

whereas recipient pups, who acquired IAV infection between day 2-3 (Fig. 1D), began

shedding virus from day 4 post contact (Fig. 3B). This pattern of virus transit suggested

that the timing of peak shedding from the index (days 1-3) corresponded with the timing

of transmission to recipient pups (days 2-4) (Fig. S3) further confirming that a key

determinant of IAV transmission in this model is shedding of virus from the secretions of

index pups. Notably, detectable shedding in the contacts lagged transmission (higher

transmission rate compared to number of contacts shedding virus), because of the

period of viral replication required prior to the detection of shed virus (Fig. S3).

Transmission efficiency of influenza viruses in mice is virus and age-dependent.

Virus strain has been shown to be important in the efficiency of transmission in

adult mice (20, 23, 26, 32). We thus tested the capacity of infant mice to support

transmission of other IAV subtypes and an influenza B virus (IBV) (Table 1).

Transmission among pups was greater for influenza A/H3N2 viruses and IBV, but lower

for A/H1N1 viruses. Notably, A/X-31 was more efficiently transmitted compared to its

parent A/PR/8/1934 virus, suggesting that either the HA and/or NA proteins are

responsible for efficient shedding and transmission of IAV in infant mice.

Surprisingly, the mean viral URT titers in index pups did not correlate with IAV

transmission (r=0.315), indicating that virus replication in the URT alone was insufficient

to mediate effective transmission. To determine the cause of the differences in

transmission efficiencies observed among virus strains, the shedding for each virus was

analyzed. We observed that virus shed from index pups correlated with IAV

transmission (r=0.8663), further supporting virus shedding as the main determinant of

IAV transmission efficiency in infant mice (Fig. S4A-B).

Given the effectiveness of the infant mouse in supporting transmission of IAV, we

evaluated the disparities of transmission efficiency previously reported in adult mice (20-

23, 26). Mice infected with A/X-31 at different ages were housed with uninfected age-

matched contacts, and transmission efficiency was assessed at 5 days p.i. (Table 2).

We observed that 100% transmission was sustained in mice up to 7-days of age.

Weaned and active adult mice (>28 days of age) failed to sustain efficient transmission

altogether. Furthermore, mouse age correlated with transmission rate among contact

mice (r=-0.8346) (Fig. S4C), confirming that in the murine model, the requirement for

young age is necessary to support efficient IAV transmission. Although the transmission

experiments in this study were done with an IAV inoculum of 250 PFU, and increasing

the inoculum size to 10^3-10^5 PFU correlated with increasing IAV titers in the URT tract

of index mice (r=0.9264), inoculum size was not associated with more efficient

transmission among adult mice (r=-0.2582) (Fig. S4D).

Humoral immunity from prior influenza virus infection limits shedding and transmission.

To further validate the relationship of viral shedding and transmission, we evaluated the role of IAV-specific immunity in this model. Because pups are infected at a young age and lack a fully functional adaptive immune response, it was necessary to provide IAV immunity via the mother (from prior IAV infection), who would then transfer immunoglobulins to her pups either pre-natally via trans-placental passage or post-natally via breastfeeding. Pups from immune mothers who were subjected to an intralitter IAV transmission experiment shed significantly fewer virus over the first 5 days of infection compared to pups from non-immune mothers (Fig. 3A left). Reduced shedding was associated with decreased transmission (20%) among immune litters (p<0.0001) (Fig. 3A below graph). To determine if the passage of anti-IAV immunoglobulins occurred pre-natally or post-natally, mothers were switched shortly after delivery such that an immune mother raised pups from a non-immune mother or vice-versa. These cross-foster experiments demonstrated that maternal passage of immunoglobulins either pre-natally or post-natally decreased IAV shedding amongst all pups, and that transmission to contact pups was more efficiently blocked when maternal antibodies were passed post-natally via breastfeeding (p<0.01) (Fig. 3A right). IAV-specific serum IgG was detected in immune mothers and pups born or cared by immune mothers, with the transfer of IgG via breastfeeding yielding higher titer of antibodies in these pups (Fig. 3B). IAV-specific serum IgA was detected in previously-infected mothers but unlike IgG was not passed to their pups in significant amounts (Fig. S4).

208 ***Streptococcus pneumoniae* colonization of the upper respiratory tract decreases**

209 **influenza virus acquisition via bacterial sialidase activity.**

There is increasing evidence of the important role of the host's gut microbiome

on IAV-specific immunity in the respiratory tract (33, 34). Yet, there is only one study

evaluating the role of the URT microbiota in IAV infection (35) and no studies on its

effect on transmission. This is surprising given that the nasopharynx, a non-sterile

environment extensively colonized by a diverse bacterial flora, is the first location

encountered by IAV. Since Spn carriage is highest in children (36, 37), Spn colonization

often precedes IAV infection in childhood (38, 39). Given that infant mice support

efficient Spn colonization in the URT (25, 40), we investigated the impact of Spn

colonization on IAV transmission. All pups in a litter were colonized with Spn prior to IAV

infection of index pups to control for the efficient pup-to-pup transmission of Spn in the

setting of IAV infection (25). IAV shedding was collected daily for each pup prior to

evaluating for IAV transmission in contact littermates at 4 days p.i (Fig. 4A). We

observed that the Spn colonized contact mice acquired IAV at a decreased rate (32%)

as compared to uncolonized mice (100%) (p<0.0001) (Fig. 5B below graph), which

corresponded to lower viral shedding among colonized contacts (Fig. 5B left). Since

index Spn colonized and uncolonized mice infected with IAV (via inoculation) shed IAV

at similar levels, this suggested an antagonistic effect of Spn colonization on IAV

transmission through decreased acquisition by contact mice (Fig. 5B left).

Previous studies showed that sialidases expressed by colonizing Spn depletes

host sialic acid (SA) from the epithelial surface of the murine URT, allowing Spn to

utilize free SA for its nutritional requirements (41). Given that IAV requires SA for

efficient attachment, we evaluated the role of Spn sialidases on IAV acquisition. We

generated a double mutant lacking two common Spn sialidases: NanA and NanB

(Spn*nanA⁻nanB⁻*), and tested its ability to alter IAV transmission in our system. These

bacterial sialidases preferentially cleave 2,3-, 2,6-, and 2,8- or only 2,3-linked SA,

respectively (42). We found that by colonizing mice with the Spn*nanA⁻nanB⁻* mutant, we

completely restored the efficiency of IAV transmission from 32% to 100% ($p<0.0001$)

(Fig. 5B middle). We then tested the single sialidase mutants (Spn*nanA*- and Spn*nanB*-)

and found that the presence of NanA (via Spn*nanB*- colonization) was sufficient to limit

IAV acquisition by contacts by 50% ($p<0.001$). Notably, there was no correlation

between the colonization density of the bacterial mutants and their effect on shedding or

transmission.

To determine the role of sialidases in general in IAV acquisition, infant index and

contact mice were treated IN twice daily with *Vibrio cholerae* neuraminidase (VCNA),

which cleaves both 2,3- and 2,6-linked sialic acids (43). We found that, like Spn NanA,

VCNA treatment was sufficient to decrease IAV acquisition (71.4%) and inhibit shedding

by the contacts (Fig. 5B right). Together, these observations suggest that sialidase

activity from colonizing bacteria has the capacity to inhibit IAV acquisition in the URT,

specifically via its cleavage of 2,3- and 2,6-linked SA.

DISCUSSION

The inability to study in detail the host's role in IAV transmission has been a

major drawback of the ferret and guinea pig animal models, and has limited our current

understanding of IAV contagion (16-19). Herein we established an efficient and

tractable infant mouse IAV transmission model with the goal of utilizing the extensive

resources of mouse biology, to explore the role that host factors, immune pathways,

and the URT flora play in IAV transmission.

Our study corroborated previous findings that infant mice support efficient and

consistent IAV transmission (20, 25), and document an age-dependent effect on the

efficiency of transmission, highlighting inefficient transmission in adult mice. This

suggests an inherent quality of younger mice (i.e. less mobility allowing closer contact,

suckling, presence/absence of a host factor, microbiota composition, immunodefficient

or developmental status) which facilitates the shedding and transmissibility of IAV. In

humans, young age correlates with increased IAV nasopharyngeal shedding (44) and

longer duration of shedding (45), increasing the potential for transmission in this age

group. Although it has been shown in the study of Edenborough (23) that 56-day old

adult mice support the transmission of A/X-31 and A/Udorn/307/72 (H3N2) viruses at

100% efficiency in BALB/c mice, we were unable to observe comparable efficiency in

transmission using our A/X-31 virus in C57BL/6J mice older than 28-days of age. The

work of Lowen (22) also failed to observe IAV transmission in adult BALB/c mice, further

supporting an inconsistent transmission phenotype observed in adult mice which has

limited its utilization as an IAV transmission model. Notably, we are the first to

demonstrate that infant mice support efficient IBV transmission, which contrasts with the

inefficient IBV transmission previously reported in adult mice (32).

Although not evaluated in this study, the difference in mouse strains could also

affect the irregular success of IAV transmission in adult mice. One mouse strain,

C57BL/6J, has been tested in infant mice and demonstrated 100% transmission

efficiency in two independent studies (present study and (25)). In contrast, several

mouse strains have been tested for IAV transmission among adult mice, with variable

efficiencies among studies (0 to 100%) despite using the similar virus strains. They

include BALB/c (22, 23, 26, 28), C57BL/6J (present study), *Mx1*-competent C57BL/6J

(24), Swiss Webster (20, 26, 46), New Colony Swiss (47), Manor Farms (MF-1) (32),

DBA/2J (26), and Kunming (27). We learned from these studies that, host traits (mouse

age, strain, microbiota composition) all contribute to the infectivity and susceptibility of

the murine species to IAV, and the host's contribution to transmission should be

explored further using an efficient and tractable model of human disease.

Several studies have demonstrated that virus strain is an important determinant

of IAV transmission in mice (20, 23, 32, 46, 48). Like more recent studies (23, 48), we

highlighted the increased efficiency of transmission of A/H3N2 over A/H1N1 viruses. In

addition, A/H2N2 viruses (not tested here) have also shown to have increased

transmission efficiency in mice over A/H1N1 viruses (32, 46). Although we have not

evaluated specific viral moieties that confer a transmissible phenotype, the viral HA has

been demonstrated to play a role in transmission in mice (23). In addition, we postulate

that the activity of some NA in combination with specific HA favors viral release from the

nasal epithelium which allows viral shedding and transmission in mice. Thus together,

our data highlights that both host and viral-specific features are important to consider to

understand the requirements for IAV transmissibility.

We demonstrated that free virus particles present in secretions of mice, and not

replicating virus in the URT, correlated with IAV transmission efficiency. A similar

observation was also reported by Schulman (32), Edenborough (23), and Carrat (31)

demonstrating that transmissibility was associated with greater shedding of virus in

index mice, higher viral titers in the saliva of index mice, and shedding of virus from

infected humans, respectively. These studies supported our conclusion that viruses

which replicate in the URT without having the ability to exit the host (via shedding)

cannot be transmitted efficiently. Additionally, studies by Milton (44) suggested that URT

symptoms was associated with nasopharyngeal shedding in humans, and that coughing

was not necessary for the spread of infectious virus. This helps explains how mice,

which lack the cough reflex, can still produce and shed infectious virions. This

emphasizes an important future role of the infant mouse IAV transmission model as a

tool to study viral shedding as a surrogate marker of IAV contagion.

Two host traits have been identified in this study that influence IAV transmission:

IAV-specific immunoglobulin and the URT microbiota. The passive transfer of maternal

immunity is transferred via the placenta pre-natally in an IgG-dependent manner (49,

50), or via breastmilk post-natally mediated by several factors: immunoglobulins,

leukocytes, and antimicrobial/anti-inflammatory factors (51-53). Our data recapitulates

the value of maternal-infant transfer of IAV-specific IgG as a correlate of infant

immunity, by demonstrating a significant inhibitory effect on viral shedding and

transmission of infant mice after experimental (inoculation) and natural infection (via

transmission). Our experiments also demonstrate that IAV-specific serum IgG is

predominantly transferred via breastfeeding in mice, as previously reported (53, 54).

The concept of maternal serum IgG passage via breastmilk in mice has not often been

recognized, even though it has been shown to occur (53, 55-57). IgG can be

synthesized locally in the mammary gland, transferred across the mammary gland

epithelium, and subsequently transported from the infant gut to the circulation via

neonatal Fc receptors (FcRn) expressed in the proximal intestine (58-60). Although this

mechanism of maternal IgG acquisition by infants has not yet been correlated in

humans, presence of FcRn in the human intestine has been confirmed (61, 62). Our

study does not address the contribution of secretory IgA, which is known to be the most

abundant immunoglobulin in breastmilk. Yet, our data suggest that adequate amounts

of IAV-specific IgG, which is known to wane within 8 weeks of birth in infants (63),

maybe necessary to maintain anti-IAV immunity in the URT and limit IAV transmission

in infants, given that at this young age, infants don't have a fully functional adaptive

immune response.

In addition to humoral immunity, our study identified an inhibitory role for the

common URT colonizer, Spn, at the step of viral acquisition during transmission of IAV.

This phenomenon has never been previously observed, although there has been some

evidence suggesting that the preceding colonization of Spn reduces IAV infection (25,

64). Notably, the study by McCullers (64) showed that preceding colonization with Spn

protected mice from mortality after IAV challenge, whereas the reverse process: prior

infection with IAV with subsequent challenge with Spn yielded the opposite effect. This

implied that the timing of pathogen encounter mattered, and the composition of the host

microbiota may serve a "prophylactic-like" protective effect. Although no studies have

evaluated the role of the respiratory microbiota on IAV transmission, we hypothesize

that the differences between the transmissibility of different IAV strains and

susceptibility of different populations (infants vs adults) to IAV may be due to a

combined effect of the virus's ability to release from SA and exit the host via shedding,

and the susceptibility to viral acquisition by contact hosts based on the composition of

their URT microbiota. Our work provides proof-of-principle and highlights the

amenability of the infant mouse model as a tool to understand the complex dynamics of

virus and host, and their combined effect in IAV transmission.

Lastly, we demonstrate the role of Spn sialidases, NanA and NanB, in

antagonizing the acquisition and shedding of IAV by contact mice. We hypothesize that

bacteria-driven de-sialylation of the host's URT glycoproteins for use as nutrient (41),

may deplete SA residues necessary for IAV adhesion and infection, thus limiting virus

susceptibility, and hence acquisition. Notably, the antagonistic effect of bacterial

sialidases on IAV shedding of the index group is not statistically different from

uncolonized controls. This is analogous to the clinical effects of NI, whereby oseltamivir

treatment of index cases alone has not been shown to reduce viral shedding (10, 11).

Only when treatment of both index and naïve contacts were partaken (as in post-

exposure prophylaxis), has the effects of NI been effective at preventing acquisition of

infection among the contact group (12, 13). Because SA is the primary recognition

moiety for many viral respiratory pathogens, the concept of utilizing bacterial sialidases

as a broad antiviral agent is currently being explored in humans, although its effect on

transmission has not yet been evaluated (65-72).

While the advantages of using murine models are evident, these can also be

drawbacks. Humans are genetically diverse, live in complex environments, and have

been exposed to a myriad of pathogens, all of which can affect transmissibility and

susceptibility to IAV, therefore findings generated in animal models of human disease

should always be cautiously interpreted. Nevertheless, studying the complexities of IAV

transmission biology in a tractable animal model such as infant mice, will allow intricate

and sophisticated investigations, which will further our understanding of IAV contagion

that may translate into better vaccines and therapeutics.

MATERIALS AND METHODS

Mice. C57BL/6J mice (Jackson Laboratories, ME) were maintained and bred in a

conventional animal facility. Pups were housed with their mother for the duration of all

experiments. Animal studies were conducted in accordance with the *Guide for the Care*

and Use of Laboratory Animals (73), and approved by the *Institutional Animal Care and*

Use Committee of NYU Langone Health (Assurance Number A3317-01). All procedures

were in compliance with *Biosafety in Microbiological and Biomedical Laboratories*.

Cells and viruses. Madin-Darby canine kidney (MDCK) cells were cultured in

Dulbecco's modified Eagle's medium with 10% fetal bovine serum and 1% penicillin-

streptomycin (Gibco).

Viruses: A/X-31 (H3N2) [HA/NA genes from A/Aichi/2/1968 and internal genes from

A/Puerto Rico/8/1934], a gift from Jan Erikson (U. Penn), whose sequence has been

deposited in GenBank (XXXXXX). The following reagents were obtained through BEI

Resources (NIAID, NIH): A/X-47 (H3N2) [HA/NA genes from A/Victoria/3/1975 and

internal genes from A/Puerto Rico/8/1934] [NR-3663], A/Hong Kong/1/1968-2 MA 21-2

(H3N2) [NR-28634], A/Puerto Rico/8/1934 (H1N1) V-301-011-000 [NR-3169],

A/WSN/1933 (H1N1) [NR-2555], A/Brisbane/59/2007 (H1N1) [NR-12282],

A/California/4/2009 (H1N1) [NR-13659], B/Lee/1940 V-302-001-000 [NR-3178]. IAV and

IBV were propagated in 8-10-day old embryonated chicken eggs (Charles River, CT) for

2 days at 37^{o}C and 33^{o}C, respectively. All viruses were tittered by standard plaque

assay in MDCK cells in the presence of TPCK (tolylsulfonyl phenylalanyl chloromethyl

ketone)-treated trypsin (Thermo Scientific) (74). Purified virus for ELISA was prepared

by harvesting allantoic fluid from eggs containing virus followed by centrifugation

(3,000×g, 30min, 4°C) to remove debris. Viruses were pelleted through a 30% sucrose

cushion (30% sucrose in NTE buffer [100mM NaCl+10mM Tris-HCl+1mM EDTA], pH

7.4) by ultracentrifugation (83,000×g, 2hrs), resuspended in phosphate-buffered saline

(PBS), and stored at -80°C.

Virus infection, shedding, and transmission. Pups in a litter (4-7 days of age) were

infected (index) with a 3µl sterile PBS inoculum without general anesthesia (to avoid

direct lung inoculation) by IN instillation of 250 PFU of IAV (unless otherwise specified),

and returned to the litter at the time of inoculation for the duration of the experiment.

Shedding of virus was collected by dipping the nares of each mouse in viral media

(PBS+0.3% BSA) daily, and samples evaluated via plaque assay. Intralitter

transmission was assessed in littermates (contact) at 4-5 days p.i. (day 10-14 of life).

Pups and mother were euthanized by CO_2 asphyxiation followed by cardiac puncture,

the URT was subjected to a retrograde lavage (flushing 300µl PBS from the trachea and

collecting through the nares), and samples were used to quantify virus (via plaque

assay or qRT-PCR) or Spn density (described below). Ratio of index to contact pups

ranged from 1:3-1:4.

Where indicated, pups were IN treated twice daily with 90 µU (3µl inoculum) of *Vibrio*

cholerae neuraminidase (VCNA, Sigma-Aldrich).

The MID_{50} was calculated by the Reed and Müench method (75).

Induction of maternal IAV immunity. Adult female mice were infected IN with 250

PFU of A/X-31 in a 6µl inoculum without anesthesia. Mice were left to recover from

infection for 7 days prior to breeding. Litters of immune mothers were used in

experiments.

Bacteria strain construction and culture. A streptomycin-resistant derivative of

capsule type-4 isolate TIGR4 (**P2406**) was used in this study and cultured on tryptic-soy

(TS)-agar-streptomycin (200µg/ml) plates (40). The *nanA*- knockout strain (**P2508**) was

constructed by transforming P2406 with genomic DNA from strain P2082 (76)

(MasterPure DNA purification kit, Illumina), and selection on TS-agar-chloramphenicol

(2.5µg/ml) plates. The *nanB*- knockout strain (**P2511**) was constructed by amplifying the

Janus cassette (77) from genomic DNA of strain P2408 (78), with flanking upstream and

downstream regions to the *nanB* gene added via isothermal assembly. Strain P2406

was transformed with the PCR product, and the transformants selected on TS-agar-

kanamycin (125µg/ml) plates. The *nanA-nanB*- double knockout strain (**P2545**) was

constructed by transforming P2511 with genomic DNA from strain P2508, and

transformants selected on TS-agar-chloramphenicol plates.

Spn strains were grown statically in TS broth (BD, NJ) at 37°C to an optical density

(OD) 620nm of 1.0. For quantitation, serial dilutions (1:10) of the inoculum or URT

lavages were plated on TS-agar-antibiotic selection plates with 100µl catalase (30,000

U/ml, Worthington Biochemical) and incubated overnight (37°C, 5% CO_2). Bacteria were

stored in 20% glycerol at −80°C. Colonization of pups was carried out by IN instillation

of 10^3 CFU in 3µl of PBS, 1 or 3 days prior to IAV infection.

qRT-PCR. Following a retrograde URT lavage with 300µl RLT lysis buffer, RNA was

isolated (RNeasy Kit, Qiagen), cDNA was generated (High-Capacity RT kit, Applied

Biosystems), and used for quantitative PCR (SYBR Green PCR Master Mix, Applied

Biosystems). Results were analyzed using the $2^{-\Delta\Delta CT}$ method (79) by comparison to

GAPDH transcription. Values represent the fold change over uninfected.

ELISA. Immulon 4 HBX plates (Thermo Scientific) were coated with 5µg/ml purified A/X-

31 in coating buffer (0.015M Na_2CO_3+0.035M $NaHCO_3$, 50µl/well), and incubated

overnight, 4°C. After three washes with PBS-T (PBS+0.1% Tween 20, 100µl/well),

plates were incubated with blocking solution (BS) (PBS-T+0.5% milk powder+3% goat

serum [ThermoFisher], 1hr, 20°C). BS was discarded, mice sera were diluted to a

starting concentration of 1:100, then serially diluted 1:2 in BS (100µl/well), and

incubated (2hr, 20°C). Three washes with PBS-T was done prior to adding secondary

antibody (horseradish peroxidase [HRP]-labeled anti-mouse IgG [whole Ab], GE

Healthcare or alkaline phosphatase [AP]-labeled anti-mouse IgA [α chain], Sigma)

diluted in BS (1:3000, 50µl/well). After incubation (1hr, 20°C) and three washes with

PBS-T, plates were developed for 10min using 100µl/well SigmaFast OPD (*o*-

phenylenediamine dihydrochloride [Sigma] and stopped with 3M HCl (50µl/well) or

developed for 1-18hr using pNPP (p-nitrophenyl phosphate) [KPL];). Plates were read at

OD490nm for the OPD substrate or 405nm for the AP substrate. The endpoint titers

were determined by calculating the dilution at which the absorbance is equal to 0.1. The

geometric mean titers (GMT) were calculated from the reciprocal of the endpoint titers.

Statistical analysis. GraphPad Prism 7 software was used for all statistical analyses.

Unless otherwise noted, data were analyzed using the Mann-Whitney U test to compare

two groups, and the Kruskal-Wallis test with Dunn's post-analysis for multiple group

comparisons.

Data Availability. The authors confirm that data will be made publicly available upon

publication upon request, without restriction.

ACKNOWLEDGMENTS

We thank Elodie Guedin (NYU) for the sequencing of A/X-31. MBO was supported by

the NYU Division of Infectious Diseases, Department of Medicine Silverman

Scholarship, and Physician Scientist Training Program Award; SB was supported by the

NYU School of Medicine and National Institute of Diabetes & Digestive & Kidney

Diseases fellowship. This project was supported by grants from the U.S. Public Health

Service to JNW (AI038446 and AI105168).

REFERENCES

1. Rolfes MA FI, Garg S, Flannery B, Brammer L, Singleton JA, et al. 2016. Estimated Influenza Illnesses, Medical Visits, Hospitalizations, and Deaths Averted by Vaccination in the United States. https://www.cdc.gov/flu/about/disease/2015-16.htm. Accessed 9/2018.

2. CDC. 9/27/2018. National Press Conference Kicks Off 2018-2019 Flu Vaccination Campaign. https://www.cdc.gov/flu/spotlights/press-conference-2018-19.htm#ref1. Accessed 9/2018.

3. Flannery B, Reynolds SB, Blanton L, Santibanez TA, O'Halloran A, Lu PJ, Chen J, Foppa IM, Gargiullo P, Bresee J, Singleton JA, Fry AM. 2017. Influenza Vaccine Effectiveness Against Pediatric Deaths: 2010-2014. Pediatrics 139.

4. Arriola C, Garg S, Anderson EJ, Ryan PA, George A, Zansky SM, Bennett N, Reingold A, Bargsten M, Miller L, Yousey-Hindes K, Tatham L, Bohm SR, Lynfield R, Thomas A, Lindegren ML, Schaffner W, Fry AM, Chaves SS. 2017. Influenza Vaccination Modifies Disease Severity Among Community-dwelling Adults Hospitalized With Influenza. Clin Infect Dis 65:1289-1297.

5. CDC. 9/19/2014. Selected Publications on Influenza Vaccine Effectiveness, *on* Centers for Disease Control and Prevention. https://www.cdc.gov/flu/about/qa/publications.htm. Accessed 9/2018.

6. Lowen AC, Steel J, Mubareka S, Carnero E, Garcia-Sastre A, Palese P. 2009. Blocking interhost transmission of influenza virus by vaccination in the guinea pig model. J Virol 83:2803-18.

7. Houser KV, Pearce MB, Katz JM, Tumpey TM. 2013. Impact of prior seasonal
H3N2 influenza vaccination or infection on protection and transmission of
emerging variants of influenza A(H3N2)v virus in ferrets. J Virol 87:13480-9.
8. CDC. 9/27/2018. 2017-18 Flu Season.
https://www.cdc.gov/flu/fluvaxview/1718season.htm. Accessed 10/2018.
9. CDC. 9/25/2018. Seasonal Influenza Vaccine Effectiveness, 2004-2018.
https://www.cdc.gov/flu/professionals/vaccination/effectiveness-studies.htm.
Accessed 10/2018.
10. Cheung DH, Tsang TK, Fang VJ, Xu J, Chan KH, Ip DK, Peiris JS, Leung GM,
Cowling BJ. 2015. Association of Oseltamivir Treatment With Virus Shedding,
Illness, and Household Transmission of Influenza Viruses. J Infect Dis 212:391-6.
11. Ng S, Cowling BJ, Fang VJ, Chan KH, Ip DK, Cheng CK, Uyeki TM, Houck PM,
Malik Peiris JS, Leung GM. 2010. Effects of oseltamivir treatment on duration of
clinical illness and viral shedding and household transmission of influenza virus.
Clin Infect Dis 50:707-14.
12. Hayden FG, Belshe R, Villanueva C, Lanno R, Hughes C, Small I, Dutkowski R,
Ward P, Carr J. 2004. Management of influenza in households: a prospective,
randomized comparison of oseltamivir treatment with or without postexposure
prophylaxis. J Infect Dis 189:440-9.
13. Welliver R, Monto AS, Carewicz O, Schatteman E, Hassman M, Hedrick J,
Jackson HC, Huson L, Ward P, Oxford JS, Oseltamivir Post Exposure
Prophylaxis Investigator G. 2001. Effectiveness of oseltamivir in preventing
influenza in household contacts: a randomized controlled trial. JAMA 285:748-54.

14.   Halloran ME, Hayden FG, Yang Y, Longini IM, Jr., Monto AS. 2007. Antiviral

528   effects on influenza viral transmission and pathogenicity: observations from

529   household-based trials. Am J Epidemiol 165:212-21.

530   15.   Hayward A. 2010. Does treatment with oseltamivir prevent transmission of

531   influenza to household contacts? Clin Infect Dis 50:715-6.

532   16.   Herfst S, Schrauwen EJ, Linster M, Chutinimitkul S, de Wit E, Munster VJ, Sorrell

533   EM, Bestebroer TM, Burke DF, Smith DJ, Rimmelzwaan GF, Osterhaus AD,

534   Fouchier RA. 2012. Airborne transmission of influenza A/H5N1 virus between

535   ferrets. Science 336:1534-41.

536   17.   Imai M, Watanabe T, Hatta M, Das SC, Ozawa M, Shinya K, Zhong G, Hanson

537   A, Katsura H, Watanabe S, Li C, Kawakami E, Yamada S, Kiso M, Suzuki Y,

538   Maher EA, Neumann G, Kawaoka Y. 2012. Experimental adaptation of an

539   influenza H5 HA confers respiratory droplet transmission to a reassortant H5

540   HA/H1N1 virus in ferrets. Nature 486:420-8.

541   18.   Schrauwen EJ, Fouchier RA. 2014. Host adaptation and transmission of

542   influenza A viruses in mammals. Emerg Microbes Infect 3:e9.

543   19.   Lowen AC, Mubareka S, Steel J, Palese P. 2007. Influenza virus transmission is

544   dependent on relative humidity and temperature. PLoS Pathog 3:1470-6.

545   20.   Eaton MD. 1940. Transmission of Epidemic Influenza Virus in Mice by Contact. J

546   Bacteriol 39:229-41.

547   21.   Schulman JL, Kilbourne ED. 1962. Airborne transmission of influenza virus

548   infection in mice. Nature 195:1129-30.

549   22.   Lowen AC, Mubareka S, Tumpey TM, Garcia-Sastre A, Palese P. 2006. The

550   guinea pig as a transmission model for human influenza viruses. Proc Natl Acad

551   Sci U S A 103:9988-92.

552   23.   Edenborough KM, Gilbertson BP, Brown LE. 2012. A mouse model for the study

553   of contact-dependent transmission of influenza A virus and the factors that

554   govern transmissibility. J Virol 86:12544-51.

555   24.   Klinkhammer J, Schnepf D, Ye L, Schwaderlapp M, Gad HH, Hartmann R,

556   Garcin D, Mahlakoiv T, Staeheli P. 2018. IFN-lambda prevents influenza virus

557   spread from the upper airways to the lungs and limits virus transmission. Elife 7.

558   25.   Diavatopoulos DA, Short KR, Price JT, Wilksch JJ, Brown LE, Briles DE,

559   Strugnell RA, Wijburg OL. 2010. Influenza A virus facilitates Streptococcus

560   pneumoniae transmission and disease. FASEB J 24:1789-98.

561   26.   Price GE, Lo CY, Misplon JA, Epstein SL. 2014. Mucosal immunization with a

562   candidate universal influenza vaccine reduces virus transmission in a mouse

563   model. J Virol 88:6019-30.

564   27.   Wu R, Sui Z, Liu Z, Liang W, Yang K, Xiong Z, Xu D. 2010. Transmission of

565   avian H9N2 influenza viruses in a murine model. Vet Microbiol 142:211-6.

566   28.   Rigoni M, Toffan A, Viale E, Mancin M, Cilloni F, Bertoli E, Salomoni A, Marciano

567   S, Milani A, Zecchin B, Capua I, Cattoli G. 2010. The mouse model is suitable for

568   the study of viral factors governing transmission and pathogenesis of highly

569   pathogenic avian influenza (HPAI) viruses in mammals. Vet Res 41:66.

570   29.   Ivinson K, Deliyannis G, McNabb L, Grollo L, Gilbertson B, Jackson D, Brown LE.

571   2017. Salivary Blockade Protects the Lower Respiratory Tract of Mice from

572   Lethal Influenza Virus Infection. J Virol 91.

573   30.   Bouvier NM, Lowen AC. 2010. Animal Models for Influenza Virus Pathogenesis

574   and Transmission. Viruses 2:1530-1563.

575   31.   Carrat F, Vergu E, Ferguson NM, Lemaitre M, Cauchemez S, Leach S, Valleron

576   AJ. 2008. Time lines of infection and disease in human influenza: a review of

577   volunteer challenge studies. Am J Epidemiol 167:775-85.

578   32.   Schulman JL. 1967. Experimental transmission of influenza virus infection in

579   mice. IV. Relationship of transmissibility of different strains of virus and recovery

580   of airborne virus in the environment of infector mice. J Exp Med 125:479-88.

581   33.   Ichinohe T, Pang IK, Kumamoto Y, Peaper DR, Ho JH, Murray TS, Iwasaki A.

582   2011. Microbiota regulates immune defense against respiratory tract influenza A

583   virus infection. Proc Natl Acad Sci U S A 108:5354-9.

584   34.   Rosshart SP, Vassallo BG, Angeletti D, Hutchinson DS, Morgan AP, Takeda K,

585   Hickman HD, McCulloch JA, Badger JH, Ajami NJ, Trinchieri G, Pardo-Manuel

586   de Villena F, Yewdell JW, Rehermann B. 2017. Wild Mouse Gut Microbiota

587   Promotes Host Fitness and Improves Disease Resistance. Cell 171:1015-1028

588   e13.

589   35.   Lee KH, Gordon A, Shedden K, Kuan G, Ng S, Balmaseda A, Foxman B. 2018.

590   The respiratory microbiome and susceptibility to influenza virus infection. bioRxiv

591   doi:10.1101/372649.

36.   Bogaert D, van Belkum A, Sluijter M, Luijendijk A, de Groot R, Rumke HC,

Verbrugh HA, Hermans PW. 2004. Colonisation by Streptococcus pneumoniae

and Staphylococcus aureus in healthy children. Lancet 363:1871-2.

37.   Adler H, Ferreira DM, Gordon SB, Rylance J. 2017. Pneumococcal Capsular

Polysaccharide Immunity in the Elderly. Clin Vaccine Immunol 24.

38.   Takano M, Ozaki K, Nitahara Y, Higuchi W, Takano T, Nishiyama A, Yamamoto

T. 2009. Streptococcus pneumoniae and Haemophilus influenzae at the initial

stage of influenza. Pediatr Int 51:687-95.

39.   Short KR, Habets MN, Hermans PW, Diavatopoulos DA. 2012. Interactions

between Streptococcus pneumoniae and influenza virus: a mutually beneficial

relationship? Future Microbiol 7:609-24.

40.   Zafar MA, Kono M, Wang Y, Zangari T, Weiser JN. 2016. Infant Mouse Model for

the Study of Shedding and Transmission during Streptococcus pneumoniae

Monoinfection. Infect Immun 84:2714-22.

41.   Siegel SJ, Roche AM, Weiser JN. 2014. Influenza promotes pneumococcal

growth during coinfection by providing host sialylated substrates as a nutrient

source. Cell Host Microbe 16:55-67.

42.   Xu G, Kiefel MJ, Wilson JC, Andrew PW, Oggioni MR, Taylor GL. 2011. Three

Streptococcus pneumoniae sialidases: three different products. J Am Chem Soc

133:1718-21.

43.   Moustafa I, Connaris H, Taylor M, Zaitsev V, Wilson JC, Kiefel MJ, von Itzstein

M, Taylor G. 2004. Sialic acid recognition by Vibrio cholerae neuraminidase. J

Biol Chem 279:40819-26.

44. Yan J, Grantham M, Pantelic J, Bueno de Mesquita PJ, Albert B, Liu F, Ehrman

S, Milton DK, Consortium E. 2018. Infectious virus in exhaled breath of

symptomatic seasonal influenza cases from a college community. Proc Natl Acad

Sci U S A 115:1081-1086.

45. Ng S, Lopez R, Kuan G, Gresh L, Balmaseda A, Harris E, Gordon A. 2016. The

Timeline of Influenza Virus Shedding in Children and Adults in a Household

Transmission Study of Influenza in Managua, Nicaragua. Pediatr Infect Dis J

35:583-6.

46. Schulman JL, Kilbourne ED. 1963. Experimental Transmission of Influenza Virus

Infection in Mice. I. The Period of Transmissibility. J Exp Med 118:257-66.

47. Schulman JL, Kilbourne ED. 1963. Experimental Transmission of Influenza Virus

Infection in Mice. Ii. Some Factors Affecting the Incidence of Transmitted

Infection. J Exp Med 118:267-75.

48. Short KR, Reading PC, Wang N, Diavatopoulos DA, Wijburg OL. 2012.

Increased nasopharyngeal bacterial titers and local inflammation facilitate

transmission of Streptococcus pneumoniae. MBio 3.

49. Puck JM, Glezen WP, Frank AL, Six HR. 1980. Protection of infants from

infection with influenza A virus by transplacentally acquired antibody. J Infect Dis

142:844-9.

50. Reuman PD, Ayoub EM, Small PA. 1987. Effect of passive maternal antibody on

influenza illness in children: a prospective study of influenza A in mother-infant

pairs. Pediatr Infect Dis J 6:398-403.

51.   Goldman AS. 1993. The immune system of human milk: antimicrobial,

638   antiinflammatory and immunomodulating properties. Pediatr Infect Dis J 12:664-

639   71.

640   52.   Slade HB, Schwartz SA. 1987. Mucosal immunity: the immunology of breast milk.

641   J Allergy Clin Immunol 80:348-58.

642   53.   Reuman PD, Paganini CM, Ayoub EM, Small PA, Jr. 1983. Maternal-infant

643   transfer of influenza-specific immunity in the mouse. J Immunol 130:932-6.

644   54.   Brambell FW. 1969. The transmission of immune globulins from the mother to

645   the foetal and newborn young. Proc Nutr Soc 28:35-41.

646   55.   Macchiaverni P, Arslanian C, Frazao JB, Palmeira P, Russo M, Verhasselt V,

647   Condino-Neto A. 2011. Mother to child transfer of IgG and IgA antibodies against

648   Dermatophagoides pteronyssinus. Scand J Immunol 74:619-27.

649   56.   Palmeira P, Costa-Carvalho BT, Arslanian C, Pontes GN, Nagao AT, Carneiro-

650   Sampaio MM. 2009. Transfer of antibodies across the placenta and in breast milk

651   from mothers on intravenous immunoglobulin. Pediatr Allergy Immunol 20:528-

652   35.

653   57.   Van de Perre P. 2003. Transfer of antibody via mother's milk. Vaccine 21:3374-6.

654   58.   Cianga P, Medesan C, Richardson JA, Ghetie V, Ward ES. 1999. Identification

655   and function of neonatal Fc receptor in mammary gland of lactating mice. Eur J

656   Immunol 29:2515-23.

657   59.   Roopenian DC, Akilesh S. 2007. FcRn: the neonatal Fc receptor comes of age.

658   Nat Rev Immunol 7:715-25.

60. Mackenzie N. 1984. Fc receptor-mediated transport of immunoglobulin across

the intestinal epithelium of the neonatal rodent. Immunol Today 5:364-6.

61. Israel EJ, Taylor S, Wu Z, Mizoguchi E, Blumberg RS, Bhan A, Simister NE.

1997. Expression of the neonatal Fc receptor, FcRn, on human intestinal

epithelial cells. Immunology 92:69-74.

62. Story CM, Mikulska JE, Simister NE. 1994. A major histocompatibility complex

class I-like Fc receptor cloned from human placenta: possible role in transfer of

immunoglobulin G from mother to fetus. J Exp Med 180:2377-81.

63. Nunes MC, Cutland CL, Jones S, Hugo A, Madimabe R, Simoes EA, Weinberg

A, Madhi SA, Maternal Flu Trial T. 2016. Duration of Infant Protection Against

Influenza Illness Conferred by Maternal Immunization: Secondary Analysis of a

Randomized Clinical Trial. JAMA Pediatr 170:840-7.

64. McCullers JA, Rehg JE. 2002. Lethal synergism between influenza virus and

Streptococcus pneumoniae: characterization of a mouse model and the role of

platelet-activating factor receptor. J Infect Dis 186:341-50.

65. Bergelson LD, Bukrinskaya AG, Prokazova NV, Shaposhnikova GI, Kocharov

SL, Shevchenko VP, Kornilaeva GV, Fomina-Ageeva EV. 1982. Role of

gangliosides in reception of influenza virus. Eur J Biochem 128:467-74.

66. Griffin JA, Basak S, Compans RW. 1983. Effects of hexose starvation and the

role of sialic acid in influenza virus release. Virology 125:324-34.

67. Air GM, Laver WG. 1995. Red cells bound to influenza virus N9 neuraminidase

are not released by the N9 neuraminidase activity. Virology 211:278-84.

68.   Thompson CI, Barclay WS, Zambon MC, Pickles RJ. 2006. Infection of human

682   airway epithelium by human and avian strains of influenza a virus. J Virol

683   80:8060-8.

684   69.   Malakhov MP, Aschenbrenner LM, Smee DF, Wandersee MK, Sidwell RW,

685   Gubareva LV, Mishin VP, Hayden FG, Kim DH, Ing A, Campbell ER, Yu M, Fang

686   F. 2006. Sialidase fusion protein as a novel broad-spectrum inhibitor of influenza

687   virus infection. Antimicrob Agents Chemother 50:1470-9.

688   70.   Nicholls JM, Moss RB, Haslam SM. 2013. The use of sialidase therapy for

689   respiratory viral infections. Antiviral Res 98:401-9.

690   71.   Triana-Baltzer GB, Gubareva LV, Nicholls JM, Pearce MB, Mishin VP, Belser JA,

691   Chen LM, Chan RW, Chan MC, Hedlund M, Larson JL, Moss RB, Katz JM,

692   Tumpey TM, Fang F. 2009. Novel pandemic influenza A(H1N1) viruses are

693   potently inhibited by DAS181, a sialidase fusion protein. PLoS One 4:e7788.

694   72.   Salvatore M, Satlin MJ, Jacobs SE, Jenkins SG, Schuetz AN, Moss RB, Van

695   Besien K, Shore T, Soave R. 2016. DAS181 for Treatment of Parainfluenza Virus

696   Infections in Hematopoietic Stem Cell Transplant Recipients at a Single Center.

697   Biol Blood Marrow Transplant 22:965-70.

698   73.   U.S. Department of Health and Human Services NIoH. 2015. Public Health

699   Service Policy on Humane Care and Use of Laboratory Animals.

700   74.   Baer A, Kehn-Hall K. 2014. Viral concentration determination through plaque

701   assays: using traditional and novel overlay systems. J Vis Exp

702   doi:10.3791/52065:e52065.

703   75.   Reed LJ, Muench H. 1938. A SIMPLE METHOD OF ESTIMATING FIFTY PER

704   CENT ENDPOINTS12. American Journal of Epidemiology 27:493-497.

705   76.   Dalia AB, Standish AJ, Weiser JN. 2010. Three surface exoglycosidases from

706   Streptococcus pneumoniae, NanA, BgaA, and StrH, promote resistance to

707   opsonophagocytic killing by human neutrophils. Infect Immun 78:2108-16.

708   77.   Sung CK, Li H, Claverys JP, Morrison DA. 2001. An rpsL cassette, janus, for

709   gene replacement through negative selection in Streptococcus pneumoniae. Appl

710   Environ Microbiol 67:5190-6.

711   78.   Lemon JK, Weiser JN. 2015. Degradation products of the extracellular pathogen

712   Streptococcus pneumoniae access the cytosol via its pore-forming toxin. MBio 6.

713   79.   Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using

714   real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods

715   25:402-8.

FIGURE LEGENDS

Figure 1. Transmission of IAV in infant mice.

(A) Schematic and timeline of experimental design. Index and contact pups were

arbitrarily assigned, maintained in the same cage, and cared for by the same mother. At

day 0 (4-7 days of age), pups were infected IN with 250 PFU of A/X-31 (index), and

cohoused with uninfected littermates (contact) for 4-5 days prior to evaluating for

transmission. (B) Transmission of IAV to contact pups was evaluated via qRT-PCR (left

panel) or plaque assay (right panel) from retrograde URT lavages after sacrifice.

URT titers are represented by a box plot extending from the minimum to maximum

values for each data set. Each symbol represent the titer measured from a single pup

with median values indicated by a line within the box. Index and contact pups are shown

in black and red symbols, respectively. (C) Window of transmission was evaluated by

separating index and contact pups for a defined period prior to contact. After infection of

index pups, uninfected contact pups were housed apart (in a separate cage) for 0 and

72 hours prior to cohousing with infected index for 5 days. Transmission to contact pups

was evaluated via plaque assay from retrograde URT lavages. URT titers are

represented by a box plot as described above. (D) Timecourse of A/X-31 transmission.

Pups in a litter were subjected to an A/X-31 transmission experiment (described above)

and transmission to contact pups was evaluated via qRT-PCR from retrograde URT

lavages at indicated day post contact. Mean URT titers ± standard deviation (SD) are

represented. (E) Morbidity of A/X-31 infection in index and contact pups over the course

of 20 days. Pups in a litter were subjected to an A/X-31 transmission experiment

(described above) and weight of each pup was measured daily. Percent of initial weight

± SD is represented (uninfected group n=9, index group n=3-4, contact group n=4-5).

Differences among group means were analyzed using the Student's *t* test.

All panels represent at least two independent experiments. * *p*<0.05, ** *p*<0.01,

IAV=Influenza A virus, URT=upper respiratory tract, NP=nucleoprotein, PFU=plaque

forming unit, LOD=Limit of detection.

Figure 2. Shedding of IAV.

(A) Image of infant mouse shedding procedure and schematic timeline of experimental

design. At day 0 (4-7 days of age), pups were infected IN with 250 PFU of A/X-31

(index), and cohoused with uninfected littermates (contact) for 10 days. Shedding of IAV

was collected by dipping the nares of each mouse in viral media daily. (B) Shedding

samples from each day were evaluated individually via plaque assay. Each symbol

represents the shedding titer measured from a single mouse at the specified day. Index

and contact pups are shown in black and red symbols, respectively. Mean values are

connected by a line. IAV=Influenza A virus, PFU=plaque forming unit, LOD=Limit of

detection.

Figure 3. Maternal-infant transfer of IAV-specific immunity limits IAV shedding

and transmission.

(A) Adult females were infected IN with 250 PFU of A/X-31 and were left to recover from

infection prior to breeding. Soon after birth, pups born from previously infected

(immune) mothers or those born from non-immune mothers were either left with their

biological mother or exchanged with a foster mother of opposite immune status. Pups

paired with their biological or foster mothers were left to acclimate until 4-5 days of life,

prior to being subjected to an IAV transmission experiment. Schematic for each

experimental condition is shown. Pups in a litter were infected IN with 250 PFU of A/X-

31 (index), and cohoused with uninfected littermates (contact) for 5 days. Shedding of

IAV was collected by gently dipping the nares of each mouse in viral media daily.

Shedding samples from each day were evaluated individually via plaque assay for each

pup. Shedding titers shown represent pooled values for days 1-5 for index pups and

days 4-5 for contact pups, representing days of maximum shedding for each group (as

per Fig. 2). Each symbol represent the shedding titer measured from a single mouse for

a specific day. Index and contact pups are shown in black and red symbols,

respectively. Median values are indicated. At the end of 5 days, pups and mothers were

sacrificed, and transmission to contact pups was evaluated via plaque assay from

retrograde URT lavages. Percentage of transmission among contact pups is displayed

below the graph. (B) Serum from mother and pups were obtained at the time of

sacrifice. Samples from individual mice were evaluated for IAV-specific IgG by ELISA.

IgG geometric mean titers (GMT) are represented by a box plot extending from the 25th

to 75th percentile for each data set. Whiskers for each box encompasses the minimum

to maximum values. Median values are indicated by a line within the box.

All panels represent at least two independent experiments. Differences in transmission

were analyzed using the Fisher's exact test. * $p<0.05$, ** $p<0.01$, *** $p<0.001$, ****

$p<0.0001$, PFU=plaque forming unit, LOD=Limit of detection, GMT=Geometric mean

titer.

Figure 4. *Streptococcus pneumoniae* sialidases limit acquisition of IAV via

transmission.

(A) Schematic timeline of experimental design. At day -1 or -3 (3-4 days of age), all

pups in a litter were colonized IN with either wild-type Spn; mutant Spn lacking NanA,

NanB, or both; or treated IN with *Vibrio cholerae* neuraminidase (VCNA) twice daily. At

day 0, pups were infected IN with 250 PFU of A/X-31 (index), and cohoused with

uninfected littermates (contact) for 4 days. Shedding of IAV was collected by dipping the

nares of each mouse in viral media daily. Transmission to contact pups was evaluated

at day 4. (B) Shedding samples from each day were evaluated individually via plaque

assay for each pup. Shedding titers shown represent pooled values for days 1-4 for

index pups and days 3-4 for contact pups. Each symbol represent the shedding titer

measured from a single mouse for a specific day with median values indicated. Index

and contact pups are shown in black and red symbols, respectively. At the end of 4

days, pups were sacrificed, and transmission to contacts was evaluated via plaque

assay from retrograde URT lavages. Percentage of transmission among contact pups is

displayed below the graph. Density of colonizing Spn was measured in URT lavage

samples of each pup. Each blue symbol represents the median Spn density ±

interquartile range for each group. Differences in transmission were analyzed using the

Fisher's exact test. * $p<0.05$, ** $p<0.01$, *** $p<0.001$, **** $p<0.0001$, PFU=plaque

forming unit, CFU=colony forming unit, LOD=Limit of detection.

Figure S1. Window of IAV transmission.

As Figure 1C, the window of transmission was evaluated by separating index and

contact pups for a defined period prior to contact. After infection of index pups,

uninfected contact pups were housed apart (in a separate cage) for additional

timepoints, including 24, 48, and 72 hours prior to cohousing with infected index for 5

days. Transmission to contact pups was evaluated via plaque assay from retrograde

URT lavages. URT titers are represented by a box plot extending from the minimum to

maximum values for each data set. Each symbol represent the titer measured from a

single pup with median values indicated by a line within the box. Index and contact pups

are shown in black and red symbols, respectively. URT=upper respiratory tract.

Figure S2. Mode of transmission is via direct pup-to-pup contact.

Upper panels represent schematic for each experimental condition. The mother or cage

contents were evaluated as possible sources of transmission (fomites). (A) All infants in

one litter (4-7 days of age) were infected IN with 250 PFU of A/X-31 (index) while the

second litter in a separate cage were left uninfected (contact). The mothers from the

infected and uninfected cage were exchanged daily without disturbing the pups or cage

contents. After 5 days, transmission to contact litter was evaluated via plaque assay

from retrograde URT lavages. (B) Index pups in a litter were infected and kept

separated from an uninfected contact litter as described above. Cages and cage

contents (bedding) from infected and uninfected litters were exchanged daily. After 5

days, transmission to contact litter was evaluated via plaque assay from retrograde URT

lavages. (C) Index pups are infected IN with 250 PFU of A/X-31, and cohoused with

uninfected littermates (contact) for 5 days prior to evaluating for transmission.

Transmission to pups and mother were evaluated via plaque assay from retrograde

URT lavages. (D) Mother was infected IN with 250 PFU of A/X-31 and placed back with

her uninfected litter. After 5 days, transmission to pups was evaluated via plaque assay

from retrograde URT lavages.

URT titers are represented by a box plot extending from the minimum to maximum

values for each data set. Each symbol represent the titer measured from a single pup

with median values indicated by a line within the box. Index and contact pups are shown

in black and red symbols, respectively. IAV=Influenza A virus, URT=upper respiratory

tract, PFU=plaque forming unit, LOD=Limit of detection.

Figure S3. Timing of IAV shedding corresponds to timing of transmission.

Overlay of shedding data (Fig. 2) (black and red symbols) with transmission data (Fig.

1D). Percent transmission was calculated for each day from raw data in Fig. 1D, and is

indicated by blue symbols connected by a blue line. Area under the curve corresponds

to the proportion of contact pups that have likely acquired IAV infection. IAV=Influenza A

virus, PFU=plaque forming unit, LOD=Limit of detection.

Figure S4. Correlation analyses of influenza virus transmission.

(A) Mean URT titers (black) and mean shedding titers (green) from index pups were

compared to transmission efficiency in contact pups. (B) Index pup shedding titers were

compared among various influenza virus strains. Median values are indicated.

Threshold of transmission is displayed with dashed line, and was calculated using best-

fit linear regression line equation in (A) when transmission X=33.3%. This represents

the minimum level of shedding likely to result in transmission of 1 out of 3 contact pups.

Each symbol represent the shedding titer measured from a single pup. Shedding titers

shown represent at least 2 independent experiments. Transmissible virus and non-

transmissible viruses are shown in green and gray symbols, respectively. (C) Age of

mice (orange) was compared to transmission efficiency in contact mice. (D) Mean URT

titers from index adult mice of 35 days of age (black) and transmission efficiency in

contact adult mice of 35 days of age (blue) were compared to index mice inoculum virus

titer.

Pearson correlation (r) was calculated for each data set. Best fit linear regression

curves were fitted on data sets with adequate correlation and goodness of fit (R^2) was

calculated. Significance of r and R^2 were calculated automatically by Graphpad Prism 7

software, * $p < 0.05$.

Figure S5. Serum IgA levels does not correspond to inhibition of shedding.

As per Fig. 3, serum from mother and pups were obtained at the time of sacrifice.

Samples from individual mice were evaluated for IAV-specific IgA by ELISA. Assay

controls include serum-deficient PBS samples (Neg) and normal mouse serum (N

Serum). IgA geometric mean titers (GMT) are represented by a box plot extending from

the 25[th] to 75[th] percentile for each data set. Whiskers for each box encompasses the

minimum to maximum values. Median values are indicated by a line within the box. All

panels represent at least two independent experiments. PFU=plaque forming unit,

LOD=Limit of detection, GMT=Geometric mean titer.

TABLE 1. Transmissibility of influenza viruses in infant mice

Virus	Index [a]			Contact [b]		Transmission [f]
	No. Infected [c]	URT Titers [d]	Shedding Titers [e]	No. Infected [c]	URT Titers [d]	
A/H3N2						
A/X-31 [g]	8/8	3.04±1.32	2.20±1.02	15/15	4.10±1.05	100
A/Hong Kong/1/1968	4/4	4.85±0.40	2.62±0.67	8/8	5.64±0.35	100
A/X-47 [h]	3/3	3.99±0.25	2.13±0.74	2/9	2.58±2.31	22.2
A/H1N1						
A/Puerto Rico/8/1934	4/4	2.85±0.56	N/A [i]	2/13	2.51±0.44	15.4
A/WSN/1933	6/6	4.18±0.63	1.43±0.64	1/10	N/A [i]	10
A/Brisbane/59/2007	5/5	2.50±1.74	1.63±0.75	1/12	N/A [i]	8.3
A/California/4/2009	5/5	3.77±0.32	1.26±0.42	0/12	N/A [i]	0
B						
B/Lee/1940	5/5	4.44±0.93	2.58±1.31	12/15	3.62±1.67	80

[a] Index pups were infected IN with 250 PFU of virus.

[b] Uninfected contact pups were housed together with infected index pups at the time of inoculation for the duration of the experiment (4-8 days).

[c] Sum of index or contact pups assayed in at least 2 independent experiments.

[d] URT titers, expressed as the mean $\log_{10}$ PFU/mL±STD, were assessed via plaque assay at time of sacrifice from retrograde tracheal lavages for each pup.

[e] Shedding titers, expressed as the mean $\log_{10}$ PFU/mL±STD, were assessed via plaque assay from daily shedding samples collected for each pup.

[f] Percentage of contact pups containing detectable virus in the URT. A/H3N2 viruses were assayed after 4 days. A/PR/8, A/WSN, and B/Lee viruses were assayed after 4 and 8 days. A/Brisbane and A/California were assayed after 8 days.

[g] HA/NA from A/Aichi/2/1968 (H3N2) + genes from A/Puerto Rico/8/1934 (H1N1)

[h] HA/NA from A/Victoria/3/1975 (H3N2) + genes from A/Puerto Rico/8/1934 (H1N1)

[i] Data not applicable for any value representing less than 2 pups

TABLE 2

Mice Age [c]	Index [a]		Contact [b]		Transmission [f]
	No. Infected [d]	URT Titers [e]	No. Infected [d]	URT Titers [e]	
4	3/3	4.17±0.77	4/4	4.66±0.50	100
7	7/7	4.40±1.23	11/11	3.94±1.04	100
14	9/9	3.12±0.95	11/17	3.41±1.00	64.7
21	4/4	3.21±0.78	6/9	2.82±0.78	66.7
Weaned [g]					
28	3/3	3.80±0.86	0/6	N/A [h]	0
35	5/5	3.15±0.52	0/8	N/A [h]	0
56	6/6	2.50±0.71	1/9	N/A [h]	11.1

[a] Index mice were infected IN with 250 PFU of virus.

[b] Uninfected age-matched contact mice were housed together with infected index mice at the time of inoculation for the duration of the experiment (5 days).

[c] Age of mice expressed in days after birth.

[d] Sum of index or contact mice assayed in at least 2 independent experiments.

[e] URT titers, expressed as the mean $\log_{10}$ PFU/mL±STD, were assessed via plaque assay at time of sacrifice from retrograde tracheal lavages for each mice.

[f] Percentage of contact mice containing detectable virus in URT after 5 days of contact.

[g] Mice weaned from breastfeeding and separated from the mother.

[h] Data not applicable for any value representing less than 2 pups