An ER-resident calcium binding protein modulates egress of malaria parasites

from erythrocytes

Manuel A. Fierro[1,2], Miryam A. HortuaTriana[1], Carrie F. Brooks[1], Beejan Asady[1],

Catherine Li[1], Silvia N.J. Moreno[1,2], and Vasant Muralidharan[1,2]

[1]Center for Tropical and Emerging Global Diseases

[2]Department of Cellular Biology

University of Georgia, Athens, GA, USA

Address correspondence to Vasant Muralidharan, vasant@uga.edu

Abstract

Calcium signaling pathways coordinate the lifecycle progression of apicomplexan parasites, specifically playing a key role in egress from host cells. The main Ca^{2+}-signaling hub in these divergent eukaryotic parasites is presumed to be the endoplasmic reticulum (ER); however, no proteins involved in Ca^{2+}-signaling have been identified in the ER of these deadly human parasites. In this study, we explored the role of the Endoplasmic Reticulum-resident Calcium-binding protein (ERC) in the lifecycle of two apicomplexan parasites, *Toxoplasma gondii* and *Plasmodium falciparum*. We find that the *Toxoplasma* ortholog (TgERC) plays a role in the storage of Ca^{2+} in the ER but is dispensable for the asexual lytic cycle. On the other hand, the *Plasmodium* ortholog (PfERC) is essential for asexual growth but not required for Ca^{2+} storage, organelle biogenesis, or protein trafficking. Instead, knockdown of PfERC inhibits parasite egress and invasion. Our results show that PfERC is critical for the rupture of the parasitophorous vacuole membrane, which is the first step in the egress of malaria parasites from erythrocytes. Surprisingly, PfERC knockdown does not affect the Ca^{2+}-dependent autoprocessing of the key egress protease, SUB1, but knockdown inhibits the second proteolytic maturation of SUB1 that occurs during its secretion from exoneme vesicles. We have therefore identified the first ER-resident protein in *Plasmodium* that modulates egress of malaria parasites.

Introduction

Members of the phylum *Apicomplexa* are responsible for severe human diseases such as malaria, toxoplasmosis, and cryptosporidiosis. Together, this group of obligate intracellular parasites causes several hundred million infections every year and remains one of the major drivers of infant mortality (1-4). In fact, malaria results in nearly half a million deaths each year and most of the mortality is attributed to one species, *Plasmodium falciparum*. All the clinical symptoms of malaria are directly correlated to the invasion and lysis of the host red blood cells by these parasites. Malaria remains a vast health and economic burden in susceptible parts of the world and, despite continuous attempts for eradication and control, there is no effective vaccine and resistance has emerged to all existing drugs (5-8). This highlights the importance of understanding the factors that control the intracellular growth and development of the parasite, which will lead to the identification of key targets for antimalarial drug development.

Egress and invasion of apicomplexan parasites are ordered and essential processes which are regulated by signaling pathways dependent upon the second messengers, cGMP and Ca^{2+} (9-13). Upon invading the host cell, the parasites create and reside within a host-derived vacuole called the parasitophorous vacuole (PV). Within this vacuole, the parasites grow and divide into daughter cells, called merozoites in *Plasmodium*, which must egress from the host cell to complete the life cycle. This event is triggered by the activation of a cGMP dependent protein kinase (PKG) and the inhibition of PKG activity blocks egress (10, 14, 15). Ca^{2+}-signaling also induces egress although it is uncertain whether this pathway works downstream (11) or synergistically with cGMP signaling (10, 16, 17). For example, studies have shown that blocking the release of Ca^{2+} from intracellular stores using cell permeable Ca^{2+} chelators blocks egress in malaria parasites (17-19). It has been suggested that this release of Ca^{2+} into the cytoplasm comes from the parasite ER; however, the parasite genome lacks identifiable orthologs of ligand-gated Ca^{2+} channels such as the inositol 1,4,5,-triphosphate or ryanodine receptors (20). Increase in cytoplasmic Ca^{2+} is thought to activate calcium dependent protein kinases (CDPKs) resulting in release of egress-related vesicles (16, 21).

In malaria parasites, these egress-related vesicles contain specific proteases that are proteolytically processed and activated, perhaps upon release into the PV (22-24).

For example, one such pivotal enzyme is the serine protease, Subtilisin 1 (SUB1), which

undergoes two cleavage events. First, the zymogen undergoes Ca^{2+} dependent

autoprocessing in the ER (25, 26) and then, it is cleaved again by the recently identified

aspartic protease, Plasmepsin X (PMX), probably during its secretion into the PV (23, 24).

The proteolytic cleavage and activation of SUB1 is understood to be the point of no-return

as the release of SUB1 into the PV results in the rapid (~10 minutes) breakdown of the

parasitophorous vacuole membrane (PVM) (22, 27). Then, substrates of SUB1 such as

merozoite surface protein 1 (MSP1) and serine-repeat antigen 6 (SERA6) help

breakdown the RBC cytoskeleton and the RBC membrane (RBCM) (28, 29). Once egress

is completed, the merozoites subsequently invade fresh RBCs to start the cycle again.

Like egress, invasion requires specific secretory events such as fusion of micronemes to

the merozoite membrane and secretion of rhoptry contents into the host cell, which

provides the ligand-receptor pair essential for driving the parasite into the host cell (30,

31).

The parasite endoplasmic reticulum (ER) is thought to play a key role in egress

and invasion of daughter merozoites. The putative functions of the parasite ER during

these lifecycle stages, include biogenesis of the specific egress and invasion related

organelles, transporting proteins to these organelles, and propagating Ca^{2+} signals

essential for egress and invasion (32, 33). However, none of the proteins responsible for

these ER-related functions during egress and invasion of apicomplexan parasites have

been identified. We hypothesized that an ER-resident calcium binding protein (ERC) that

is conserved in both *P. falciparum* (PfERC: PF3D7_1108600) and *T. gondii* (TgERC:

TGME49_229480) plays a role in the Ca^{2+} signaling function of the parasite ER. In malaria

parasites, this protein has been shown to bind calcium (34). However, the biological

function of ERC is unknown in either apicomplexan parasite.

To address this, we used a ribozyme-based conditional knockdown system to

study the role of PfERC in the asexual lifecycle *P. falciparum*, and we used a knock-out

approach to study the role of TgERC in the biology of *T. gondii*. We characterized the role

of both Ca^{2+} binding proteins, PfERC and TgERC, and propose a model for their potential

roles in Ca^{2+} signaling and in the life cycle of both parasites. This is the first ER resident

protein identified in *P. falciparum* that is critical for the egress of malaria parasites from
human red blood cells.

Results

<u>PfERC and TgERC are EF-hand containing proteins localized in the ER</u>

PfERC (PF3D7_1108600) and TgERC (TgME49_229480) are proteins related to
the CREC (<u>C</u>alumenin, <u>R</u>eticulocalbin 1, <u>E</u>RC-55, <u>C</u>ab-45) family of proteins, which are
characterized by the presence of multiple EF-hands and localize in various parts of the
secretory pathway (35) (Figure 1A and Supplementary Figure 1). Like CREC members in
other organisms, both PfERC and TgERC contain a signal peptide, multiple EF-hands,
and an ER-retention signal (Figure 1A). However, TgERC differs from its *Plasmodium*
homolog in that it only contains 5 EF-hands while PfERC contains and extended C-
terminus that may contain a 6^{th}, though degenerate, EF-hand (residues 314-325)
(Supplementary Figure 1) (34). Since both proteins have an ER-retention signal they are
predicted to reside within the lumen of the ER.

<u>Generating conditional mutants of PfERC</u>

In order to determine the biological role of PfERC, we used CRISPR/Cas9 gene
editing to generate mutant parasite lines where the endogenous locus of PfERC was
tagged with the conditional knockdown system employing the inducible ribozyme *glmS* or
the inactive version of the ribozyme, *M9* (termed PfERC-*glmS* and PfERC-*M9*
respectively) (Figure 1B and 1C) (36). Western blot and PCR analysis of PfERC-*glmS*
and PfERC-*M9* parasite clones from two independent transfections demonstrate the
correct insertion of the HA tag and the *glmS/M9* ribozymes at the PfERC locus (Figure
1D and 1E). Co-localization of PfERC with an ER marker also confirmed its sub-cellular
localization to this organelle (Figure 1F).

<u>PfERC is essential for asexual growth and for schizont to ring transition</u>

To determine if PfERC was essential for intraerythrocytic survival, we grew
asynchronous PfERC-*glmS* and PfERC-*M9* parasites in the presence of glucosamine
(GlcN), which activates the *glmS* ribozyme leading to mRNA cleavage (Figure 1B).

Western blot analysis shows that this leads to a reproducible reduction of PfERC expression in PfERC-*glmS* parasites while there is no reduction of protein levels in PfERC-*M9* parasites grown under identical conditions (Figure 2A-B). Importantly, this reduction in PfERC levels inhibited the asexual expansion of PfERC-*glmS* parasites, while there was no effect on the growth of PfERC-*M9* parasites (Figure 2C). The response of PfERC-*glmS* parasites was dose-dependent upon GlcN (Supplementary figure 2A)

Since our data show that PfERC was essential for growth within the host RBC, we used synchronous parasites to determine which asexual stage was affected by knockdown. We added GlcN to synchronized schizonts and took blood smears at regular intervals during the intraerythrocytic life cycle to detect morphological development of the asexual stages (Figure 2D). Our data show that all intracellular stages developed normally in both PfERC-*glmS* and PfERC-*M9* parasites grown with GlcN (Figure 2D and Supplementary Figure 2B). However, 55hrs after addition of GlcN, the PfERC-*glmS* parasites remained either as morphologically normal schizonts or were observed as daughter merozoites in the extracellular space as well as some that were attached to RBCs (Figure 2D). On the other hand, PfERC-*M9* parasites were able to egress and re-invade fresh RBCs and developed into ring stage parasites (Figure 2D and Supplementary figure 2B).

These data suggest that the knockdown of PfERC resulted in a defect in the conversion from schizonts to rings. Therefore, we induced knockdown and observed the conversion of schizonts into rings via flow cytometry at 44, 48, and 56 h post-addition of GlcN. The data show that over the course of 12 hours, PfERC-*M9* parasites transitioned from schizonts to rings as determined by the ring:schizont ratio while PfERC-*glmS* parasites were unable to convert from schizonts into rings resulting in a drastically reduced ratio (Figure 2E). These data show that there was a delay in the disappearance of the morphologically normal PfERC-*glmS* schizonts over the final few hours of the asexual life cycle compared to PfERC-*M9* schizonts, suggesting that knockdown of PfERC led to a defect in egress (Figure 2F and 2G). Consequently, the delayed egress may lead to reduced numbers of ring stage parasites in PfERC-*glmS* parasites unlike PfERC-*M9* parasites (Figure 2F and 2H).

 <u>PfERC is required for PVM breakdown</u>

Egress of daughter merozoites from the infected RBC is an ordered and rapid process where the PVM breakdown precedes the disruption of RBCM (Figure 3A). Therefore, we analyzed how egress of PfERC-*glmS* parasites was failing during knockdown. We took synchronized PfERC-*glmS* and PfERC-*M9* schizonts and initiated knockdown with addition of GlcN. These schizonts were allowed to reinvade fresh RBCs and proceed through the asexual stages for 48 hours until they developed into schizonts again. Then, these second cycle schizonts were incubated with inhibitors that block key steps during egress of *P. falciparum* (Figure 3A). To ensure synchronized egress, we used reversible inhibitors of PKG, Compound 1 (C1) or Compound 2 (C2), because inhibition of PKG allows merozoites to develop normally but prevents them from initiating egress (Figure 3A) (10, 15). We used flow cytometry to observe PfERC-*glmS* and PfERC-*M9* schizonts after washing off C1 and saw that there was a delay in the egress of PfERC-*glmS* schizonts while the majority (>60%) of the PfERC-*M9* schizonts were able to complete egress within two hours after washout of C1 (Figure 3B). Removal of C1 initiates the breakdown of the PVM followed by RBCM rupture (Figure 3A), suggesting that PfERC-*glmS* parasites fail to breach one of these membranes down despite removal of the PKG inhibitor.

Therefore, we tested whether PfERC knockdown prevented rupture of PVM or if PfERC functioned during RBCM breakdown (Figure 3A). PfERC-*glmS* and PfERC-*M9* schizonts (where knockdown had been initiated in the previous cycle) were incubated with C2 (10, 15) and observed by scanning electron microscopy (SEM) (Figure 3A and 3C). We observed that parasites treated with C2 were morphologically identical and had developed into mature schizonts within the PVM inside the RBC (Figure 3C). Then, we washed C2 from the parasites and observed these schizonts after 30 mins by SEM (Figure 3C). During this time period, PfERC-*M9* schizonts were able to initiate egress after removal of C2 and we observed free merozoites attached to the RBC as well as clusters of merozoites that had broken out of the PVM but were contained by a collapsed RBCM wrapped around them (Figure 3C). In contrast, the PfERC-*glmS* schizonts were still stuck within the RBC and looked identical to the C2 arrested schizonts, suggesting that they had not initiated egress even though PKG was no longer inhibited (Figure 3C).

These data suggest that knockdown of PfERC blocks egress at an early step, perhaps
blocking the rupture of the PVM (Figure 3C).

We directly observed if breakdown of the PVM was impacted by knockdown of
PfERC using transmission electron microscopy (TEM) (Figure 3D). Knockdown was
induced by adding GlcN to PfERC-*glmS* and PfERC-*M9* schizonts and these parasites
were allowed to go through one asexual cycle and develop into schizonts again 48hrs
later. These schizonts were prevented from completing egress using the irreversible
cysteine protease inhibitor, E64 (Figure 3A). This inhibitor blocks the breakdown of the
RBCM but allows both the breakdown of PVM and poration of the RBCM, which results
in the loss of the electron dense contents of the infected RBC (Figure 3A) (15, 29, 37).
Our results show that the PfERC-*M9* schizonts were able to break down the PVM as well
as proceed with the poration of the RBCM after an 8-hour incubation with E64, while the
PfERC-*glmS* mutants were unable to proceed through the first step of egress and failed
to rupture the PVM (Figure 3D). Overall, these data demonstrate that PfERC function is
critical for the breakdown of the PVM (Figure 3C and 3D).

<u>Generating knockouts of TgERC</u>

We hypothesized that the ortholog of PfERC in the related apicomplexan parasite,
T. gondii, may play a similar in the egress of this parasite from its host cell. Therefore, we
studied the biological role of TgERC by disrupting the transcription of the *tgerc* gene using
CRISPR/Cas9 gene editing to insert the selection marker gene, dihydrofolate reductase
(DHFR), in the *tgerc* locus (38, 39) (Figure 4A). Insertion of this gene into the *tgerc* locus
interfered with the synthesis of the correct transcript and allowed the selection of clonal
populations with pyrimethamine. The mutant *Δtgerc* was isolated by limiting dilution, and
a PCR product of 6.1 kb obtained from the mutant parasites DNA indicated the insertion
of the marker in the predicted locus (Figure 4B). This fragment was only amplified from
DNA of mutants while a fragment of 2.9kb was amplified from parental cells, which
correspond to the intact locus of *tgerc* (Figure 4B).

In order to verify the localization of TgERC, we cloned and expressed TgERC in
E. coli and the purified recombinant protein to immunize mice for the generation of specific
antibodies. Western blot analysis using this antibody showed the presence of a ~38kDa

band corresponding to TgERC in the parental (RH) parasites but absent in the Δ*tgerc*

mutants (Figure 4C). Furthermore, IFAs using this antibody showed that TgERC localized

predominantly to the endoplasmic reticulum in both extracellular and intracellular

tachyzoites (Figure 4D).

TgERC is required for Ca^{2+} storage but PfERC is not

To determine whether TgERC was essential for the lytic cycle of the parasites, we

performed plaque assays comparing growth of RH wild type with Δ*tgerc* (Figure 4E). Our

data show that the mutant parasites were able to form plaques about the same size as

the controls (Figure 4F). These data suggest that unlike PfERC, TgERC was not required

for egress from the host cell nor was it essential for the lytic cycle of *T. gondii*.

We next tested the if TgERC had any role in the storage of Ca^{2+} by testing for

increases in cytoplasmic Ca^{2+} in response to thapsigargin. This inhibits the SERCA-

Ca^{2+} ATPase, exposing leakage of Ca^{2+} from the ER into the cytoplasm as detected by

fluorescence changes in parasites loaded with the Ca^{2+} indicator Fura-2AM (Figure 4G-

J) (40, 41). We observed that the cytoplasmic Ca^{2+} increase in the Δ*tgerc* mutants after

adding thapsigargin was not significantly different than the response obtained under

identical conditions when using wild type parasites (Figure 4G). However, when we

performed the same experiment in the presence of extracellular Ca^{2+} (Figure 4I), the

response to thapsigargin in Δ*tgerc* tachyzoites was significantly reduced. The reduced

response in the Δ*tgerc* parasites indicates that the ER of these parasites is unable to

store Ca^{2+} as efficiently as the wild type. This indicates that TgERC is part of the ER

machinery involved in storing Ca^{2+}. However, it is evident that other mechanisms are in

play since the parasites are still able to grow normally (Figure 4E-F).

Since knockout of TgERC resulted in defects in Ca^{2+} storage in the ER of *T. gondii*

(Figure 4I and 4J)*,* we hypothesized that PfERC is required for egress because it plays a

similar role in *P. falciparum* and does not have redundant mechanisms like *T. gondii* that

compensate for the lack of TgERC. To test this hypothesis, synchronized PfERC-*glmS*

and PfERC-*M9* schizonts were incubated with GlcN and allowed to proceed through one

asexual cycle until they formed schizonts again. The second cycle schizonts were isolated

using saponin lysis and loaded with Fluo-4AM to measure cytosolic Ca^{2+} (Supplementary

Figure 3A). To assess if the storage of Ca^{2+} in the ER of the parasite was affected by

knockdown of PfERC, we added the SERCA-Ca^{2+} ATPase inhibitor Cyclopiazoic acid

(CPA) to these saponin-isolated parasites (Supplementary Figure 3A and 3B) (42).

Inhibiting the SERCA-Ca^{2+} ATPase allows Ca^{2+} stored in the ER to leak into the

cytoplasm, which results in a detectable change in the fluorescence of Fluo-4AM

(Supplementary Figure 3B). Our measurements show that there was no difference in the

amount of Ca^{2+} that leaked from the parasite ER, upon SERCA-Ca^{2+} ATPase inhibition,

between PfERC-*glmS* and PfERC-*M9* schizonts (Supplementary Figure 3B).

To test if there was a defect in Ca^{2+} storage in neutral stores, we used the

ionophore, Ionomycin, which releases Ca^{2+} from all neutral stores in the cell and

measured the release of Ca^{2+} into the cytoplasm of PfERC-*glmS* and PfERC-*M9*

schizonts. The parasites were isolated as described above and the changes in

cytoplasmic Ca^{2+} were measured using Fluo-4AM (Supplementary Figure 3A and 3C).

Again, we did not observe any difference in the amount of Ca^{2+} released into the

cytoplasm of PfERC-*glmS* and PfERC-*M9* schizonts treated with ionomycin

(Supplementary Figure 3C). These data suggest that the availability of free Ca^{2+} in the

ER of *P. falciparum* is not affected by knockdown of PfERC. Furthermore, these data

suggest that the observed egress defect upon PfERC knockdown was not a result of

disequilibrium of Ca^{2+} in the parasite ER.

<u>Efficient SUB1 activation requires PfERC</u>

Electron microscopy data show that knockdown of PfERC prevents the breakdown

of the PVM (Figure 3). A key event required for PVM breakdown is the proteolytic

processing of SUB1, which starts a cascade that ends in the release of merozoites from

the infected RBC (22, 29). Therefore, we tested if knockdown of PfERC affects processing

of PfSUB1. This protease is processed twice, once in the ER, where it undergoes a Ca^{2+}-

dependent autocatalytic processing from its zymogen form (83-kDa) into a 54-kDa semi-

proenzyme form (p54) (25, 26, 43). From the ER, SUB1 is transported to the egress-

related secretory vesicles, the exonemes, which are secreted into the PV to initiate

breakdown of the PV membrane. It is proposed that during secretion of SUB1, it is

processed by PMX from its semi-proenzyme form (p54) to its mature form (p47) (23, 24).

The secretion of mature p47 form of SUB1 initiates the breakdown of the PVM (22, 43).

We hypothesized that PfERC is required for the first Ca^{2+}-dependent autoprocessing of

SUB1, in the ER, from the 82-kDa zymogen into the p54 semi-proenzyme.

To test this hypothesis, PfERC-*glmS* and PfERC-*M9* schizonts were incubated with

GlcN and allowed to progress through one asexual growth cycle (48 hours) to develop

into schizonts again. Lysates from these synchronized schizonts were separated on a

Western blot and probed with antibodies against SUB1 (Figure 5A and Supplementary

Figure 4A). No change was observed in the Ca^{2+}-dependent autoprocessing of the

zymogen-form of SUB1 into the semi-proenzyme (p54) form (Figure 5A and

Supplementary Figure 4A). Surprisingly, we observed a reproducible and significant

decrease in the processing of SUB1 from p54 to the p47 form in PfERC-*glmS* parasites

(Figure 5A and 5B). Compared to PfERC-*M9* parasites, there was a >50% decrease in

the amount of processed SUB1 (p47) in PfERC-*glmS* parasites (Figure 5B). These data

suggest that PfERC is required for the secretion-dependent processing of SUB1 in

exoneme vesicles.

Since we observed the presence of some mature SUB1 in PfERC-*glmS* parasites

(Figure 5A), we tested if the activity of SUB1 was affected upon knockdown of PfERC by

assaying for the processing of a known substrate of SUB1, the merozoite surface protein

1 (MSP1). MSP1 is required for the initial attachment of merozoites onto RBCs and it has

been shown that correct processing of MSP1, by SUB1, is also required for efficient

egress as it plays a role in breakdown of the RBC cytoskeleton after release from the

PVM (28, 44-46). Lysates from synchronized second-cycle PfERC-*glmS* and PfERC-*M9*

schizonts, treated as above, were separated on a Western blot and probed using anti-

MSP1 antibodies (Figure 5C and Supplementary Figure 4B). Our data show that the there

was a ~60% reduction in MSP1 processing in our PfERC-*glmS* parasites as compared to

PfERC-*M9* parasites after knockdown (Figure 5C). In fact, we observed that PfERC-*glmS*

schizonts had 3 times more unprocessed MSP1 in contrast to PfERC-*M9* schizonts

(Figure 5D). These data reveal that knockdown of PfERC leads to defects in SUB1

processing and activity, and consequently, MSP1 processing (Figure 5).

<u>Knockdown of PfERC does not affect protein trafficking or organelle biogenesis</u>

MSP1 is a GPI-anchored merozoite membrane protein that is presumably
processed by SUB1 once the protease is secreted into the PV (28, 29). Therefore, we
wanted to verify that knockdown of PfERC led to a specific defect in the egress cascade
and is not due to a block in protein trafficking via the ER or defects in the biogenesis of
egress and invasion related organelles. To address this, we used super resolution
structured illumination microscopy (SR-SIM) to observe if there was a difference in the
surface expression of MSP1 between PfERC-*glmS* and PfERC-*M9* schizonts upon
knockdown of PfERC (Figure 5E). As before, knockdown was initiated in synchronized
PfERC-*glmS* and PfERC-*M9* schizonts and after 48 hours, these schizonts were stained
with anti-MSP1 antibodies. Our data shows that there was no difference in the trafficking
of MSP1 to the surface of developing PfERC-*glmS* or PfERC-*M9* merozoites after
knockdown (Figure 5E). In addition, we also tested whether the localization of other
egress-and invasion-related proteins that traffic through the ER where affected after
knockdown by staining similarly treated schizonts for the micronemal protein AMA1 and
the rhoptry-bulb protein RAP1. Our data show that, again, there was no difference in the
localization of these proteins in schizonts between PfERC-*glmS* and PfERC-*M9* parasites
suggesting that the knockdown of PfERC does not cause a generalized defect in the
secretory pathway (Figure 5F&G).

As the ER generates the lipid membranes required for generating organelles
essential for egress and invasion, we wanted to see if their biogenesis was inhibited upon
knockdown. If true, this could explain the processing defects that we observe for SUB1
and MSP1. We observed E64-treated schizonts (Figure 3A) by transmission electron
microscopy and these data show that both micronemes and rhoptries remain
morphologically intact in both PfERC-*glmS* and PfERC-*M9* parasites after knockdown
was induced (Figure 5H). Successful knockdown was confirmed by the lack of PVM
membrane fragments in PfERC-*glmS* parasites, which are present in the PfERC-*M9*
parasites, suggesting that PVM rupture occurred normally in the presence of PfERC but
not when the protein is knocked down (Figure 5H). These data suggest that the
knockdown of PfERC does not lead to defects in organelle biogenesis (Figure 3D and
Figure 5H).

<u>PfERC is required for invasion of merozoites</u>

The synchronized growth assays suggest that knockdown of PfERC inhibits the

invasion of merozoites into RBCs (Figure 2D, H and Figure 5C, D). To assess if invasion

was reduced upon knockdown, PfERC-*glmS* and PfERC-*M9* schizonts in the second

cycle after 48 hours in GlcN, were incubated with the PKG inhibitor, C1, for four hours

(Figure 3A). The inhibitor was then washed off and the formation of ring stages was

observed over two hours by flow cytometry (Figure 6A). We observed that there was a

delay in the formation of ring stages as well as a drastic decrease in the numbers of ring

stage parasites formed in PfERC-*glmS* parasites compared to the PfERC-*M9* control

(Figure 6A). This could be due to loss of egress or could be a combination of effects on

egress and invasion due to PfERC knockdown.

To decouple the egress and invasion phenotypes, we directly measured the

efficiency of merozoite invasion (Figure 6B). This was accomplished by incubating second

cycle PfERC-*glmS* and PfERC-*M9* schizonts with E-64 and then mechanically releasing

the merozoites (Figure 6B) (47). Incubation with E-64 for 8 hours allows for the completion

of schizogony and formation of invasion-competent merozoites. These purified

merozoites were then allowed to invade fresh RBCs and the invasion rate was quantified

using flow cytometry as described previously (Figure 6C) (47). These data show that there

was a drastic reduction in the invasion efficiency of PfERC-*glmS* merozoites as compared

to the control PfERC-*M9* merozoites, thus demonstrating that knockdown of PfERC led

to a defect in invasion as well (Figure 6C).

<u>PfERC knockdown affects proteolytic processing of AMA1</u>

The reduced invasion of PfERC-*glmS* merozoites could be explained by the

reduction in processing of MSP1, which is known to be required for the initial attachment

of merozoites to the RBC (44, 45, 48). Invasion of RBCs by *P. falciparum* merozoites

requires secretion of contents from another apical organelle, the rhoptries, into the RBC

(49-51). Proteins in the rhoptries, like the rhotry-bulb protein, RAP1, require proteolytic

processing for their activity (23, 24). Once in the rhoptry, RAP1 is processed by the

aspartic protease, Plasmepsin IX (PMIX), from a pro-form (p83) to a mature form (p67)

(23, 24, 52, 53). Therefore, we tested if RAP1 processing was affected by knockdown of

PfERC using Western blot analysis (Figure 6F and Supplementary Figure 4C). Our data

show that the proteolytic processing of RAP1 was not affected by the knockdown of

PfERC (Figure 6G), showing that knockdown does not lead to a generalized defect in the

processing of all proteins that traverse through the secretory pathway.

Another key and essential step in invasion of merozoites is the formation of a tight

junction between the parasite and the RBC and AMA1 is critical for the formation of this

tight junction (30, 31). AMA1 is trafficked from micronemes to the merozoite surface and

there it is processed from its pro-form (p83) to its mature form (p66) by an unknown

protease (54-56). The proteolytic processing of AMA1 is critical for its function during

invasion (57). Therefore, we tested if the processing of the AMA1 was affected upon

knockdown of PfERC. As before, after initiating knockdown in synchronized schizonts,

we separated lysates from second cycle PfERC-*glmS* and PfERC-*M9* schizonts on a

Western blot and probed it with anti-AMA1 antibodies (Figure 6D and Supplementary

Figure 4D). We observed a significant reduction in the proteolytic processing of AMA1

upon knockdown (Figure 6D). Indeed, there was a >40% decrease in the processing of

AMA1 in PfERC-*glmS* mutants compared to the PfERC-*M9* control (Figure 6E). These

data suggest that PfERC is required for the correct processing of AMA1 and therefore,

essential for invasion of merozoites into the host RBC.

Discussion

In this study, we addressed the biological role of a conserved Ca^{2+}-binding protein

that resides in the lumen of the ER of apicomplexan parasites. We have shown that the

Toxoplasma homolog, TgERC, is dispensable for *in vitro* growth, and the *Plasmodium*

homolog, PfERC, is essential for asexual replication (Figures 2 and 3). Our data show

that knockdown of PfERC did not affect the ring and trophozoite development but had a

clear effect on the subsequent schizont-to-ring transition (Figures 2D-H, 3B, 6A, and

Supplementary Figure 2). Specifically, these data show that PfERC is required for both

egress from infected RBCs (Figures 2D, 2G, and 3) and invasion into host erythrocytes

(Figures 2D, 2H and 6A-C). This is consistent with data that suggest PfERC may be

transcriptionally controlled by the invasion-specific transcription factor PfAP2-I (58). Our

data show that knockdown of PfERC leads to defects in the processing of proteins critical

for invasion of merozoites into the host RBC, namely, MSP1 and AMA1 (Figures 5C-D
and 6 D-E). However, the observed invasion defect is likely a secondary effect because
several proteins critical for invasion are processed during egress (28, 29). Given the
limitations of the conditional knockdown system, we cannot tease out a specific role for
PfERC in invasion. As invasion occurs rapidly (<2mins), a potential specific invasion-
related function of PfERC could be tested using a small molecule that specifically targets
PfERC (59). Overall, these data show that PfERC is essential for egress of merozoites
from the infected RBC and for invasion of merozoites into the host erythrocyte.

It has been observed previously that the mechanisms at play during egress of
daughter parasites from the host cells differs between *Toxoplasma* and *Plasmodium.* This
is highlighted by studies showing that orthologs of apicomplexan proteins can serve
different functions in their respective organisms. For example, TgCDPK3 is required for
egress of *T. gondii* tachyzoites while its *Plasmodium* ortholog, PfCDPK1, is essential for
both egress and invasion (60). Similarly, another calcium dependent protein kinase,
TgCDPK1, plays a vital role in egress and invasion of *Toxoplasma* tachyzoites but the
Plasmodium ortholog, PfCDPK4, was shown to be required for the development of sexual
stages (60). Additionally, TgSUB1 localizes to micronemes and is used for processing of
micronemal proteins that are required for invasion of *T. gondii* into host cells (61).
However, PfSUB1 is critical for egress of *Plasmodium* merozoites, localizes to exonemes,
and upon secretion into the PV, processes both micronemal and non-micronemal proteins
(22, 62). These differences could be related to their development within different host
cells. *Plasmodium* prefers the erythrocyte for its asexual expansion and *Toxoplasma* is
able to make its home within any nucleated cell. Another factor is the difference in the
primary structure of TgERC and PfERC, which possesses an additional putative EF-hand
domain compared to its homolog in *T. gondii.* Therefore, it is not surprising that the
biological role of TgERC may be different from the function of PfERC. This was evident
in the data that suggest that TgERC plays a role in ER Ca^{2+} storage but PfERC may not
(Figures 4I, 4J and Supplementary Figure 3). It is interesting to note that there are no
orthologues of many Ca^{2+} entry channels such as store-operated channels (ORAI), or
their partners such as the ER sensor protein stromal interaction molecule (STIM) in the
genome of apicomplexan parasites. Future work on identifying the interacting partners of

TgERC may provide some clues as to the identity of putative ER Ca^{2+} sensors that could
regulate the function Ca^{2+} entry channels in these divergent parasites.

During the formation of daughter merozoites in schizonts, several key egress and
invasion related organelles essential for propagation of the infection are generated. The
ER is thought to play a key role in the biogenesis of these organelles and the ER is
responsible for transporting the essential proteins to these organelles (32, 33). As an ER-
resident protein, knockdown of PfERC could affect several ER functions such as protein
trafficking, organellar biogenesis, and Ca^{2+} signaling. Therefore, we tested if PfERC
functions in the trafficking of proteins required for schizont to ring transition such as MSP1,
AMA1, and RAP1. A defect in the secretory pathway would explain the observed
deficiencies in the proteolytic processing of SUB1, MSP1 and AMA1, as transport out of
the ER is required for their maturation (10, 26, 52, 63). However, super-resolution and
electron microscopy experiments showed that trafficking of proteins to the merozoite
surface, micronemes, and rhoptries or their biogenesis are not affected (Figures 3D and
5E-5H). Likewise, Western blot analysis showed that the proteolytic processing of a
rhoptry protein, RAP1, which is processed after transport to the organelle, is not affected
by knockdown of PfERC (Figures 5G, and 6F-G) (23, 24). These data show that
knockdown of PfERC does not result in a generalized defect in protein trafficking via the
ER or in organelle biogenesis.

Immunofluorescence microscopy, electron microscopy, and Ca^{2+} measurements
ruled out a role for PfERC in Ca^{2+} storage, protein trafficking, or organelle biogenesis
(Figures 3D, 5E-G, 6F, 6G and Supplementary Figure 3). Another interesting possibility
is that PfERC may play a role the signal-dependent release of Ca^{2+} from the ER. PfERC
is a member of the CREC family of proteins and other members of this family are known
to play key roles in the function of ligand-gated Ca^{2+} channels. For example, calumenin
and reticulocalbin 1 have been shown to interact with the Ryanodine and IP_3 receptors,
respectively, and regulate their function (64, 65). It has also been suggested that some
CREC members serve as Ca^{2+} sensors in the ER as they exhibit calcium-dependent
structural changes (66, 67). This is difficult to test in *Plasmodium* since there are no clear
homologs for a ligand-dependent Ca^{2+} channel in its genome (20). At present, Ca^{2+}
measurements in *P. falciparum* require lysis of the RBCM that results in loss of the RBC

contents and this complicates any conclusions reached from egress-related experiments
since it is unknown if components in the RBC are essential for the signaling pathway
required for egress. However, this may be addressed by using less invasive approaches
that do not require lysis of the RBCM such as genetically-encoded Ca^{2+} indicators (GECI)
to measure changes in ER Ca^{2+} during egress or by studying the egress-related roles of
proteins that interact with PfERC.

Consistent with a role for PfERC in signal-dependent release of Ca^{2+} from the ER,
we observed that knockdown resulted in a clear defect in egress (Figures 2G, 3, and 5A-
D). Egress is a rapid process and the genetic mechanisms in the ER of apicomplexan
parasites that control the signal-dependent release of Ca^{2+} from the ER are unknown.
Our data supports a model where PfERC functions in the signaling pathway that releases
Ca^{2+} from the ER required for the egress of merozoites from the infected RBC. This model
would also predict that PfERC plays a role in the release of Ca^{2+} from the ER, such that
in the absence of PfERC, the ER is unable to release Ca^{2+} upon the requisite signal. Such
a putative role for PfERC is supported by data which show that knockdown of PfERC
inhibits egress by preventing the breakdown of the PVM (Figure 3C-D).

A key enzyme that is required for initiating egress is the protease SUB1. Another
hypothesis we tested was if PfERC was required for the Ca^{2+} dependent activity of SUB1.
This serine proteases plays a critical role in disruption of both the PVM and RBCM. It is
produced as an 82-kDa zymogen in the ER, where it rapidly self-processes into a 54-kDa
semi-proenzyme that remains associated with the cleaved prodomain, thus inhibiting its
activity (43). The autocatalytic activity of SUB1 is Ca^{2+} dependent and therefore, it was
possible that PfERC was required for this autoprocessing event by providing the required
Ca^{2+} to SUB1. This would explain the phenotypes observed during knockdown of PfERC
as they are similar to that seen when SUB1 is conditionally knocked out (29). However,
our data show only the second processing step of SUB1 that produces the mature, active
form of the protease (p54 to p47), is affected (Figure 5A-B and Supplementary Figure
4A). There is no effect on the Ca^{2+} dependent autoprocessing of SUB1 (p83 to p54) that
occurs in the ER (Figure 5A-B and Supplementary Figure 4A). The processing of SUB1
from p54 to p47 occurs once it is trafficked out of the ER and putatively during secretion
of exoneme vesicles (25). This processing defect reduces the activity of SUB1 as it is

defective in processing its substrate, MSP1 (Figure 5A-D). This processing event

happens after the secretion of SUB1 into the PV as MSP1 is a GPI-anchored protein on

the membrane of merozoites (28). These data suggest that PfERC is required for

secretion of exonemes, which is consistent with a role for PfERC in signal-dependent

release of Ca^{2+} from the ER.

A principal finding of these studies is the identification of the first protein in the ER

of *P. falciparum* with a specific role in egress of malaria parasites from RBCs and

potentially in the invasion of parasites into the RBC (Figures 2, 3, 5 and 6). These studies

lay the foundation for understanding the vital and key role that ER-resident proteins play

in the egress of human malaria parasites from their infected RBC and their re-entry into

the host cell. Some studies have suggested that a key class of antimalarials containing

endoperoxides, which includes the frontline antimalarial artemisinin, may target PfERC

(59) and one of the transcriptomic responses of artemisinin-resistant parasites is the

overexpression of PfERC (68). These data suggest that targeting PfERC, and thus

egress, is a viable strategy for antimalarial drug development.

Material and Methods

Cell culture and transfections. *Plasmodium* parasites were cultured in RPMI 1640

medium supplemented with Albumax I (Gibco) and transfected as described earlier (69-

72). *T. gondii* tachyzoites were maintained in vitro by serial passage in Dulbecco's

modified minimal essential media (DMEM) with 1% FBS, 2.5 µg/ml amphotericin B, and

100 µg/ml streptomycin. Host cells were human telomerase reverse transcriptase

immortalized foreskin fibroblasts (hTERT) (73).

To generate PfERC-*glmS* and PfERC-*M9* parasites, a mix of two plasmids (50µg

of each) was transfected in duplicate into 3D7 parasites. The plasmid mix contained

pUF1-Cas9-guide (74) which contains the DHOD resistance gene, and pPfERC-HA-

SDEL-*glmS* or pPfERC-HA-SDEL-*M9*, which are marker-free. Drug pressure was applied

48hrs after transfection, using 1µM (DSM1) (75), selecting only for Cas9 expression.

DSM1 was then removed from the culturing medium once the parasites were detected in

the culture, around 3 weeks post-transfection.

The *Δtgerc* knock-out line was generated by inserting the dihydrofolate reductase
(DHFR) gene into the first ~300 nucleotides of the *tgerc* gene to disrupt the coding
sequence. The CRISPR/Cas9 gene knockout method previously reported was used (39).
A fragment corresponding to the single guide RNA (sgRNA) using primers 1+2 was
cloned into pSAG1:CAS9::U6:sgUPRT (Addgene #54467) using the Q5 site-directed
mutagenesis kit (NEB). The resulting pSAG1:CAS9::U6:sgTgERC was co-transfected
with the DHFR cassette (in proportion 3:1) using primers 3+4 to tachyzoites of the RH
strain. After pyrimethamine selection (1 µM), parasites were sub-cloned by limiting dilution
and screened for positive clones by PCR using primers 5+6 shown in Table S1.

Construction of PfERC plasmids. Genomic DNA was isolated from *P. falciparum*
cultures using the QIAamp DNA blood kit (Qiagen). Constructs utilized in this study were
confirmed by sequencing. PCR products were inserted into the respective plasmids using
the In-Fusion cloning system (Clontech), the sequence- and ligation-independent cloning
(SLIC) method (71, 72), T4-ligation (New England BioLabs), or site-directed mutagenesis
using QuickChange (Agilent). To generate the pHA-SDEL-*glmS*/*M9* plasmid, primers 7+8
were used to add an SDEL sequence at the end of the HA tag in pHA-*glmS* and pHA-*M9*
plasmids (71, 72).

For the generation of the PfERC-*glmS*/*M9* conditional mutants, pHA-SDEL-
glmS/*M9* plasmid was used as a donor DNA template consisting of two homology regions
flanking the HA-SDEL tag and the *glmS* or *M9* sequence. To allow efficient genomic
integration of the pHA-SDEL-*glmS* and pHA-SDEL-*M9* donor plasmids, 800-bp
sequences were used for each homology region. The C-terminus of the *pferc* coding
region was PCR amplified from genomic DNA using primers 9+10 (containing the shield
mutation) and was inserted into pHA-SDEL-*glmS* and pHA-SDEL-*M9* using restriction
sites SacII and AfeI. The 3'UTR of *pferc* was PCR amplified from genomic DNA using
primers 11+12 and was inserted into pHA-SDEL-*glmS* and pHA-SDEL-*M9* (already
containing the C-terminus region) using restriction sites HindIII and NheI. For expression
of PfERC guide RNA, oligos 13+14 were inserted into pUF1-Cas9-guide as previously
described (74, 76). Briefly, pUF1-Cas9-guide was digested with BtgZI and annealed

oligos were inserted using the SLIC method. Primers 9+12 and primers 9+15 (which
recognizes the *glmS*/*M9* sequence) were used for clone verification.

***Plasmodium* growth assays**. Asynchronous growth assays were done as described
previously (71, 72). Briefly, 5mM glucosamine (GlcN) (Sigma) was added to the growth
medium and parasitemia was monitored every 24hrs via flow cytometry using a CyAn
ADP (Beckman Coulter) or CytoFLEX (Beckman Coulter) instrument and analyzed by
FlowJo software (Treestar, Inc.). As required, parasites were subcultured to avoid high
parasite density, and relative parasitemia at each time point was back-calculated based
on actual parasitemia multiplied by the relevant dilution factors. One hundred percent
parasitemia was determined as the highest relative parasitemia and was used to
normalize parasite growth. Data were fit to exponential growth equations using Prism
(GraphPad Software, Inc.).

To determine the ring:schizont ratio of PfERC-*glmS* and PfERC-*M9* parasites,
7.5mM GlcN was added to percoll isolated schizont-stage parasites and parasites were
allowed to egress and reinvade fresh RBCs. Two hours later, 5% sorbitol +7.5mM GlcN
was added to the invaded culture to lyse any remaining schizonts and isolate two-hour
rings. The ring-stage parasites were grown again in media supplemented with GlcN. Then
samples were taken at 44hrs, 48hrs, and 56hrs, and read by flow cytometry to determine
the population of rings and schizonts present at those times using FlowJo software
(Treestar, Inc.). To determine the development of each life cycle stage during the asexual
lifecycle of PfERC-*glmS* and PfERC-*M9* parasites, 7.5mM was added to percoll isolated
schizont-stage parasites and parasites were allowed to egress and reinvade fresh RBCs.
Time points were then taken by blood smears and the percentage of the specific lifecycle
stage was calculated as: $\% \ of \ stage = \frac{number \ of \ specific \ stage}{total \ number \ of \ parasites}$. Time 0hr is when GlcN was
added. To determine the % amount of egressing or invading parasites, we took the %
values of schizonts or rings at 44hrs and subtracted that number from % values of
schizonts or rings at 12hrs later.

For growth assays using Compound 1, synchronized schizonts were grown with
7.5mM GlcN for about 48hrs. Then, schizonts were percoll isolated and incubated with
Compound 1 for 4hrs and then removed by gently washing parasites twice with 1mL of

warm, complete RPMI. Parasites were resuspended with fresh media and RBCs and

samples were taken and read by flow cytometry. DNA content was determined using

Hoechst staining (ThermoFisher).

Plaque assays. Plaque assays were done as previously described (77). Two hundred

freshly egressed tachyzoites were used to infect a confluent monolayer of hTERT

fibroblasts followed by 7 days of growth. Monolayers were fixed and stained with 5x

crystal violet. Plaque sizes were analyzed with FIJI by measuring the area of fifteen

plaques per biological replicate. At least 3 biological experiments were done for wild type

control RH and TgERC knockout.

Generation of TgERC antibody. The *tgerc* gene (TgGT1_229480) was identified from

ToxoDB. Total RNA of the wild type strain (RH) was extracted from freshly lysed parasites

using Trizol® reagent (Sigma) following the manufacturer's instructions. The RNA sample

was further treated with DNAse I for 10 min at 37 °C (New England Biolabs) to remove

contaminating DNA. Four micrograms of purified mRNA were used to synthesize cDNA

using the superscript III first-strand synthesis system according to the manufacturer's

protocol (ThermoFisher Scientific-Life Technologies). *T. gondii* cDNA was used as a

template for amplifying the *tgerc* gene (~ 1 Kb) with the primers 16+17 (Table S1). The

PCR product was gel purified and cloned using the pET32 Ek/LIC vector system

(Novagen). The construct was sequenced and transformed into *E. coli* BL21(DE3)-

CodonPlus. Expression of the TgERC protein was induced for 3 h at 37°C with 0.4 mM

isopropyl-β-D-thiogalactopyranoside (IPTG). Cells were pelleted and resuspended in

equilibration/binding buffer (50 mM Na_3PO4, 300 mM NaCl, 10 mM Imidazole, and 1:500

protease inhibitor cocktail P-8849). Cells were then sonicated on ice for 80 seconds total

with amplitude set to 30% (Branson Digital Sonifier Model 250) and centrifuged at 12,000

rpm at 4°C for 20 min. The supernatant was filtered through a 0.45 μm cellulose acetate

filter (VWR) and the protein was purified using HisPur Ni-NTA chromatography following

instructions from the manufacturer. Unbound proteins were washed off with a total volume

of 12 ml of wash buffer (50 mM Na_3PO_4, 300 mM NaCl and 20 mM Imidazole) and the

TgERC recombinant protein was eluted with 5 ml of elution buffer (50 mM Na_3PO_4 ,300

mM NaCl and 250 mM Imidizole). Elution fractions were verified by SDS-PAGE.

Antibodies against the TgERC recombinant protein were generated in mice. Six

CD-1 mice (Charles River, Inc.) were intraperitoneally injected with 100 µg of purified

TgERC recombinant protein mixed 1:1 with complete Freund adjuvant. Mice were

boosted twice with 50 µg of antigen and the final serum was collected by cardiac puncture

after CO_2 euthanasia (IACUC protocol A2018 02-021-Y1-A0)

Western blotting. Western blotting for *Plasmodium* parasites was performed as

described previously (71, 72). Briefly, parasites were permeabilized selectively by

treatment with ice-cold 0.04% saponin in PBS for 10 min and pellets were collected for

detection of proteins with the parasite. For detection of MSP1, schizonts were isolated on

a Percoll gradient (Genesee Scientific) and whole-cell lysates were generated by

sonication. The antibodies used in this study were rat anti-HA (3F10; Roche, 1:3,000),

rabbit anti-HA (715500; Invitrogen, 1:100), rabbit anti-PfEF1α (from D. Goldberg,

1:2,000), mouse anti-plasmepsin V (from D. Goldberg, 1:400), rabbit anti-SUB1 (from Z.

Dou and M. Blackman, 1:10,000), rat anti-AMA1 (28G2; Alan Thomas via BEI Resources,

NIAID, NIH 1:500), mouse anti-MSP1 (12.4; European Malaria Reagent Repository,

1:500) and mouse anti-RAP1 (2.29; European Malaria Reagent Repository, 1:500). The

secondary antibodies that were used are IRDye 680CW goat anti-rabbit IgG and IRDye

800CW goat anti-mouse IgG (LICOR Biosciences) (1:20,000). The Western blot images

were processed using the Odyssey Clx LICOR infrared imaging system software (LICOR

Biosciences). Calculation of knockdown and processing ratios was determined by both

the Odyssey infrared imaging system software and ImageJ 1.8 (NIH).

For *T. gondii*, lysates were prepared by resuspending a pellet of 1 x 10^8 tachyzoites

in 50 µl of Cell Lytic^M lysis buffer containing 12.5 U benzonase and 1X protease cocktail

inhibitor (P8340 Sigma). The reaction was stopped with one volume of 2% SDS and 1

mM EDTA. Total lysates were boiled in Laemmli sample buffer (BioRad) (78).

Immunoblots were done following established protocols using polyclonal mouse anti-

TgERC serum (1:1,000). The Odyssey Clx LICOR system was used for detection, and

the secondary antibody was goat anti-mouse IRDye 800WC (1:10,000).

Immunofluorescence microscopy. For IFAs, cells were fixed as described previously (71, 72). The antibodies used for IFA were: rat anti-HA antibody (clone 3F10; Roche, 1:100), rat anti-PfGRP78 (MRA-1247; BEI Resources, NIAID, NIH 1:100), mouse anti-MSP1 (12.4; European Malaria Reagent Repository, 1:500), rat anti-AMA1 (28G2; Alan Thomas via BEI Resources, NIAID, NIH 1:500), and mouse anti-RAP1 (2.29; European Malaria Reagent Repository, 1:500). Secondary antibodies used were anti-rat antibody conjugated to Alexa Fluor 488 or 546 and anti-rabbit antibody conjugated to Alexa Fluor 488, (Life Technologies, 1:100). Cells were mounted on ProLong diamond with 4',6'-diamidino-2-phenylindole (DAPI) (Invitrogen) and imaged using a Delta-Vision II microscope system with an Olympus IX-71 inverted microscope using a 100x objective or an Elyra S1 SR-SIM microscope (Zeiss, Jena, Germany). Image processing, analysis, and display were performed using SoftWorx or Axiovision and Adobe Photoshop. Adjustments to brightness and contrast were made for display purposes.

For *T. gondii* IFAs, extracellular parasites were collected and purified as previously described (77). Parasites were washed once with buffer A with glucose (BAG, 116 mM NaCl, 5.4 mM KCl, 0.8 mM MgSO₄, 5.5 mM glucose and 50 mM HEPES, pH 7.4) and an aliquot with 2 x 10⁴ parasites was overlaid on a coverslip previously treated with poly-L-Lysine. Intracellular tachyzoites were grown on hTERT cells on coverslips previously infected with freshly lysed parasites. Both extracellular and intracellular parasites preparations were fixed with 2.5% paraformaldehyde for 20 min at room temperature (RT), permeabilized with 0.25% Triton X-100 and blocked with 3% bovine serum albumin (BSA). For co-localization studies parasites were exposed to primary antibodies mouse anti-TgERC 1:1000 and rabbit anti-GFP 1:800 after transfection with the P30-GFP-HDEL plasmid (79). The secondary antibodies were Alexa-Fluor488 goat-anti mouse and Alexa-Flour594 goat-anti rabbit at 1:1,000. Slides were examined using an Olympus IX-71 inverted fluorescence microscope with a photometric CoolSNAP HQ charge-coupled device (CCD) camera driven by DeltaVision software (Applied Precision).

Invasion Rate Quantification. To calculate the invasion rate, parasites were treated as described previously (47). Briefly, 7.5mM GlcN was added to percoll isolated schizont-

stage parasites and parasites were allowed to egress and reinvade fresh RBCs. 48hrs

later, schizonts were percoll purified and incubated with 20µM E-64 (Sigma) at 37°C in

an incubator for 7-8hrs. Once incubation was done, merozoites were isolated by gently

resuspending and passing the schizonts through a 1.2µm Acrodisic Syringe Filter (PALL).

Merozoites were spun at 2000xg for 5min, and then resuspended in 100µL of complete

RMPI medium and added to a 1mL culture of uninfected RBCs at 2% hematocrit. Cultures

were grown in a FluoroDish cell culture dish (World Precision Instruments) and gassed in

a chamber for 20-24hrs. Invasion rate was then measured by the following equation: $IR =$

$iRBC \cdot \left[\frac{RBC/\mu L}{Mz/\mu L} \right]$ where "*iRBC*" is the parasitemia 20-24hrs later, "*RBC/µL*" are the free

RBCs used before addition of merozoites and "*Mz/µL*" are the merozoites found in the

100µL suspension used before adding to fresh RBCs. Values for these variables were

acquired by flow cytometry (CytoFLEX Beckman Coulter) with cells stained with acridine

orange. The data were normalized using the IR values for PfERC-*M9* merozoites as

100%.

Transmission Electron Microscopy. 7.5mM GlcN was added to percoll isolated

schizont-stage parasites and parasites were allowed to egress and reinvade fresh RBCs.

48hrs later, parasites were percoll-isolated and then incubated with 20µM E-64 for 8hrs.

After incubation, parasites were washed with 1X PBS and gently resuspended in 2.5%

glutaraldehyde in 0.1M sodium cacodylate-HCl (Sigma) buffer pH 7.2 for 1hr at room

temperature. Parasites were then rinsed in 0.1M Cacodylate-HCl buffer before agar-

enrobing the cells in 3% Noble agar. Parasites were post fixed in 1% osmium

tetroxide/0.1M Cacodylate-HCl buffer for 1 hour and rinsed in buffer and deionized water.

Dehydration of the parasite samples was done with an ethanol series and then exposed

to Propylene oxide before infiltration with Epon-Araldite. The blocks of parasites were

trimmed, and sections were obtained using a Reichert Ultracut S ultramicrotome (Leica,

Inc., Deerfield, IL). 60-70nm sections were placed on 200-mesh copper grids and post-

stained with ethanolic uranyl acetate and Reynolds Lead Citrate. Grids were viewed with

a JEOL JEM-1011 Transmission Electron Microscope (JEOL USA, Inc., Peabody, MA)

using an accelerating voltage of 80 KeV. Images were acquired using an AMT XR80M

Wide-Angle Multi-Discipline Mid-Mount CCD Camera (Advanced Microscopy
Techniques, Woburn, MA).

Scanning Electron Microscopy. 7.5mM GlcN was added to percoll isolated schizont-
stage parasites and parasites were allowed to egress and reinvade fresh RBCs. 48hrs
later, parasites were percoll-isolated and then incubated with 2μM Compound 2 for 4
hours without shaking at 37°C in an incubator. After incubation, parasites were washed
twice with warm, complete RPMI + 7.5mM GlcN. Samples were taken immediately after
washing off Compound 2 and then 30min after and fixed as with TEM samples. Parasites
were rinsed with 0.1M Cacodylate-HCl buffer before placing on glass coverslips prepared
with 0.1% Poly-L-lysine. Parasites were allowed to settle onto the glass coverslips in a
moist chamber overnight and then post fixed in 1% osmium tetroxide/0.1M Cacodylate-
HCl buffer for 30 minutes. Cells on coverslips were rinsed three times in deionized water
and then dehydrated with an ethanol series. The glass coverslips were critical point dried
in an Autosamdri-814 Critical Point Dryer (Tousimis Research Corporation, Rockville,
MD), mounted onto aluminum pin stubs with colloidal paint, and sputter coated with gold-
palladium with a Leica EM ACE600 Coater (Leica Microsystems Inc., Buffalo Grove, IL).
Stubs were examined with the FE-SEM FEI Teneo (FEI, Inc., Hillsboro, OR) using the
secondary electron detector to obtain digital images.

Calcium Measurements. To measure Ca^{2+} in PfERC mutants, knockdown was induced
on synchronized schizonts. After 48hrs, schizonts were percoll purified and permeabilized
selectively by treatment with ice-cold 0.04% saponin in PBS for 10 min. Isolated parasites
were then washed 2X with BAG Buffer (116mM NaCl, 5.4mM KCl, 0.8mM $MgSO_4·7H_2O$,
50mM HEPES, 5.5mM Glucose) + 7.5mM GlcN and incubated with 10μM Fluo-4AM
(ThermoFisher) for 45min while rocking at 37°C. After incubation, cells were washed 2X
with BAG buffer + 7.5mM GlcN and immediately taken for fluorimetric measurements.
Fluorescence measurements were carried out in a cuvette (Sarstedt) containing parasites
suspended in 2.5 ml of BAG buffer and 100uM EGTA (Sigma). The cuvette was placed
in a Hitachi F-4500 fluorescence spectrophotometer and Fluo-4AM excitation was done
at 505 nm with emission read at 530 nm (80). Drugs and reagents were added via a

Hamilton syringe. Final concentration of CPA (Sigma) was 3 µM, and Ionomycin (Sigma)

at 2µM.

Loading of *Toxoplasma gondii* tachyzoites with Fura-2AM (ThermoFisher) was

done as previously described (40). Briefly, freshly collected parasites were washed twice

with BAG by centrifugation (706 g for 10 min) and resuspended at a final density of 1 x

10^9 parasites/ml in loading buffer (BAG plus 1.5% sucrose, and 5µM Fura-2AM). The

suspension was incubated for 26 min at 26 °C with mild agitation. Subsequently, parasites

were washed twice with BAG to remove extracellular dye, re-suspended to a final density

of 1 x 10^9/ml in BAG and kept on ice. For fluorescence measurements, 2 x 10^7

parasites/ml were placed in a cuvette with 2.5 ml of BAG buffer. Fluorescence

measurements were done in a Hitachi F-7000 fluorescence spectrophotometer using the

Fura-2AM conditions for excitation (340 and 380 nm) and emission (510 nm). The Fura-

2AM fluorescence response to Ca^{2+} was calibrated from the ratio of 340/380 nm

fluorescence values after subtraction of the background fluorescence of the cells at 340

and 380 nm as described previously (81). The Ca^{2+} release rate is the change in Ca^{2+}

concentration during the initial 20 s after compound addition (41).

Acknowledgments

We thank Michael Reese for comments on the manuscript; Dan Goldberg for comments

on the manuscript, anti Ef1α and anti-PMV anitbodies; The European Malaria Reagent

Repository for anti-MSP1 12.4 and anti-RAP1 2.29, antibodies; Alan Thomas and BEI

Resources NIAID, NIH for anti-AMA1 28G2 and anti-BiP antibodies; Zhicheng Dou and

Michael Blackman for anti-SUB1 antibody; Purnima Bhanot for Compound 1 and

Compound 2; Muthugapatti Kandasamy at the University of Georgia Biomedical

Microscopy Core, Julie Nelson at the CTEGD Cytomtetry Shared Resource Lab, and

Mary Ard from the Georgia Electron Microscopy for technical assistance; Michael

Cipriano for assistance with protein alignment. This work was supported by UGA Startup

funds and UGA Faculty Research Grant (FRG-SE0031) to V.M., and the US National

Institutes of Health (R21AI133322) to V.M. and S.N.J.M., and (T32AI060546) to M.A.F.

References

1. Torgerson PR, Mastroiacovo P. The global burden of congenital toxoplasmosis: a systematic review. Bull World Health Organ. 2013;91(7):501-8.

2. Checkley W, White J,A.Clinton, Jaganath D, Arrowood MJ, Chalmers RM, Chen X, et al. A review of the global burden, novel diagnostics, therapeutics, and vaccine targets for cryptosporidium. Lancet Infect Dis. 2015; 15:85-94.

3. Liu L, Oza S, Hogan D, Chu Y, Perin J, Zhu J, et al. Global, regional, and national causes of under-5 mortality in 2000-15: an updated systematic analysis with implications for the Sustainable Development Goals. Lancet. 2016;388(10063):3027-35.

4. WHO. World Malaria Report 2017. Geneva: World Health Organization 2017

5. Dondorp AM, Nosten F, Yi P, Das D, Phyo AP, Tarning J, et al. Artemisinin resistance in Plasmodium falciparum malaria. N Engl J Med. 2009;361(5):455-67.

6. Mita T, Venkatesan M, Ohashi J, Culleton R, Takahashi N, Tsukahara T, et al. Limited geographical origin and global spread of sulfadoxine-resistant "dhps" alleles in Plasmodium falciparum populations. J Infect Dis. 2011(12):1980.

7. Roper C, Pearce R, Nair S, Sharp B, François Nosten, Anderson T. Intercontinental spread of pyrimethamine-resistant malaria. Science. 2004(5687):1124.

8. Gosling R, Lorenz vS. The future of the RTS,S/AS01 malaria vaccine: an alternative development plan. PLoS Medicine, 2016;13 :e1001994.

9. Donald RGK, Allocco J, Singh SB, Nare B, Salowe SP, Wiltsie J, et al. Toxoplasma gondii cyclic GMP-dependent kinase: chemotherapeutic targeting of an essential parasite protein kinase. Eukaryot Cell. 2002;1(3):317-28.

10. Christine RC, Hackett F, Strath M, Penzo M, Chrislaine Withers-Martinez, David AB, et al. Malaria parasite cGMP-dependent protein kinase regulates blood stage merozoite secretory organelle discharge and egress. PLoS Pathogens, 2013; 9(5):e1003344.

11. Brochet M, Collins MO, Smith TK, Thompson E, Sebastian S, Volkmann K, Schwach F, et al. Phosphoinositide metabolism links cGMP-dependent protein kinase G to essential Ca^{2+} signals at key decision points in the life cycle of malaria parasites. PloS Biology, 2014; 12(3): e1001806

12. Hortua Triana MA, Márquez-Nogueras KM, Vella SA, Moreno SNJ. Calcium
signaling and the lytic cycle of the Apicomplexan parasite Toxoplasma gondii. BBA -
Molecular Cell Research. 2018; 1865: 1846-1856

13. Brochet M, Billker O. Calcium signalling in malaria parasites. Mol Microbiol. 2016;
100(3):397-408.

14. Holder A, Baker D, Baker DA, Taylor HM, Sicard A, Grainger M, et al. The malaria
parasite cyclic GMP-dependent protein kinase plays a central role in blood-stage
schizogony. Eukaryot Cell. 2010;9(1):37-45.

15. Hale VL, Watermeyer JM, Hackett F, Vizcay G, Ooij Cv, Thomas JA, et al.
Parasitophorous vacuole poration precedes its rupture and rapid host erythrocyte
cytoskeleton collapse in Plasmodium falciparum egress. Proc Natl Acad Sci USA.
2017; 114(13):3439-3444.

16. Absalon S, Blomqvist K, Rudlaff RM, DeLano TJ, Pollastri MP, Dvorin JD. Calcium-
dependent protein kinase 5 Is required for release of egress-specific organelles in
Plasmodium falciparum. MBio. 2018;9(1):e00130.

17. Glushakova S, Beck JR, Garten M, Busse BL, Nasamu AS, Tenkova-Heuser T, et
al. Rounding precedes rupture and breakdown of vacuolar membranes minutes
before malaria parasite egress from erythrocytes. Cell Microbiol. 2018;
20(10):e12868.

18. Garg S, Agarwal S, Chitnis CE, Singh S, Kumar S, Yazdani SS. Calcium-dependent
permeabilization of erythrocytes by a perforin-like protein during egress of malaria
parasites. Nat Commun. 2013;4: 1736

19. Lourido S, Moreno SNJ. The calcium signaling toolkit of the Apicomplexan parasites
Toxoplasma gondii and Plasmodium spp. Cell Calcium. 2015;57(3):186-93.

20. Lourido S, Tang K, Sibley LD. Distinct signalling pathways control Toxoplasma
egress and host-cell invasion. EMBO J. 2012;31(24):4524-34.

21. Yeoh S, O'Donnell RA, Koussis K, Dluzewski AR, Ansell KH, Osborne SA, et al.
Subcellular discharge of a serine protease mediates release of invasive malaria
parasites from host erythrocytes. Cell. 2007;131(6):1072-83.

22. Nasamu AS, Glushakova S, Russo I, Oksman BVA, Kim AS, Fremont DH, et al.
Plasmepsins IX and X are essential and druggable mediators of malaria parasite
egress and invasion. Science. 2017; 358(6362):518-522.
23. Pino P, Caldelari R, Mukherjee B, Vahokoski J, Klages N, Maco B, et al. A
multistage antimalarial targets the plasmepsins IX and X essential for invasion and
egress. Science. 2017; 358(6362):522-528.
24. Blackman MJ, Fujioka H, Stafford WHL, Sajid M, Clough B, Fleck SL, et al. A
subtilisin-like protein in secretory organelles of Plasmodium falciparum merozoites. J
Biol Chem. 1998; 273(36):23398-409.
25. Sajid M, Withers-Martinez C, Blackman MJ. Maturation and specificity of
Plasmodium falciparum subtilisin-like protease-1, a malaria merozoite subtilisin-like
serine protease. J Biol Chem. 2000; 275(1):631-41.
26. Tawk L, Kent R, Gorgette O, Mercereau-Puijalon O, Barale J, Lacroix C, et al. A key
role for plasmodium subtilisin-like SUB1 protease in egress of malaria parasites from
host hepatocytes. J Biol Chem. 2013;288(46):33336-46.
27. Das S, Hertrich N, Perrin AJ, Withers-Martinez C, Collins CR, Jones ML, et al.
Processing of Plasmodium falciparum merozoite surface protein MSP1 activates a
spectrin-binding function enabling parasite egress from RBCs. Cell Host Microbe.
2015;18(4):433-44.
28. Thomas JA, Tan MSY, Hackett F, Blackman MJ, Bisson C, Umrekar TR, et al. A
protease cascade regulates release of the human malaria parasite Plasmodium
falciparum from host red blood cells. Nat Microbiol. 2018;3(4):447-55.
29. Besteiro S, Lebrun M, Lamarque M, Besteiro S, Papoin J, Roques M, et al. The
RON2-AMA1 interaction is a critical step in moving junction-dependent invasion by
Apicomplexan Parasites. PLoS Pathogens. 2011;7(2): e1001276.
30. Paul AS, Egan ES, Duraisingh MT. Host-parasite interactions that guide red blood
cell invasion by malaria parasites. Curr Opin Hematol. 2015;22(3):220-226.
31. Tomavo S, Slomianny C, Meissner M, Vern BC. Protein trafficking through the
endosomal system prepares intracellular parasites for a home invasion. PLoS
Pathogens. 2013; 9(10):e1003629.

32. Coffey MJ, Jennison C, Tonkin CJ, Boddey JA. Role of the ER and Golgi in protein
export by Apicomplexa. Curr Opin Cell Biol. 2016;41:18-24.

33. Greca N, Hibbs AR, Riffkin C, Foley M, Tilley L. Identification of an endoplasmic
reticulum-resident calcium-binding protein with multiple EF-hand motifs in asexual
stages of Plasmodium falciparum. Mol Biochem Parasitol. 1997;89(2):283-93.

34. Honoré B. The rapidly expanding CREC protein family: members, localization,
function, and role in disease. Bioessays. 2009;31(3):262-77.

35. Prommana P, Uthaipibull C, Wongsombat C, Kamchonwongpaisan S, Yuthavong Y,
Knuepfer E, et al. Inducible knockdown of Plasmodium gene expression using the
glmS ribozyme. PLoS ONE. 2013;8(8): e73783.

36. Salmon BL, Oksman A, Goldberg DE. Malaria parasite exit from the host
erythrocyte: a two-step process requiring extraerythrocytic proteolysis. Proc Natl
Acad Sci U S A. 2001; 98(1):271-6.

37. Sidik SM, Hackett CG, Tran F, Westwood NJ, Lourido S. Efficient genome
engineering of Toxoplasma gondii using CRISPR/Cas9. PLoS ONE. 2014;9(6):
e100450.

38. Shen B, Brown KM, Lee TD, Sibley LD. Efficient gene disruption in diverse strains of
Toxoplasma gondii using CRISPR/CAS9. MBio. 2014;5(3): e01114-14.

39. Moreno SNJ, Zhong L. Acidocalcisomes in Toxoplasma gondii tachyzoites. Biochem
J. 1996; 313(Pt2):655-9.

40. Pace DA, McKnight CA, Liu J, Jimenez V, Moreno SNJ. Calcium entry in
Toxoplasma gondii and its enhancing effect of invasion-linked traits. J Biol Chem.
2014;289(28):19637-47.

41. Alleva LM. Calcium regulation in the intraerythrocytic malaria parasite Plasmodium
falciparum. Molecular and biochemical parasitology. 2001;117(2):121-8.

42. Giganti D, Bouillon A, Tawk L, Robert F, Martinez M, Crublet E, et al. A novel
Plasmodium-specific prodomain fold regulates the malaria drug target SUB1
subtilase. Nat Commun. 2014;5:4833.

43. Goel VK, Li X, Chen H, Shih-Chun Liu, Chishti AH, Steven SO. Band 3 Is a host
receptor binding merozoite surface protein 1 during the Plasmodium falciparum
invasion of erythrocytes. Proc Natl Acad Sciences USA. 2003; 100(9):5164-9.

44. Li X, Chen H, Oo TH, Daly TM, Bergman LW, Shih-Chun Liu, et al. A co-ligand
complex anchors Plasmodium falciparum merozoites to the erythrocyte invasion
receptor b and 3. J Biol Chem. 2004;279(7):5765-71.

45. Baldwin MR, Li X, Hanada T, Liu S, Chishti AH. Merozoite surface protein 1
recognition of host glycophorin A mediates malaria parasite invasion of red blood
cells. Blood. 2015; 125(17):2704-11.

46. Boyle MJ, Wilson DW, Richards JS, Riglar DT, Tetteh K, Conway DJ, et al. Isolation
of viable Plasmodium falciparum merozoites to define erythrocyte invasion events
and advance vaccine and drug development. Proc Natl Acad Sciences USA. 2010;
107(32): 14378-83.

47. Lin CS, Uboldi AD, Epp C, Bujard H, Tsuboi T, Czabotar PE, et al. Multiple
Plasmodium falciparum merozoite surface protein 1 complexes mediate merozoite
binding to human erythrocytes. J Biol Chem. 2016;291(14):7703-15.

48. Bannister LH, Mitchell GH, Butcher GA, Dennis ED. Lamellar membranes
associated with rhoptries in erythrocytic merozoites of Plasmodium knowlesi: a clue
to the mechanism of invasion. Parasitology. 1986;92(pt2):291-303.

49. Riglar DT, Richard D, Wilson DW, Boyle MJ, Dekiwadia C, Turnbull L, et al. Super-
resolution dissection of coordinated events during malaria parasite invasion of the
human erythrocyte. Cell Host Microbe. 2011;9(1):9-20.

50. Cowman AF, Tonkin CJ, Tham W, Duraisingh MT. The molecular basis of
erythrocyte invasion by malaria parasites. Cell Host Microbe. 2017;22:232-45.

51. Howard RF, Schmidt CM. The secretary pathway of Plasmodium falciparum
regulates transport of p82/RAP-1 to the rhoptries. Mol Biochem Parasitol.
1995;74(1):43-54.

52. Howard RF, Narum D, Blackman M, Thurman J. Analysis of the processing of
Plasmodium falciparum rhoptry-associated protein 1 and localization of Pr86 to
schizont rhoptries and p67 to free merozoites. Mol Biochem Parasitol. 1998;
92(1):111-22.

53. Narum DL, Thomas AW. Differential localization of full-length and processed forms
of PF83/AMA-1 an apical membrane antigen of Plasmodium falciparum merozoites.
Mol Biochem Parasitol. 1994;67(1):59-68.

54. Howell SA, Withers-Martinez C, Kocken C, Thomas AW, Blackman MJ. Proteolytic
processing and primary structure of Plasmodium falciparum apical membrane
antigen-1. J Biol Chem. 2001;276(33):31311-20.

55. Bannister LH, Hopkins JM, Dluzewski AR, Margos G, Williams IT, Blackman MJ, et
al. Plasmodium falciparum apical membrane antigen 1 (PfAMA-1) is translocated
within micronemes along subpellicular microtubules during merozoite development.
J Cell Sci. 2003;116(18):3825-34.

56. Healer J, Tiriglia T, Hodder AN, Gemmill AW, Cowman AF. Functional analysis of
Plasmodium falciparum apical membrane antigen 1 utilizing interspecies domains.
Infect Immun. 2005; 73(4):2444-2451.

57. Santos JM, Josling G, Ross P, Joshi P, Orchard L, Campbell T, et al. Red blood cell
invasion by the malaria parasite is coordinated by the PfAP2-I transcription factor.
Cell Host Microbe. 2017;21(6): 731-741.e10.

58. Kim H, Morita M, Sanai H, Hiramoto A, Sato A, Wataya Y, et al. Plasmodium
falciparum endoplasmic reticulum-resident calcium binding protein is a possible
target of synthetic antimalarial endoperoxides, N-89 and N-251. J Proteome Res.
2012;11(12):5704-11.

59. Billker O, Lourido S, Sibley LD. Calcium-Dependent Signaling and Kinases in
Apicomplexan Parasites. Cell Host & Microbe. 2009;5(6):612-22.

60. Lagal V, Binder EM, Huynh M, Kafsack BFC, Harris PK, Diez R, et al. Toxoplasma
gondii protease TgSUB1 is required for cell surface processing of micronemal
adhesive complexes and efficient adhesion of tachyzoites. Cell Microbiol.
2010;12(12):1792-808.

61. Monerri NCS, Flynn HR, Campos MG, Hackett F, Koussis K, Withers-Martinez C, et
al. Global identification of multiple substrates for Plasmodium falciparum SUB1, an
essential malarial processing protease. Infect Immun. 2011;79(3):1086-1097.

62. Knuepfer E, Rug M, Klonis N, Tilley L, Cowman AF. Trafficking of the major
virulence factor to the surface of transfected P falciparum-infected erythrocytes.
Blood. 2005;105(10):4078-87.

63. Jung DH, Mo SH, Kim DH. Calumenin, a multiple EF-hands Ca2+-binding protein,
interacts with ryanodine receptor-1 in rabbit skeletal sarcoplasmic reticulum.
Biochem Biophys Res Commun. 2006;343(1):34-42.
64. Xu S, Xu Y, Chen L, Fang Q, Song S, Chen J, et al. RCN1 suppresses ER stress-
induced apoptosis via calcium homeostasis and PERK-CHOP signaling.
Oncogenesis. 2017 Mar 20,;6(3):e304.
65. Mazzorana M, Hussain R, Sorensen T. Ca-dependent folding of human calumenin.
PLoS ONE. 2016;11(3): e0151547.
66. Yonezawa N, Suzuki N, Ban S, Itoh E, Chen S, Yonezawa N, et al. Calcium-
dependent structural changes in human reticulocalbin-1. J Biochem.
2014;155(5):281-93.
67. Mok S, Ashley EA, Ferreira PE, Zhu L, Lin Z, Yeo T, et al. Population
transcriptomics of human malaria parasites reveals the mechanism of artemisinin
resistance. Science. 2015;347(6220):431-435.
68. Drew ME, Banerjee R, Uffman EW, Gilbertson S, Rosenthal PJ, Goldberg DE.
Plasmodium food vacuole plasmepsins are activated by falcipains. J Biol Chem.
2008(19):12870-12876.
69. Russo I, Oksman A, Goldberg DE. Fatty acid acylation regulates trafficking of the
unusual Plasmodium falciparum calpain to the nucleolus. Mol Microbiol. 2009;
72(1):229-245.
70. Cobb DW, Florentin A, Fierro MA, Krakowiak M, Moore JM, Muralidharan V. The
exported chaperone PfHsp70x is dispensable for the Plasmodium falciparum
intraerythrocytic life cycle. mSphere. 2017 Oct 25,;2(5):e00363-17.
71. Florentin A, Cobb DW, Fishburn JD, Cipriano MJ, Kim PS, Fierro MA, et al. PfClpC
is an essential Clp chaperone required for plastid integrity and Clp protease stability
in Plasmodium falciparum. Cell Reports. 2017; 21(7):1746-56.
72. Farwell DG, Shera KA, Koop JI, Bonnet GA, Matthews CP, Reuther GW, et al.
Genetic and epigenetic changes in human epithelial cells immortalized by
telomerase. Am J Pathol. 2000; 156(5):1537-47.

73. Spillman NJ, Beck JR, Ganesan SM, Niles JC, Goldberg DE. The chaperonin TRiC forms an oligomeric complex in the malaria parasite cytosol. Cell Microbiol. 2017; 19(6): e12719.

74. Ganesan SM, Morrisey JM, Ke H, Painter HJ, Laroiya K, Phillips MA, et al. Yeast dihydroorotate dehydrogenase as a new selectable marker for Plasmodium falciparum transfection. MolBiochem Parasitol. 2011;177(1):29-34.

75. Ghorbal M, Gorman M, Macpherson CR, Martins RM, Scherf A, Lopez-Rubio J. Genome editing in the human malaria parasite Plasmodium falciparum using the CRISPR-Cas9 system. Nat Biotechnol. 2014;32(8):819-21.

76. Liu J, Pace D, Dou Z, King TP, Guidot D, Li ZH, et al. A vacuolar-H+-pyrophosphatase (TgVP1) is required for microneme secretion, host cell invasion, and extracellular survival of Toxoplasma gondii. Mol Microbiol. 2014;93(4):698-712.

77. Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. Nature. 1970;227(5259):680-5.

78. Hager KM, Striepen B, Tilney LG, Roos DS. The nuclear envelope serves as an intermediary between the ER and Golgi complex in the intracellular parasite Toxoplasma gondii. J Cell Sci. 1999;112(16):2631.

79. Budu A, Gomes MM, Melo PM, El CM, Bagnaresi P, Azevedo MF, et al. Calmidazolium evokes high calcium fluctuations in Plasmodium falciparum. Cell Signal. 2016;28(3):125-35.

80. Grynkiewicz G, Poenie M, Tsien RY. A new generation of Ca2+ indicators with greatly improved fluorescence properties. J Biol Chem. 1985;260(6):3440-50.

81. Waterhouse AM, Procter JB, Martin DM, Clamp M, Barton GJ. Jalview Version 2-a multiple sequence alignment editor and analysis workbench. Bioinformatics. 2009; 25(9):1189-91.

Figure 1: Generating PfERC-*glmS/M9* mutant parasites. (A) Schematic representation of the domain structure of PfERC and TgERC. Both proteins contain a signal peptide, 5 (TgERC) or 6 (PfERC) EF hands, and an ER-retention signal. (B) Mechanism of the conditional expression of PfERC using the *glmS* ribozyme system. This is an inactive ribozyme that is transcribed, but not translated, with the mRNA of a protein of interest. Addition of glucosamine (GlcN) leads to its phosphorylation within the

cell to glucosamine-6-phosphate (GlcN-6P). GlcN-6P binds to the transcribed PfERC-

glmS mRNA and the *glmS* ribozyme is activated and cleaves itself from the mRNA. This

leads to disassociation of the mRNA from its poly-A tail and leads to the degradation of

target specific mRNA. The resulting decline in mRNA levels leads to reduced protein

levels and, thus, loss of gene expression. As a control, we generated parasite lines

containing a mutated version of the *glmS* ribozyme, called *M9*, which cannot cleave

itself upon binding of GlcN. (C) Using the CRISPR/Cas9 system, we induced a double-

stranded break in the PfERC locus that was repaired by a donor plasmid containing

homology templates to the PfERC locus and appended a C-terminal 3XHA tag and a

stop codon followed by the *glmS* or *M9* sequence to the targeted gene. The location of

diagnostic primers used to demonstrate the repair of the locus via double cross-over

homologous integration are also shown (P5, P8 and P11). (D) Western blot of lysates

isolated from two independent clones and the parental line (3D7) probed with anti-HA

antibodies show that the PfERC gene was tagged with HA in the mutants but not the

parental line. PfBiP was the loading control. (E) PCR analysis of the generated mutants

using specific primers (P5+P8; Table S1) in the C-terminus and 3'UTR of PfERC shows

integration of the HA tag and *glms*/*M9* ribozymes into the PfERC locus. Modification of

PfERC gene introduces an AfeI restriction enzyme site in this locus that is absent in the

parental line. Digesting the PCR products (using AfeI) resulting from amplification using

primers P5+P8 shows that AfeI is able to digest the PCR products from our mutants but

not the parental line. PCR analysis using another primer pair (P5+P11) that sits on the

glmS/*M9* sequence shows that amplification only occurs in the mutants but not in the

parental line. (F) Representative IFA of PfERC-*M9* parasites showing that tagged

PfERC localizes to the ER as shown with co-localization with the ER chaperone BiP in

all asexual stages of the parasite. From left to right, the images are phase-contrast

micrographs of parasites, parasites stained with anti-BiP antibody (green), parasites

stained with anti-HA antibody (red), parasite nucleus stained with DAPI (blue), and

fluorescence merge images of the parasites. Abbreviations: R, rings; T, trophozoites; S,

schizonts. Scale bar, 5µm.

Figure 2: PfERC mutants fail to transition from schizonts to rings. (A) Western blot of parasite lysates isolated from PfERC-*glmS* and PfERC-*M9* parasites grown in the presence of 7.5 mM GlcN and probed with antibodies against HA (red) and against the ER-resident protease Plasmepsin V (PMV; green). One representative experiment out of

four is shown. (B) Quantification of the overall reduction in PfERC in PfERC-*glmS* and PfERC-*M9* parasites over time after addition of GlcN, as shown in (A). Data were normalized to the loading control (PMV) and are represented as mean ± SEM (n=4 biological replicates). (C) Growth curve of asynchronous PfERC-*glmS* and PfERC-*M9* clones grown in the presence of GlcN shows that PfERC is essential for asexual reproduction. Data are normalized to parasites grown without GlcN and are represented as the mean ± SEM (n=3 biological replicates). (D) Representative blood smears of synchronous PfERC-*glmS* and PfERC-*M9* parasites grown in the presence of GlcN (n=2 biological replicates). (E) Ring:schizont ratio of synchronous PfERC-*glmS* and PfERC-*M9* parasites taken at 40-52hrs post-addition of GlcN. Data are represented as the mean ± SEM (n=3 biological replicates). (F) The parasites shown in blood smears (D) were counted. The amount of each lifecycle stage (ring, trophozoite, schizont) was determined as a percentage of the total number of parasites for each time point. (G) The egress of PfERC-*glmS* or PfERC-*M9* parasites was quantified by determining the amount of schizonts in blood smears as in (D) present at 44hrs and at 56hrs. Egress was quantified using the following equation: (# of schizonts at 56h / # schizonts at 44h) x100. Data are represented as mean ± SEM (n=2 biological replicates; *$P<0.05$, unpaired *t*-test). (H) Invasion of PfERC-*glmS* or PfERC-*M9* parasites was quantified by determining the amount of ring stages in blood smears as in (D) present at 44hrs and at 56hrs. This was quantified using the following equation: (# of rings at 44h / # rings at 56h) x100. Data are represented as mean ± SEM (n=2 biological replicates; *$P<0.05$, unpaired *t*-test).

Figure 3: PfERC knockdown prevents PVM breakdown. (A) Schematic showing the experimental layout to study the effect of PfERC knockdown using compounds that inhibit egress in parasites. Abbreviations: C1/2- PKG inhibitors, compound 1 or compound 2;

SEM- Scanning Electron Microscopy; TEM- Transmission Electron Microscopy. (B) As

shown in (A), synchronized PfERC-*glmS* and PfERC-*M9* schizonts were observed by flow

cytometry after removal of C1 (time 0hr). Schizonts were quantified as a percentage of

the total amount of parasites as determined by flow cytometry. Data are represented as

the mean ± SEM (n=3 biological replicates). (C) Representative SEM images of C2

arrested PfERC-*glmS* (n=2 biological replicates) and PfERC-*M9* (n=2 biological

replicates) mutants fixed immediately after washing off C2 and after 30mins, as shown in

(A). Scale bar, 800nm. (D) Representative TEM images of PfERC-*glmS* (n=2 biological

replicates) and PfERC-*M9* (n=2 biological replicates) schizonts incubated with E-64, as

shown in (A). PVM is marked by black arrowheads.

1099

1100

Figure 4: Generation and characterization of TgERC mutants. (A) Using
CRISPR/Cas9, we generated *Δtgerc* mutants by introducing a DHFR cassette that

disrupted the open-reading frame of TgERC. (B) PCR analysis using primers that sit on
the homology regions used for insertion showing amplification of the WT band (~3Kb) and
insertion of the DHRF cassette as demonstrated by a higher product (~6Kb) in the
mutants. (C) Western blot using generated antibodies against TgERC showing the
presence of TgERC in the RH (WT) parasites but not in the $\Delta tgerc$ mutants. αTubulin was
used as the loading control. (D) IFAs of either extracellular or intracellular $\Delta tgerc$ mutants
transiently transfected with the P30-GFP-HDEL plasmid (79) which contains an ER-
localized GFP. Left panel shows parasites probed with anti-TgERC antibodies, middle
panel shows presence of transfected GFP, and right panel shows overlay of staining with
DIC. Scale bar, 5μm. (E) Representative image of one plaque assay out of three biological
replicates. (F) Quantification of plaque sizes between RH and $\Delta tgerc$ parasites (n=3
biological replicates; n.s=non-significant, unpaired t-test). (G) Representative Ca^{2+}
tracings of RH and $\Delta tgerc$ parasites treated with the extracellular Ca^{2+} chelator EGTA and
the SERCA pump inhibitor thapsigargin. (H) Quantification of the rate of Ca^{2+} release of
RH and $\Delta tgerc$ parasites from (G) (n=3 biological replicates; n.s=non-significant, unpaired
t-test). (I) Representative Ca^{2+} tracings of RH and $\Delta tgerc$ parasites treated with the
extracellular Ca^{2+} and thapsigargin. (J) Quantification of the rate of Ca^{2+} release of RH
and $\Delta tgerc$ from (I) (n= 3 biological replicates, *$P<0.05$, unpaired t-test).

1121

Figure 5: PfERC knockdown inhibits SUB1 and MSP1 processing. (A) Western blot of parasite lysates isolated from PfERC-*glmS* and PfERC-*M9* schizonts grown in the presence of GlcN for 48 hours and probed with anti-SUB1 antibodies (top panel) and against the ER-resident protease PMV (loading control, bottom panel). One representative experiment out of four is shown. The protein marker sizes that co-migrated

with the probed protein are shown on the left. (B) Quantification of the overall reduction

in processing of SUB1 in PfERC-*glmS* and PfERC-*M9* parasites over time after addition

of GlcN, as shown in (A). Data were normalized to the total SUB1 (processed +

unprocessed) and are represented as mean ± SEM (n=4 biological replicates; **P<0.005

unpaired *t*-test). (C) Western blot of parasite lysates isolated from PfERC-*glmS* and

PfERC-*M9* schizonts grown in the presence of GlcN for 48 hours and probed with anti-

MSP1 antibodies (top panel) and against the anti-EF1α (loading control, bottom panel).

One representative experiment out of two is shown. The protein marker sizes that co-

migrated with the probed protein are shown on the left. (D) Quantification of the overall

reduction in processing of MSP1 in PfERC-*glmS* and PfERC-*M9* parasites over time after

addition of GlcN. Quantification of the overall reduction in MSP1 in PfERC-*glmS* and

PfERC-*M9* parasites over time after addition of GlcN, as shown in (B). Data were

normalized to the loading control (EF1α) and are represented as mean ± SEM (n=2

biological replicates; **P<0.005 unpaired *t*-test). (E-G) Representative Super-Resolution

SIM images of PfERC-*glmS* and PfERC-*M9* schizonts stained with antibodies against

MSP1, AMA1, and RAP1. From top to bottom, the images are parasites stained with anti-

MSP1 (E), anti-AMA1 (F), anti-RAP1 (G) antibodies (green), parasite nucleus stained with

DAPI (blue), and fluorescence merge images of the schizonts incubated with GlcN for

48hrs (n=2 biological replicates). Scale bar 2μm. (H) Representative TEM images of

PfERC-*glmS* and PfERC-*M9* schizonts grown for 48hrs with GlcN and incubated with E-

64 for 8 hours, as shown in Figure 3A (n=2 biological replicates). Small arrowheads point

to rhoptries, large arrowheads to micronemes, and white arrowheads to PVM fragments

(15). Scale bar, 200nm.

Figure 6: PfERC knockdown prevents invasion of RBCs. (A) As shown in Figure 4A,
synchronized PfERC-*glmS* and PfERC-*M9* parasites were observed by flow cytometry
after removal of C1 (time 0hr). Rings were quantified as a percentage of the total amount
of parasites as determined by flow cytometry. Data are represented as the mean ± SEM
(n=3 biological replicates). (B) Schematic showing the experimental layout to study the

effect of PfERC knockdown on invasion of merozoites into host RBCs. Merozoites were

purified using mechanical lysis after 8hr incubation with E64. (C) Invasion rates of

mechanically purified merozoites from PfERC-*glmS* and PfERC-*M9* parasites, as shown

in (B), were quantified using flow cytometry (47). All replicates were normalized to PfERC-

M9 merozoites. Data are represented as mean ± SEM (n=7 biological replicates;

****$P<0.0001$, unpaired *t*-test). (D) Western blot of parasite lysates isolated from PfERC-

glmS and PfERC-*M9* schizonts grown in the presence of GlcN for 48 hours and probed

with anti-AMA1 antibodies (top panel) and anti- EF1αantibodies (loading control, bottom

panel). One representative experiment out of eight is shown. The protein marker sizes

that co-migrated with the probed protein are shown on the left. (E) Quantification of the

overall reduction in processing of AMA1 in PfERC-*glmS* and PfERC-*M9* parasites over

time after addition of GlcN, as shown in (D). Data were normalized to the total AMA1

(processed + unprocessed) and are represented as mean ± SEM (n=8 biological

replicates; *$P<0.05$ unpaired *t*-test). (F) Western blot of parasite lysates isolated from

PfERC-*glmS* and PfERC-*M9* schizonts grown in the presence of GlcN for 48 hours and

probed with anti-RAP1 antibodies (top panel) and anti- EF1α antibodies (loading control,

bottom panel). One representative experiment out of five is shown. The protein marker

sizes that co-migrated with the probed protein are shown on the left. (G) Quantification of

the overall reduction in processing of RAP1 in PfERC-*glmS* and PfERC-*M9* parasites over

time after addition of GlcN, as shown in (F). Data were normalized to the total RAP1

(processed + unprocessed) and are represented as mean ± SEM (n=5 biological

replicates; n.s=non-significant, unpaired *t*-test).

 Supplementary Figure 1: Sequence alignment of TgERC and PfERC to other members
 of the CREC family of proteins using MUSCLE alignment, viewed using JalView Software
 (http://www.jalview.org/) and BOXSHADE (82). Alignment was done using the human
 homologs: Cab-45, ERC-55 (RCN2), Calumenin, and Reticulocalbin1 (RCN1). Identical
 residues are shaded in black, similar residues are shaded in gray, and EF-hands are
 highlighted in red.

1184

1185

1186

1187

1188

1189

1190

1191

1192

1193

1194

1195

1196

1197

1198

1199

1200

PfERC-*glmS* —— PfERC-*M9* - - -

Supplementary Figure 2: Effect of PfERC knockdown on parasite growth. (A)

Representative graph (one out of three) showing dose-dependent response of PfERC-

*glmS*2 to GlcN. Asynchronous PfERC-*glmS* parasites were incubated in different

concentrations of GlcN and growth after three days was assessed by flow cytometry. Data

are represented as mean ± SEM of n=3 biological replicates. (B) Representative life cycle

graph showing all stages from Figures 3D and 3F (n=2 biological replicates).

1207

1208

1209

1210

1211

1212

1213

1214

Supplementary Figure 3: Effect of PfERC knockdown on Ca_2^{2+} in the ER. (A) Experimental schematic showing how Ca^{2+} measurements were done in PfERC-*glmS* and PfERC-*M9* mutants. Synchronized PfERC-*glmS* and PfERC-*M9* schizonts were incubated with GlcN for 48 hours and isolated using saponin lysis, which lyses the RBC membrane but leaves the PV intact. Abbreviations: CPA- cyclopiazoic acid (B) Representative fluorescence tracings after CPA addition to PfERC-*glmS* and PfERC-*M9* schizonts, isolated as in (A). Quantification was done by calculating the difference in fluorescence between the basal to the highest peak of fluorescence. Data are represented as the combined mean ± SEM (PfERC-*glmS;* n=15 biological replicates; PfERC-*M9;* n=9 biological replicates; n.s- non-significant, unpaired *t*-test). (C)

1225   Representative fluorescence tracings after Ionomycin addition to PfERC-*glmS* and

1226   PfERC-*M9* schizonts, isolated as in (A). Quantification was done by calculating the

1227   difference in fluorescence between the basal to the highest peak of fluorescence. Data

1228   are represented as the combined mean ± SEM (PfERC-*glmS;* n=9 biological replicates;

1229   PfERC-*M9;* n=5 biological replicates; n.s- non-significant, unpaired *t*-test).

A

αSUB1

B

αMSP1

C

αRAP1

D

αAMA1

1230

1231   **Supplementary Figure 4:** Representative whole-blot images of Western blots of lysates

1232   from PfERC-*glmS* and PfERC-*M9* schizonts incubated with GlcN for 48 hours, probed

1233   with anti-SUB1, anti-MSP1, anti-AMA1, and anti-RAP1 antibodies from Figures 5 and 6.

1234   The protein marker sizes that co-migrated with the probed protein are shown on the left.

1235

1236

1237

1238

1239

1240

1241

1242

1243

1244

1245

1246

1247

1248

1249

1250

1251

1252

1253

1254   **Table S1:** Primers used in this study.

Primer	Primer Sequence
1	GCGGCGATGCCCAAGTTGTCGTTTTAGAGCTAGAAATAGC
2	AACTTGACATCCCCATTTAC
3	CGACTGTTCCCCCCAATCAATTCTACTTTCGTCGCTTATCTCGAGAAAACAAAGCTTCGCCAGGCTGTAAATCC
4	CTCCTGTCCGCAGAAGCTCTGTACGTGTTCTCAACAACTAGCTCTTCATGTTCATCCTGCAAGTGCATAGAAGGAAAGTTG
5	TCCGACTCTCCAACGTGGTTGC
6	GCTGCTACACAGCACTCCCTG
7	CTGCAGGTCTGGACATTTAAAGTTCATCACTAGCGTAATCTGGAACATCG
8	CGATGTTCCAGATTACGCTAGTGATGAACTTTAAATGTCCAGACCTGCAG
9	AATTCGCCCTTTCCGCGGAGAATAGAAAAATTATTTCATTTGATAGATAAAAACAATGAT
10	TGGGTAACTAGTAGCGCTTAATTCATCAATTGCTGGGGATTTTTGTTGCGATGCATCGTC
11	ATGATCTTGCCGGCAAGCTTTTTATATAAACATATTTTTTTTTTTTTAACATAAAGGG
12	CCTTGAGCTCGCTAGCGACAAATTGGATAGATAATAGGGGGTACAAATATACATAC
13	AAGTATATAATATTCAATTGCTGGGGATTTTTGTGTTTAGAGCTAGAA
14	TTCTAGCTCTAAAACACAAAAATCCCCAGCAATTGAATATTATATACTTA
15	GCAAGATCATGTGATTTCTCTTTGTTCAAGGAGTCACCCCC
16	GACGACGACAAGATAATGGCGGCCTTCGCAGACAG
17	GAGGAGAAGCCCGGTCAGTTCATCGTGTTTGTCTTCGTCGTCG

1255

1256