Genetic Heterogeneity Profiling by Single Cell RNA Sequencing

Zilu Zhou, Graduate Group in Genomics and Computational Biology, University of Pennsylvania, Philadelphia, PA, USA. zhouzilu@pennmedicine.upenn.edu

Bihui Xu, Department of Radiation Oncology, Parker Institute for Cancer Immunotherapy, Abramson Family Cancer Research Institute, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA.

Andy Minn, Department of Radiation Oncology, Parker Institute for Cancer Immunotherapy, Abramson Family Cancer Research Institute, Graduate Group in Cell and Molecular Biology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA.

Nancy R Zhang*, Department of Statistics, The Wharton School, University of Pennsylvania, Philadelphia, PA, USA. nzh@wharton.upenn.edu

* Corresponding author

Abstract

Detection of genetically distinct subclones and profiling the transcriptomic differences between them is important for studying the evolutionary dynamics of tumors, as well as for accurate prognosis and effective treatment of cancer in the clinic. For the profiling of intra-tumor transcriptional heterogeneity, single cell RNA-sequencing (scRNA-seq) is now ubiquitously adopted in ongoing and planned cancer studies. Detection of somatic DNA mutations and inference of clonal membership from scRNA-seq, however, is currently unreliable. We propose DENDRO, an analysis method for scRNA-seq data that detects genetically distinct subclones, assigns each single cell to a subclone, and reconstructs the phylogenetic tree describing the tumor's evolutionary history. DENDRO utilizes information from single nucleotide mutations in transcribed regions and accounts for technical noise and expression stochasticity at the single cell level. The accuracy of DENDRO was benchmarked on spike-in datasets and on scRNA-seq data with known subpopulation structure. We applied DENDRO to delineate subclonal expansion in a mouse melanoma model in response to immunotherapy, highlighting the role of neoantigens in treatment response. We also applied DENDRO to primary and lymph-node metastasis samples in breast cancer, where the new approach allowed us to better understand the relationship between genetic and transcriptomic intratumor variation.

Keywords

single-cell RNA sequencing, intratumor heterogeneity, cancer genomics, phylogeny inference, multi-omics analysis

Background

DNA alterations, especially single nucleotide alteration (SNA) and epigenetic modulation both contribute to intratumor heterogeneity [1], which mediates tumor initiation, progression, metastasis and relapse [2, 3]. Intratumor genetic and transcriptomic variation underlie patients' response to treatment, as natural selection can lead to the emergence of subclones that are drug resistant [4]. Thus, identifying subclonal DNA alterations and assessing their impact on intratumor transcriptional dynamics can elucidate the mechanisms of tumor evolution and, further, uncover potential targets for therapy. To characterize intratumor genetic heterogeneity, most prior studies have used bulk tumor DNA sequencing [5-12], but these approaches have limited resolution and power [13].

Breakthroughs in single-cell genomics promise to reshape cancer research by allowing comprehensive cell type classification and rare subclone identification. For example, in breast cancer, single-cell DNA sequencing (scDNA-seq) was used to distinguish normal cells from malignant cells, the latter of which were further classified into subclones [14-16]. For the profiling of intra-tumor transcriptional heterogeneity, single cell RNA-sequencing (scRNA-seq), such as Smart-seq2 [17], Drop-seq [18], and 10X Genomics Chromium^{TM}, is now ubiquitously adopted in ongoing and planned cancer studies. ScRNA-seq studies have already led to novel insights into cancer progression and metastasis, as well as into tumor prognosis and treatment response, especially response variability in immune checkpoint blockade (ICB) [19-26]. Characterization of intratumor genetic heterogeneity and identification of subclones using scRNA-seq is challenging, as SNAs derived from scRNA-seq reads are extremely noisy and most studies have relied on the detection of chromosome-level copy number aberrations through smoothed gene expression profiles. Yet, as intratumor transcriptomic variation is partially driven by intratumor genetic variation, the classification of cells into subclones and the characterization of each subclone's genetic alterations should ideally be an integral step in any scRNA-seq analysis.

The appeal of subclone identification in scRNA-seq data is compounded by the shortage of technology

for sequencing the DNA and RNA molecules in the *same* cell with acceptable accuracy, throughput, and

cost. Although one can apply both scDNA-seq and scRNA-seq to a given cell population, the mutation

analysis and RNA quantification cannot be conducted in the same set of cells. Although there are now

technologies for deep targeted sequencing of select transcripts matched with same-cell whole

transcriptome sequencing [27, 28], these methods are still, in effect, profiling DNA-level variation by

sequencing expressed transcripts, and are thus subject to the technical issues, especially dropout due to

transcriptional stochasticity, that we modeled in DENDRO.

Subclone detection using scRNA-seq is difficult mainly because only a small portion of the SNAs of each

cell is expected to be seen in the read output of scRNA-seq. This is because to be sequenced, an SNA

needs to fall in a transcribed region of the genome, at a location within the transcript that will

eventually be read by the chosen sequencing protocol. Even for SNAs that satisfy these requirements,

the mutated allele are often missing in the read output due to *dropout*, especially in the heterozygous

case. This is due, in part, to the bursty nature of gene transcription in single cells [29-31], where in any

given cell, a substantial fraction of the genes are only expressed from one of the alleles. Thus, an SNA

residing in a gene that is expressed at the bulk tissue level may not be observed in a particular cell,

simply because the mutated allele, by chance, is not expressed in the given cell. We refer to alleles that

are not captured due to expression stochasticity as *biological dropouts*. Even for a mutated allele that is

expressed, it has to be successfully converted to cDNA and then sequenced to be represented in the

final read output; we refer to alleles lost due to technical reasons as *technical dropouts*. In addition to

dropout events, post-transcriptional modification, such as RNA editing, and sequencing errors impede

both the sensitivity and the specificity of SNA discovery. As a result, methods developed for single cell

SNA detection using scDNA-seq, such as Monovar [32], as well as methods designed for SNA detection in

bulk DNA or RNA sequencing data do not yield accurate results in the scRNA-seq setting [33-38].

Here we present a new statistical and computational framework – DNA based EvolutionNary tree

preDiction by scRNA-seq technOlogy (DENDRO) - that reconstructs the phylogenetic tree for cells

sequenced by scRNA-seq based on genetic divergence calculated from DNA-level mutations. DENDRO

assigns each cell to a leaf in the tree representing a subclone, and, for each subclone, infers its mutation

profile. DENDRO can detect genetically divergent subclones by addressing challenges unique to scRNA-

seq, including transcriptional variation and technical noise. A DENDRO clustering of scRNA-seq data

allows joint genetic and transcriptomic analysis on the same set of cells.

We evaluate DENDRO against existing approaches, through spike-in data sets and a metastasized renal

cell carcinoma dataset with known subpopulation labels, and show that DENDRO improved the accuracy

of subclone detection. We then demonstrate the DENDRO to biological discovery through two

applications. The first application profiles the treatment response in a melanoma model to immune

checkpoint blockade therapy. DENDRO identified a subclone that contracted consistently in response to

ICB therapy, and revealed that the contraction was driven by the high mutation burden and increased

availability of predicted neoantigens. Transcriptional divergence between the subclones in this model

was very weak, and thus the neoantigen-driven sub-clonal dynamics would not have been detected

without extracting DNA-level information. In the second application to a breast tumor dataset, DENDRO

detected subclones and allowed for the joint characterization of transcriptomic and genetic divergence

between cells in lymph-node metastasis and cells in primary resections.

The DENDRO package, implemented in R, is available at https://github.com/zhouzilu/DENDRO, where

we also provide a power calculation toolkit, DENDROplan, to aid in the design of scRNA-seq experiments

for subclonal mutation analysis using DENDRO.

Results

Method overview

Overview of DENDRO model and pipeline

Figure 1A shows an overview of DENDRO's analysis pipeline. Per cell counts of total read coverage (N

matrix) and mutation allele read coverage (X matrix) at SNA locations are extracted after read alignment

and SNA detection (details in Methods, Figure S1). Based on these matrices, DENDRO then computes a

cell-to-cell genetic divergence matrix, where entry (c, c') of the matrix is a measure of the genetic

divergence between cells c and c'. Details of this genetic divergence evaluation will be given in the next

section. DENDRO then clusters the cells into genetically distinct subclones based on this pairwise

divergence matrix, and selects the number of subclones based on inspection of the intra-cluster

divergence curve. Reads from the same subclone are then pooled together, and the SNA profile for each

subclone is re-estimated based on the pooled reads, which improves upon the previous SNA profiles

computed at the single cell level. Finally, DENDRO generates a parsimony tree using the subclone-level

mutation profiles to more accurately reflect the evolutionary relationship between the subclones.

Genetic divergence evaluation

Due to the high rates of biological and technical dropout, SNA detection within each individual cell lacks

sensitivity. We also expect low specificity due to the high base error rate in scRNA-seq protocols. Thus,

simple distance measures such as the Hamming or Euclidean distances evaluated on the raw SNA

genotype matrix or the raw allele frequency matrix do not accurately reflect the genetic divergence

between cells.

To more accurately estimate the cell-to-cell genetic divergence, we have developed a statistical model

that accounts for technical dropout, sequencing error and expression stochasticity. Consider two cells, c

and c', and let I_c and $I_{c'}$ index the clonal group to which the cells belong. That is, $I_c = I_{c'}$ if cells c and c'

come from the same subclone and thus share the same SNA profile. Let $X_c = (X_{c1}, \ldots, X_{cm})$ be the

mutation allele read counts for this cell at the m SNA sites profiled, and $N_c = (N_{c1}, \ldots, N_{cm})$ be the total

read counts at these sites. We define the genetic divergence between the two cells as

129
$$d_{cc'} = -\log P(X_c, X_{c'} | N_c, N_{c'}, I_c = I_{c'}) = \sum_{g=1}^{m} d_{cc'}^g$$

130
$$\text{where } d_{cc'}^g = -\log P\left(X_{cg}, X_{c'g} \middle| N_{cg}, N_{c'g}, I_c = I_{c'}\right).$$

In other words, $d_{cc'}$ is the negative log likelihood of the mutation allele counts of cells c and c', given

the total read counts and the event that the two cells belong to the same subclone. If c and c' have

mutations in mismatched positions, this likelihood for X_c, $X_{c'}$ conditioned on $I_c = I_{c'}$ would be small,

giving a large value for $d_{cc'}$. By the assumption of independence between sites, $d_{cc'}$ is the sum of $d_{cc'}^g$,

where $d_{cc'}^g$ is the contribution of mutation site g to the divergence measure. In characterizing the

conditional distribution for X_{cg} and $X_{c'g}$, we use a Beta-Binomial distribution to model expression

stochasticity and a Binomial model to capture sequencing errors and rare RNA-editing events. Referring

to Figure 1b, mutations residing in bursty genes, such as gene g, would tend to have U-shaped allele

frequency distributions and are more likely to be "dropped" due to low or zero expression. In contrary,

mutations residing in constitutive (non-bursty) genes, such as gene g' in Figure 1b, would have bell-

shaped allele frequency distributions and can be genotyped more reliably. Thus, even if the read counts

for the mutation loci residing in genes g and g' are identical across two cells (c_1 and c_2 in Figure 1b),

the locus in g' would contribute a higher value, compared to the locus in g, to the divergence between

cells c_1 and c_2. Please see Methods for details.

Accuracy assessment

 Accuracy assessment with in silico spike-in experiment

First, we designed a spike-in procedure to assess the accuracy of DENDRO versus existing approaches

and to make realistic power projections for subclone detection (Figure 2a). Since DENDRO is currently

the only method for SNA-based subclone detection using scRNA-seq data alone, we benchmarked

against more straightforward approaches such as hierarchical clustering based on mutation allele

frequencies. The spike-in procedure starts with an assumed evolutionary tree, where the leaves are

subclones and mutations can be placed on the branches. In the absence of prior information, a simple

tree structure is used, such as the one shown in Figure 2A. Parameters of simulation are (1) total

number of mutations, (2) total number of cells, (3) the proportion of cells in each clade, (4) the

proportion of mutations along each branch, and (5) mean read coverage across loci. Some of these

parameters can be determined using bulk DNA-seq and/or bulk RNA-seq data if available (Methods).

Parameters (1-4) determine the mutation profile matrix Z. To get the matrix of alternative and total

read counts for each mutation loci in each cell (X_{cg} and N_{cg}, respectively), we overlay the mutation

matrix onto the scRNA-seq data. This allows the spike-in framework to retain the expression

stochasticity and sequencing error of real scRNA-seq data. DENDRO is then applied to the read count

matrices to obtain the subclone clusters, which is then compared with the known labels. Accuracy is

evaluated by three metrics: adjusted Rand index, capture rate and purity (Supplementary Materials).

Such spike-in procedure can also facilitate experiment design, as it predicts the expected clustering

accuracy by DENDRO given sequencing parameters and available bulk data for the tumor (See

DENDROplan in Methods).

Using the above framework, we conducted a systematic evaluation of DENDRO's subclone detection

accuracy on an example scRNA-seq dataset with allelic information [39]. The results, compiled in Figure

2b, Supplementary Materials and Figure S3, show that DENDRO has better performance than simply

clustering on mutation allele frequencies and quantify how accuracy depends on the mutation burden,

mutation read depth, mutation distribution, subclone cell proportion, and cell populations. Even when

there are only 100 mutations with relatively low average coverage (read depth equals to 1), DENDRO

can still extract meaningful clustering results (average ARI $\approx$ 0.8). More importantly, variation in total

expression of genes does not influence DENDRO's divergence measure. DENDRO shows consistent

results in spike-in analysis between populations of single cell type and multiple cell types (Figure S3).

This is due to DENDRO's reliance only on the distribution of the mutation allele frequency conditioned

on the total read coverage, as illustrated by the simulation study (Supplementary Material and Figure

S2). The divergence evaluation reflects solely genetic distance not transcriptomic difference, allowing for

easy interpretation. A more extensive spike-in analysis can be found in the Supplementary Materials.

Accuracy assessment on a renal cell carcinoma and its metastasis

We also benchmarked DENDRO against existing methods on the renal cell carcinoma dataset from Kim

et al [21] (Figure 2C). This dataset contained 116 cells sequenced using the Smart-seq technology [17],

obtained from three tumors derived from one patient: a patient-derived xenograft (PDX) from the

primary renal cell carcinoma (PDX_pRCC), a biopsy of the metastasis to the lung 1 year after treatment

of primary site (Pt_mRCC), and a PDX of the lung metastasis renal cell carcinoma (PDX_mRCC) (Figure

S4A). The cells should share common early driver mutations due to their shared origin from the same

patient, but the metastasis and the cultivation of each tumor in separate medium (human or mouse)

should have allowed for the accumulation of new mutations. Thus, we expect the three tumors to be

clonally distinct. This knowledge allows us to use this dataset to benchmark accuracy and to illustrate

how DENDRO enables joint analysis of the genetic and transcriptomic heterogeneity at single cell

resolution.

We compared 4 different clustering methods: (1) DENDRO, (2) hierarchical clustering based on the

primary genotype matrix Z generated by GATK ($Z_{cg} = 1$ when a mutation g is detected for cell c, $Z_{cg} =$

0 otherwise), (3) hierarchical clustering based on the X/N matrix that preserve the variant allele

information and (4) hierarchical clustering based on gene expression ($\log TPM$). DENDRO gives the

cleanest separation between the three populations with adjusted Rand Index of 0.932 (1.0 indicates

perfect clustering, Figure 2C panel 1), as compared to 0.754 for Z matrix (Figure 2C panel 2), 0.519 for

X/N (Figure 2C panel 3) and 0.489 for expression (Figure 2C panel 4). Inspection of the tree shows that,

as expected, divergence between primary tumor and metastasis exceeds divergence between patient

sample and PDX sample, as PDX_mRCC clusters with Pt_mRCC rather than PDX_pRCC. All of the other

three methods successfully separated the primary sample from the metastatic samples, but could not

differentiate between the two metastasis samples.

For DENDRO, the intra-cluster divergence curve flattened at 3, and thus we stopped splitting at 3

clusters (Figure S4E and Methods). We annotated the clusters as PDX_mRCC, PDX_pRCC and Pt_mRCC

by their cell compositions (Table S3A). DENDRO found minimal sharing of subclones among the tumors

derived from three sources, and low genetic heterogeneity within each tumor. This is unsurprising since

relapsed metastasis consists of cells that have already undergone selection, and since the PDX tumors

are each seeded by a small subsample of cells from the original tumor, each tumor consists of unique

subclones not detected in other sites [40-42]. Additional joint analysis of transcriptome and DNA

mutations can be found in the Supplementary Materials.

DENDRO analysis of melanoma model in response to immune checkpoint blockade highlights

the role of neoantigens

Immune checkpoint blockade (ICB) of the inhibitory receptors CTLA4 and PD1 can result in durable

responses in multiple cancer types [43]. Features intrinsic to cancer cells that can impact ICB treatment

outcome include their repertoire of neoantigens [44], tumor mutational burden (TMB) [45], and

expression of PDL1 [46]. DENDRO analysis of scRNA-seq data allows joint DNA-RNA analysis of single

cells, thus enabling the simultaneous quantification of tumor mutational burden, the prediction of

neoantigen repertoire, and the characterization of gene expression profile at subclonal resolution. Thus,

to demonstrate the power of DENDRO and to better understand the relationship between ICB response

and intratumor heterogeneity, we profiled the single cell transcriptomes across three conditions derived

from 2 melanoma cell lines (Figure 3A): B16 melanoma cell line, which has shown modest initial

response to ICB treatment but eventually grows out, and Res 499 melanoma cell line (R499), which was

derived from a relapsed B16 tumor after combined treatment of radiation and anti-CTLA4 and is fully

resistant to ICB [47]. B16 was evaluated with and without anti-PD1 treatment, as we wanted a tumor

model that captures a transient ICB response. A total of 600 tumor cells were sequenced with Smart-seq

technology from six mice across three conditions: two mice with B16 without treatment (B16), two mice

with B16 after anti-PD1 treatment (B16PD1) and two mice with R499 without treatment (R499) (Figure

3A and Methods). The existence of multiple subclones in B16 and R499 was suggested by bulk WES

analysis [47, 48]. Our goal here is to determine whether the subclones differ in anti-PD1 response, and if

so, what are the subclonal differences.

A DENDRO analysis of 4059 putative mutation sites across 460 cells retained after QC (see Methods and

Figure S9A, B, C) yields the clustering displayed in Figure 3B, with four subclones suggested by the intra-

cluster divergence curve (Figure S9D). All subclones are shared among the three conditions, which is not

unexpected given that all tumor cells were derived from the same parental cell line. However, the sub-

clonal proportions vary significantly between conditions (Figure 3B). The subclonal proportions of

B16PD1 are approximately intermediate between that of B16 and R499. This is expected as R499 had

gone through immune editing whereas B16PD1, at the time of harvest, was still undergoing immune

editing and was at the transient response state. Furthermore, the selective pressure of radiation plus

anti-CTLA4 is likely more than that of anti-PD1 treatment, as the former but not the latter results in

complete responses in our B16 model [47]. The frequency of Clone 2 is lower in B16PD1 and R499,

indicating sensitivity to anti-PD1 treatment, while the frequencies of Clone 3 and Clone 4 increase after

treatment and are the highest in R499, indicating resistance to therapy (Figure 3C, S10A).

To explore why subclones vary in sensitivity to anti-PD1 treatment, we compared the mutation profile of

Clone 2 to the other subclones. We pooled cells in each of the four subclones and re-estimated their

mutation profiles, which were then used to construct a phylogenetic tree (Figure 3D). The phylogeny

suggests that Clone 3 and Clone 4 are genetically closer to each other than to Clone 2, and thus, their

similarity in treatment response may be in part due to similarity in their mutation profiles. The re-

estimated mutation profiles show that Clone 2 has the highest tumor mutation burden, which has been

associated with increased likelihood of ICB response [49, 50]. We then predicted the quantity of high-

affinity ($\leq$ 100 nm) neoantigens in each subclone given its mutation profile [48]. As shown in Figure 3E,

Clone 2 has twice as many high-affinity neoantigens as the other three subclones. The high level of

neoantigens can lead to better T cell recognition, resulting in increased efficacy of anti-PD1 treatment

[51].

Analysis of gene expression, on the other hand, did not yield detectable known signatures associated

with anti-PD1 treatment sensitivity. Projections based on the expression of highly variable genes, as

shown in PCA and t-SNE plots (Figure S8), did not yield meaningful clusters. Differential expression

analysis between each subclone and the other subclones found few genes with adjusted P-value < 0.05,

indicating similar expression across sub-clones that is concordant with the lack of structure in the

expression PCA and tSNE plots. Expressions of *Pdl1* (aka. *Cd274*) showed no differences between

subclones (KS-test: P-value > 0.42, Figure S10B). In addition, there were no detectable chromosome-

level differences in smoothed gene expression, indicating that there are no large CNV events that

distinguish the subclones (Figure S11). DENDRO, detecting exonic mutations from scRNA-seq data,

enabled the finding of subclones in this data, the prediction of neoantigen load of each subclone, and

the analysis of subclonal dynamics due to treatment. Our analysis suggests that the genetic

heterogeneity, rather than transcriptomic heterogeneity, contributes to treatment efficacy in this tumor

model.

Simultaneous analysis of genetic and transcriptomic variation in single cell breast cancer

We next applied DENDRO to the analysis of data from a study of primary and metastasized breast cancer

[20]. We focused on two tumors (BC03 and BC09) that had the most cells sequenced (Table S5 and

Figure S12). BC03 included cells from the primary tumor (here after BC03P) as well as cells from regional

metastatic lymph nodes (here after BC03LN), whereas BC09 included cells only from the primary

resection. 132 single cell transcriptomes were profiled by Smart-seq protocol [17]. We first assess

whether DENDRO separated BC03 and BC09, and then examine the transcriptomic and genetic

heterogeneity within each tumor.

GATK [52] detected a total of 2,364,823 mutation sites across the 132 cells, 353,647 passed QC and were

retained for downstream analysis (Figure S12A, B, C). Figure 4 shows the clustering determined by

DENDRO. DENDRO separates BC09 cells from BC03 cells with 100% accuracy (Figure 4A). The intra-

cluster divergence curve flattened at five subclones: three subclones for BC03 and two for BC09 (Figure

4A, Figure S12D and Table S3B). Within BC03, Clone Mix_1 and Clone Mix_2 contained a mixture of cells

from the primary tumor and lymph nodes, and Clone LN_1 contained mostly cells from the lymph nodes.

This suggests that tumor cells that have metastasized to the lymph nodes are genetically

heterogeneous, with some cells remaining genetically similar to the primary population and others

acquiring new genetic mutations. In comparison, hierarchical clustering based on expression (using log

transcripts-per-million values) did not separate BC03 from BC09, and gave a negative adjusted Rand

index within BC03, indicating effectively random assignment of cells to the two spatial locations (Figure

4B).

We then pooled cells within each of the 5 clusters and re-estimated their mutation profiles with

DENDRO. We defined a variant as subclonal if it was not present in all of the subclones within a tumor.

Based on detection marginal likelihood, we picked the top 10,000 most confident variants to construct a

phylogenetic tree (Figure 4C). As expected, the two BC09 clusters are far from the three BC03 clusters.

Within BC03, the length of the branches shows that the subclone containing mostly cells from lymph

nodes (labeled BC03LN_1) is genetically more similar to Clone Mix_2 compared to Clone Mix_1 (Figure

4C). In addition, window-smoothed expression plot with cells grouped by DENDRO clustering shows

broad chromosome-level shifts in expression patterns between subclones, most likely due to copy

number aberrations that are consistent with SNAs (Figure S13) [22].

A comparison of the transcriptomes of the subclones revealed substantial differences in the expression

of PAM50 genes, which are prognostic markers for breast cancer (Figure 4D) [53]. DENDRO detected one

rare subclone, BC09_2, with only six cells (<5% of the total number of cells) which had a strong basal-like

signature. Interestingly, in BC03, Clone LN_1 has the TNBC/basal-like subtype with an invasive gene

signature, while Clone Mix_2 has the $ESR1^+$ subtype. Thus, the genetic divergence of Clone LN_1 from

Clone Mix_2 is accompanied by its acquisition of an invasive metastatic expression signature. In a direct

comparison between cells from the primary site and cells from the lymph node without distinguishing

subclones, these expression differences would be much weaker since the subclones do not cleanly

separate by site. Compared with the original analysis that assigned each tumor to one specific breast

cancer subtype, this analysis identifies subclones with different expression phenotypes, potentially

allowing for better therapy design that targets all subclone phenotypes to reduce the risk of tumor

relapse.

Existing scRNA-seq studies of cancer tissue cluster cells based on total gene expression or copy number

profiles derived from smoothed total expression, making it difficult to separate the effects of sub-clonal

copy number aberrations from transcriptomic variation [19, 22, 24]. Differential expression analysis

based on clusters derived from total expression is prone to self-fulfilling prophecy, as there would

indeed be differentially expressed genes because this is the clustering criteria. Because DENDRO's

subclone identification is based solely on genetic divergence, and not on expression profile, the

downstream differential gene expression analysis can be precisely attributed to transcriptional

divergence between subclones.

Hence, we conducted a transcriptome-wide search for pathways that have differential expression

between subclones (Methods and Table S6), and assessed their overlap with pathways that are

differentially mutated between subclones. Focusing on tumor BC03, pathways for G2M checkpoint and

KRAS signaling are up-regulated in lymph node metastasis Clone BC03LN_1, while pathways for estrogen

response and apoptosis are down-regulated, indicating a more invasive phenotype (Table S6E). In

addition, *GAPDH* is up-regulated in the metastatic subclone (BC03LN_1) and down-regulated in the two

mix-cell subclones, consistent with previous findings [54, 55] (Figure S14D). Differentially expressed

genes between other subclone pairs in BC03 are also enriched in estrogen response, apoptosis, and DNA

repair (Table S6C, D). In parallel, subclone-specific mutated genes are highly enriched in cancer-related

pathways including MYC target, G2M checkpoints and mitotic spindle, and immune related pathways

such as, interferon response, TNF-a signaling and inflammatory response (Table S6). Interestingly, few of

the differentially mutated genes are associated with estrogen and androgen responses, suggesting that

the differential expression of hormone related genes is not mediated directly by genetic mutations in

these pathways. This is consistent with the recent studies that epigenetic alteration, such as histone

acetylation and methylation, regulate hormones receptor signaling in breast cancer [56-59]. DNA-RNA

joint analysis between other subclones are included in the Table S6 and Figure S14. Overall, this example

illustrates how DENDRO enables the joint assessment of genetic and transcriptomic contributions to

clonal diversity at single-cell resolution.

Discussion

We have described DENDRO, a statistical framework to reconstruct intratumor DNA-level heterogeneity

using scRNA-seq data. DENDRO starts with mutations detected directly from the scRNA-seq reads, which

are very noisy due to a combination of factors: (1) errors are introduced in reverse-transcription,

sequencing and mapping, (2) low sequencing depth and low molecule conversion efficiency leading to

technical dropouts, and (3) expression burstiness at the single cell level leading to biological dropouts.

DENDRO overcomes these obstacles through the statistical modeling of each component. Given noisy

mutation profiles and allele-specific read counts, DENDRO computes a distance between each pair of

cells that quantifies their genetic divergence after accounting for transcriptional bursting, dropout and

sequencing error. Then, DENDRO clusters the cells based on this distance as subclone and re-estimates a

more robust subclone-specific mutation profile by pooling reads across cells within the same cluster.

These re-estimated mutations profiles are then passed to downstream mutation analysis and

phylogenetic tree reconstruction.

Importantly, the genetic divergence used by DENDRO for cell clustering is based solely on allelic

expression ratios and do not reflect the difference in total expression between cells at mutation sites.

Thus, DENDRO differs from, and complements, existing tools that cluster cells based on total expression.

In fact, as shown by spike-in analysis, DENDRO clusters the cells based on true underlining mutation

profiles, and is robust to changes in total gene expression. As expected, the numbers of cells, the depth

of sequencing, the actual number of subclonal mutations and the phylogenetic tree structure all

influence the power of DENDRO. To aid researchers in experiment design, we developed DENDROplan,

which predicts DENDRO's clustering accuracy given basic experimental parameters and the expected

informative mutation count, which can be obtained from bulk DNA sequencing.

Ideally, joint sequencing of the DNA and RNA on the same cells would allow us to relate genomic profiles

to transcriptomic variations. Currently, there is yet no scalable technology for doing this. Separately

performing scDNA-seq and scRNA-seq on different batches of cells within the same tumor would meet

the nontrivial challenge of matching the subclones between the two data sets. DENDRO takes advantage

of the central dogma and utilizes computational methods to extract genetic divergence information

from noisy mutation calls in coding regions. Through two case studies, we illustrate the insights gained

from the subclonal mutation and expression joint analysis that DENDRO enables.

We have demonstrated that proper computational modeling can excavate the DNA-level heterogeneity

in scRNA-seq data. Yet, there is always limitation in working with RNA. While rare RNA editing events are

absorbed by the parameter ϵ, DENDRO cannot distinguish subcluster-specific constituent RNA editing

events from subcluster-specific DNA mutations. In the extreme and unlikely scenario where RNA editing

events are common and pervasive, DENDRO's cluster would reflect RNA editing. In such cases, we

recommend using matched bulk DNA-seq of the same tumor to filter the loci detected in the first step of

DENDRO, keeping only those that are supported by at least on read in the bulk DNA-seq data.

Conclusions

We have developed DENDRO, a statistical method for tumor phylogeny inference and clonal

classification using scRNA-seq data. DENDRO accurately infers the phylogeny relating the cells and

assigns each single cell from the scRNA-seq data set to subclone. DENDRO allows us to (1) cluster cells

based on genetic divergence while accounting for transcriptional bursting, technical dropout and

sequencing error, as benchmarked by in silico mixture and a spike-in analysis, (2) characterize the

transcribed mutations for each subclone, and (3) perform single-cell multi-omics analysis by examining

the relationship between transcriptomic variation and mutation profile with the same set of cells. We

evaluate the performance of DENDRO through a spike-in analysis and a data set with known subclonal

structure. We further illustrate DENDRO through two case studies. In the first case study of relationship

between intratumor heterogeneity and ICB treatment response, DENDRO estimates tumor mutation

burden and predicts repertoire of high-affinity neoantigens in each subclone from scRNA-seq. In the

second case study on a primary breast tumor dataset, DENDRO brought forth new insights on the

interplay between intratumor transcriptomic variation and subclonal divergence.

386 Methods

scRNA-seq alignment and SNA calling pipeline

Figure S1 illustrates the SNA calling pipeline. Raw scRNA-seq data is aligned by STAR 2-pass method,

which accounts for splicing junctions and achieve higher mapping quality [60]. In the next step, raw

variants calling is made using the Haplotype Caller (GATK tool) on the BAM files after sorting, joining

read groups, removing duplicated reads, removing overhangs into intronic regions, realigning and

recalibration [61]. Conventionally, there are two methods from GATK tools for mutation detection:

haplotype caller and mutect2. Haplotype caller has a RNA-seq setting which handle splice junctions

correctly, but assumes VAF around 50%, while mutect2 can detect mutations with low VAF but does

account for splice junction. The reason we select haplotype caller instead of mutect2 is that we extract

allele read counts for all cells as long as one of the cells is listed as carrying the mutation. Thus, as long

as one cell has VAF reaching 50%, this mutation would be detected. Calls with stand_call_conf greater

than 20 and population frequency greater than 5% but less than 95% were preserved for further

analysis. Admittedly, such lenient filtering typically introduces false positive sites. However, our priority

at this step is to minimize false negative rate, while the genetic divergence matrix in the following step

robustly estimates cell population substructure. Both the coverage of the alternative allele and the total

read coverage are extracted for each site for further analysis.

Genetic Divergence and Beta-Binomial framework

Consider two cells: c and c'. Let I_c and $I_{c'}$ denote the clonal group to which the cells belong, i.e. $I_c = I_{c'}$

if and only if cells c and c' come from the same subclone. We define the genetic divergence at loci

g, by $d_{cc'}^g$:

$$d_{cc'}^g = \log \frac{1}{P(X_{cg}, X_{c'g} | N_{cg}, N_{c'g}, I_c = I_{c'})}$$

$$= \log \frac{P(X_{cg}, X_{c'g} | N_{cg}, N_{c'g}, I_c = I_{c'}) + P(X_{cg}, X_{c'g} | N_{cg}, N_{c'g}, I_c \neq I_{c'})}{P(X_{cg}, X_{c'g} | N_{cg}, N_{c'g}, I_c = I_{c'})})$$

where $X_c = (X_{c1}, X_{c2}, \dots X_{cg}, \dots X_{cm})$ are the mutation allele read counts for cell c and $N_c =$

$(N_{c1}, N_{c2}, \dots N_{cg}, \dots N_{cm})$ are the total read counts at these sites. More intuitively, if cells c and c' are not

from the same clonal group, the numerator has larger value compared to denominator. Thus $d_{cc'}^g$ is

large, indicating bigger divergence between the two cells. With further derivation (Supplementary

Materials), $d_{cc'}^g$ is a function of the five following probabilities:

$d_{cc'}^g$

$= f\left(P_g; P(X_{cg}|N_{cg}, Z_{cg} = 0); P(X_{cg}|N_{cg}, Z_{cg} = 1); P(X_{c'g}|N_{c'g}, Z_{c'g} = 0); P(X_{c'g}|N_{c'g}, Z_{c'g} = 1)\right)$

where $Z_{cg} \in \{0,1\}$ is SNA indicator for cell c at site g and $P_g = P(Z_g = 1)$ is mutation frequency across

the cells estimated by GATK calls.

In the above formula for $d_{cc'}^g$, $P(X_{cg}|N_{cg}, Z_{cg} = 0)$ and $P(X_{c'g}|N_{cg}, Z_{cg} = 0)$ reflect reverse-

transcription/sequencing/mapping errors and rare RNA editing events, because when there is no

mutation (i.e. $Z_{cg} = 0$), all mutation reads reflect such technical errors or RNA editing. Let ϵ denote the

combined rate of technical error and RNA editing, we have

$$P(X_{cg}|N_{cg}, Z_{cg} = 0) \sim \text{Binomial}(X_{cg}|N_{cg}, \epsilon)$$

For cases where there are mutations (i.e. $Z_{cg} = 1$), the distribution of mutated read counts given total

read counts is modeled with a Beta Binomial distribution, which is capable of modeling technical

dropout and transcriptional bursting, and is supported by previous allele specific expression studies [30,

62] .

$$P(X_{cg}|N_{cg}, Z_{cg} = 1) \sim \int_0^1 \text{Binomial}(X_{cg}|N_{cg}, Q_{cg} = q) dF(q),$$

$$q \sim \text{Beta}(\alpha_g, \beta_g)$$

where Q_{cg} indicates proportion of mutated alleles expressed in cell c at site g, with Beta distribution as

prior. Respectively, α_g and β_g represent gene activation and deactivation rate, which are estimated

empirically.

Kernel based clustering and optimal cluster assignment

We cluster the cells using a kernel-based algorithm, such as hierarchical clustering. Given that there are

multiple sorting schemes, we leave the user to choose it. For the default-sorting scheme, we

recommend "ward.D" [63]. This is because $d_{cc'}$ behaves like a log likelihood ratio, which should follow a

χ^2 distribution when the two cells share the same subclone. The "ward.D" method has been shown to

work well in Euclidian space. Empirically, among different hierarchical clustering algorithms on the renal

cell carcinoma dataset (Figure S5) "ward.D" based hierarchical clustering performs the best.

To determine the number of clusters we use an intra-cluster divergence curve computed from the

divergence matrix. Existing software rely on AIC, BIC, or another model selection metric [64, 65].

However, since we only have the "distance" matrix, these traditional methods cannot be applied. Let N_k

be the number of cell pairs in cluster C_k and N be the total number of pairs between cells for all

clusters. Let K be the number of clusters. The weighted sum of intra-cluster distance W_K is

$$W_K = \sum_{k=1}^{K} N_k \sum_{(i,j)\in C_k} \frac{d_{ij}}{N}$$

Note that small clusters are naturally down-weighted in the above metric. DENDRO relies on visual

examination of the intra-cluster divergence curve (W_K plotted against K) to find the "elbow point",

which can be taken as a reasonable clustering resolution.

Power analysis toolkit and experimental design

Before conducting a single cell RNA-seq experiment on a tumor sample, it is important to project how

subclone detection power depends on the number of cells sequenced and the coverage per cell. To

facilitate experiment design, we have developed a tool, DENDROplan (Figure 2a), that predicts the

expected clustering accuracy by DENDRO given sequencing parameters and available bulk data for the

tumor. Given an assumed tree structure and a target accuracy, DENDROplan computes the necessary

read depth and number of cells needed.

As shown in Figure 2a, if bulk DNA sequencing and/or RNA sequencing data are available for the tumor

being studied, these data can be harnessed to make more realistic power calculations. For example, if

SNAs have been profiled using bulk DNA sequencing data, the set of mutations that lie in the exons of

annotated genes can be retrieved and used directly in constructing the spike-in data. Furthermore,

phylogeny construction algorithms for bulk DNA-seq data can be used to infer a putative tree structure

that can be used as input to DENDROplan [5, 65]. If bulk RNA-seq data is available, the bulk expression

level of the mutation-carrying genes can be used to predict the expression level of the mutation in the

single cell data. For details, see Methods. The power analysis tool is also available at

https://github.com/zhouzilu/DENDRO.

Spike-in analysis

In our spike-in analysis, we adopt a dataset from Deng et al, which has allele specific read counts [39].

We further added binomial noise to mimic sequencing error. When analyzing DENDRO performance in

terms of various number mutation site, number of cells, proportion of cells in each clade and proportion

of mutations along each branch, we only take a subset of cells with similar cell types. On the other hand,

we utilize a mixture cell population to test the robustness of DENDRO performance with regard to

various expression profiles.

SNA inference in "bulk" and phylogenetic tree construction

As stated previously, DENDRO further inferred SNA after pooling the reads from all cells within each

cluster. Because, with our choice of thresholds, we identify SNAs in single cells with high sensitivity, the

"bulk" level SNAs should be a subset of the SNAs in single cells, and mutation allele counts and total

allele counts should provide us with enough information for SNA detection using a maximum likelihood

framework [66], which accounts for both sequencing error and rare RNA-editing events. Suppose g is

the genotype (number of reference allele) at a site and assume m, the ploidy, equals to 2. Then the

likelihood is:

484
$$\mathcal{L}(g) = \frac{1}{m^k} \prod_{j=1}^{l} [(m-g)\epsilon + g(1-\epsilon)] \prod_{j=l+1}^{k} [(m-g)(1-\epsilon) + g\epsilon]$$

485
$$g^* = \underset{g}{\operatorname{argmax}} -\mathcal{L}(g)$$

where k is number of reads at a site and the first l bases ($l \leq k$) be the same to reference and the rests

are same to alternative allele. ϵ is the sequencing error and rare RNA-editing combined rate. Given

mutation profiles, DENDRO then constructs a phylogenetic tree with the neighbor-joining method,

which can more accurately capture the evolutionary relationship between different subclones [67] than

the initial tree given by hierarchical clustering.

Differential gene expression, mutation annotation and gene ontology analysis

We use Seurat and scDD to identify differentially expressed genes between tumors and between tumor

subclones [68-70]. For each comparison, we apply two different methods: MAST implemented by Seurat

and scDD. Genes with adjusted p-value < 0.05 count as significant differentially expressed gene for each

method. We further intersect these two sets of differentially expressed genes to increase robustness.

Subclonal mutations are annotated by ANNOVAR with default parameters and variants associated with

intergenic regions were discarded for downstream analysis [71]. For GO analysis, we apply Gene Set

Enrichment Analysis tool [52]. Hallmark gene sets serve as fundamental database with FDR q-value <

0.05 as significant.

Single cell RNA-seq of Tumor Model Derived from B16.

Six C57bl/6 mice were injected on both flanks with either B16 or R499: four with B16 and two with R499.

Two of the mice implanted with B16 were treated with 200 ug of anti-PD1 per mouse on Days 5, 8 and

11. On Day 15, all tumors were harvested and made into single cell suspension. 100,000 CD45 negative

tumor cells were sorted on Aria to enrich for live tumor cells and loaded on SMARTer ICELL8 cx Single-

Cell System prior to full length single cell RNA-sequencing library preparation using Smart-seq following

manufacturer's recommendations. 460 cells and 11531 genes passed standard QC and were retained for

downstream analysis.

Neoantigen prediction.

Based on gene expression from RNA-seq data, variants from unexpressed transcripts are removed. The

MHC-I binding affinities of variants are then predicted using NetMHC version 4.0 for H-2-Kb and H-2-Db

using peptide lengths from 8 to 11 [72]. Given subclonal mutation profile, we further assign the

neoantigens to each subclone.

List of abbreviations

SNA: Single nucleotide alteration

scDNA-seq: Single-cell DNA sequencing

scRNA-seq: Single-cell RNA sequencing

PDX: Patient-derived xenograft

TPM: Transcripts per kilobase million

ICB: Immune checkpoint blockade

TMB: Tumor mutational burden

Declarations

Acknowledgements

We thank Dr. Nan Lin for informing useful tools for biological insight analysis, and Dr. Kai Tan and Dr.

Mingyao Li for helpful comments and suggestions.

Funding

This work was supported by National Institutes of Health (NIH) grant 1P01CA210944-01 to AM and BX,

5R01-HG006137-07 to ZZ and to NRZ, and 1U2CCA233285-01 to NRZ.

Availability of data and materials

DENDRO is an open-source R package available at https://github.com/zhouzilu/DENDRO with license

GPL-3.0.

Authors' contributions

ZZ and NRZ formulated the model. ZZ developed and implemented the algorithm and conducted all

computational analyses. BX and AM designed the immunotherapy case study. BX performed the

experiments of the immunotherapy case. ZZ, BX and NRZ wrote the manuscript. All authors read and

approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Consent for publication

Not applicable.

Ethics approval and consent to participate

Not applicable.

1.   Gamazon, E.R. and B.E. Stranger, *The impact of human copy number variation on gene*
expression. Briefings in Functional Genomics, 2015. **14**(5): p. 352-357.
2.   Hanks, S., et al., *Constitutional aneuploidy and cancer predisposition caused by biallelic*
mutations in BUB1B. Nat Genet, 2004. **36**(11): p. 1159-61.
3.   Vicente-Duenas, C., et al., *Epigenetic Priming in Cancer Initiation.* Trends Cancer, 2018.
4(6): p. 408-417.
4.   Burrell, R.A., et al., *The causes and consequences of genetic heterogeneity in cancer*
evolution. Nature, 2013. **501**(7467): p. 338-45.

5.   Jiang, Y., et al., *Assessing intratumor heterogeneity and tracking longitudinal and spatial*
clonal evolutionary history by next-generation sequencing. Proc Natl Acad Sci U S A,
2016. **113**(37): p. E5528-37.
6.   Deshwar, A.G., et al., *PhyloWGS: reconstructing subclonal composition and evolution*
from whole-genome sequencing of tumors. Genome Biol, 2015. **16**: p. 35.
7.   Zare, H., et al., *Inferring clonal composition from multiple sections of a breast cancer.*
PLoS Comput Biol, 2014. **10**(7): p. e1003703.
8.   Carter, S.L., et al., *Absolute quantification of somatic DNA alterations in human cancer.*
Nat Biotechnol, 2012. **30**(5): p. 413-21.
9.   Li, B. and J.Z. Li, *A general framework for analyzing tumor subclonality using SNP array*
and DNA sequencing data. Genome Biol, 2014. **15**(9): p. 473.
10.   Oesper, L., A. Mahmoody, and B.J. Raphael, *THetA: inferring intra-tumor heterogeneity*
from high-throughput DNA sequencing data. Genome Biol, 2013. **14**(7): p. R80.
11.   Ha, G., et al., *TITAN: inference of copy number architectures in clonal cell populations*
from tumor whole-genome sequence data. Genome Res, 2014. **24**(11): p. 1881-93.
12.   Miller, C.A., et al., *SciClone: inferring clonal architecture and tracking the spatial and*
temporal patterns of tumor evolution. PLoS Comput Biol, 2014. **10**(8): p. e1003665.
13.   Navin, N.E., *The first five years of single-cell cancer genomics and beyond.* Genome Res,
2015. **25**(10): p. 1499-507.
14.   Navin, N., et al., *Tumour evolution inferred by single-cell sequencing.* Nature, 2011.
472(7341): p. 90-4.
15.   Wang, Y., et al., *Clonal evolution in breast cancer revealed by single nucleus genome*
sequencing. Nature, 2014. **512**(7513): p. 155-60.
16.   Gao, R., et al., *Punctuated copy number evolution and clonal stasis in triple-negative*
breast cancer. Nat Genet, 2016. **48**(10): p. 1119-30.
17.   Picelli, S., et al., *Smart-seq2 for sensitive full-length transcriptome profiling in single cells.*
Nat Methods, 2013. **10**(11): p. 1096-8.
18.   Klein, A.M., et al., *Droplet barcoding for single-cell transcriptomics applied to embryonic*
stem cells. Cell, 2015. **161**(5): p. 1187-1201.
19.   Patel, A.P., et al., *Single-cell RNA-seq highlights intratumoral heterogeneity in primary*
glioblastoma. Science, 2014. **344**(6190): p. 1396-401.
20.   Chung, W., et al., *Single-cell RNA-seq enables comprehensive tumour and immune cell*
profiling in primary breast cancer. Nat Commun, 2017. **8**: p. 15081.
21.   Kim, K.T., et al., *Application of single-cell RNA sequencing in optimizing a combinatorial*
therapeutic strategy in metastatic renal cell carcinoma. Genome Biol, 2016. **17**: p. 80.
22.   Tirosh, I., et al., *Dissecting the multicellular ecosystem of metastatic melanoma by*
single-cell RNA-seq. Science, 2016. **352**(6282): p. 189-96.
23.   Jerby-Arnon, L., et al., *A Cancer Cell Program Promotes T Cell Exclusion and Resistance to*
Checkpoint Blockade. Cell, 2018. **175**(4): p. 984-997 e24.
24.   Tirosh, I., et al., *Single-cell RNA-seq supports a developmental hierarchy in human*
oligodendroglioma. Nature, 2016. **539**(7628): p. 309-313.
25.   Venteicher, A.S., et al., *Decoupling genetics, lineages, and microenvironment in IDH-*
mutant gliomas by single-cell RNA-seq. Science, 2017. **355**(6332).

26. Li, H., et al., *Reference component analysis of single-cell transcriptomes elucidates*
cellular heterogeneity in human colorectal tumors. Nat Genet, 2017. **49**(5): p. 708-718.
27. van Galen, P., et al., *Single-Cell RNA-Seq Reveals AML Hierarchies Relevant to Disease*
Progression and Immunity. Cell, 2019. **176**(6): p. 1265-1281 e24.
28. Nam, A.S., et al., *Somatic mutations and cell identity linked by Genotyping of*
Transcriptomes. Nature, 2019. **571**(7765): p. 355-360.
29. Raj, A., et al., *Stochastic mRNA synthesis in mammalian cells.* PLoS Biol, 2006. **4**(10): p.
e309.
30. Jiang, Y., N.R. Zhang, and M. Li, *SCALE: modeling allele-specific gene expression by single-*
cell RNA sequencing. Genome Biol, 2017. **18**(1): p. 74.
31. Padovan-Merhar, O., et al., *Single mammalian cells compensate for differences in*
cellular volume and DNA copy number through independent global transcriptional
mechanisms. Mol Cell, 2015. **58**(2): p. 339-52.
32. Zafar, H., et al., *Monovar: single-nucleotide variant detection in single cells.* Nat
Methods, 2016. **13**(6): p. 505-7.
33. Piskol, R., G. Ramaswami, and J.B. Li, *Reliable identification of genomic variants from*
RNA-seq data. Am J Hum Genet, 2013. **93**(4): p. 641-51.
34. Brennecke, P., et al., *Accounting for technical noise in single-cell RNA-seq experiments.*
Nat Methods, 2013. **10**(11): p. 1093-5.
35. Pierson, E. and C. Yau, *ZIFA: Dimensionality reduction for zero-inflated single-cell gene*
expression analysis. Genome Biol, 2015. **16**: p. 241.
36. Vallejos, C.A., J.C. Marioni, and S. Richardson, *BASiCS: Bayesian Analysis of Single-Cell*
Sequencing Data. PLoS Comput Biol, 2015. **11**(6): p. e1004333.
37. Ding, B., et al., *Normalization and noise reduction for single cell RNA-seq experiments.*
Bioinformatics, 2015. **31**(13): p. 2225-7.
38. Qiu, X., et al., *Single-cell mRNA quantification and differential analysis with Census.* Nat
Methods, 2017. **14**(3): p. 309-315.
39. Deng, Q., et al., *Single-cell RNA-seq reveals dynamic, random monoallelic gene*
expression in mammalian cells. Science, 2014. **343**(6167): p. 193-6.
40. Eirew, P., et al., *Dynamics of genomic clones in breast cancer patient xenografts at*
single-cell resolution. Nature, 2015. **518**(7539): p. 422-6.
41. Gerlinger, M., et al., *Intratumor heterogeneity and branched evolution revealed by*
multiregion sequencing. N Engl J Med, 2012. **366**(10): p. 883-892.
42. Shi, Y.J., et al., *Intratumoral Heterogeneity in Breast Cancer: A Comparison of Primary*
and Metastatic Breast Cancers. Oncologist, 2017. **22**(4): p. 487-490.
43. Ribas, A. and J.D. Wolchok, *Cancer immunotherapy using checkpoint blockade.* Science,
2018. **359**(6382): p. 1350-1355.
44. Schumacher, T.N. and R.D. Schreiber, *Neoantigens in cancer immunotherapy.* Science,
2015. **348**(6230): p. 69-74.
45. Rizvi, N.A., et al., *Cancer immunology. Mutational landscape determines sensitivity to*
PD-1 blockade in non-small cell lung cancer. Science, 2015. **348**(6230): p. 124-8.
46. Tumeh, P.C., et al., *PD-1 blockade induces responses by inhibiting adaptive immune*
resistance. Nature, 2014. **515**(7528): p. 568-71.

47. Twyman-Saint Victor, C., et al., *Radiation and dual checkpoint blockade activate non-*
redundant immune mechanisms in cancer. Nature, 2015. **520**(7547): p. 373-7.
48. Benci, J.L., et al., *Opposing Functions of Interferon Coordinate Adaptive and Innate*
Immune Responses to Cancer Immune Checkpoint Blockade. Cell, 2019. **178**(4): p. 933-
948 e14.
49. Patel, S.A. and A.J. Minn, *Combination Cancer Therapy with Immune Checkpoint*
Blockade: Mechanisms and Strategies. Immunity, 2018. **48**(3): p. 417-433.
50. Goodman, A.M., et al., *Tumor Mutational Burden as an Independent Predictor of*
Response to Immunotherapy in Diverse Cancers. Mol Cancer Ther, 2017. **16**(11): p. 2598-
2608.
51. Rosenthal, R., et al., *Neoantigen-directed immune escape in lung cancer evolution.*
Nature, 2019. **567**(7749): p. 479-485.
52. Subramanian, A., et al., *Gene set enrichment analysis: a knowledge-based approach for*
interpreting genome-wide expression profiles. Proc Natl Acad Sci U S A, 2005. **102**(43): p.
15545-50.
53. Parker, J.S., et al., *Supervised risk predictor of breast cancer based on intrinsic subtypes.* J
Clin Oncol, 2009. **27**(8): p. 1160-7.
54. Zhang, J.Y., et al., *Critical protein GAPDH and its regulatory mechanisms in cancer cells.*
Cancer Biol Med, 2015. **12**(1): p. 10-22.
55. Tarrado-Castellarnau, M., et al., *Glyceraldehyde-3-phosphate dehydrogenase is*
overexpressed in colorectal cancer onset. Translational Medicine Communications, 2017.
2(1): p. 6.
56. Mann, M., V. Cortez, and R.K. Vadlamudi, *Epigenetics of estrogen receptor signaling: role*
in hormonal cancer progression and therapy. Cancers (Basel), 2011. **3**(3): p. 1691-707.
57. Green, K.A. and J.S. Carroll, *Oestrogen-receptor-mediated transcription and the influence*
of co-factors and chromatin state. Nat Rev Cancer, 2007. **7**(9): p. 713-22.
58. Dreijerink, K.M., et al., *Menin links estrogen receptor activation to histone H3K4*
trimethylation. Cancer Res, 2006. **66**(9): p. 4929-35.
59. Kim, H., et al., *Requirement of histone methyltransferase SMYD3 for estrogen receptor-*
mediated transcription. J Biol Chem, 2009. **284**(30): p. 19867-77.
60. Dobin, A., et al., *STAR: ultrafast universal RNA-seq aligner.* Bioinformatics, 2013. **29**(1):
p. 15-21.
61. McKenna, A., et al., *The Genome Analysis Toolkit: a MapReduce framework for analyzing*
next-generation DNA sequencing data. Genome Res, 2010. **20**(9): p. 1297-303.
62. Skelly, D.A., et al., *A powerful and flexible statistical framework for testing hypotheses of*
allele-specific gene expression from RNA-seq data. Genome Res, 2011. **21**(10): p. 1728-
37.
63. Ward, J.H., *Hierarchical Grouping to Optimize an Objective Function.* Journal of the
American Statistical Association, 1963. **58**(301): p. 236-&.
64. Goutte, C., et al., *Feature-space clustering for fMRI meta-analysis.* Hum Brain Mapp,
2001. **13**(3): p. 165-83.
65. Urrutia, E., et al., *Integrative pipeline for profiling DNA copy number and inferring tumor*
phylogeny. Bioinformatics, 2018. **34**(12): p. 2126-2128.

66.   Li, B., et al., *A likelihood-based framework for variant calling and de novo mutation
685   detection in families.* PLoS Genet, 2012. **8**(10): p. e1002944.
686   67.   Schliep, K.P., *phangorn: phylogenetic analysis in R.* Bioinformatics, 2011. **27**(4): p. 592-3.
687   68.   Finak, G., et al., *MAST: a flexible statistical framework for assessing transcriptional
688   changes and characterizing heterogeneity in single-cell RNA sequencing data.* Genome
689   Biol, 2015. **16**: p. 278.
690   69.   Korthauer, K.D., et al., *A statistical approach for identifying differential distributions in
691   single-cell RNA-seq experiments.* Genome Biol, 2016. **17**(1): p. 222.
692   70.   Butler, A., et al., *Integrating single-cell transcriptomic data across different conditions,
693   technologies, and species.* Nat Biotechnol, 2018. **36**(5): p. 411-420.
694   71.   Wang, K., M. Li, and H. Hakonarson, *ANNOVAR: functional annotation of genetic
695   variants from high-throughput sequencing data.* Nucleic Acids Res, 2010. **38**(16): p.
696   e164.
697   72.   Karosiene, E., et al., *NetMHCcons: a consensus method for the major histocompatibility
698   complex class I predictions.* Immunogenetics, 2012. **64**(3): p. 177-86.
699

Figure 1. DENDRO analysis pipeline and genetic divergence evaluation. a DENDRO analysis pipeline overview. **b** Statistical model for genetic divergence evaluation function. In this example, we show relationship of cell c_1 and c_2 with gene g and g' in terms of their genetic divergence. Middle panels shows the stochasticity of cell expression and bottom panels illustrates the distribution of variants allele frequency across cells for these two genes, indicating that gene g is a bursty gene and g' is a constitutive gene.

Figure 2. DENDRO accuracy assessment. a The overall spike-in analysis pipeline. Three statistics measure the accuracy of DENDRO - adjusted Rand index (global accuracy, 1 indicating perfect classification), capture rate (subclone specific accuracy, out of all cells from the clade, how many of them are correctly assigned) and purity (subclone specific accuracy, out of all cells from the detected cluster, how many are actually from the clade). **b** Cluster accuracy via spike-in studies.Various parameters show effects on cluster accuracy (measured by adjusted Rand index) based on tree structure on the right. Left panel: effect of mutation read depth. Right panel: effect of total mutation burden. **c** Evaluation of DENDRO on a renal cell carcinoma and its metastasis. (Left to right) (1) DENDRO clustering result from primary and metastatic renal cell carcinoma dataset. Background colors represent DEDRO clustering result. (2) Clustering of the same dataset using Z matrix (indicator matrix, $z_{ij}=1$ when detected a mutation for cell i at site j by GATK tool). (3) Clustering of the same dataset using X/N matrix (mutation allele frequency matrix) (4) Clustering of the same dataset using expression (logT-PM).

Figure 3. Clonal composition alternations with anti-PD1 treatments and cell lines. a Experimental overview. For each condition at Day 15, we have two biological replicates. There are total 600 cells from 6 tumors sequenced. **b** DENDRO cluster result. No clone is exclusively associated with any tumor condition. **c** Frequencies of the subclonal populations in B16, B16PD1 and R499. **d** Neighbor joining phylogenetic tree given detected subclones. **e** Number of high affinity neoantigen predicted for each clone. Clone 2 have the highest number of neoantigen.

Figure 4. Analysis of scRNA-seq dataset of primary breast cancer. a DENDRO cluster result for primary breast cancer dataset (Chung et al., 2017). **b** Hierarchical clustering result for the same dataset based on expression (*log*TPM). (dashlines indicate cluster boundaries). **c** Neighbor joining phylogenetic tree given detected subclones for breast cancer dataset. **d** PAM50 gene panel expression shows breast cancer subtypes of each subclone.