Genetic Heterogeneity Profiling by Single Cell RNA Sequencing

Zilu Zhou, Graduate Group in Genomics and Computational Biology, University of Pennsylvania,

Philadelphia, PA, USA. zhouzilu@pennmedicine.upenn.edu

Nancy R Zhang*, Department of Statistics, The Wharton School, University of Pennsylvania, Philadelphia,

PA, USA. nzh@wharton.upenn.edu

* Corresponding author

Abstract

Detection of genetically distinct subclones and profiling the transcriptomic differences between subclones is important for studying the evolutionary dynamics of tumors, as well as for accurate prognosis and effective treatment of cancer in the clinic. Single cell RNA-sequencing (scRNA-seq) is now ubiquitously adopted in ongoing and planned cancer studies for the profiling of intra-tumor transcriptional heterogeneity. Inference of clonal membership from scRNA-seq is currently unreliable. We propose DENDRO, an analysis method for scRNA-seq data that detects genetically distinct subclones, assigns each single cell to a subclone, and reconstructs the evolutionary tree. DENDRO utilizes information from single nucleotide mutations in transcribed regions and accounts for single-cell level expression stochasticity and technical noise. The accuracy of DENDRO is benchmarked on spike-in datasets and on scRNA-seq data with known subpopulation structure. We applied DENDRO to two scRNA-seq data sets, where the new approach allowed us to elucidate the interplay between genetic and transcriptomic intratumor variation.

Keywords

single-cell RNA sequencing, intratumor heterogeneity, clonal reconstruction, statistical modeling

Background

Tumor initiation, progression, metastasis and relapse are mediated by DNA alterations[1]. Single nucleotide alteration (SNA) is a basic type of DNA alteration that can affect the transcriptome and

contribute to intratumor heterogeneity [2]. Intratumor heterogeneity is a key mediator of patients'

response to treatment, as clonal selection can lead to the emergence of subclones that are drug

resistant [3]. Thus, identifying subclonal DNA alterations and assessing their impact on subclonal

transcriptional dynamics can elucidate the mechanisms of tumor evolution and determine potential

gene targets for therapy. To characterize intratumor genetic heterogeneity, most prior studies have

used bulk tumor DNA sequencing [4-11], but these approaches have limited resolution and power [12].

Breakthroughs in single-cell genomics promise to reshape cancer research by allowing comprehensive

cell type classification and rare subclone identification. For example, in breast cancer, single-cell DNA

sequencing (scDNA-seq) was used to distinguish normal cells from malignant cells, the latter of which

were further classified into subclones [13-15]. However, scDNA-seq still suffers from low coverage,

intensive labor and low consistency, hindering its widespread adoption in the cancer field [16]. Single-

cell RNA sequencing (scRNA-seq), on the other hand, is comparatively more mature and cost effective.

ScRNA-seq technologies such as Smart-seq2 [17], Drop-seq [18], and 10X Genomics Chromium^{TM}, which

can simultaneously profile thousands of cells with reasonable cost, have advanced tumor transcriptome

profiling to single cell resolution [19]. Previous single-cell studies revealed novel insights into the

developmental and immunological programs of gene expression in glioblastoma and breast cancer [20,

21]. Kim et al. and Tirosh et al. used scRNA-seq to elucidate drug target pathways and drug response

variability across samples and tumor metastasis [22, 23]. More recently, studies have focused on tumor

cellular architecture and tumor cell lineage in gliomas, leading to new expression signatures that are

predictive of prognosis [24-26]. However, despite the recognized importance of clonal DNA mutations in

tumor pathogenesis, this crucial aspect has not been explored in detail by these scRNA-seq studies, due

to the lack of appropriate computational methods for analyzing DNA-level mutation heterogeneity using

scRNA-seq data.

Only a small portion of the SNAs of each cell are expected to be seen in the read output of scRNA-seq.

To be sequenced, a SNA needs to fall in a transcribed region of the genome, at a location within the

transcript that will eventually be read. Gene transcription in single cells is typically bursty [27-29], and

thus an SNA residing in a gene that is expressed at the bulk tissue level may not be observed in a

particular cell, simply because the mutated allele, by chance, is not expressed in the given cell. We refer

to alleles that are not captured due to expression stochasticity as *biological dropouts*. Even if a mutated

allele is expressed, it has to be successfully converted to cDNA and then sequenced to be represented in

the final read output; we refer to alleles lost at this step as *technical dropouts*. In addition to dropout

events, post-transcriptional modification, such as RNA editing, and sequencing errors impede both

sensitivity and specificity of SNA discovery. As a result, methods developed for single cell SNA detection

using scDNA-seq, such as Monovar [30], as well as methods designed for SNA detection in bulk DNA or

RNA sequencing data do not yield accurate results in the scRNA-seq setting [31-36].

Intratumor genetic heterogeneity gives rise to intratumor transcriptomic heterogeneity, and cell-to-cell

transcriptional variation is driven by inter-clonal competition. Thus, in the analysis of scRNA-seq data of

tumor samples, the accurate classification of cells into clones and the characterization of each clone's

genetic alterations should be an integral step. The potential of identifying subclonal DNA mutations

using scRNA-seq data and the prospect of using these alterations for clonal inference has not been

extensively explored. This analysis gap is compounded by the lack of technology for profiling DNA

mutations and RNA expression in the *same* cell with acceptable accuracy, throughput, and cost.

Although one can apply scDNA-seq and scRNA-seq to the same tumor, the mutation analysis and RNA

quantification could not be conducted in the same set of cells.

Here we develop a new statistical and computational framework – DNA based EvolutionNary tree

preDiction by scRNA-seq technOlogy (DENDRO) - that reconstructs the phylogenetic tree for cells

sequenced by scRNA-seq based on genetic divergence calculated from DNA-level mutations. DENDRO

assigns each cell to a clone in the tree, and, for each clone, infers its mutation profile. DENDRO achieves

high power to detect genetically divergent subclones by addressing challenges unique to scRNA-seq,

including transcriptional variation and technical noise. A DENDRO clustering of scRNA-seq data allows

joint genetic and transcriptomic analysis on the same set of cells. We evaluate the accuracy of DENDRO

through spike-in benchmark data sets, and then illustrate its application through two case studies. In the

first case study on a metastasized renal cell carcinoma dataset, we validate DENDRO's results using

known subpopulation labels, and further elucidate the genetic-transcriptomic changes in a PDX tumor

model. In the second case study on a primary breast tumor dataset, DENDRO detects unbiased

subclones and jointly characterizes the transcriptomic and genetic changes between cells in lymph

nodes and primary resections, illustrating the clinical essence of studying intratumor heterogeneity. The

DENDRO package, implemented in R, is available at https://github.com/zhouzilu/DENDRO, where we

also provide a power calculation toolkit, DENDROplan, to aid in the design of scRNA-seq experiments for

subclonal mutation analysis using DENDRO.

98   Results

Overview of DENDRO model and pipeline

Figure 1A shows an overview of DENDRO's analysis pipeline. Input data can be in fastq or BAM format.

For fastq files, we recommend STAR 2-pass method for alignment, which accounts for splicing junctions

and achieve higher mapping quality [37]. In the next step, GATK tools are used for SNA detection [38];

parameters for this step are chosen to allow high sensitivity but low specificity, since the SNA loci will be

further refined in downstream analysis (see Methods and Figure S1 for details). Per cell counts of total

read coverage (N matrix) and mutation allele read coverage (X matrix) at SNA locations are extracted.

Based on these matrices, DENDRO then computes a cell-to-cell genetic divergence matrix, where entry

(c, c') of the matrix is a measure of the genetic divergence between cells c and c'. Details of this genetic

divergence evaluation are given in the next section. DENDRO next clusters the cells into genetically

distinct clones based on this pairwise divergence matrix, and selects the number of clones based on

inspection of the intra-cluster divergence curve. Reads from the same clone are then pooled together,

and the SNA profile for each clone is re-estimated based on the pooled reads, significantly improving

upon the previous SNA profiles computed at the single cell level. Finally, DENDRO generates a parsimony

tree using the clone-level mutation profiles, to more accurately reconstruct their evolutionary

relationship.

Genetic divergence evaluation

Due to the high rate of biological and technical dropout, SNA detection by GATK within each cell lacks

sensitivity. We also expect a high false positive rate due to the known high base error rate in scRNA-seq

protocols. Given this, simple approaches such as the Hamming or Euclidean distance computed using

the raw SNA genotype matrix or the allele frequency matrix do not accurately measure the genetic

divergence between cells.

To more accurately estimate the cell-to-cell genetic divergence, we developed a statistical model that

accounts for technical dropout, sequencing error and expression stochasticity. Consider two cells, c

and c', and let I_c and $I_{c'}$ index the clonal group to which the cells belong. That is, $I_c = I_{c'}$ if cells c and c'

come from the same subclone and thus share the same SNA profile. Let $X_c = (X_{c1}, \dots, X_{cm})$ be the

mutation allele read counts for this cell at the m SNA sites profiled, and let $N_c = (N_{c1}, \ldots, N_{cm})$ be the

total read counts at these sites. We define the genetic divergence between the two cells as

$$d_{cc'} = \log P(X_c, X_{c'} | N_c, N_{c'}, I_c = I_{c'}) = \sum_{g=1}^{m} d_{cc'}^g$$

$$\text{where } d_{cc'}^g = -\log P(X_{cg}, X_{c'g} | N_{cg}, N_{c'g}, I_c = I_{c'})$$

In other words, $d_{cc'}$ is the negative log likelihood of the mutation allele counts of cells c and c', given

the total read counts and the event that the two cells belong to the same clone. If c and c' have

mutations in mismatched positions, the likelihood for $X_c, X_{c'}$ conditioned on $I_c = I_{c'}$ would be small,

giving a large value for $d_{cc'}$. By the assumption of independence between sites, $d_{cc'}$ is the sum of $d_{cc'}^g$,

where $d_{cc'}^g$ is the contribution of mutation site g to the divergence measure. In characterizing the

conditional distribution for X_{cg} and $X_{c'g}$, we use a Beta-Binomial distribution to model expression

stochasticity and a Binomial model to capture sequencing errors and rare RNA-editing events, please see

Methods for details.

To illustrate how transcriptional stochasticity affects the divergence calculation, consider the toy

example shown in Figure 1B, where we compare the contributions of two mutation sites, g and g', to

the pairwise divergence between cell c_1 and cell c_2. First, note that site g is identical to site g' in

mutation allele frequency and total read count in both cells. However, they contribute different values

of $d_{cc'}$ due to their different expression stochasticity: Gene g is bursty, that is, expression of both alleles

is sporadic leading to a U-shaped mutation allele frequency distribution across cells. In contrast, gene g'

is constitutive, that is, expression of both alleles is stable, leading to a bell shaped allele-frequency

distribution across cells. Thus, biological dropout is more likely for gene g than for gene g', which means

that, despite the identical mutation allele frequency and total coverage, $d_{c_1 c_2}^g < d_{c_1 c_2}^{g'}$.

Accuracy assessment using in silico spike-ins

We designed a spike-in procedure to assess the accuracy of DENDRO and to make realistic power

projections for subclone detection (Figure 2a). The procedure, in its basic mode, starts with an assumed

evolutionary tree, where the leaves are clades and mutations can be placed on the branches. In the

absence of prior information, a simple tree structure is used, such as the one shown in Figure 2A.

Parameters of simulation are (1) total number of mutations, (2) total number of cells, (3) the proportion

of cells in each clade, (4) the proportion of mutations along each branch, and (5) mean read coverage

across loci. Some of these parameters can be determined using bulk DNA-seq and/or bulk RNA-seq data,

if available, see our description of DENDROplan in next section. Parameters (1-4) determine the

mutation matrix Z. To get the matrix of alternative and total read counts for each mutation loci in each

cell (X_{cg} and N_{cg}, respectively), we overlay the mutation matrix onto the scRNA-seq data, see details in

Methods. This allows the spike-in framework to retain the expression stochasticity and sequencing error

of real scRNA-seq data. DENDRO is applied to the spike-in data to obtain the clonal clusters, which is

then compared with the known labels. Subclone detection is evaluated by three metrics: adjusted Rand

index, capture rate and purity. Details on these metrics are given in the Supplementary Material.

Using the above framework, we conducted a systematic evaluation of DENDRO's subclone detection

accuracy on an example scRNA-seq dataset (Figure 2). The results show: (1) As mutation burden of a

clade increases, the adjusted Rand index and clade-specific capture rate decrease, but cluster purity of

interested clade increases (Figure 2B), because there are few mutations remaining to distinguish the

other two subclones; (2) similarly, as cell proportion of a clade increases, its adjusted Rand index and

capture rate decrease, but cluster purity increases (Figure 2C); (3) more mutation site increases DENDRO

performance; (4) lower sequencing depth decreases accuracy; (5) As false positive rate of mutation calls

increasing, DENDRO performs worse; (6) lower sequencing error rate improves DENDRO accuracy; and

(7) variation in total expression of genes does not influence DENDRO's divergence measure (Figure S3).

The last point is due to DENDRO's reliance only on the distribution of the mutation allele frequency

conditioned on the total read coverage, and further illustrated by mutation spike-ins in a mixer cell

population with different expression pattern. This indicates that the divergence evaluation reflects

solely genetic distance not transcriptomic difference. We further illustrate this property by a simulation

study. See Supplementary Material and Figure S2 for details.

As a result, DENDRO's subclone identification is based solely on genetic divergence, and not on

expression profile. Previous scRNA-seq studies identify subclones based on total gene expression,

making it difficult to separate genetic and transcriptomic variations. Clustering based on total expression

counts and then performing differential gene expression analysis is prone to self-fulfilling prophecy, as

there will indeed be differential expressed genes because this was the clustering criteria. DENDRO's

approach makes the downstream differential gene expression analysis between subclones more

interpretable.

Power analysis toolkit and experimental design

Before conducting a single cell RNA-seq experiment on a tumor sample, it is important to project how

subclone detection power depends on the number of cells sequenced and the coverage per cell. To

facilitate experiment design, we developed a tool, DENDROplan (Figure 2a), that predicts the expected

clustering accuracy by DENDRO given sequencing parameters and available bulk data for the tumor.

Given an assumed tree structure and a target accuracy, DENDROplan computes the necessary read

depth and number of cells needed.

As shown in Figure 2a, if bulk DNA sequencing and/or RNA sequencing data are available for the tumor

being studied, these data can be harnessed to make more realistic power calculations. For example, if

SNAs have been profiled using bulk DNA sequencing data, the set of mutations that lie in the exons of

annotated genes can be retrieved and used directly in constructing the spike-in data. Furthermore,

phylogeny construction algorithms for bulk DNA-seq data can be used to infer a putative tree structure

that can be used as input to DENDROplan [4, 39]. If bulk RNA-seq data is available, the bulk expression

level of the mutation-carrying genes can be used to predict the expression level of the mutation in the

single cell data. For details, see Methods. The power analysis tool is also available at

https://github.com/zhouzilu/DENDRO.

Evaluation of DENDRO on a renal cell carcinoma and its metastasis

We applied DENDRO to the analysis of the renal cell carcinoma dataset from Kim et al [22] (Figure 3).

This dataset is composed of 116 cells from three different tumors derived from one single patient: a

patient-derived xenograft (PDX) from the primary renal cell carcinoma (PDX_pRCC), a biopsy of the

metastasis to the lung 1 year after treatment of primary site (Pt_mRCC), and a PDX of the lung

metastasis renal cell carcinoma (PDX_mRCC) (Figure S4A). Transcriptomic profiles of each individual cell

were generated using the Smart-seq protocol [17]. The cells should share common early driver

mutations due to their shared origin from the same patient, but the metastasis and the cultivation of

each tumor in separate medium (human or mouse) should have allowed for the accumulation of new

mutations. Thus, we expect the three tumors to be clonally distinct. Given this prior expectation, we use

this dataset to examine the accuracy of DENDRO and to illustrate how DENDRO enables joint analysis of

the genetic and transcriptomic heterogeneity within the same group of cells.

GATK detected 2,867,029 mutation sites across all cells [38]. Mutations that are detected in less than 5%

(too rare) or more than 95% (too common) of the cells were removed, which leaves 72,206 mutations.

On average, 10801 mutations are detected in each cell and 17.35 cells possess the same mutation for

each loci (Figure S4B, C). For majority sites, only few cells have nonzero read coverage, highlighting the

fact that many mutations are missed due to technical and biological dropout (Figure S4D) [32-36].

We compared 4 different clustering methods: DENDRO, hierarchical clustering based on the primary

genotype matrix Z generated by GATK ($Z_{cg} = 1$ when a mutation g is detected for cell c, $Z_{cg} =$

0 otherwise), hierarchical clustering based on the X/N matrix that preserve the variant allele

information and hierarchical clustering based on gene expression ($logTPM$). DENDRO gives the cleanest

separation between the three populations with adjusted Rand Index of 0.932 (1.0 indicates perfect

clustering, Figure 3A), as compared to 0.754 for Z matrix (Figure 3B), 0.519 for X/N (Figure 3C) and 0.489

for expression (Figure 3D). Inspection of the graph shows that, as expected, divergence between

primary tumor and metastasis exceeds divergence between patient sample and PDX sample, as

PDX_mRCC clusters with Pt_mRCC rather than PDX_pRCC. All of the other three methods successfully

separated the primary sample from the metastatic samples, but cannot differentiate between the two

metastasis samples. For DENDRO, the intra-cluster divergence curve kinks sharply at 3, and thus we

stopped at 3 clusters (Figure 3E and Methods). We annotated the clusters as PDX_mRCC, PDX_pRCC and

Pt_mRCC by their cell compositions (Table S1A). The fact that DENDRO found little mixing of cells

between the three tumors, and low genetic heterogeneity within each tumor, is unsurprising given how

these tumors were derived: Since relapsed metastasis consists of cells that have already underwent a

selection bottleneck, and since the PDX tumor is seeded by a small subsample of cells from the original

tumor, it is not surprising that no subclones are detected in each of the three tumors [40-42]. By

maximum likelihood, we called a total of 13,454 mutations after pooling the cells within each cluster

(Figure 3F and Methods). The unrooted phylogenetic tree computed with the revised mutation profiles

confirms that genetic divergence between metastasis and primary is greater than divergence between

PDX and primary sample.

DENDRO enables simultaneous clonal assignment and transcriptomic profiling of the same set of cells.

Plot of smoothed expression ordered by DENDRO shows unique expression patterns within each

subclone (Figure S6). We focused on the comparison of the two metastasized cell populations

(metastasis to lung and patient derived mouse xenograph). Even though PDX_mRCC was derived from

Pt_mRCC, the DENDRO analysis found substantial genetic divergence between the two cell populations.

To investigate further, we performed a differential expression analysis between PDX_mRCC and

Pt_mRCC with scDD and MAST, detecting 74 significant differentially expressed genes (Methods, Figure

S7E, Table S2A) [43-45]. Gene ontology analysis classified these 74 genes into two subgroups: immune-

related genes and cancer-related genes (Table 1, Table S2F) [46]. Immune-related differentially

expressed genes are enriched for the terms TNF-α signaling, complement system and allograft rejection.

On the other hand, cancer related differentially expressed genes overlap with the pathways including

hypoxia, KRAS signaling, mTORC1 signaling and epithelial mesenchymal transition.

Simultaneously, we compare the mutation profiles of these two clones. 9521 locus have different

mutated allele counts between these two populations and were further annotated by ANNOVAR [47].

After filtering, the preserved variants associated with 24 out of 74 differential expressed genes (Table

S2A). Next, we performed a similar GSEA on variants associated genes to identify mutation-related

pathway [46]. Interestingly, variant annotated genes are enriched in cancer-related pathways, including

mitotic spindle, mTORC1 signaling, EMT and hypoxia, overlapping substantially with the cancer-related

pathways identified by differential expression analysis; in comparison, none of the differentially

expressed genes from immune pathways showed up in this mutated gene analysis (Table 2, Table S2F).

In another word, cancer-related transcriptomic divergence between PDX_mRCC and Pt_mRCC is driven

directly by genetic alterations in the same genes, but immune-related differential expression is

influenced by non-DNA factors. This makes sense, since implantation of tumor cells from human to mice

alters their immune microenvironment [48-50], and thus is expected to alter immune-related signaling

within the implanted tumor cells. This illustrates how DENDRO extricates DNA variation from RNAs

allowing heir joint analysis. Differential expression and differential mutation analysis for the other

subclone pairs can be found in Table S2 and Figure S7.

Table 1. GO analysis on Differential Expressed Genes between Pt_mRCC and PDX_mRCC

Gene Set Name [# Genes (K)]			p-value	FDR q-value
Color key:	Cancer-related pathway Immune-related pathway Other			
HALLMARK_TNFA_SIGNALING_VIA_NFKB [200]			1.86 e-8	9.28 e-7
HALLMARK_HYPOXIA [200]*			5.04 e-7	1.26 e-5
HALLMARK_MTORC1_SIGNALING [200]*			1.15 e-5	1.92 e-4
HALLMARK_COMPLEMENT [200]			2.15 e-4	1.79 e-3
HALLMARK_GLYCOLYSIS [200]*			2.15 e-4	1.79 e-3
HALLMARK_KRAS_SIGNALING_UP [200]			2.15 e-4	1.79 e-3
HALLMARK_ALLOGRAFT_REJECTION [200]			3.17 e-3	1.58 e-2
HALLMARK_EPITHELIAL_MESENCHYMAL_TRANSINSITION [200]*			3.17 e-3	1.58 e-2
HALLMARK_ESTROGEN_RESPONSE_EARLY [200]			3.17 e-3	1.58 e-2
HALLMARK_INFLAMMATORY_RESPONSE [200]			3.17 e-3	1.58 e-2

266   Table 2. GO analysis on Differential Mutated Genes between Pt_mRCC and PDX_mRCC

Gene Set Name [# Genes (K)]			p-value	FDR q-value
Color key:	Cancer-related pathway Immune-related pathway Other			
HALLMARK_UV_RESPONSE_DN [144]			1.39 e-29	6.93 e-28
HALLMARK_MYC_TARGETS_V1 [200]			2.75 e-27	6.87 e-26
HALLMARK_MITOTIC_SPINDLE [200]			7.97 e-24	1.33 e-22
HALLMARK_MTORC1_SIGNALING [200]*			9.49 e-20	1.19 e-18
HALLMARK_EPITHELIAL_MESENCHYMAL_TRANSINSITION [200]*			1.91 e-17	1.91 e-16
HALLMARK_OXIDATIVE_PHOSPHORYLATION [200]			1.06 e-16	8.81 e-16
HALLMARK_HYPOXIA [200]*			5.69 e-16	4.06 e-15
HALLMARK_GLYCOLYSIS [200]*			2.97 e-15	1.86 e-14
HALLMARK_ANDROGEN_RESPONSE [101]			1.07 e-14	5.96 e-14
HALLMARK_HEME_METABOLISM [200]			7.4 e-14	3.7 e-13

Analysis of scRNA-seq dataset of primary breast cancer

We next demonstrate the application of DENDRO to a primary breast cancer dataset [21]. We focused

on two tumors (BC03 and BC09) that have the most cell sequenced (Table S3 and Figure S8). BC03

includes cells from primary tumor (here after BC03P) and from regional metastatic lymph nodes (here

after BC03LN). 132 single cell transcriptomes were profiled by Smart-seq protocol [17]. For this analysis,

we will first examine whether DENDRO can separate BC03 and BC09 based on their genetic differences,

and then, jointly characterize the transcriptomic and genetic changes between cells in lymph nodes and

primary resections.

GATK [46] detected a total of 2,364,823 mutation sites across the 132 cells, 353,647 of which have

mutation frequency greater than 5% and smaller than 95%, and were retained for downstream analysis

(Figure S8A, B, C). Figure 4 shows the clustering given by DENDRO. DENDRO separates BC09 cells from

BC03 cells with 100% accuracy (Figure 4A). The intra-cluster divergence curve kinks at 5 clones between

BC03 and BC09, yielding 3 subclones for BC03 and 2 for BC09 (Figure 4A, Figure S8D and Table S1B).

Within BC03, clones Mix_1 and Mix_2 contain a mixture of cells from the primary tumor and lymph

nodes, and clone LN_1 contains mostly cells from the lymph nodes. This suggests that tumor cells that

have transitioned to the lymph nodes are genetically heterogeneous, with some cells remaining

genetically similar to the primary population and others acquiring new genetic mutations. In

comparison, hierarchical clustering based on expression (using log transcripts-per-million values) does

not separate BC03 from BC09, and give a negative adjusted Rand index within BC03, indicating worse

than random classification (Figure 4B).

DENDRO pooled cells within each of the 5 clusters and re-estimated their mutation profiles. We define a

variant as subclonal if it is not present in all of the subclones within a tumor. Based on marginal

likelihood, we pick the top 10,000 variants to construct a phylogenetic tree (Figure 4C). As expected, the

two BC09 clusters are far separate from the BC03 clusters. Within BC03, the length of the branches

shows that the subclone containing mostly cells from lymph nodes (labeled BC03LN_1) is genetically

similar to the Mix_2 clone, and distant from the Mix_1 clone (Figure 4C).

To profile clone-specific gene expression patterns, we first plot window-smoothed expression with cells

grouped by DENDRO clustering [23]. Unsurprisingly, there are broad chromosome-level differences in

expression patterns between groups (Figure S9), most likely due to copy number aberrations and/or

epigenetic remodeling. Next, we focused on the expression of the PAM50 genes (Figure 4D), and found

substantial differences in their expression between subclones. DENDRO detected one rare subclone,

BC09_2, with only six cells (<5% of the total number of cells) which has strong basal-like signature.

Interestingly, in BC03, the subclone LN_1 has the TNBC/basal-like subtype with invasive gene signature,

while the subclone Mix_2 has the $ESR1^+$ subtype. Thus, the genetic divergence of LN_1 from Mix_2 is

accompanied by its acquisition of an invasive metastatic expression signature. If the comparison were

made directly between cells from the primary site and cells from the lymph node without DENDRO's

subclone assignment, these expression patterns would be much weaker since the clones do not cleanly

separate by site. Comparing with original paper assigning each tumor to one specific breast cancer

subtype, with the aid of DEDNRO, subtype identification in sub-tumor resolution allows better therapy

design to potentially target all subclones and reduce tumor relapse risk.

We next conducted a transcriptome-wide search for pathways that have differential expression

between clones (Methods and Table S4), and assessed their overlap with pathways that are

differentially mutated between clones. Focusing on tumor BC03, pathways for G2M checkpoint and

KRAS signaling are up-regulated in lymph node metastasis subclone BC03LN_1, while pathways for

estrogen response and apoptosis are down-regulated, indicating more invasive phenotype (Table S4E).

In addition, *GAPDH* is up-regulated in the metastatic clone (BC03LN_1) and down-regulated in the two

mix-cell clones, consistent with previous findings [51, 52] (Figure S10D). Differentially expressed genes

between other subclone pairs in BC03 are also enriched in estrogen response, apoptosis, and DNA

repair. (Table S4C, D). In parallel, subclone-specific mutated genes are highly enriched in cancer-related

pathways including MYC target, G2M checkpoints and mitotic spindle, and immune related pathways

such as, interferon response, TNF-a signaling and inflammatory response (Table S4). Interestingly, few of

the differentially mutated genes are associated with estrogen and androgen responses, suggesting that

differential expression of hormone related genes are not mediated directly by genetic mutations in

these pathways. This is consistent with the recent studies that found epigenetic alteration, such as

histone acetylation and methylation, to underlie hormones receptor signaling in breast cancer [53-56].

DNA-RNA joint analysis between other subclones are included in the Table S4 and Figure S10. Overall,

this example illustrates how DENDRO enables the joint assessment of genetic and transcriptomic

contributions to clonal diversity at single-cell resolution.

Discussion

We have described DENDRO, a statistical framework to study intratumor DNA-level heterogeneity using

scRNA-seq data. DENDRO starts with mutations detected directly from the scRNA-seq reads, which are

very noisy due to a combination of factors: Errors are introduced in the reverse-transcription,

sequencing and mapping steps in the experiment, low sequencing depth and molecule conversion

efficiency lead to technical dropouts, and expression burstiness at the single cell level lead to biological

dropouts. DENDRO overcomes these obstacles by utilizing the fact that cells from the same subclone

possess almost identical mutation profile. Given noisy mutation profiles and allele specific read counts,

DENDRO first reconstructs clonal substructure by computing a distance between each pair of cells that

quantifies their genetic divergence after accounting for transcriptional bursting, dropout and sequencing

error. Then, DENDRO re-estimates subclone-specific mutation profiles by pooling reads across cells

within the same subclone. These re-estimated mutations profiles are then passed to downstream

mutation analysis and phylogenetic tree reconstruction.

Importantly, the genetic divergence used by DENDRO for cell clustering is based solely on allelic

expression ratios and do not reflect the difference in total expression between cells at mutation sites.

Thus, DENDRO differs from and complements existing tools that cluster cells based on total expression.

In fact, as shown by spike-in analysis, DENDRO clusters the cells based on true underlining mutation

profiles, and are robust to changes in total gene expression. As expected, the numbers of cells, the

depth of sequencing, the actual number of subclonal mutations and the phylogenetic tree structure all

influence the power of DENDRO. To aid researchers in experiment design, we developed DENDROplan,

which predicts DENDRO's clustering accuracy given basic experimental parameters and the expected

informative mutation count, which can be obtained from bulk DNA sequencing.

Ideally, joint profiling of the DNA and RNA on the same cells would allow us to relate genomic profiles to

transcriptomic variations. Currently, there is yet no scalable technology for doing this. Separately

performing scDNA-seq and scRNA-seq on different batches of cells within the same tumor would meet

the nontrivial challenge of matching the subclones between the two data sets. DENDRO takes advantage

of the central dogma and utilizes computational methods to extract genetic divergence information

from noisy mutation calls in coding regions. Through two case studies, we illustrate the accuracy of

DENDRO compared to more ad hoc methods and the DNA-RNA joint analysis that DENDRO enables.

We have demonstrated that proper computational modeling can excavate the DNA-level heterogeneity

in scRNA-seq data. Yet, there is always limitation in working with RNA. While rare RNA editing events are

absorbed by the parameter ϵ, DENDRO cannot distinguish subcluster-specific constituent RNA editing

events from subcluster-specific DNA mutations. In the extreme and unlikely scenario where RNA editing

events are common and pervasive, DENDRO's cluster would reflect RNA editing. In such cases, we

recommend using matched bulk DNA-seq of the same tumor to filter the loci detected in the first step of

DENDRO, keeping only those that are supported by at least on read in the bulk DNA-seq data.

Conclusions

We have developed DENDRO, a statistical method for tumor phylogeny inference and clonal

classification using scRNA-seq data. DENDRO accurately infers the phylogeny relating the cells and

assigns each single cell from the scRNA-seq data set to subclone. DENDRO allows us to (1) cluster cells

based on genetic divergence while accounting for transcriptional bursting, technical dropout and

sequencing error, as benchmarked by in silico mixture and a spike-in analysis, (2) characterize the

transcribed mutations for each subclone, and (3) perform DNA-RNA joint analysis with the same set of

cells and examine the relationship between transcriptomic variation and mutation profile. We illustrate

DENDRO through two case studies. In the first case study on a metastasized renal cell carcinoma

dataset, we validated DENDRO's results using known subpopulation labels and elucidated the genetic-

transcriptomic changes in the PDX tumor model. In the second case study on a primary breast tumor

dataset, DENDRO brought forth new insights on the interplay between intratumor transcriptomic

variation and subclonal divergence.

Methods

SNA calling pipeline

Figure S1 illustrates the SNA calling pipeline. Raw scRNA-seq data is aligned by STAR 2-pass method [37].

Raw variants calling was made using the Haplotype Caller (GATK tool) on the BAM files after sorting,

joining read groups, removing duplicated reads, removing overhangs into intronic regions, realigning

and recalibration [38]. Conventionally, there are two methods from GATK tools for mutation detection:

haplotype caller and mutect2. Haplotype caller has a RNA-seq setting which handle splice junctions

correctly, but assumes VAF around 50%, while mutect2 can detect mutations with low VAF but does

account for splice junction. The reason we select haplotype caller instead of mutect2 is that we extract

allele read counts for all cells as long as one of the cells is listed as carrying the mutation. Thus, as long

as one cell has VAF reaching 50%, this mutation would be detected. Calls with stand_call_conf greater

than 20 and population frequency greater than 5% but less than 95% were preserved for further

analysis. Admittedly, such lenient filtering typically introduces false positive sites. However, our priority

at this step is to minimize false negative rate, while the genetic divergence matrix in the following step

robustly estimates cell population substructure. Both the coverage of the alternative allele and the total

read coverage are extracted for each site for further analysis.

Genetic Divergence and Beta-Binomial framework

Consider two cells: c and c'. Let I_c and $I_{c'}$ denote the clonal group to which the cells belong, i.e. $I_c = I_{c'}$

if and only if cells c and c' come from the same subclone. We define the genetic divergence at loci

g, by $d_{cc'}^g$:

399
$$d_{cc'}^g = \log \frac{1}{P(X_{cg}, X_{c'g} | N_{cg}, N_{c'g}, I_c = I_{c'})}$$

400
$$= \log \frac{P(X_{cg}, X_{c'g} | N_{cg}, N_{c'g}, I_c = I_{c'}) + P(X_{cg}, X_{c'g} | N_{cg}, N_{c'g}, I_c \neq I_{c'})}{P(X_{cg}, X_{c'g} | N_{cg}, N_{c'g}, I_c = I_{c'})})$$

where $X_c = (X_{c1}, X_{c2}, \dots X_{cg}, \dots X_{cm})$ are the mutation allele read counts for cell c and $N_c =$

$(N_{c1}, N_{c2}, \dots N_{cg}, \dots N_{cm})$ are the total read counts at these sites. More intuitively, if cells c and c' are not

from the same clonal group, the numerator has larger value compared to denominator. Thus $d^g_{cc'}$ is

large, indicating bigger divergence between the two cells. With further derivation (see Methods), $d^g_{cc'}$ is

a function of the five following probabilities:

$d^g_{cc'}$

$= f\left(P_g; P\left(X_{cg}\middle|N_{cg}, Z_{cg} = 0\right); P\left(X_{cg}\middle|N_{cg}, Z_{cg} = 1\right); P\left(X_{c'g}\middle|N_{c'g}, Z_{c'g} = 0\right); P\left(X_{c'g}\middle|N_{c'g}, Z_{c'g} = 1\right)\right)$

where $Z_{cg} \in \{0,1\}$ is SNA indicator for cell c at site g and $P_g = P(Z_g = 1)$ is mutation frequency across

the cells estimated by GATK calls.

In the above formula for $d^g_{cc'}$, $P\left(X_{cg}\middle|N_{cg}, Z_{cg} = 0\right)$ and $P\left(X_{c'g}\middle|N_{cg}, Z_{cg} = 0\right)$ reflect reverse-

transcription/sequencing/mapping errors and rare RNA editing events, because when there is no

mutation (i.e. $Z_{cg} = 0$), all mutation reads reflect such technical errors or RNA editing. Let ϵ denote the

combined rate of technical error and RNA editing, we have

$$P\left(X_{cg}\middle|N_{cg}, Z_{cg} = 0\right) \sim \text{Binomial}(X_{cg}|N_{cg}, \epsilon)$$

For cases where there are mutations (i.e. $Z_{cg} = 1$), the distribution of mutated read counts given total

read counts is modeled with a Beta Binomial distribution, which is capable of modeling technical

dropout and transcriptional bursting, and is supported by previous allele specific expression studies [28,

60] .

$$P\left(X_{cg}\middle|N_{cg}, Z_{cg} = 1\right) \sim \int_0^1 \text{Binomial}\left(X_{cg}\middle|N_{cg}, Q_{cg} = q\right) dF(q),$$

$$q \sim \text{Beta}(\alpha_g, \beta_g)$$

where Q_{cg} indicates proportion of mutated alleles expressed in cell c at site g, with Beta distribution as

prior. Respectively, α_g and β_g represent gene activation and deactivation rate, which are estimated

empirically. See Supplementary Material for detailed formula derivation.

Kernel based clustering and optimal cluster assignment

We cluster the cells using a kernel-based algorithm, such as hierarchical clustering. Given that there are

multiple sorting schemes, we leave the user to choose it. For the default-sorting scheme, we

recommend "ward.D" [61]. This is because $d_{cc'}$ behaves like a log likelihood ratio , which should follow a

χ^2 distribution when the two cells share the same subclone. The "ward.D" method has been shown to

work well in Euclidian space,. Empirically, among different hierarchical clustering algorithms on the renal

cell carcinoma dataset (Figure S5) "ward.D" based hierarchical clustering performs the best.

To determine the number of clusters we use an intra-cluster divergence curve computed from the

divergence matrix. Existing software rely on AIC, BIC, or another model selection metric, [39, 62].

However, since we only have the "distance" matrix, these traditional methods cannot be applied. Let N_k

be the number of cell pairs in cluster C_k and N be the total number of pairs between cells for all

clusters. Let K be the number of clusters. The weighted sum of intra-cluster distance W_K is

$$W_K = \sum_{k=1}^{K} N_k \sum_{(i,j) \in C_k} \frac{d_{ij}}{N}$$

Note that small clusters are naturally down-weighted in the above metric. DENDRO relies on visual

examination of the intra-cluster divergence curve (W_K plotted against K) to find the "elbow point",

which can be taken as a reasonable clustering resolution.

Spike-in analysis

In our spike-in analysis, we adopt a dataset from Deng et al, which has allele specific read counts [63].

We further added binomial noise to mimic sequencing error. When analyzing DENDRO performance in

terms of various number mutation site, number of cells, proportion of cells in each clade and proportion

of mutations along each branch, we only take a subset of cells with similar cell types. On the other hand,

we utilize a mixture cell population to test the robustness of DENDRO performance with regard to

various expression profiles.

SNA inference in "bulk" and phylogenetic tree construction

As stated previously, DENDRO further inferred SNA after pooling the reads from all cells within each

cluster. Because, with our choice of thresholds, we identify SNAs in single cells with high sensitivity, the

"bulk" level SNAs should be a subset of the SNAs in single cells, and mutation allele counts and total

allele counts should provide us with enough information for SNA detection using a maximum likelihood

framework [64], which accounts for both sequencing error and rare RNA-editing events. Suppose g is

the genotype (number of reference allele) at a site and assume m, the ploidy, equals to 2. Then the

likelihood is:

457
$$\mathcal{L}(g) = \frac{1}{m^k} \prod_{j=1}^{l} [(m-g)\epsilon + g(1-\epsilon)] \prod_{j=l+1}^{k} [(m-g)(1-\epsilon) + g\epsilon]$$

458
$$g^* = \underset{g}{\operatorname{argmax}} -\mathcal{L}(g)$$

where k is number of reads at a site and the first l bases ($l \leq k$) be the same to reference and the rests

are same to alternative allele. ϵ is the sequencing error and rare RNA-editing combined rate. Given

mutation profiles, DENDRO then constructs a phylogenetic tree with the neighbor-joining method,

which can more accurately capture the evolutionary relationship between different subclones [65] than

the initial tree given by hierarchical clustering.

Differential gene expression, mutation annotation and gene ontology analysis

We used Seurat and scDD to identify differentially expressed genes between tumors and between tumor

subclones [43-45]. For each comparison, we used two different methods: MAST implemented by Seurat

and scDD. Genes with adjusted p-value < 0.05 counts as significant differentially expressed gene for each

method. We further intersect these two sets of differentially expressed genes to increase robustness.

Subclonal mutations are annotated by ANNOVAR with default parameters and variants associated with

intergenic regions were discarded for downstream analysis [47]. For GO analysis, we applied Gene Set

Enrichment Analysis tool [46]. Hallmark gene sets were used as fundamental database with FDR q-value

< 0.05 as significant.

List of abbreviations

SNA: Single nucleotide alteration

scDNA-seq: Single-cell DNA sequencing

scRNA-seq: Single-cell RNA sequencing

PDX: Patient-derived xenograft

TPM: Transcripts per kilobase million

Declarations

Acknowledgements

484   We thank Dr. Nan Lin for informing useful tools for biological insight analysis, and Dr. Kai Tan, Dr.

485   Mingyao Li, and Dr. Andy Minn for helpful comments and suggestions.

Funding

487   This work was supported by National Institutes of Health (NIH) grant 5R01-HG006137-07 to ZZ and to

488   NRZ.

Availability of data and materials

490   DENDRO is an open-source R package available at https://github.com/zhouzilu/DENDRO with license

491   GPL-3.0.

Authors' contributions

493   NRZ initiated and envisioned the study. ZZ and NRZ formulated the model. ZZ developed and

494   implemented the algorithm and conducted all analyses. ZZ and NRZ and wrote the manuscript. All

495   authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Consent for publication

Not applicable.

Ethics approval and consent to participate

Not applicable.

1.   Hanks, S., et al., *Constitutional aneuploidy and cancer predisposition caused by biallelic*
503   *mutations in BUB1B.* Nat Genet, 2004. **36**(11): p. 1159-61.
504   2.   Gamazon, E.R. and B.E. Stranger, *The impact of human copy number variation on gene*
505   *expression.* Briefings in Functional Genomics, 2015. **14**(5): p. 352-357.
506   3.   Burrell, R.A., et al., *The causes and consequences of genetic heterogeneity in cancer*
507   *evolution.* Nature, 2013. **501**(7467): p. 338-45.
508   4.   Jiang, Y., et al., *Assessing intratumor heterogeneity and tracking longitudinal and spatial*
509   *clonal evolutionary history by next-generation sequencing.* Proc Natl Acad Sci U S A,
510   2016. **113**(37): p. E5528-37.
511   5.   Deshwar, A.G., et al., *PhyloWGS: reconstructing subclonal composition and evolution*
512   *from whole-genome sequencing of tumors.* Genome Biol, 2015. **16**: p. 35.
513   6.   Zare, H., et al., *Inferring clonal composition from multiple sections of a breast cancer.*
514   PLoS Comput Biol, 2014. **10**(7): p. e1003703.
515   7.   Carter, S.L., et al., *Absolute quantification of somatic DNA alterations in human cancer.*
516   Nat Biotechnol, 2012. **30**(5): p. 413-21.
517   8.   Li, B. and J.Z. Li, *A general framework for analyzing tumor subclonality using SNP array*
518   *and DNA sequencing data.* Genome Biol, 2014. **15**(9): p. 473.
519   9.   Oesper, L., A. Mahmoody, and B.J. Raphael, *THetA: inferring intra-tumor heterogeneity*
520   *from high-throughput DNA sequencing data.* Genome Biol, 2013. **14**(7): p. R80.
521   10.   Ha, G., et al., *TITAN: inference of copy number architectures in clonal cell populations*
522   *from tumor whole-genome sequence data.* Genome Res, 2014. **24**(11): p. 1881-93.
523   11.   Miller, C.A., et al., *SciClone: inferring clonal architecture and tracking the spatial and*
524   *temporal patterns of tumor evolution.* PLoS Comput Biol, 2014. **10**(8): p. e1003665.
525   12.   Navin, N.E., *The first five years of single-cell cancer genomics and beyond.* Genome Res,
526   2015. **25**(10): p. 1499-507.
527   13.   Navin, N., et al., *Tumour evolution inferred by single-cell sequencing.* Nature, 2011.
528   **472**(7341): p. 90-4.
529   14.   Wang, Y., et al., *Clonal evolution in breast cancer revealed by single nucleus genome*
530   *sequencing.* Nature, 2014. **512**(7513): p. 155-60.
531   15.   Gao, R., et al., *Punctuated copy number evolution and clonal stasis in triple-negative*
532   *breast cancer.* Nat Genet, 2016. **48**(10): p. 1119-30.
533   16.   Gawad, C., W. Koh, and S.R. Quake, *Single-cell genome sequencing: current state of the*
534   *science.* Nat Rev Genet, 2016. **17**(3): p. 175-88.

535   17.   Picelli, S., et al., *Smart-seq2 for sensitive full-length transcriptome profiling in single cells.*
536   Nat Methods, 2013. **10**(11): p. 1096-8.
537   18.   Klein, A.M., et al., *Droplet barcoding for single-cell transcriptomics applied to embryonic*
538   *stem cells.* Cell, 2015. **161**(5): p. 1187-1201.
539   19.   Cloney, R., *Cancer genomics: Single-cell RNA-seq to decipher tumour architecture.* Nat
540   Rev Genet, 2017. **18**(1): p. 2-3.
541   20.   Patel, A.P., et al., *Single-cell RNA-seq highlights intratumoral heterogeneity in primary*
542   *glioblastoma.* Science, 2014. **344**(6190): p. 1396-401.
543   21.   Chung, W., et al., *Single-cell RNA-seq enables comprehensive tumour and immune cell*
544   *profiling in primary breast cancer.* Nat Commun, 2017. **8**: p. 15081.
545   22.   Kim, K.T., et al., *Application of single-cell RNA sequencing in optimizing a combinatorial*
546   *therapeutic strategy in metastatic renal cell carcinoma.* Genome Biol, 2016. **17**: p. 80.
547   23.   Tirosh, I., et al., *Dissecting the multicellular ecosystem of metastatic melanoma by*
548   *single-cell RNA-seq.* Science, 2016. **352**(6282): p. 189-96.
549   24.   Tirosh, I., et al., *Single-cell RNA-seq supports a developmental hierarchy in human*
550   *oligodendroglioma.* Nature, 2016. **539**(7628): p. 309-313.
551   25.   Venteicher, A.S., et al., *Decoupling genetics, lineages, and microenvironment in IDH-*
552   *mutant gliomas by single-cell RNA-seq.* Science, 2017. **355**(6332).
553   26.   Li, H., et al., *Reference component analysis of single-cell transcriptomes elucidates*
554   *cellular heterogeneity in human colorectal tumors.* Nat Genet, 2017. **49**(5): p. 708-718.
555   27.   Raj, A., et al., *Stochastic mRNA synthesis in mammalian cells.* PLoS Biol, 2006. **4**(10): p.
556   e309.
557   28.   Jiang, Y., N.R. Zhang, and M. Li, *SCALE: modeling allele-specific gene expression by single-*
558   *cell RNA sequencing.* Genome Biol, 2017. **18**(1): p. 74.
559   29.   Padovan-Merhar, O., et al., *Single mammalian cells compensate for differences in*
560   *cellular volume and DNA copy number through independent global transcriptional*
561   *mechanisms.* Mol Cell, 2015. **58**(2): p. 339-52.
562   30.   Zafar, H., et al., *Monovar: single-nucleotide variant detection in single cells.* Nat
563   Methods, 2016. **13**(6): p. 505-7.
564   31.   Piskol, R., G. Ramaswami, and J.B. Li, *Reliable identification of genomic variants from*
565   *RNA-seq data.* Am J Hum Genet, 2013. **93**(4): p. 641-51.
566   32.   Brennecke, P., et al., *Accounting for technical noise in single-cell RNA-seq experiments.*
567   Nat Methods, 2013. **10**(11): p. 1093-5.
568   33.   Pierson, E. and C. Yau, *ZIFA: Dimensionality reduction for zero-inflated single-cell gene*
569   *expression analysis.* Genome Biol, 2015. **16**: p. 241.
570   34.   Vallejos, C.A., J.C. Marioni, and S. Richardson, *BASiCS: Bayesian Analysis of Single-Cell*
571   *Sequencing Data.* PLoS Comput Biol, 2015. **11**(6): p. e1004333.
572   35.   Ding, B., et al., *Normalization and noise reduction for single cell RNA-seq experiments.*
573   Bioinformatics, 2015. **31**(13): p. 2225-7.
574   36.   Qiu, X., et al., *Single-cell mRNA quantification and differential analysis with Census.* Nat
575   Methods, 2017. **14**(3): p. 309-315.
576   37.   Dobin, A., et al., *STAR: ultrafast universal RNA-seq aligner.* Bioinformatics, 2013. **29**(1):
577   p. 15-21.

38. McKenna, A., et al., *The Genome Analysis Toolkit: a MapReduce framework for analyzing*
next-generation DNA sequencing data. Genome Res, 2010. **20**(9): p. 1297-303.

39. Urrutia, E., et al., *Integrative pipeline for profiling DNA copy number and inferring tumor*
phylogeny. Bioinformatics, 2018. **34**(12): p. 2126-2128.

40. Eirew, P., et al., *Dynamics of genomic clones in breast cancer patient xenografts at*
single-cell resolution. Nature, 2015. **518**(7539): p. 422-6.

41. Gerlinger, M., et al., *Intratumor heterogeneity and branched evolution revealed by*
multiregion sequencing. N Engl J Med, 2012. **366**(10): p. 883-892.

42. Shi, Y.J., et al., *Intratumoral Heterogeneity in Breast Cancer: A Comparison of Primary*
and Metastatic Breast Cancers. Oncologist, 2017. **22**(4): p. 487-490.

43. Finak, G., et al., *MAST: a flexible statistical framework for assessing transcriptional*
changes and characterizing heterogeneity in single-cell RNA sequencing data. Genome
Biol, 2015. **16**: p. 278.

44. Korthauer, K.D., et al., *A statistical approach for identifying differential distributions in*
single-cell RNA-seq experiments. Genome Biol, 2016. **17**(1): p. 222.

45. Butler, A., et al., *Integrating single-cell transcriptomic data across different conditions,*
technologies, and species. Nat Biotechnol, 2018. **36**(5): p. 411-420.

46. Subramanian, A., et al., *Gene set enrichment analysis: a knowledge-based approach for*
interpreting genome-wide expression profiles. Proc Natl Acad Sci U S A, 2005. **102**(43): p.
15545-50.

47. Wang, K., M. Li, and H. Hakonarson, *ANNOVAR: functional annotation of genetic*
variants from high-throughput sequencing data. Nucleic Acids Res, 2010. **38**(16): p.
e164.

48. Straussman, R., et al., *Tumour micro-environment elicits innate resistance to RAF*
inhibitors through HGF secretion. Nature, 2012. **487**(7408): p. 500-4.

49. Michor, F. and V.M. Weaver, *Understanding tissue context influences on intratumour*
heterogeneity. Nat Cell Biol, 2014. **16**(4): p. 301-2.

50. Augsten, M., *Cancer-associated fibroblasts as another polarized cell type of the tumor*
microenvironment. Front Oncol, 2014. **4**: p. 62.

51. Zhang, J.Y., et al., *Critical protein GAPDH and its regulatory mechanisms in cancer cells.*
Cancer Biol Med, 2015. **12**(1): p. 10-22.

52. Tarrado-Castellarnau, M., et al., *Glyceraldehyde-3-phosphate dehydrogenase is*
overexpressed in colorectal cancer onset. Translational Medicine Communications, 2017.
2(1): p. 6.

53. Mann, M., V. Cortez, and R.K. Vadlamudi, *Epigenetics of estrogen receptor signaling: role*
in hormonal cancer progression and therapy. Cancers (Basel), 2011. **3**(3): p. 1691-707.

54. Green, K.A. and J.S. Carroll, *Oestrogen-receptor-mediated transcription and the influence*
of co-factors and chromatin state. Nat Rev Cancer, 2007. **7**(9): p. 713-22.

55. Dreijerink, K.M., et al., *Menin links estrogen receptor activation to histone H3K4*
trimethylation. Cancer Res, 2006. **66**(9): p. 4929-35.

56. Kim, H., et al., *Requirement of histone methyltransferase SMYD3 for estrogen receptor-*
mediated transcription. J Biol Chem, 2009. **284**(30): p. 19867-77.

57. Han, L., et al., *The Genomic Landscape and Clinical Relevance of A-to-I RNA Editing in*
Human Cancers. Cancer Cell, 2015. **28**(4): p. 515-528.

58.   Mayba, O., et al., *MBASED: allele-specific expression detection in cancer tissues and cell
623   lines.* Genome Biol, 2014. **15**(8): p. 405.
624   59.   Rhee, J.K., et al., *Allelic imbalance of somatic mutations in cancer genomes and
625   transcriptomes.* Sci Rep, 2017. **7**(1): p. 1653.
626   60.   Skelly, D.A., et al., *A powerful and flexible statistical framework for testing hypotheses of
627   allele-specific gene expression from RNA-seq data.* Genome Res, 2011. **21**(10): p. 1728-
628   37.
629   61.   Ward, J.H., *Hierarchical Grouping to Optimize an Objective Function.* Journal of the
630   American Statistical Association, 1963. **58**(301): p. 236-&.
631   62.   Goutte, C., et al., *Feature-space clustering for fMRI meta-analysis.* Hum Brain Mapp,
632   2001. **13**(3): p. 165-83.
633   63.   Deng, Q., et al., *Single-cell RNA-seq reveals dynamic, random monoallelic gene
634   expression in mammalian cells.* Science, 2014. **343**(6167): p. 193-6.
635   64.   Li, B., et al., *A likelihood-based framework for variant calling and de novo mutation
636   detection in families.* PLoS Genet, 2012. **8**(10): p. e1002944.
637   65.   Schliep, K.P., *phangorn: phylogenetic analysis in R.* Bioinformatics, 2011. **27**(4): p. 592-3.
638

Figure 1. DENDRO analysis pipeline and genetic divergence evaluation. a DENDRO analysis pipeline overview. **b** Statistical model for genetic divergence evaluation function. In this example, we show relationship of cell c_1 and c_2 with gene g and g' in terms of their genetic divergence. Middle panels shows the stochasticity of cell expression and bottom panels illustrates the distribution of variants allele frequency across cells for these two genes, indicating that gene g is a bursty gene and g' is a constitutive gene.

Figure 2. DENDRO accuracy assessment by spike-in analysis. a The overall spike-in analysis pipeline. Three statistics measure the accuracy of DENDRO - adjusted Rand index (global accuracy, 1 indicating perfect classification), capture rate (subclone specific accuracy, out of all cells from the clade, how many of them are correctly assigned) and purity (subclone specific accuracy, out of all cells from the detected cluster, how many are actually from the clade). **b** Statistics under different SNV load of a given clade of interest (i.e. as fraction of total mutation counts). **c** Statistics under different cell proportion of a given clade of interest. (Mutation counts: number of mutation identified; Avg mutation read depth: average read depth of all the mutation sites.)

Figure 3. Evaluation of DENDRO on a renal cell carcinoma and its metastasis. a DENDRO clustering result from primary and metastatic renal cell carcinoma dataset. Background colors represent DEDRO clustering result. **b** Clustering of the same dataset using *Z* matrix (indicator matrix, $z_{ij}=1$ when detected a mutation for cell *i* at site *j* by GATK tool). **c** Clustering of the same dataset using *X/N* matrix (mutation allele frequency matrix) **d** Clustering of the same dataset using expression (*log*TPM). **e** Intra-cluster divergence curve to select optimal number of cluster by DENDRO. Here, *optK=3* **f** Phylogenetic tree of three cell populations identified by DEDNRO. (In **a - d**, dashlines indicate cluster boundaries)

Figure 4. Analysis of scRNA-seq dataset of primary breast cancer. a DENDRO cluster result for primary breast cancer dataset (Chung et al., 2017). **b** Hierarchical clustering result for the same dataset based on expression (*log*TPM). (dashlines indicate cluster boundaries). **c** Neighbor joining phylogenetic tree given detected subclones for breast cancer dataset. **d** PAM50 gene panel expression shows breast cancer subtypes of each subclone.