Fluctuating pathogen-mediated selection drives the maintenance of innate

immune gene polymorphism in a widespread wild ungulate

Erwan Quéméré[1*], Pauline Hessenauer[2], Maxime Galan[3], Marie Fernandez[1], Joël Merlet[1], Yannick Chaval[1],

Aidan J Mark Hewison[1], Nicolas Morellet[1], Hélène Verheyden[1], Emmanuelle Gilot-Fromont[4,5], Nathalie

Charbonnel[3]

[1]CEFS, INRA, Université de Toulouse, Castanet-Tolosan Cedex F-31326, France

[2]Département de Foresterie, Université Laval, Quebec, Canada

[3]CBGP, INRA, CIRAD, IRD, Montpellier SupAgro, Univ Montpellier, Montpellier, France

[4]Université de Lyon, Université Lyon 1, UMR CNRS 5558, Villeurbanne, France,

[5]Université de Lyon, VetAgro Sup, Marcy l'Etoile, France

* Corresponding author:

erwan.quemere@ inra.fr. Phone: (+33) 561 285 497. Fax: (+33) 561 285 500.

Abstract (246 words)

Toll-like Receptors (TLR) play a central role in recognition and host frontline defence against a wide range of pathogens. A number of recent studies have shown that *Tlrs* often exhibit a large polymorphism in natural populations. Yet, there is little knowledge on how this polymorphism is maintained and how it influences disease susceptibility in the wild. We investigated the selective mechanisms by which multiple-pathogens drive the genetic diversity of *Tlr* genes, as well as the commonly-used *Mhc -Drb* in European roe deer (*Capreolus capreolus*), an abundant and widespread ungulate in Europe. In a previous work, we showed that some *Tlrs* exhibit contemporary signature of balancing selection in roe deer. To assess the relative contribution of "heterozygous advantage", "negative-frequency dependence" and "fluctuating selection", we tested associations between *Mhc-Drb* or *Tlr* variants and infection or co-infection with *Toxoplasma* and *Chlamydia*, two intracellular pathogens known to cause reproductive failure in ungulates. We showed that *Toxoplasma* and *Chlamydia* exposures vary significantly across space and time with few co-infection events and that the two pathogens act antagonistically on *Tlr* polymorphism leading to fluctuating selection. By contrast, we only observed *Tlr* heterozygous advantage for a single pathogen and we did not find evidence of negative-frequency dependence. We did not detect any association with *Mhc-Drb.* Our study confirmed the importance of looking beyond *Mhc* genes in wildlife immunogenetic studies. It also emphasized the necessity to consider multiple pathogen challenges and their spatio-temporal variations to improve our understanding of vertebrate defence evolution against pathogens.

Keywords: balancing selection, habitat heterogeneity, Toll-like genes, antagonistic effects, roe deer

Running title: Fluctuating selection on TLR genes in deer

1. INTRODUCTION

While evolutionary theory predicts that selection and random drift should deplete genetic diversity, a number of genes and genomic regions have maintained abundant polymorphism in natural populations with exceptionally high ratio of non-synonymous (protein-altering) to synonymous substitutions (Hedrick, 2006; Trowsdale & Parham, 2004). Balancing selection is a collective term for evolutionary processes that adaptively maintain variation in populations. It is a widespread process driving the diversity of genes involved in numerous biological functions including pathogen resistance (Sommer, 2005), color polymorphism (Wellenreuther, 2017) or mate choice/male competiveness (Johnston et al., 2013; Moore & Moore, 1999). However, identifying the mechanisms driving balancing selection has been a major challenge. At least three not-mutually exclusive processes (putting aside sexual selection) can act in synergy: the "heterozygote advantage" or "overdominance" hypothesis (Doherty & Zinkernagel, 1975; Lewontin & Hubby, 1966) proposes that heterozygous individuals have a higher relative fitness, on average, than homozygous individuals. The "negative frequency-dependent selection" (also called rare-allele advantage) hypothesis suggests that the fitness-advantage of alleles is inversely proportional to their frequency in the population as a result of the host-pathogen (Takahata & Nei, 1990) or prey-predator co-evolutionary dynamics (Cook, 2003). Lastly, the "fluctuating selection hypothesis" (Hill, 1991; Levene, 1953) proposes that high genetic diversity may be maintained when spatially and/or temporally varying directional selection occurs with sometime antagonistic effects on fitness (Wittmann, Bergland, Feldman, Schmidt, & Petrov, 2017).

Immunity-related genes are ideal candidates for studying how adaptive genetic diversity is maintained in wild populations because of their direct effect on survival (Møller & Saino, 2004) or reproductive success (Pedersen & Greives, 2008). The genes of the major histocompatibility complex (MHC) are often used as a model since they show an exceptional polymorphism in vertebrates (Spurgin & Richardson, 2010). Their structure and mechanisms of interaction with antigens are now well described (J. Klein & Figueroa, 1986). Furthermore, their ability to trigger a rapid adaptive response to varying pathogen-mediated selective pressures (PMS) has been demonstrated experimentally (Eizaguirre, Lenz, Kalbe, & Milinski, 2012; Phillips et al., 2018). Over the last few years, a growing body of literature identified PMS at *Mhc* genes in a large range of taxa (see Spurgin & Richardson, 2010 for a reviews) but few of them convincingly elucidated the relative influence of the three mechanisms mentioned above (but see Oliver et al., 2009; Philips, 2018). One likely reason is that most studies have an unrealistic view of the functioning of immune system and of the ecological context in which it evolves, by modelling associations between a single gene (most often a *Mhc* gene) and only one pathogen.

Mhc, although important, corresponds to a small fraction of the genetic variation in pathogen

resistance. Before the adaptive immunity has an opportunity to intervene in the host response,

invading pathogens interact with different effectors of innate immunity that recognize specific

structures (Acevedo-Whitehouse & Cunningham, 2006). Therefore many genes encoding proteins

involved in pathogen recognition or immune regulation may be associated with the resistance

against the same pathogen (Turner, Begon, Jackson, Bradley, & Paterson, 2011). In particular, genes

encoding Toll-like receptors (TLR) play a key role in the host frontline defence against a wide range of

microparasites such as fungi, bacteria, protozoa. This family of genes conserved in both invertebrate

and vertebrate lineages are critical for the stimulation of many immune pathways such as

inflammation, innate or acquired immunity (Akira, Takeda, & Kaisho, 2001). A multitude of studies

have revealed associations between *Tlr* gene polymorphism and infectious diseases in humans and

domestic species (Mun et al., 2003; Yang & Joyee, 2008). However, very few (most focusing on birds

or rodents) have investigated their role in host-pathogen interactions in natural environment and

their contemporary mechanisms of evolution at a population level (but see Turner et al., 2011;

Tschirren et al., 2013; Grueber et al., 2013).

Moreover, co-infection with multiple pathogens is a real world rule (Maizels & Nussey, 2013) and

some selective mechanisms such as heterozygous advantage or fluctuating selection may better be

detected when accounting for the multi-pathogen context (Sin et al., 2014; Wegner, Kalbe, Kurtz,

Reusch, & Milinski, 2003). Most immunity-related genes have pleiotropic effects on resistance

against several pathogens (Hedrick, 1999) and the maintenance of 'susceptible' (deleterious) alleles

in a wild population can be due to antagonist effects of a given immunity-related gene on several

pathogens (*i.e.* antagonistic pleiotropy). To date, only few studies have examined the selective

effects of multiple pathogens on host immunogenetic patterns and they have only focused on *Mhc*

genes (Tollenaere et al. 2008 on voles, Loiseau et al. 2008 on house sparrows, Kamath et al. 2014 in

plain zebras, Froeschke & Sommer 2012 on striped mouse).

Here, we investigated the selective mechanisms by which multiple pathogens could drive the

genetic diversity of immunity-related genes in European roe deer (*Capreolus capreolus*), the most

abundant and widespread ungulate in Europe. We studied a free-ranging population of roe deer

inhabiting an agricultural landscape in Southwestern France. This population is heavily exposed to a

range of pathogens that also infect livestock, including the abortive *Toxoplasma gondii* or *Chlamydia*

abortus (Candela et al., 2014). These two pathogens are known to have negative influence on the

survival and reproductive performances of ungulates (Dubey, 2009; Pioz et al., 2008). They may

therefore exert selective pressures on immunity-related genes. We focused on four genes encoding

pattern-recognition receptors (PRRs) that detect specific microbes or pathogen-associated molecular

patterns: the widely-used *Mhc class II-Drb* gene and three non-*Mhc* genes encoding Toll-like receptors genes (*Tlr2, Tlr4, Tlr5*). *Mhc* and *Tlr* genes may have pleiotropic effects on the resistance/susceptibility to pathogens harboring similar pathogen-associated molecular patterns (Kaisho & Akira, 2004; Loiseau et al., 2008). Moreover, *Tlr* genes have been shown to play a major role in host defence against multiple pathogens, including *Toxoplasma* (Miller, Boulter, Ikin, & Smith, 2009) and *Chlamydia* (Beckett et al., 2012). In a recent work, we showed that these genes exhibit high levels of amino acid diversity within French roe deer populations including the population studied here, hence suggesting an important functional repertoire (Quéméré et al., 2015). *Tlr2* and *Tlr4* also exhibited contemporary signatures of balancing selection: several haplotypes were observed at moderate frequency and showed very low levels of differentiation among roe deer populations compared to the patterns obtained with neutral markers. These findings support the hypothesis that the polymorphism of these genes is shaped by multiple pathogen-mediated selection.

More specifically, we assessed the role of "heterozygous advantage", "negative-frequency dependence" and "fluctuating" selection in shaping the signatures of balancing selection observed in the European roe deer. We analyzed whether the prevalence of *Toxoplasma* and *Chlamydia* or a combination of these pathogens (*i.e.* co-infection status) was associated with specific patterns of genetic diversity observed at the four immunity-related genes (e.g. heterozygosity/allelic divergence, presence of rare or common alleles), while accounting for ecological predictors (age, sex, spatial and temporal variation in infection level). The "heterozygote advantage" hypothesis has received limited support when authors focused on a single pathogen species (but see Olivier et al., 2009) but a growing number of case studies has revealed *Mhc* heterozygous superiority in animals co-infected with multi-pathogens (Hughes & Nei, 1992; McClelland et al., 2003; Froeschke & Sommer, 2012 but see Sin et al., 2014). In agreement, we expected associations between immunity-related gene heterozygosity and the co-infection status of deer, in particular because *Toxoplasma* and *Chlamydia* employ similar strategies to invade host cells (Romano & Coppens, 2013). Negative-frequency dependence, through host-pathogen "Red Queen" co-evolution, may confer selective advantage to novel alleles as recently demonstrated experimentally in wild populations (Phillips et al., 2018). Therefore, under this hypothesis, we expected negative relationships between rare *Mhc* or *Tlr* alleles and pathogen infection. Lastly, given the high temporal and spatial heterogeneity in pathogen exposure in the study area (Candela et al., 2014; Sevila et al., 2014), balancing selection was also predicted to act on *Mhc-Drb* and *Tlr* genes through spatial and/or temporal fluctuations in pathogen-mediated selection. We thus expected to find various associations between particular *Mhc-Drb* or *Tlr* alleles and pathogen prevalence.

2. MATERIAL AND METHODS

2.1 Study population and data collection

The study focused on a roe deer population inhabiting a heterogeneous agricultural landscape in Southern France (43°13′N, 0°52′E). This area is covered by a mix of open fields (including 38% meadows and pastures for cattle and sheep, and 32% crops), hedges (16%), small woodlands patches (14%) and two larger forest blocks (5%). Individuals were caught by drive-netting during winter (from January to March) between 2008 and 2016 in seven sampling sites with varying vegetation cover (Figure S1). Individual sex, age class (juveniles: ≤ 1 year of age versus adults: > 1 year of age) and body mass were recorded. Blood samples were collected for pathogen screening (see below) and ear punches for genetic analyses. In total, we gathered samples from 433 annual captures corresponding to 328 different individuals (190 females and 138 males, 164 juveniles and 164 adults). All applicable institutional and European guidelines for the care and use of animals were followed. The study was permitted by the land manager. All the procedures involving animals were approved by the Ethical Committee 115 of Toulouse and authorized by the French Ministry in charge of ethical evaluation (n° APAFIS#7880-2016120209523619v5).

2.2 *Tlr* and *Mhc-Drb* genotyping

DNA was extracted from alcohol-preserved tissues using the DNeasy Blood and Tissue kit (QIAGEN). The second exon of the *Mhc-Drb* class II gene encoding the ligand-binding domain of the protein was amplified and sequenced for 293 deer samples using Illumina MiSeq system following the procedure detailed in Quéméré et al. (2015). Haplotypes and individual genotypes were identified using the SESAME barcode software (Piry, Guivier, Realini, & Martin, 2012). *Tlr* genes (*Tlr2*, *Tlr4* and *Tlr5*) were genotyped using the two-step procedure described in Quéméré et al. (2015). Previously, a pilot study on 32 individuals was performed to identify polymorphic sites (SNPs) and Linkage-disequilibrium (LD) groups. We screened almost the entire coding region of the three *Tlr* genes (82% in average) including the leucine-rich extracellular region of receptors involved in antigen-binding. Detailed on primer sequences, SNP positions and codon changes are provided in Quéméré et al. (2015). We then selected one SNP per LD group (primarily targeting non-synonymous sites) that was genotyped for all individuals (N=328) using the KASPar allele-specific genotyping system provided by KBiosciences (Hoddesdon, UK). A total of 13 SNPs were typed including 5, 3 and 5 SNPs for *Tlr2*, *Tlr4* and *Tlr5* respectively. Details on SNP position and codon change can be found in Table S2. Haplotypes were then reconstructed from the phased SNPs using the procedure implemented in DNASP v5 (Librado & Rozas, 2008). All sequences have been submitted to NCBI Genbank (Accession nos. are in Table S1, Supporting information).

2.3 Pathogen screening

We determined infection by *Toxoplasma gondii* and/or *Chlamydia abortus* by analysing the serological status of roe deer with regard to these pathogens using classical enzyme-linked immunosorbent assays (ELISA) with specific commercial kits (Sevila et al., 2014). According to the manufacturer's instructions, blood samples with antibody recognition level (ARL) > 30% (respectively 40%) were considered positive for *T. gondii* (respectively *C. abortus*) (see Sevila et al., 2014 for further details). In total, we obtained the *Toxoplasma* and *Chlamydia* serological status of 277 (with 74 repeated measures) and 196 (36 repeated measures) roe deer respectively (caught between 2008 and 2016). The proportion of co-infected individuals was examined in the years where both pathogens were screened (between 2008 and 2013). The importance of interactions between *Toxoplasma* and *Chlamydia* in roe deer was analyzed using a screening association method (Vaumourin et al., 2014). This method tests whether some pathogen associations are significantly more or less detected than expected under the null hypothesis that these associations are random.

2.4 Statistical analyses

We analyzed the relationships between the prevalence of *T. gondii* or *C. abortus* infection (binary response variable – presence/absence) and immunity-related gene variation using generalized linear mixed-effect models (GLMM) with a binomial family (using a logit link function) and the *glmer* function implemented in the lme4 package (Bates, Maechler, & Bolker, 2012). We also tested whether immunity-related gene variation affected the number of pathogens co-infecting roe deer (0, 1 or 2) using a GLMM with a Poisson family (using a log link function).

In a first step, we built for each pathogen and for the co-infection status a reference "non-genetic" model containing biological and environmental factors that possibly affect pathogen exposure. We tested the influence of sex (male *versus* female) and age-class (juvenile *versus* adult) because both behaviour and host susceptibility may vary between sexes and across an individual's lifetime (Klein, 2000) and because infection persistence is expected to result in a positive age-prevalence relationship (Gotteland et al., 2014). We also considered the spatial and inter-annual variation in environmental conditions that may affect pathogen survival and thus encounter probability (e.g. climate, food resources, population density, etc.). For each individual and year, we calculated an annual probability of infection (API) corresponding to the number of infected individuals in a given year (except the focal individual) divided by the number of animals sampled this year (except the focal individual). Similarly, we calculated a spatial probability of infection (SPI) as the number of infected individuals in a given sampling site (except the focal individual) divided by the number of animals sampled within this site (except the focal individual). In the co-infection models,

we tested the effect of sex, age, API and SPI of both pathogens. In all models, we included individual

identity as a random effect to account for correlation amongst different observations of the same

individual.

In step two, using the best 'non-genetic' model as reference, we tested, independently for each

pathogen/co-infection and gene, the three mechanisms potentially underlying balancing selection.

The "heterozygote advantage" hypothesis was tested either by i) adding heterozygosity as a binary

fixed effect (heterozygote vs homozygote) or ii) fitting the number of SNPs (allelic distance) among

the two alleles with an individual's genotype. The "negative frequency dependence" hypothesis was

investigated by including the presence or absence of rare haplotype (<10%) as explanatory variable.

Finally, the "fluctuating selection" hypothesis was examined by considering associations between

infection status and the presence/absence (0/1) or the number of copies (0, 1 or 2) of the most or

the second most frequent haplotype. Each genetic effect was tested either as a simple additive effect

or in interaction with API or SPI. These latter enabled to test the hypotheses that the genetic effect

on pathogen prevalence varies across time or space, according to the intensity of exposition.

In total, 28 models were tested for each combination of gene and pathogen or gene and co-

infection status (Table S3). We used a model selection procedure using Akaike information criterion

with a correction for small sample sizes (AIC_C) to determine which model best explained variation in

pathogen infection (models with lowest AIC) . We only examined models with ΔAIC< 2 relative to the

best model. Statistical analyses were performed using the software R version 3.3.3 (R Development

Core Team, 2017).

3. RESULTS

3.1 Immunogenetic variation

Among the 293 tested individuals, we found nine functional alleles for *Mhc* class II-*Drb* gene, coding

for different amino acid sequences. One haplotype (CacaDRB*0302) was common (60%), two other

(CacaDRB*0201 and CacaDRB*0102) had intermediate frequency (18% and 10% respectively) while

the six others were rare (<5%). Observed heterozygosity (H_O) reached 0.55. We isolated six different

haplotypes for *Tlr2* in 327 genotyped individuals, all corresponding to different amino acids

sequences. Two haplotypes ("CATCG" and "TGCCG") showed particularly high frequency (35% and

51% respectively) while the four others were rare (< 5%). The two most common haplotypes

belonged to two highly divergent genetic clusters and were separated by seven substitutions

(including four non-synonymous ones) (Quéméré et al., 2015). The *Tlr4* gene exhibited four

functional haplotypes in 327 individuals including two haplotypes ("CGG" and "GAG") with moderate

to high frequency (26% and 55% respectively) and two haplotypes with weak frequencies (10% and

7%). Lastly, we isolated six haplotypes (in 324 individuals) for the *Tlr5* among which two ("GCTCG",

"ACCTG") had high frequencies (38% and 31% respectively), one ("ATCCG") had intermediate

frequency (18%) and the three others were rare (<5%). Observed heterozygosity was relatively high

for the three *Tlr* genes (H_O = 0.62, 0.64 and 0.69, *Tlr2*, *Tlr4* and *Tlr5* respectively).

3.2 Ecological patterns of parasitism

Seroprevalence of *Toxoplasma* and *Chlamydia* reached 35% and 16% respectively (Table 1). Among

the 232 annual captures for which both pathogens were screened (196 different deer), only 18

individuals (7%) were co-infected by two pathogens simultaneously while 93 (40%) and 36 (15%)

animals were infected solely by *Toxoplasma* and *Chlamydia* respectively. We did not reveal any

significant pathogen association using the screening analysis (*P* = 0.39) (Figure S2).

For the two studied pathogens, we found a significant positive effect of annual and spatial

variation in environmental conditions (modeled as a linear effect of API and SPI) on the prevalence of

the two pathogens: API ranged between 0.16 (in 2014) and 0.83 (in 2009) for *Toxoplasma*; between 0

(in 2012 and 2013) and 0.37 (in 2010) for *Chlamydia*. SPI ranged between 0.15 (in site 9) and 0.55 (in

site 1) for *Toxoplasma;* between 0.11 (in site 9) and 0.22 (in site 7) for *Chlamydia.* Additionally, the

Toxoplasma reference (non-genetic) model revealed a significant effect of age class, with lower

antibody prevalence in juvenile individuals compared to adults (OR = 0.38 [CI 0.15-0.91]). Lastly, we

found a significant positive effect of API_{TOXO}, API_{CHLA} and negative effect of SPI_{CHLA} on co-infection

status. Most co-infections (16 out of 18) occurred in 2009 and 2010 (Table 1).

3.3 Genetic effects on pathogen infection

Toxoplasma gondii

The inclusion of *Tlr2* genotype significantly improved the *Toxoplasma* infection non-genetic model.

The best-fitted model (delta AICc = -4.68 with non-genetic model) (Table S3) included an effect of

individual age-class, the yearly prevalence (API) and the number of copies of the "TGCCG" haplotype

of *Tlr2*. Roe deer carrying two copies of this frequent haplotype (51%) showed a decreased

Toxoplasma seroprevalence (odd ratio OR = 0.24 [0.075-0.76]) compared to individuals without this

haplotype (Figure 1a). The model with the same haplotype fitted as a binary effect

(presence/absence) received a slightly lower support (delta AICc = -2.86). We also observed a strong

relationship between *Toxoplasma* prevalence and *Tlr4* heterozygosity (delta AICc = -3.86) or *Tlr4*

allelic distance (delta AICc = -4.19), each time in interaction with the annual probability of infection

(Figure 2). Roe deer carrying two different or distant copies of *Tlr4* had a significantly lower

probability of being infected with *Toxoplasma* than other animals, in years when population

exposure was high. The opposite pattern was observed when population exposure was low. Other
models testing rare allele advantage received a weaker support than the reference non-genetic
model. We did not detect any association between *Toxoplasma* infection and *Tlr5* or *Mhc-Drb*
genetic variation.

Chlamydia abortus

We revealed a strong relationship between *Tlr2* genotype and *Chlamydia* prevalence. The best-
fitted model (delta AIC = -6.15 with reference model) included a positive effect of API, SPI and the
number of copies of the most-common haplotype ("TGCCG"). Homozygous individuals for this allele
had an increased probability to be infected (OR = 9 [1.11-80]) (Table S3, Figure 1). Alternatively, the
second best-fitted model (delta AIC = -5.48) indicated that deer carrying one or two copies of the
second most-frequent haplotype ("CATCG") were less infected with *Chlamydia* (OR = 0.46[0.13-1.68]
for deer carrying a single copy ; OR = 0.019 [0.004-0. 95] for deer carrying two copies). We also found
a significant association between *Tlr5* genotype and *Chlamydia* infection (delta AIC = -4 with
reference model). Individuals carrying rare alleles (<5%) were more likely to be infected (OR= 5.44
[1.02-28.9]) (Table S3). All models testing directional selection or heterozygosity advantage received
a weaker support than the minimal non-genetic model. No association was observed when
considering *Mhc-Drb* and *Tlr4* genes.

Co-infections

We did not find any relationship between the co-infection status of individuals and their
immunogenetic profiles (Table S3).

4. DISCUSSION

So far, very few studies, focusing on a limited number of taxa (most exclusively birds or rodents),
examined the mechanisms behind pathogen-mediated selective processes contemporarily acting on
immunity-related genes within natural populations (Grueber et al., 2013; Barbara Tschirren, Råberg,
& Westerdahl, 2011; Turner, Begon, Jackson, & Paterson, 2012). Being evolutionary ancient and
under strong functional constraints, genes involved in innate immune responses are expected to
evolve primarily under purifying selection and to exhibit limited polymorphism (Parham, 2003). Yet, it
is now quite clear that some of these genes, including *Tlr* genes encoding surface recognition
receptors, may have relaxed selective constraints and show large nucleotide diversity, sometimes
comparable to *Mhc* genes (Seabury et al., 2011). In a previous work, we have shown that the high
sequence and allelic diversity observed at roe deer *Tlr* genes could be maintained through balancing
selection (Quéméré et al., 2015). In this study, we developed a multi-gene/multi-pathogen

association approach to bring new insights into the mechanisms underlying such balancing selection. Most of the pathogen-gene associations found here suggested a key role of fluctuating directional selection and, to a lesser extent, of heterozygous advantage. We only found a limited influence of negative-frequency dependence process. We showed that different *Tlr* genes may be associated with resistance to the same microparasite (here *Tlr2* and *Tlr4* with *Toxoplasma* or *Tlr2* and *Tlr5* with *Chlamydia*) in agreement with the general idea that pathogens interact with many immune recognition receptors. We also revealed that different pathogens may act antagonistically on the polymorphism of a given gene (here *Tlr2* with *Toxoplasma* and *Chlamydia*).

4.1 Predominant role of directional and fluctuating selection on *Tlr* gene variation

We revealed that the polymorphism of *Tlr2* is predominantly shaped by directional selection exerted by both *Toxoplasma* and *Chlamydia*. Roe deer carrying the most or second most frequent haplotype were less likely to be infected either by *Toxoplasma* or *Chlamydia*. We know that these two abortive pathogens may have significant impact on the annual reproductive success of ungulates (Pioz et al., 2008). TLRs have been shown to activate the complement system, a component of innate immunity known to be important for resistance to microparasites during early infection (Raby et al., 2011). A higher affinity of a *Tlr* haplotype to *Toxoplasma* or *Chlamydia* ligands may result in a stronger complement response and thus in an increased resistance to these pathogens (Tschirren et al., 2013).

Our results also revealed a high temporal and spatial heterogeneity in both pathogen prevalence probably owing to variation in exposure. Annual variation in pathogen prevalence may be related to both climate and host factors. Climatic conditions such as temperature or humidity are known to affect pathogen survival in the environment and may influence host transmission *via* the quantity of infected aerosols for *Chlamydia* (Tang, 2009) or oocysts for *Toxoplasma* (Gotteland et al., 2014). For example, Sevila et al. (2014) showed that *Toxoplasma* prevalence in roe deer was higher in mild and wet years. The high turnover of individuals in the population in relation with a strong harvest pressure may also partly explain why we observed a so high inter-annual variability (Candela et al., 2014). Moreover, habitat heterogeneity and land use within roe deer's home range may explain the observed spatial variation in pathogen exposure. For example, *Toxoplasma* prevalence increased with the proportion of human dwellings (a proxy of the cat presence) within home range.

Spatial and temporal variation in pathogen-mediated directional selection means that *Tlr* haplotypes inheritance may vary among years and according to where females established their home range. This supported the hypothesis that fluctuating selection is a key process in the maintenance of balanced polymorphism in *Tlr* genes in this population. This finding is fully consistent with the Simpsonian 'dynamic' Adaptive Landscape concept (Simpson, 1944) that states that genetic variance

and evolution of fitness characters in the wild is mainly shaped by spatial and temporal fluctuations

in selection, specially variation in biotic interactions (Calsbeek, Gosden, Kuchta, & Svensson, 2012;

Siepielski, DiBattista, & Carlson, 2009). This general concept can be easily applied to host-pathogen

interactions and immunity-related gene evolution. Different habitats in terms of biotic and abiotic

factors are likely to support distinct pathogen communities and so distinct pathogen-mediated-

selection regime (Eizaguirre & Lenz, 2010). Using simulation-based models, Hedrick (2002)

demonstrated that variation in space and time in pathogen resistance can in itself be a strong

mechanism for the maintenance of polymorphism at immunity-related genes. In agreement, several

empirical studies found that *Mhc* genetic diversity varies across space in a mosaic of habitats likely

due to differences in pathogen communities (Alcaide, Edwards, Negro, Serrano, & Tella, 2008;

Eizaguirre & Lenz, 2010; Landry & Bernatchez, 2001). We noted that few studies have convincingly

elucidated the ecological drivers of this immunogenetic variation and that all focused on *Mhc* genes.

4.2 Antagonistic pleiotropy and potential immunogenetic trade-off

Interestingly, we observed that different haplotypes of a same *Tlr* gene (here *Tlr2*) may be associated

with the infection status of the two pathogens. Pleiotropic effects in *Tlr2* were expected since this

receptor is known to be implicated in the recognition of many bacterial, fungal, protozoa (de Oliviera

Nascimento, Massari, & Wetzler, 2012) including *Toxoplasma* (Yarovinsky, 2008) and *Chlamydia*

(Beckett et al., 2012; Darville et al., 2003). Here, the most common *Tlr2* haplotype ("TGCCG") (51%)

was associated with decreased *Toxoplasma* infection and increased *Chlamydia* infection while the

concurrent haplotype ("CATCG") (35%) showed opposite trends. This result provides an example of

antagonistic pleiotropy, which has been proposed as a widespread mechanism by which genetic

variation is maintained within populations (Roff & Fairbairn, 2007; Rose, 1982), in particular at

immunity-related genes (Aderem & Ulevitch, 2000; Carter & Nguyen, 2011). Antagonistic pleiotropy

has been rarely evidenced in natural populations (Turner et al., 2011) and, to our knowledge, this

study is the first demonstration with regard to a *Tlr* gene. Two non-exclusive mechanisms may be

invoked to explain this antagonistic effect: decrease of intracellular pathogen competition (Loiseau et

al., 2008) and host immunogenetic trade-off (Kamath et al., 2014). *Toxoplasma* and *Chlamydia* use

similar strategies to invade cells route and compete for the same nutrient pools in co-infected cells

(Romano & Coppens, 2013). Therefore, antagonistic effects might arise because *Tlr2* gene alters the

competitive interactions between the two pathogens. In other words, because individuals carrying

the most frequent *Tlr2* haplotype are less infected by *Toxoplasma*, this may indirectly promote

infection by *Chlamydia* without a direct role of the second most frequent haplotype (and *vice versa*).

In a study on house sparrows, Loiseau et al. (2008) found antagonistic effects of a *Mhc* class I gene on

multiple malarial parasite strains. In this particular case, deleterious 'susceptibility alleles could be

maintained in the population due to within-host competition between malaria parasites. Here, we

did not observe fewer co-infections than expected given the prevalence of the two pathogens.

Moreover *Tlr2* genotype had no effect on individual co-infection status. So it is unlikely that

antagonistic effect may result from the competitive interactions between *Toxoplasma* and

Chlamydia.

The most likely hypothesis is the presence of a host immunogenetic trade-off (Kamath et al.,

2014). In this case, pathogens do not directly compete but *TLr2* genotype has a direct effect on the

resistance/susceptibility to both pathogens. The allele inferred to be beneficial for decreasing

Toxoplasma infection is also associated with increased susceptibility to *Chlamydia* (and *vice versa*).

This scenario is reinforced by the fact that the two concurrent *Tlr2* haplotypes exhibit very distinct

DNA and amino-acid sequences, what suggests marked functional differences. Similar patterns and

processes have previously been described for ticks and nematodes infections (Kamath et al., 2014;

Turner et al., 2011). To our knowledge, our study is the second one showing evidence of antagonistic

selection at non-*Mhc* gene in a wild population (see for the other one Turner et al., 2011).

4.3 Marginal role of heterozygous advantage

A negative relationship between gene heterozygosity and deer co-infection status was expected

under the heterozygosity advantage hypothesis (McClelland et al., 2003). Instead, we found only an

association between *Tlr4* heterozygosity or allelic divergence and the infection to a single pathogen

(*Toxoplasma*). This finding must be interpreted with caution since we analyzed two pathogens only.

However, Sin et al. (2014) also revealed several single gene-single pathogen associations in European

badger (*Meles meles*) and did not find any support for *Mhc* heterozygosity being associated with

fewer co-infecting pathogens despite examining 13 pathogenic species from a wide variety of taxa. In

our study, the absence of relationship between heterozygosity at immunity-related genes and co-

infection could be due to the low individual's probability of being exposed to both pathogens

Toxoplasma and *Chlamydia* across their life-time, explaining the low fitness benefit of heterozygous

genotypes. The lack of association between Mhc-Drb and the two pathogens supported the general

idea that MHC class II molecules principally bind exogenous antigens and are primarily involved in

the immune response to extracellular pathogens (Hughes & Yeager, 1998)

The strong association between *Tlr4* allelic divergence and *Toxoplasma* infection supported the

divergent allele advantage hypothesis (Wakeland et al., 1990). Individuals carrying highly distant

copies of *Tlr4* appeared less likely to be infected by *Toxoplasma* than those carrying one haplotype or

two genetically closer haplotypes, but in years of high *Toxoplasma* exposure only. This genetic effect

in interaction with inter-annual variation in pathogen exposure again emphasizes the importance of

fluctuating selection on immunogenetic polymorphism. We also found an advantage for homozygotes or individuals carrying two genetically close haplotypes when *Toxoplasma* exposure was low. However, further information on the genetic diversity of *Toxoplasma* in this deer population and on the co-evolutionary dynamics with *Tlr4* is needed to fully understand this complex pattern.

5. Conclusion

In conclusion, our study illustrates the importance of looking beyond *Mhc* genes (Acevedo-Whitehouse & Cunningham, 2006) and single snapshot gene/pathogen association (Maizels & Nussey, 2013) in wildlife immunogenetic studies. In the present case, we did not find any influence of *Mhc* diversity on the resistance to two microparasites. Moreover, we highlighted the importance of antagonistic effects and fluctuating selection on the maintenance of innate immunogenetic diversity, what could not have been possible using a classical single gene-single pathogen approach.

ACKNOWLEDGMENTS

This work was supported by the French National Institute for Agricultural Research (INRA). We thank the local hunting associations, the Fédération Départementale des Chasseurs de Haute-Garonne. We thank numerous co-workers and volunteers for their assistance in field data collection, in particular J-M. Angibault, B. Cargnelutti, J-L. Rames, J. Joachim, B. Lourtet, D. Picot and N. Cebe. We also thank Karine Laroucau and Rachid Aaziz from ANSES the laboratory for animal health in Maison-Alfort (France) for her support and her expertise on *C. abortus* results. We also thank Monica G Candela, Julie Sevila and co-workers for their support on laboratory work (serology). Genetic data used in this work were produced through molecular genetic analysis technical facilities of the labex "Centre Méditerranéen de l'Environnement et de la Biodiversité".

REFERENCES

Acevedo-Whitehouse, K., & Cunningham, A. (2006). Is MHC enough for understanding wildlife immunogenetics? *Trends Ecol Evol*, *21*, 433–438.

Aderem, A., & Ulevitch, R. J. (2000). Toll-like receptors in the induction of the innate immune response. *Nature*, *406*(6797), 782.

Akira, S., Takeda, K., & Kaisho, T. (2001). Toll-like receptors: critical proteins linking innate and acquired immunity. *Nature Immunology*, *2*(8), 675–680.

Alcaide, M., Edwards, S. V., Negro, J. J., Serrano, D., & Tella, J. L. (2008). Extensive polymorphism and geographical variation at a positively selected MHC class II B gene of the lesser kestrel (Falco naumanni). *Molecular Ecology*, *17*(11), 2652–2665.

Bates, D., Maechler, M., & Bolker, B. (2012). lme4: Linear mixed-effects models using S4 classes.

435   Beckett, E. L., Phipps, S., Starkey, M. R., Horvat, J. C., Beagley, K. W., Foster, P. S., & Hansbro, P. M. (2012).
436   TLR2, but not TLR4, is required for effective host defence against Chlamydia respiratory tract infection
437   in early life. *PLoS One*, *7*(6), e39460.

438   Calsbeek, R., Gosden, T. P., Kuchta, S. R., & Svensson, E. I. (2012). Fluctuating selection and dynamic adaptive
439   landscapes. *The Adaptive Landscape in Evolutionary Biology*, 89–109.

440   Candela, M. G., Serrano, E., Sevila, J., León, L., Caro, M. R., & Verheyden, H. (2014). Pathogens of zoonotic and
441   biological importance in roe deer (Capreolus capreolus): Seroprevalence in an agro-system population
442   in France. *Research in Veterinary Science*, *96*(2), 254–259.

443   Carter, A. J., & Nguyen, A. Q. (2011). Antagonistic pleiotropy as a widespread mechanism for the maintenance
444   of polymorphic disease alleles. *BMC Medical Genetics*, *12*(1), 160.

445   Cook, L. M. (2003). The rise and fall of the carbonaria form of the peppered moth. *The Quarterly Review of*
446   *Biology*, *78*(4), 399–417.

447   Darville, T., O'Neill, J. M., Andrews, C. W., Nagarajan, U. M., Stahl, L., & Ojcius, D. M. (2003). Toll-like receptor-
448   2, but not Toll-like receptor-4, is essential for development of oviduct pathology in chlamydial genital
449   tract infection. *The Journal of Immunology*, *171*(11), 6187–6197.

450   de Oliviera Nascimento, L., Massari, P., & Wetzler, L. M. (2012). The role of TLR2 in infection and immunity.
451   *Frontiers in Immunology*, *3*, 79.

452   Doherty, P. C., & Zinkernagel, R. M. (1975). Enhanced immunological surveillance in mice heterozygous at the
453   H-2 gene complex. *Nature*, *256*(5512), 50.

454   Dubey, J. P. (2009). Toxoplasmosis in sheep—the last 20 years. *Veterinary Parasitology*, *163*(1–2), 1–14.

455   Eizaguirre, C., & Lenz, T. L. (2010). Major histocompatibility complex polymorphism: dynamics and
456   consequences of parasite-mediated local adaptation in fishes. *Journal of Fish Biology*, *77*(9), 2023–
457   2047.

458   Eizaguirre, C., Lenz, T. L., Kalbe, M., & Milinski, M. (2012). Rapid and adaptive evolution of MHC genes under
459   parasite selection in experimental vertebrate populations. *Nature Communications*, *3*, 621.

460   Froeschke, G., & Sommer, S. (2012). Insights into the complex associations between MHC class II DRB
461   polymorphism and multiple gastrointestinal parasite infestations in the striped mouse. *PLoS One*, *7*(2),
462   e31820.

463   Gotteland, C., Aubert, D., Gibert, P., Moinet, M., Klein, F., Game, Y., … Gilot-Fromont, E. (2014). Toxoplasmosis
464   in natural populations of ungulates in France: prevalence and spatiotemporal variations. *Vector-Borne*
465   *and Zoonotic Diseases*, *14*(6), 403–413.

466   Grueber, C. E., Wallis, G. P., & Jamieson, I. G. (2013). Genetic drift outweighs natural selection at toll-like
467   receptor (TLR) immunity loci in a re-introduced population of a threatened species. *Molecular Ecology*,
468   *22*(17), 4470–4482.

469   Hedrick, P. W. (1999). Antagonistic pleiotropy and genetic polymorphism: a perspective. *Heredity*, *82*(2), 126–
470   133.

471   Hedrick, P. W. (2002). Pathogen resistance and genetic variation at MHC loci. *Evolution*, *56*(10), 1902–1908.

472   Hedrick, P. W. (2006). Genetic polymorphism in heterogeneous environments: the age of genomics. *Annu. Rev.*
473   *Ecol. Evol. Syst.*, *37*, 67–93.

474   Hill, A. V. (1991). HLA associations with malaria in Africa: some implications for MHC evolution. In *Molecular*
475   *evolution of the major histocompatibility complex* (pp. 403–420). Springer.

476   Hughes, A. L., & Nei, M. (1992). Models of host-parasite interaction and MHC polymorphism. *Genetics*, *132*(3),
477   863.

478   Hughes, A. L., & Yeager, M. (1998). Natural selection and the evolutionary history of major histocompatibility
479   complex loci. *Frontiers in Bioscience: A Journal and Virtual Library*, *3*, d509-516.

480   Johnston, S. E., Gratten, J., Berenos, C., Pilkington, J. G., Clutton-Brock, T. H., Pemberton, J. M., & Slate, J.
481   (2013). Life history trade-offs at a single locus maintain sexually selected genetic variation. *Nature*,
482   *502*(7469), 93.

483   Kaisho, T., & Akira, S. (2004). Pleiotropic function of Toll-like receptors. *Microbes and Infection*, *6*(15), 1388–
484   1394.

485   Kamath, P. L., Turner, W. C., Küsters, M., & Getz, W. M. (2014). Parasite-mediated selection drives an
486   immunogenetic trade-off in plains zebras (Equus quagga). *Proceedings of the Royal Society of London*
487   *B: Biological Sciences*, *281*(1783), 20140077.

488   Klein, J., & Figueroa, F. (1986). Evolution of the major histocompatibility complex. *Critical Reviews in*
489   *Immunology*, *6*(4), 295–386.

490   Klein, S. L. (2000). The effects of hormones on sex differences in infection: from genes to behavior.
491   *Neuroscience & Biobehavioral Reviews*, *24*(6), 627–638.

492   Landry, C., & Bernatchez, L. (2001). Comparative analysis of population structure across environments and
493   geographical scales at major histocompatibility complex and microsatellite loci in Atlantic salmon
494   (Salmo salar). *Molecular Ecology*, *10*(10), 2525–2539.

495   Levene, H. (1953). Genetic equilibrium when more than one ecological niche is available. *The American*
496   *Naturalist*, *87*(836), 331–333.

497   Lewontin, R. C., & Hubby, J. L. (1966). A molecular approach to the study of genic heterozygosity in natural
498   populations. II. Amount of variation and degree of heterozygosity in natural populations of Drosophila
499   pseudoobscura. *Genetics*, *54*(2), 595.

500   Loiseau, C., Zoorob, R., Garnier, S., Birard, J., Federici, P., Julliard, R., & Sorci, G. (2008). Antagonistic effects of a
501   Mhc class I allele on malaria-infected house sparrows. *Ecology Letters*, *11*(3), 258–265.

502   Maizels, R. M., & Nussey, D. H. (2013). Into the wild: digging at immunology's evolutionary roots. *Nature*
503   *Immunology*, *14*(9), 879.

504   McClelland, E. E., Penn, D. J., & Potts, W. K. (2003). Major histocompatibility complex heterozygote superiority
505   during coinfection. *Infection and Immunity*, *71*(4), 2079–2086.

506   Miller, C. M., Boulter, N. R., Ikin, R. J., & Smith, N. C. (2009). The immunobiology of the innate response to
507   Toxoplasma gondii. *International Journal for Parasitology*, *39*(1), 23–39.

Møller, A. P., & Saino, N. (2004). Immune response and survival. *Oikos*, *104*(2), 299–304.

Moore, A. J., & Moore, P. J. (1999). Balancing sexual selection through opposing mate choice and male
competition. *Proceedings of the Royal Society of London B: Biological Sciences*, *266*(1420), 711–716.

Mun, H.-S., Aosai, F., Norose, K., Chen, M., Piao, L.-X., Takeuchi, O., … Yano, A. (2003). TLR2 as an essential
molecule for protective immunity against Toxoplasma gondii infection. *International Immunology*,
15(9), 1081–1087.

Oliver, M. K., Telfer, S., & Piertney, S. B. (2009). Major histocompatibility complex (MHC) heterozygote
superiority to natural multi-parasite infections in the water vole (Arvicola terrestris). *Proceedings of*
the Royal Society of London B: Biological Sciences, *276*(1659), 1119–1128.

Parham, P. (2003). Innate immunity: the unsung heroes. *Nature*, *423*(6935), 20–20.

Pedersen, A. B., & Greives, T. J. (2008). The interaction of parasites and resources cause crashes in a wild
mouse population. *Journal of Animal Ecology*, *77*(2), 370–377.

Phillips, K. P., Cable, J., Mohammed, R. S., Herdegen-Radwan, M., Raubic, J., Przesmycka, K. J., … Radwan, J.
(2018). Immunogenetic novelty confers a selective advantage in host–pathogen coevolution.
Proceedings of the National Academy of Sciences, 201708597.

Pioz, M., Loison, A., Gauthier, D., Gibert, P., Jullien, J.-M., Artois, M., & Gilot-Fromont, E. (2008). Diseases and
reproductive success in a wild mammal: example in the alpine chamois. *Oecologia*, *155*(4), 691–704.

Piry, S., Guivier, E., Realini, A., & Martin, J. (2012). SESAME Barcode: NGS-oriented software for amplicon
characterization–application to species and environmental barcoding. *Molecular Ecology Resources*,
12(6), 1151–1157.

Quéméré, E., Galan, M., Cosson, J.-F., Klein, F., Aulagnier, S., Gilot-Fromont, E., … Charbonnel, N. (2015).
Immunogenetic heterogeneity in a widespread ungulate: the European roe deer (Capreolus
capreolus). *Molecular Ecology*. Retrieved from
http://onlinelibrary.wiley.com/doi/10.1111/mec.13292/full

Raby, A.-C., Holst, B., Davies, J., Colmont, C., Laumonnier, Y., Coles, B., … Labéta, M. O. (2011). TLR activation
enhances C5a-induced pro-inflammatory responses by negatively modulating the second C5a
receptor, C5L2. *European Journal of Immunology*, *41*(9), 2741–2752. doi:10.1002/eji.201041350

Roff, D. A., & Fairbairn, D. J. (2007). The evolution of trade-offs: where are we? *Journal of Evolutionary Biology*,
20(2), 433–447.

Romano, J. D., & Coppens, I. (2013). Host Organelle Hijackers: a similar modus operandi for Toxoplasma gondii
and Chlamydia trachomatis: co-infection model as a tool to investigate pathogenesis. *Pathogens and*
Disease, *69*(2), 72–86.

Rose, M. R. (1982). Antagonistic pleiotropy, dominance, and genetic variation. *Heredity*, *48*(1), 63.

Seabury, C. M., Bhattarai, E. K., Taylor, J. F., Viswanathan, G. G., Cooper, S. M., Davis, D. S., … Seabury, P. M.
(2011). Genome-wide polymorphism and comparative analyses in the white-tailed deer (Odocoileus
virginianus): a model for conservation genomics. *Plos One*, *6*(1), e15811.

Sevila, J., Richomme, C., Hoste, H., Candela, M. G., Gilot-Fromont, E., Rodolakis, A., … Verheyden, H. (2014).
Does land use within the home range drive the exposure of roe deer (Capreolus capreolus) to two
abortive pathogens in a rural agro-ecosystem? *Acta Theriologica*, *59*(4), 571–581.

Siepielski, A. M., DiBattista, J. D., & Carlson, S. M. (2009). It's about time: the temporal dynamics of phenotypic
selection in the wild. *Ecology Letters*, *12*(11), 1261–1276.

Simpson, G. G. (1944). Tempo and Mode in Evolution (Columbia Classics in Evolution Series.).

Sin, Y. W., Annavi, G., Dugdale, H. L., Newman, C., Burke, T., & MacDonald, D. W. (2014). Pathogen burden, co-
infection and major histocompatibility complex variability in the European badger (Meles meles).
Molecular Ecology, *23*(20), 5072–5088. doi:10.1111/mec.12917

Sommer, S. (2005). The importance of immune gene variability (MHC) in evolutionary ecology and
conservation. *Front Zool*, *2*(1), 16.

Spurgin, L., & Richardson, D. (2010). How pathogens drive genetic diversity: MHC, mechanisms and
misunderstandings. *Proc Biol Sci*, *277*, 979–988.

Takahata, N., & Nei, M. (1990). Allelic genealogy under overdominant and frequency-dependent selection and
polymorphism of major histocompatibility complex loci. *Genetics*, *124*(4), 967–978.

Tang, J. W. (2009). The effect of environmental parameters on the survival of airborne infectious agents.
Journal of The Royal Society Interface, *6*(Suppl 6), S737–S746. doi:10.1098/rsif.2009.0227.focus

Tollenaere, C., Bryja, J., Galan, M., Cadet, P., Deter, J., Chaval, Y., … Laakkonen, J. (2008). Multiple parasites
mediate balancing selection at two MHC class II genes in the fossorial water vole: insights from
multivariate analyses and population genetics. *Journal of Evolutionary Biology*, *21*(5), 1307–1320.

Trowsdale, J., & Parham, P. (2004). Mini-review: defense strategies and immunity-related genes. *European
Journal of Immunology*, *34*(1), 7–17.

Tschirren, B, Andersson, M., Scherman, K., Westerdahl, H., Mittl, P., & Raberg, L. (2013). Polymorphisms at the
innate immune receptor TLR2 are associated with Borrelia infection in a wild rodent population. *Proc
Biol Sci*, *280*, 20130364.

Tschirren, Barbara, Råberg, L., & Westerdahl, H. (2011). Signatures of selection acting on the innate immunity
gene Toll-like receptor 2 (TLR2) during the evolutionary history of rodents. *Journal of Evolutionary
Biology*, *24*(6), 1232–1240.

Turner, A. K., Begon, M., Jackson, J. A., Bradley, J. E., & Paterson, S. (2011). Genetic Diversity in Cytokines
Associated with Immune Variation and Resistance to Multiple Pathogens in a Natural Rodent
Population. *PLoS Genet*, *7*(10), e1002343. doi:10.1371/journal.pgen.1002343

Turner, A. K., Begon, M., Jackson, J. A., & Paterson, S. (2012). Evidence for selection at cytokine loci in a natural
population of field voles (Microtus agrestis). *Molecular Ecology*, *21*(7), 1632–1646.

Vaumourin, E., Vourc'h, G., Telfer, S., Lambin, X., Salih, D., Seitzer, U., … Gasqui, P. (2014). To be or not to be
associated: power study of four statistical modeling approaches to identify parasite associations in
cross-sectional studies. *Frontiers in Cellular and Infection Microbiology*, *4*, 62.

Wakeland, E. K., Boehme, S., She, J. X., Lu, C.-C., McIndoe, R. A., Cheng, I., … Potts, W. K. (1990). Ancestral
polymorphisms of MHC class II genes: divergent allele advantage. *Immunologic Research*, *9*(2), 115–
122.

Wegner, K. M., Kalbe, M., Kurtz, J., Reusch, T. B., & Milinski, M. (2003). Parasite selection for immunogenetic
optimality. *Science*, *301*(5638), 1343–1343.

Wellenreuther, M. (2017). Balancing selection maintains cryptic colour morphs. *Molecular Ecology*, *26*(22),

6185–6188.

Wittmann, M. J., Bergland, A. O., Feldman, M. W., Schmidt, P. S., & Petrov, D. A. (2017). Seasonally fluctuating

selection can maintain polymorphism at many loci via segregation lift. *Proceedings of the National*

Academy of Sciences, *114*(46), E9932–E9941.

Yang, X., & Joyee, A. (2008). Role of toll-like receptors in immune responses to chlamydial infections. *Current*

Pharmaceutical Design, *14*(6), 593–600.

Yarovinsky, F. (2008). Toll-like receptors and their role in host resistance to Toxoplasma gondii. *Immunology*

Letters, *119*(1–2), 17–21.

Data Accessibility:

597   Haplotype DNA sequences: Primers and genBank accessions are in Table S1. SNP positions and
598   characteristics are in Table S2. Genotypes of all individuals at each *Tlr* and *Mhc-Drb* locus and
599   environmental/parasitological data can be found on Dryad: Dryad entry doi:10.5061/dryad.qt0079b.

Authors' contributions

601   EQ, EGM and NC designed the study and interpreted the data; AJMH, HV and YC led the data
602   collection process on the field; MG, JM and EQ carried out the *Tlr* and *Mhc-Drb* genotyping; EQ, PH,
603   NM and MF performed the statistical analyses; EQ led the writing, all authors contributed to the
604   manuscript and gave final approval for publication.

Tables and Figures.

606 **Table 1 - Summary of the pathogen** data (number of samples per year in brackets). NA indicates the years for
607 which no data were available.

Pathogen	Number of samples	Number of individuals	Overall seroprevalence	Seroprevalence evaluated per year								
				2008	2009	2010	2011	2012	2013	2014	2015	2016
Toxoplasma	351	277	0.35	0.23 (40)	0.83 (42)	0.56 (43)	0.24 (50)	0.21 (42)	0.24 (38)	0.16 (31)	0.3 (49)	0.43 (16)
Chlamydia	232	196	0.16	0.00 (40)	0.14 (41)	0.37 (43)	0.28 (50)	0.05 (42)	0.00 (16)	NA	NA	NA
coïnfection	232	196	0.08	0 (40)	5 (41)	11 (43)	2 (50)	0 (42)	0 (16)	NA	NA	NA

Figure 1. Genetic polymorphisms at *Tlr2* are associated with both *Chlamydia* and *Toxoplasma*
serological status. Predicted prevalence of *Toxoplasma* and *Chlamydia* infection in roe deer with the
number of copies (0/1/2) of the most common haplotype ("TGCCG", 51%).

Figure 2. Infection to *Toxoplasma* is associated with *Tlr4* allelic distance. Predicted prevalence of
Toxoplasma infection in roe deer in function of the distance (in base pairs) between the two alleles of
the *Tlr4* gene with a positive interaction with the annual probability of infection (Low API = 0.3,
Intermediate = 0.5 and high = 0.7)

Supporting information

Table S1. Details of primer sequences, product size and genbank accessions.

Table S2. Summary of the 13 *Tlr* SNPs genotyped using the KASPar SNP genotyping system.

Table S3. Details of the mixed linear models of *Toxoplasma*, *Chlamydia* infection or co-infection for
each gene (*Trl2*, *Tlr4*, *Tlr5* and *MHC-Drb*).

Figure S1. Map of the study area.

Figure S2. Outputs of the screening association analysis.