Fluctuating pathogen-mediated selection favours the maintenance of innate
immune gene polymorphism in a widespread wild ungulate

Erwan Quéméré[1*], Pauline Hessenauer[2], Maxime Galan[3], Marie Fernandez[1], Joël Merlet[1], Yannick Chaval[1],

Aidan J Mark Hewison[1], Nicolas Morellet[1], Hélène Verheyden[1], Emmanuelle Gilot-Fromont[4,5], Nathalie

Charbonnel[3]

[1]CEFS, INRA, Université de Toulouse, Castanet-Tolosan Cedex F-31326, France

[2]Département de Foresterie, Université Laval, Quebec, Canada

[3] CBGP, INRA, CIRAD, IRD, Montpellier SupAgro, Univ Montpellier, Montpellier, France

[4]Université de Lyon, Université Lyon 1, UMR CNRS 5558, Villeurbanne, France,

[5]Université de Lyon, VetAgro Sup, Marcy l'Etoile, France

12
13   * Corresponding author:

14   erwan.quemere@inra.fr. Phone: (+33) 561 285 497. Fax: (+33) 561 285 500.

Abstract (233 words)

Toll-like Receptors (TLR) play a central role in recognition and host frontline defence against a wide range of pathogens. A number of recent studies have shown that *Tlrs* often exhibit a large polymorphism in natural populations. Yet, there is little knowledge on how this polymorphism is maintained and how it influences disease susceptibility in the wild. We investigated the selective mechanisms by which multiple-pathogens drive the genetic diversity of *Tlr* genes, as well as the commonly-used *Mhc -Drb* in European roe deer (*Capreolus capreolus*), an abundant and widespread ungulate in Europe. In a previous work, we showed that some *Tlrs* exhibit high levels of genetic diversity and contemporary signatures of balancing selection in roe deer. To assess the relative contribution of "heterozygous advantage", and "fluctuating selection", we evaluated associations between *Mhc-Drb* or *Tlr* variants and infection with *Toxoplasma* and *Chlamydia*, two intracellular pathogens known to cause reproductive failure in ungulates. We showed that *Toxoplasma* and *Chlamydia* exposures vary significantly across space and time with few co-infection events detected, and that the two pathogens act antagonistically on *Tlr* polymorphism leading to fluctuating selection. By contrast, we found no evidence of association with *Mhc-Drb* and *Tlr* heterozygosity. Our study confirmed the importance of looking beyond *Mhc* genes in wildlife immunogenetic studies. It also emphasized the necessity to consider multiple pathogen challenges and their spatio-temporal variations to improve our understanding of vertebrate defence evolution against pathogens.

Keywords: balancing selection, habitat heterogeneity, Toll-like genes, antagonistic effects, roe deer

Running title: Fluctuating selection on *Tlr* genes in deer

1. INTRODUCTION

While evolutionary theory predicts that selection and random drift should deplete genetic diversity, a number of genes and genomic regions have maintained abundant polymorphism in natural populations with exceptionally high ratio of non-synonymous (protein-altering) to synonymous substitutions (Hedrick, 2006; Trowsdale & Parham, 2004). Balancing selection is a collective term for evolutionary processes that adaptively maintain variation in populations. It is a widespread process driving the diversity of genes involved in numerous biological functions including pathogen resistance (Sommer, 2005), color polymorphism (Wellenreuther, 2017) or mate choice/male competiveness (Johnston et al., 2013; Moore & Moore, 1999). However, identifying the mechanisms driving balancing selection has been a major challenge. At least three not-mutually exclusive processes (putting aside sexual selection) can act in synergy: the "heterozygote advantage" or "overdominance" hypothesis (Doherty & Zinkernagel, 1975; Lewontin & Hubby, 1966) proposes that heterozygous individuals have a higher relative fitness, on average, than homozygous individuals. The "fluctuating selection hypothesis" (Hill, 1991; Levene, 1953) proposes that high genetic diversity may be maintained when spatially and/or temporally varying directional selection occurs with sometime antagonistic effects on fitness (Wittmann, Bergland, Feldman, Schmidt, & Petrov, 2017). Third, the "negative frequency-dependent selection" (also called rare-allele advantage) hypothesis suggests that the fitness-advantage of alleles is inversely proportional to their frequency in the population as a result of the host-pathogen (Phillips et al., 2018; Takahata & Nei, 1990) or prey-predator co-evolutionary dynamics (Cook, 2003). This third mechanism is particularly challenging to evidence in natural systems as it requires long time series of species prevalence and allele frequencies of genes involved in "Red Queen" co-evolution (Spurgin & Richardson, 2010, but see Charbonnel & Pemberton, 2005).

Immunity-related genes are ideal candidates for studying how adaptive genetic diversity is maintained in wild populations because of their direct effect on survival (Møller & Saino, 2004) or reproductive success (Pedersen & Greives, 2008). The genes of the major histocompatibility complex (MHC) are often used as a model since they show an exceptional polymorphism in vertebrates (Spurgin & Richardson, 2010). Their structure and mechanisms of interaction with antigens are well described (Klein & Figueroa, 1986). Furthermore, their ability to trigger a rapid adaptive response to varying pathogen-mediated selective pressures (PMS) has been demonstrated experimentally (Eizaguirre, Lenz, Kalbe, & Milinski, 2012; Phillips et al., 2018). Over the last few years, a growing body of literature identified PMS at *Mhc* genes in a large range of taxa (see Spurgin & Richardson, 2010 for a review) but few of them convincingly elucidated the relative influence of the three mechanisms mentioned above (but see Oliver et al., 2009; Philips et al., 2018). One likely reason is

that most studies modelling associations between a single gene (most often a *Mhc* gene) and only
one pathogen. However, decreasing sequencing costs now offer new opportunities to examine
multiple genes and diseases in a single study, leading to a more realistic view of the functioning of
the immune system and of the ecological context in which it evolves.

Mhc, although important, only corresponds to a fraction of the genetic variation in pathogen
resistance (Acevedo-Whitehouse & Cunningham, 2006). Before the adaptive immunity has an
opportunity to intervene in the host response, invading pathogens interact with different effectors of
innate immunity that recognize specific structures (Jepson et al., 1997; Lam-Yuk-Tseung & Gros,
2003). Therefore many genes encoding proteins involved in pathogen recognition or immune
regulation may be associated with the resistance against the same pathogen (Turner, Begon, Jackson,
Bradley, & Paterson, 2011). In particular, genes encoding Toll-like receptors (TLR) play a key role in
the host frontline defence against a wide range of microparasites such as fungi, bacteria, protozoa.
This family of genes conserved in both invertebrate and vertebrate lineages are critical for the
stimulation of many immune pathways such as inflammation, innate or acquired immunity (Akira,
Takeda, & Kaisho, 2001). A multitude of studies have revealed associations between *Tlr* gene
polymorphism and infectious diseases in humans and domestic species (Mun et al., 2003; Yang &
Joyee, 2008). However, very few (most focusing on birds or rodents) have investigated their role in
host-pathogen interactions in natural environment and their contemporary mechanisms of evolution
at a population level (but see Kloch et al., 2018; Tschirren et al., 2013).

Moreover, infection with multiple pathogens is a real world rule (Maizels & Nussey, 2013) and
some selective mechanisms such as heterozygous advantage or fluctuating selection may better be
detected when accounting for the multi-pathogen context (Sin et al., 2014b; Wegner, Kalbe, Kurtz,
Reusch, & Milinski, 2003). Most immunity-related genes have pleiotropic effects on resistance
against several pathogens (Hedrick, 1999) and the maintenance of 'susceptible' (deleterious) alleles
in a wild population can be due to antagonist effects of a given immunity-related gene on several
pathogens (*i.e.* antagonistic pleiotropy). To date, only few studies have examined the selective
effects of multiple pathogens on host immunogenetic patterns and the great majority focused on
Mhc genes (Tollenaere et al., 2008 on voles; Loiseau et al., 2008 on house sparrows ; Kamath et al.,
2014 in plain zebras; Froeschke & Sommer, 2012 on striped mouse but see Antonides et al., 2019).

Here, we investigated the selective mechanisms by which multiple pathogens could drive the
genetic diversity of immunity-related genes in European roe deer (*Capreolus capreolus*), the most
abundant and widespread ungulate in Europe. We studied a free-ranging population of roe deer
inhabiting an agricultural landscape in Southwestern France. This population is heavily exposed to a

range of pathogens that also infect livestock, including the abortive *Toxoplasma gondii* or *Chlamydia*

abortus (Candela et al., 2014). These two pathogens are known to have negative influence on the

survival and reproductive performances of ungulates (Dubey, 2009; Pioz et al., 2008). They may

therefore exert selective pressures on immunity-related genes. We focused on four genes encoding

pattern-recognition receptors (PRRs) that detect specific microbes or pathogen-associated molecular

patterns: the widely-used *Mhc class II-Drb* gene and three non-*Mhc* genes encoding Toll-like

receptors genes (*Tlr2, Tlr4, Tlr5*). *Mhc* and *Tlr* genes may have pleiotropic effects on the

resistance/susceptibility to pathogens harboring similar pathogen-associated molecular patterns

(Kaisho & Akira, 2004; Loiseau et al., 2008). Moreover, *Tlr* genes have been shown to play a major

role in host defence against multiple pathogens, including protozoans and gram-negative bacteria

(Beckett et al., 2012; Gazzinelli & Denkers, 2006; Miller, Boulter, Ikin, & Smith, 2009; Netea,

Wijmenga, & O'Neill, 2012; Yarovinsky, 2008). In a recent work, we showed that these genes exhibit

high levels of amino acid diversity within French roe deer populations including the population

studied here, hence suggesting an important functional repertoire (Quéméré et al., 2015). In

particular, *Tlr2* and *Tlr4* exhibited several haplotypes at moderate frequency and very low levels of

differentiation among roe deer populations compared to the patterns obtained with neutral markers.

These findings support the hypothesis that the polymorphism of these genes is shaped by balancing

selection although the underlying mechanism(s) remains to be investigated.

In this work, we assessed the role of "heterozygous advantage" and "fluctuating" selection in

shaping the signature of balancing selection observed in the European roe deer. We analyzed

whether the prevalence of *Toxoplasma* and *Chlamydia* was associated with specific patterns of

genetic diversity observed at the four immunity-related genes (e.g. heterozygosity/allelic divergence,

presence of dominant haplotypes), while accounting for ecological predictors (age, sex, year,

sampling site). Support for the *Mhc* heterozygote advantage hypothesis has been found in many

studies of mammals (Hedrick, 2012; Oliver et al., 2009; Sin et al., 2014b), including wild ungulates

(Brambilla, Keller, Bassano, & Grossen, 2018; Paterson, Wilson, & Pemberton, 1998). However, its

importance for innate immunity-related genes has rarely been assessed (but see Antonides et al.,

2019; Grueber et al., 2013). A recent study in birds (Grueber, Wallis, & Jamieson, 2013) has found

evidence for survival advantage associated with *Tlr4* heterozygote genotypes. Here, we expected

associations between *Tlr* and/or *Mhc-Drb* heterozygosity and infection with *Toxoplasma* and

Chlamydia, two pathogens employing similar strategies to invade host cells (Romano & Coppens,

2013). Moreover, there is a high temporal and spatial heterogeneity in pathogen exposure in the

study area (Candela et al., 2014; Sevila et al., 2014), and TLR and MHC receptors have pleiotropic

functions (Akira et al., 2001; Hedrick, 1999). As such we expected to find fluctuations in pathogen-

mediated directional selection acting on different *Mhc-Drb* or *Tlr* alleles. Such fluctuations could

underlie the signatures of balancing selection previously detected on these genes (Quéméré et al.,

2015).

2. MATERIAL AND METHODS

2.1 Study population and data collection

The study focused on a roe deer population inhabiting a heterogeneous agricultural landscape in

Southern France (43°13⬚N, 0°52⬚E). This area called "Vallons et Côteaux de Gascogne" (VCG) is part

of the Aterlier Pyrénées Garonne(https://pygar.omp.eu/). The area is covered by a mix of open fields

(including 38% meadows and pastures for cattle and sheep, and 32% crops), hedges (16%), small

woodlands patches (14%) and two larger forest blocks (5%). Individuals were caught by drive-netting

during winter (from January to March) between 2008 and 2016 in seven neighboring sites located a

few kilometers apart, with varying vegetation cover (Figure S1). Deer from these sites have been

shown to belong to the same panmictic population (Coulon et al., 2006; Gervais et al., 2019).

Individual sex, age class (juveniles: ≤ 1 year of age versus adults: > 1 year of age) and body mass were

recorded. Blood samples were collected for pathogen screening (see below) and ear punches for

genetic analyses. In total, we gathered samples from 433 annual captures corresponding to 328

different individuals (190 females and 138 males, 164 juveniles and 164 adults) among which 157

deer had been included in a previous immunogenetic studies (Quéméré et al., 2015). All applicable

institutional and European guidelines for the care and use of animals were followed. The study was

permitted by the land manager. All the procedures involving animals were approved by the Ethical

Committee 115 of Toulouse and authorized by the French Ministry in charge of ethical evaluation (n°

APAFIS#7880-2016120209523619v5).

2.2 *Tlr* and *Mhc-Drb* genotyping

The *Mhc-Drb* class II and *Tlr* (*Tlr2*, *Tlr4* and *Tlr5*) genes of 157 roe deer from VCG had been genotyped

in a previous study (Quéméré et al., 2015). We completed this dataset by genotyping 171 new

individuals using exactly the same procedure. DNA was extracted from alcohol-preserved tissues

using the DNeasy Blood and Tissue kit (QIAGEN). *Tlr* genes were genotyped using a two-step

approach: a pilot study on 32 individuals was first performed to identify polymorphic sites (SNPs) and

linkage-disequilibrium (LD) groups. We screened almost the entire coding region of the three *Tlr*

genes (82% in average) including the leucine-rich extracellular region of receptors involved in

antigen-binding. Detailed on primer sequences, SNP positions and codon changes are provided in

Quéméré et al. (2015). We then selected one SNP per LD group (primarily targeting non-synonymous

sites) that was genotyped for all individuals using the KASPar allele-specific genotyping system

provided by KBiosciences (Hoddesdon, UK). A total of 13 SNPs were typed including 5, 3 and 5 SNPs

for *Tlr2*, *Tlr4* and *Tlr5* respectively. Details on SNP position and codon change can be found in Table

S2. Lastly, haplotypes were reconstructed from the phased SNPs using the procedure implemented in

DNASP v5 (Librado & Rozas, 2008). The second exon of the *Mhc-Drb* class II gene encoding the ligand-

binding domain of the protein was amplified and sequenced using Illumina MiSeq system as detailed

in Quéméré et al. (2015). Haplotypes and individual genotypes were identified using the SESAME

barcode software (Piry, Guivier, Realini, & Martin, 2012). All sequences have been submitted to NCBI

Genbank (Accession nos. are in Table S1, Supporting information).

2.3 Pathogen screening

The serological status of roe deer for *Toxoplasma gondii* and/or *Chlamydia abortus* was analyzed

using classical enzyme-linked immunosorbent assays (ELISA) with specific commercial kits (Sevila et

al., 2014). The specificity and sensitivity of these kits were respectively 97.4 and 99.4 % for *T. gondii*

and 92.2 and 89 % for *C. abortus*. Although the kits were developed for domestic ruminants, they

were shown to be reliable and efficient in wild ungulates (Gotteland et al., 2014) with a good

concordance with classical test (e.g. high concordance (0.8) between ELISA and the Modified

Agglutination Test, a reference test for *Toxoplasma* in roe deer). According to the manufacturer's

instructions, blood samples with antibody recognition level (ARL) > 30% (respectively 40%) were

considered positive for *T. gondii* (respectively *C. abortus*) (see Sevila et al., 2014 for further details).

In total, we obtained the *Toxoplasma* and *Chlamydia* serological status of 277 (with 74 repeated

measures) and 196 (36 repeated measures) roe deer respectively (caught between 2008 and 2016,

Table S3). The individual repeatability (R) and its standard error (SE) for *Toxoplasma* and *Chlamydia*

seroprevalence were calculated using the R package 'rptR' (Stoffel, Nakagawa, & Schielzeth, 2017).

The proportion of individuals seropositive for both *Toxoplasma* and *Chlamydia* was examined in the

years where both pathogens were screened (between 2008 and 2013).

2.4 Statistical analyses

We analyzed the influence of immunity-related gene variation, environmental and life-history factors

on the seroprevalence of *T. gondii* or *C. abortus* (binary response variable – presence/absence) using

generalized linear mixed-effect models (GLMM) with a binomial family (using a logit link function).

We used multimodel inference to establish which explanatory variables had an effect, averaged over

plausible models (Burhnam & Anderson 2002; Grueber et al., 2011). Analyses were performed using

the *glmer* function implemented in the 'lme4' package (Bates, Maechler, & Bolker, 2012), 'MuMin'

v1.7.7 (Barton, 2009), and 'AICcmodavg' v1.25 (Mazerolle, 2013) in R version 3.3.3 (R Development

Core Team, 2017). Multimodel inference was conducted in two steps to reduce the number of
factors in a single model and the total number of models evaluated (Sin et al., 2014b).

In the first step, we investigated the effects of non-genetic biological and environmental factors
that possibly affect pathogen exposure. The starting model included roe deer sex (male *versus*
female) and age-class (juvenile *versus* adult) because both behaviour and host susceptibility may vary
between sexes and across an individual's lifetime (Klein, 2000) and because infection persistence is
expected to result in a positive age-prevalence relationship (Gotteland et al., 2014). We also included
the capture year (9 levels) and site (7 levels) as fixed effects to account for the spatial and inter-
annual variation in environmental conditions that may affect pathogen survival and thus encounter
probability (e.g. climate, food resources, population density). Roe deer classically inhabit a single
home range across the entire adult life that is relatively small (mean 0.76 km²) (Morellet et al., 2013)
and generally included in only one capture site. As such, we considered capture sites as the spatial
units shared by individuals captured at the same location. In all models, we included individual
identity as a random effect to account for correlation amongst different observations of the same
individual. We conducted an initial exploration of our data to ascertain their distribution and identify
potential outliers (Zuur, Ieno, Walker, Saveliev, & Smith, 2009). Model averaged parameter estimates
are provided with their unconditional standard errors (SE), 95% confidence intervals and relative
importance, after model averaging (ΔAIC< 5 relative to the best model) (Burnham, Anderson, &
Huyvaert, 2011). Significant variables were retained in a reduced model for use in step 2.

In the second step, we investigated the "heterozygote advantage hypothesis" by fitting a binary
fixed effect (heterozygote *vs* homozygote) and the "haplotype advantage" hypothesis by considering
associations between the infection status and the number of copies (0, 1 or 2) of the most or second
most frequent haplotype of *Tlr2*, *Tlr4*, *Tlr5* and *MHC-Drb*. We used variance inflation factors (VIF) to
measure collinearity among predictors and retained variables with VIF values <3 (Zuur et al., 2009).
To minimize multicollinearity, the effects of the two most frequent haplotypes of a gene and its
heterozygosity were evaluated in separate models. The identity of the site and/or the year were
included as random effects in all models because (i) individuals at a site/year all experience similar
conditions of exposure (see the results of step 1) and (ii) to account for spatial/temporal
autocorrelation in allele frequencies that may lead to spurious associations with pathogen
prevalence. In total, 256 models were evaluated for both *Toxoplasma* and *Chlamydia* (see Table S4
for the full list of models).

3. RESULTS

3.1 Immunogenetic variation

Among the 293 tested individuals, we found nine functional alleles for *Mhc* class II-*Drb* gene, coding
for different amino acid sequences. One haplotype (CacaDRB*0302) was common (60%), two other
(CacaDRB*0201 and CacaDRB*0102) had intermediate frequency (18% and 10% respectively) while
the six others were rare (<5%). Observed heterozygosity (H_O) reached 0.55. We isolated six different
haplotypes for *Tlr2* in 327 genotyped individuals, all corresponding to different amino acids
sequences. Two haplotypes ("CATCG" and "TGCCG") showed particularly high frequency (35% and
51% respectively) while the four others were rare (< 5%). The two most common haplotypes
belonged to two highly divergent genetic clusters and were separated by seven substitutions
(including four non-synonymous ones) (Quéméré et al., 2015). The *Tlr4* gene exhibited four
functional haplotypes in 327 individuals including two haplotypes ("CGG" and "GAG") with moderate
to high frequency (26% and 55% respectively) and two haplotypes with low frequencies (10% and
7%). Lastly, we isolated six haplotypes (in 324 individuals) for the *Tlr5* among which two ("GCTCG",
"ACCTG") had high frequencies (38% and 31% respectively), one ("ATCCG") had intermediate
frequency (18%) and the three others were rare (<5%). Observed heterozygosity was relatively high
for the three *Tlr* genes (H_O = 0.62, 0.64 and 0.69, *Tlr2*, *Tlr4* and *Tlr5* respectively).

3.2 Ecological patterns of parasitism

Seroprevalence of *Toxoplasma* reached 35% and was moderately repeatable among years at the
individual level (R=0.21, SE=0.06). By comparison, seroprevalence of *Chlamydia* was lower (16%) but
showed a higher individual consistency among capture events (R=0.70, SE=0.05). Among the 232
annual captures for which both pathogens were screened (196 different deer), only 18 individuals
(7%) had anti-*Toxoplasma* and anti-*Chlamydia* antibodies simultaneously while 93 (40%) and 36
(15%) animals were seropositive solely for *Toxoplasma* and *Chlamydia* respectively (Table 1). We
found a strong effect of "the year of capture" for the two pathogens with a significant increased
probability of being seropositive for *Toxoplasma* in 2009 and 2010 (Seroprevalence = 83% and 56%
respectively) and a decreased seroprevalence in 2009 for *Chlamydia* (16%) (Table 1). Additionally, we
revealed a significant relationship between *Toxoplasma* infection status, roe deer age and capture
site: antibody prevalence was lower in site 9 and in juveniles compared to adults (Table S5, Figure
S2).

3.3 Genetic effects on pathogen infection

We revealed a strong relationship between *Toxoplasma* seroprevalence and the number of copies of
specific *Tlr2* haplotypes. Roe deer with two copies of the most frequent "TGCCG" haplotype (*Tlr2-1*)
showed a decreased *Toxoplasma* seroprevalence (model-averaged parameter estimates (AES) = -1.03
[95% CI -1.91, -0.14]) while individuals carrying the second most frequent haplotype ("CATCG", *Tlr2-*

2) were more likely to be *Toxoplasma* seropositive (Figure 1) (AES = 0.71 [0.04, 1.38]). We also found

a negative relationship between *Toxoplasma* seroprevalence and individual heterozygosity at *Tlr2*

gene (AES = -0.63 [-1.24, -0.02]). *Toxoplasma* seroprevalence was not associated with *Tlr4*, *Tlr5* or

Mhc-Drb genetic variation (neither heterozygosity status nor specific alleles).

Similarly to *Toxoplasma*, we found a strong relationship between *Tlr2* genotype and *Chlamydia*

seroprevalence. However, we observed the opposite pattern of association: homozygous individuals

for the most-common *Tlr2* haplotype (*Tlr2-1*) had an increased probability to be seropositive (AES =

1.28 [0.06, 2.51]) while individuals carrying *Tlr2-2* were less likely to be seropositive for *Chlamydia*

(AES = -2.30 [-4.48, -0.12] for "CATCG" homozygous deer). We also revealed an association between

Tlr5 genotype and *Chlamydia* seroprevalence: roe deer with the "ACCTG" haplotype were less likely

to be Chlamydia seropositive (AES = -1.22 [-2.18, -0.25]). No association was observed when

considering *Mhc-Drb* and *Tlr4* genotypes (neither number of alleles nor heterozygosity status).

Figure 1. Model averaged parameter estimates and their 95% confidence intervals for genetic (*Tlr2, Tlr4, Tlr5 and Mhc-

DRB* genotype) predictors associated with *Toxoplasma* and *Chlamydia* seroprevalence (predictors included in the top model

sets with ΔAic < 5). * indicate a parameter with a significant effect. //* indicates extremely wide 95% confidence intervals.

4. DISCUSSION

So far, very few studies, focusing on a limited number of taxa (most exclusively birds or rodents),

examined the mechanisms behind pathogen-mediated selective processes contemporarily acting on

innate immunity-related genes within natural populations (Grueber et al., 2013; Barbara Tschirren,

Råberg, & Westerdahl, 2011; Turner, Begon, Jackson, & Paterson, 2012). Being evolutionary ancient

and under strong functional constraints, genes involved in innate immune responses are expected to

evolve primarily under purifying selection and to exhibit limited polymorphism (Parham, 2003). Yet, it
is now quite clear that some of these genes, including *Tlr* genes encoding surface recognition
receptors, may have relaxed selective constraints and show large nucleotide diversity, sometimes
comparable to *Mhc* genes (Seabury et al., 2011). In a previous work, we have shown that the high
sequence and allelic diversity observed at roe deer *Tlr* genes could be maintained through balancing
selection (Quéméré et al., 2015). In this study, we developed a multi-gene/multi-pathogen
association approach to bring new insights into the mechanisms underlying such balancing selection.
Our results suggested a key role of directional selection on specific *Tlr* haplotypes rather than
heterozygote advantage. We showed that different *Tlr* genes may be associated with resistance to
the same microparasite (here *Tlr2* and *Tlr5* with *Chlamydia*) in agreement with the general idea that
pathogens interact with many immune recognition receptors. We also revealed that different
pathogens may act antagonistically on the polymorphism of a given gene (here *Tlr2* with *Toxoplasma*
and *Chlamydia*). Given the high temporal and spatial heterogeneity in pathogen exposure in the
study area, this supports the hypothesis that fluctuations in pathogen-mediated directional selection
may promote the maintenance of *Tlr* genetic diversity.

4.1. High temporal and spatial variation in pathogen exposure

Our results revealed a high temporal heterogeneity in both pathogen seroprevalence at individual
and population levels. In contrast with the traditional view of lifelong persistence of *Toxoplasma*
infection (Tenter, Heckeroth, & Weiss, 2000), we observed low individual repeatability of
Toxoplasma seroprevalence at the within-individual level (R=0.21). This is in line with a previous
study on the same population that reported frequent seroconversion with initially seropositive
individuals becoming seronegative within 1 to 3 years (Sevila et al., 2014). There is no clear
consensus on the persistence rate of antibodies against *T. gondii* in domestic and wild ungulates.
Toxoplasma seroprevalence is known to increase with age in wild ungulate populations, but it is most
likely due to repeated exposure from the same environment rather than antibody persistence
(Gotteland et al., 2014; Opsteegh, Swart, Fonville, Dekkers, & Van Der Giessen, 2011). Overall, this
suggests that the presence of antibodies in the study population would reflect relatively recent
rather than long-term infection. The observed annual variation in pathogen prevalence at population
level may be related to both climate and host factors. Climatic conditions such as temperature or
humidity are known to affect pathogen survival in the environment and may influence host
transmission *via* the quantity of infected aerosols for *Chlamydia* (Tang, 2009) or oocysts for
Toxoplasma (Gotteland et al., 2014). For example, Sevila et al. (2014) showed that *Toxoplasma*
prevalence in roe deer was higher in mild and wet years and that *Chlamydia* prevalence increased in
cold years. The high turnover of individuals in the population in relation with a strong harvest

pressure may also partly explain why we observed a so high inter-annual variability in multi-cohort
samples (Candela et al., 2014). Moreover, we also observed a strong variation in *Toxoplasma*
exposure across sampling sites that most likely results from habitat heterogeneity and land use
within roe deer's small home range (e.g. proportion of human dwellings within home range as a
proxy of the cat presence, Sevila et al., 2014).

4.2 Predominant role of directional and fluctuating selection on *Tlr* gene variation

We revealed that *Tlr2* gene polymorphism is predominantly shaped by directional selection exerted
by both *Toxoplasma* and *Chlamydia*. Roe deer carrying the most frequent *Tlr2* haplotype were less
likely to be infected by *Toxoplasma* but more likely infected by *Chlamydia*. The opposite pattern was
revealed for the second most frequent haplotype. These two abortive pathogens may have
significant impact on the annual reproductive success of ungulates (Pioz et al., 2008). TLRs have been
shown to activate the complement system, a component of innate immunity known to be important
for resistance to microparasites during early infection (Raby et al., 2011). A higher affinity of a *Tlr*
haplotype to *Toxoplasma* or *Chlamydia* ligands may result in a stronger complement response and
thus in an increased resistance to these pathogens (Tschirren et al., 2013).

Spatial and temporal variations in pathogen-mediated directional selection suggest that *Tlr*
haplotypes inheritance may vary among years and according to sites where females established their
home range. This supported the hypothesis that fluctuating selection is a key process in the
maintenance of *Tlr* genes' balanced polymorphism in this population. This finding is fully consistent
with the Simpsonian 'dynamic' Adaptive Landscape concept (Simpson, 1944) stating that genetic
variance and evolution of fitness characters in the wild are mainly shaped by spatial and temporal
fluctuations in selection, specially variation in biotic interactions (Calsbeek, Gosden, Kuchta, &
Svensson, 2012; Siepielski, DiBattista, & Carlson, 2009). This general concept can be easily applied to
host-pathogen interactions and immunity-related gene evolution. Different habitats in terms of biotic
and abiotic factors are likely to support distinct pathogen communities and so distinct pathogen-
mediated-selection regime (Eizaguirre & Lenz, 2010). Using simulation-based models, Hedrick (2002)
demonstrated that variation in space and time in pathogen resistance can in itself be a strong
mechanism for the maintenance of polymorphism at immunity-related genes. In agreement, several
empirical studies found that *Mhc* genetic diversity varies across space in a mosaic of habitats likely
due to differences in pathogen communities (Alcaide, Edwards, Negro, Serrano, & Tella, 2008;
Eizaguirre & Lenz, 2010; Landry & Bernatchez, 2001). We noted that few studies have convincingly
elucidated the ecological drivers of this immunogenetic variation and that all of them focused on
Mhc genes.

4.2 Antagonistic pleiotropy and potential immunogenetic trade-off

Interestingly, we observed that different haplotypes of a same *Tlr* gene (here *Tlr2*) may be associated with the infection status of the two pathogens. Pleiotropic effects in *Tlr2* were expected since this receptor is known to be implicated in the recognition of many bacterial, fungal, protozoa (de Oliviera Nascimento, Massari, & Wetzler, 2012) including *Toxoplasma* (Yarovinsky, 2008) and *Chlamydia* (Beckett et al., 2012; Darville et al., 2003). Here, the most common *Tlr2* haplotype ("TGCCG") (51%) was associated with decreased *Toxoplasma* infection and increased *Chlamydia* infection while the concurrent haplotype ("CATCG") (35%) showed the opposite trend. This result provides an example of antagonistic pleiotropy, which has been proposed as a widespread mechanism favouring the maintenance of genetic variation within populations (Roff & Fairbairn, 2007; Rose, 1982), in particular at immunity-related genes (Aderem & Ulevitch, 2000; Carter & Nguyen, 2011). Antagonistic pleiotropy has rarely been evidenced in natural populations (Turner et al., 2011) and, to our knowledge, this study is the first demonstration with regard to a *Tlr* gene. Two non-exclusive mechanisms may be invoked to explain this antagonistic effect: decrease of intracellular pathogen competition (Loiseau et al., 2008) and host immunogenetic trade-off (Kamath et al., 2014). *Toxoplasma* and *Chlamydia* use similar strategies to invade cells route and compete for the same nutrient pools in co-infected cells (Romano & Coppens, 2013). Therefore, antagonistic effects might arise because *Tlr2* gene alters the competitive interactions between the two pathogens. In other words, because individuals carrying the most frequent *Tlr2* haplotype are less infected by *Toxoplasma*, this may indirectly promote infection by *Chlamydia* without a direct role of the second most frequent haplotype (and *vice versa*). In a study of house sparrows, Loiseau et al. (2008) found antagonistic effects of a *Mhc* class I gene on multiple malarial parasite strains. In this particular case, deleterious 'susceptibility alleles' could be maintained in the population due to within-host competition between malaria parasite strains. Here, because we observed relatively few co-infections, it is unlikely that antagonistic effect may result from the competitive interactions between *Toxoplasma* and *Chlamydia*.

The most likely hypothesis is the presence of a host immunogenetic trade-off (Kamath et al., 2014). In this case, pathogens do not directly compete but *TLr2* genotype has a direct effect on the resistance/susceptibility to both pathogens. The allele inferred to be beneficial for decreasing *Toxoplasma* infection is also associated with increased susceptibility to *Chlamydia* (and *vice versa*). This scenario is reinforced by the fact that the two concurrent *Tlr2* haplotypes exhibit very distinct DNA and amino-acid sequences, what suggests marked functional differences. Similar patterns and processes have previously been described for ticks and nematodes infections (Kamath et al., 2014;

Turner et al., 2011). To our knowledge, our study is the second one showing evidence of antagonistic
selection at non-*Mhc* gene in a wild population (see for the other one Turner et al., 2011).

4.3 Limited role of other genes and selective mechanisms

While many empirical studies in a variety of organisms revealed signatures of pathogen-mediated
directional or balancing in *Mhc* loci (see Bernatchez & Landry, 2003; Sommer, 2005; Spurgin &
Richardson, 2010 for reviews), we did not find any association between *Mhc-Drb* and *Toxoplasma* or
Chlamydia. This is congruent with the findings of Quémeré et *al.* (2015), who showed clear evidence
of historical positive selection on *Mhc-Drb* in roe deer at the sequence level but no signature of
contemporary selection (in contrast to *Tlr* genes). This study suggests that genetic drift and migration
would the predominant contemporary forces shaping *Mhc* variation in roe deer. Moreover, the few
studies showing associations between MHC class II genetic diversity and pathogen infection in other
wild ungulates focused on strongyle parasites (Charbonnel & Pemberton, 2005; Kamath et al., 2014;
Paterson et al., 1998) while we looked at interactions with micro-pathogens. This supports the
general idea that MHC class II molecules principally bind exogenous antigens and are primarily
involved in the immune response to extracellular pathogens (Hughes & Yeager, 1998).

Our data provided only weak support for the "heterozygous advantage hypothesis". *Tlr2* is the only
gene that exhibited a relationship between heterozygosity status and *Toxoplasma* seroprevalence.
However, this pattern most likely resulted from the correlation between *Tlr2* heterozygosity and the
number of copies of *Tlr2-2* haplotype (Pearson's r = 0.3; *P*-value <0.001), this later being strongly
associated with *Toxoplasma* infection status. This illustrates the complexity of disentangling different
mechanisms underlying pathogen-mediated selection (Sin et al., 2014a; Spurgin & Richardson, 2010).
We noted that the "heterozygous advantage hypothesis" only received a limited support when
authors focused on a single pathogen species (but see Oliver et al., 2009), while a growing number of
case studies has revealed *Mhc* heterozygous advantage in animals co-infected with multiple
pathogens (Hughes & Nei, 1992; McClelland et al., 2003; Froeschke & Sommer, 2012). In our case,
the lack of association with roe deer heterozygosity status could be due to the low individual's
probability of being exposed to both pathogens *Toxoplasma* and *Chlamydia* across their life-time,
explaining the low fitness benefit of heterozygous genotypes. A limitation of this work is that we
could not investigate the importance of other mechanisms such as negative frequency-dependent
selection, because of insufficient statistical power to test the role of rare variants and the lack of long
time series (see Charbonnel et al., 2005 for an example).

5. Conclusion

In conclusion, our study illustrates the importance of looking beyond *Mhc* genes (Acevedo-
Whitehouse & Cunningham, 2006) and single snapshot gene/pathogen association (Maizels &
Nussey, 2013) in wildlife immunogenetic studies. In the present case, we did not find any influence of
Mhc diversity on the resistance to two microparasites. Moreover, we highlighted the importance of
antagonistic effects and fluctuating selection on the maintenance of innate immunogenetic diversity,
what could not have been possible using a classical single gene-single pathogen approach.

ACKNOWLEDGMENTS

This work was supported by the French National Institute for Agricultural Research (INRA). We thank
the local hunting associations, the Fédération Départementale des Chasseurs de Haute-Garonne. We
thank numerous co-workers and volunteers for their assistance in field data collection, in particular J-
M. Angibault, B. Cargnelutti, J-L. Rames, J. Joachim, B. Lourtet, D. Picot and N. Cebe. We also thank
Karine Laroucau and Rachid Aaziz from ANSES the laboratory for animal health in Maison-Alfort
(France) for her support and her expertise on *C. abortus* results. We also thank Monica G Candela,
Julie Sevila and co-workers for their support on laboratory work (serology). Genetic data used in this
work were produced through molecular genetic analysis technical facilities of the labex "Centre
Méditerranéen de l'Environnement et de la Biodiversité".

REFERENCES

Acevedo-Whitehouse, K., & Cunningham, A. (2006). Is MHC enough for understanding wildlife
immunogenetics? *Trends Ecol Evol*, *21*, 433–438.
Aderem, A., & Ulevitch, R. J. (2000). Toll-like receptors in the induction of the innate immune
response. *Nature*, *406*(6797), 782.
Akira, S., Takeda, K., & Kaisho, T. (2001). Toll-like receptors: critical proteins linking innate and
acquired immunity. *Nature Immunology*, *2*(8), 675–680.
Alcaide, M., Edwards, S. V., Negro, J. J., Serrano, D., & Tella, J. L. (2008). Extensive polymorphism and
geographical variation at a positively selected MHC class II B gene of the lesser kestrel (Falco
naumanni). *Molecular Ecology*, *17*(11), 2652–2665.
Antonides, J., Mathur, S., Sundaram, M., Ricklefs, R., & DeWoody, J. A. (2019). Immunogenetic
response of the bananaquit in the face of malarial parasites. *BMC Evolutionary Biology*, *19*(1),
107.
Barton, K. (2009). MuMIn: multi-model inference. R package version 1. 0. 0. *Http://R-Forge. r-Project.*
Org/Projects/Mumin/.
Bates, D., Maechler, M., & Bolker, B. (2012). *lme4: Linear mixed-effects models using S4 classes.*

Beckett, E. L., Phipps, S., Starkey, M. R., Horvat, J. C., Beagley, K. W., Foster, P. S., & Hansbro, P. M.
(2012). TLR2, but not TLR4, is required for effective host defence against Chlamydia
respiratory tract infection in early life. *PLoS One*, *7*(6), e39460.

Bernatchez, L., & Landry, C. (2003). MHC studies in nonmodel vertebrates: what have we learned
about natural selection in 15 years? *Journal of Evolutionary Biology*, *16*(3), 363–377.

Brambilla, A., Keller, L., Bassano, B., & Grossen, C. (2018). Heterozygosity–fitness correlation at the
major histocompatibility complex despite low variation in Alpine ibex (Capra ibex).
Evolutionary Applications, *11*(5), 631–644.

Burnham, K. P., Anderson, D. R., & Huyvaert, K. P. (2011). AIC model selection and multimodel
inference in behavioral ecology: some background, observations, and comparisons.
Behavioral Ecology and Sociobiology, *65*(1), 23–35.

Calsbeek, R., Gosden, T. P., Kuchta, S. R., & Svensson, E. I. (2012). Fluctuating selection and dynamic
adaptive landscapes. *The Adaptive Landscape in Evolutionary Biology*, 89–109.

Candela, M. G., Serrano, E., Sevila, J., León, L., Caro, M. R., & Verheyden, H. (2014). Pathogens of
zoonotic and biological importance in roe deer (Capreolus capreolus): Seroprevalence in an
agro-system population in France. *Research in Veterinary Science*, *96*(2), 254–259.

Carter, A. J., & Nguyen, A. Q. (2011). Antagonistic pleiotropy as a widespread mechanism for the
maintenance of polymorphic disease alleles. *BMC Medical Genetics*, *12*(1), 160.

Charbonnel, N., & Pemberton, J. (2005). A long-term genetic survey of an ungulate population
reveals balancing selection acting on MHC through spatial and temporal fluctuations in
selection. *Heredity*, *95*(5), 377.

Cook, L. M. (2003). The rise and fall of the carbonaria form of the peppered moth. *The Quarterly
Review of Biology*, *78*(4), 399–417.

Coulon, A., Guillot, G., Cosson, J. F., Angibault, J. M. A., Aulagnier, S., Cargnelutti, B., … Hewison, A. J.
M. (2006). Genetic structure is influenced by landscape features: empirical evidence from a
roe deer population. *Molecular Ecology*, *15*(6), 1669–1679.

Darville, T., O'Neill, J. M., Andrews, C. W., Nagarajan, U. M., Stahl, L., & Ojcius, D. M. (2003). Toll-like
receptor-2, but not Toll-like receptor-4, is essential for development of oviduct pathology in
chlamydial genital tract infection. *The Journal of Immunology*, *171*(11), 6187–6197.

de Oliviera Nascimento, L., Massari, P., & Wetzler, L. M. (2012). The role of TLR2 in infection and
immunity. *Frontiers in Immunology*, *3*, 79.

Doherty, P. C., & Zinkernagel, R. M. (1975). Enhanced immunological surveillance in mice
heterozygous at the H-2 gene complex. *Nature*, *256*(5512), 50.

Dubey, J. P. (2009). Toxoplasmosis in sheep—the last 20 years. *Veterinary Parasitology*, *163*(1–2), 1–
14.

Eizaguirre, C., & Lenz, T. L. (2010). Major histocompatibility complex polymorphism: dynamics and
consequences of parasite-mediated local adaptation in fishes. *Journal of Fish Biology*, *77*(9),
2023–2047.

Eizaguirre, C., Lenz, T. L., Kalbe, M., & Milinski, M. (2012). Rapid and adaptive evolution of MHC
genes under parasite selection in experimental vertebrate populations. *Nature*
Communications, *3*, 621.

Froeschke, G., & Sommer, S. (2012). Insights into the complex associations between MHC class II DRB
polymorphism and multiple gastrointestinal parasite infestations in the striped mouse. *PLoS*
One, *7*(2), e31820.

Gazzinelli, R. T., & Denkers, E. Y. (2006). Protozoan encounters with Toll-like receptor signalling
pathways: implications for host parasitism. *Nature Reviews Immunology*, *6*(12), 895–906.

Gervais, L., Perrier, C., Bernard, M., Merlet, J., Pemberton, J. M., Pujol, B., & Quéméré, E. (2019).
RAD-sequencing for estimating genomic relatedness matrix-based heritability in the wild: A
case study in roe deer. *Molecular Ecology Resources*. doi: 10.1111/1755-0998.13031

Gotteland, C., Aubert, D., Gibert, P., Moinet, M., Klein, F., Game, Y., … Gilot-Fromont, E. (2014).
Toxoplasmosis in natural populations of ungulates in France: prevalence and spatiotemporal
variations. *Vector-Borne and Zoonotic Diseases*, *14*(6), 403–413.

Grueber, C. E., Wallis, G. P., & Jamieson, I. G. (2013). Genetic drift outweighs natural selection at
toll-like receptor (TLR) immunity loci in a re-introduced population of a threatened species.
Molecular Ecology, *22*(17), 4470–4482.

Hedrick, P. W. (1999). Antagonistic pleiotropy and genetic polymorphism: a perspective. *Heredity*,
82(2), 126–133.

Hedrick, P. W. (2002). Pathogen resistance and genetic variation at MHC loci. *Evolution*, *56*(10),
1902–1908.

Hedrick, P. W. (2006). Genetic polymorphism in heterogeneous environments: the age of genomics.
Annual Review of Ecology, Evolution and Systematics, *37*, 67–93.

Hedrick, P. W. (2012). What is the evidence for heterozygote advantage selection? *Trends in Ecology*
& Evolution, *27*(12), 698–704.

Hill, A. V. (1991). HLA associations with malaria in Africa: some implications for MHC evolution. In
Molecular evolution of the major histocompatibility complex (pp. 403–420). Springer.

Hughes, A. L., & Nei, M. (1992). Models of host-parasite interaction and MHC polymorphism.
Genetics, *132*(3), 863.

Hughes, A. L., & Yeager, M. (1998). Natural selection and the evolutionary history of major
histocompatibility complex loci. *Frontiers in Bioscience: A Journal and Virtual Library*, *3*, d509-
516.

Jepson, A., Banya, W., Sisay-Joof, F., Hassan-King, M., Nunes, C., Bennett, S., & Whittle, H. (1997).
Quantification of the relative contribution of major histocompatibility complex (MHC) and
non-MHC genes to human immune responses to foreign antigens. *Infection and Immunity*,
65(3), 872–876.

Johnston, S. E., Gratten, J., Berenos, C., Pilkington, J. G., Clutton-Brock, T. H., Pemberton, J. M., &
Slate, J. (2013). Life history trade-offs at a single locus maintain sexually selected genetic
variation. *Nature*, *502*(7469), 93.

Kaisho, T., & Akira, S. (2004). Pleiotropic function of Toll-like receptors. *Microbes and Infection*, *6*(15),
1388–1394.

Kamath, P. L., Turner, W. C., Küsters, M., & Getz, W. M. (2014). Parasite-mediated selection drives an
immunogenetic trade-off in plains zebras (Equus quagga). *Proceedings of the Royal Society of*
London B: Biological Sciences, *281*(1783), 20140077.

Klein, J., & Figueroa, F. (1986). Evolution of the major histocompatibility complex. *Critical Reviews in*
Immunology, *6*(4), 295–386.

Klein, S. L. (2000). The effects of hormones on sex differences in infection: from genes to behavior.
Neuroscience & Biobehavioral Reviews, *24*(6), 627–638.

Kloch, A., Wenzel, M. A., Laetsch, D. R., Michalski, O., Welc-Falęciak, R., & Piertney, S. B. (2018).
Signatures of balancing selection in toll-like receptor (TLRs) genes–novel insights from a free-
living rodent. *Scientific Reports*, *8*(1), 8361.

Lam-Yuk-Tseung, S., & Gros, P. (2003). Genetic control of susceptibility to bacterial infections in
mouse models. *Cellular Microbiology*, *5*(5), 299–313.

Landry, C., & Bernatchez, L. (2001). Comparative analysis of population structure across
environments and geographical scales at major histocompatibility complex and microsatellite
loci in Atlantic salmon (Salmo salar). *Molecular Ecology*, *10*(10), 2525–2539.

Levene, H. (1953). Genetic equilibrium when more than one ecological niche is available. *The*
American Naturalist, *87*(836), 331–333.

Lewontin, R. C., & Hubby, J. L. (1966). A molecular approach to the study of genic heterozygosity in
natural populations. II. Amount of variation and degree of heterozygosity in natural
populations of Drosophila pseudoobscura. *Genetics*, *54*(2), 595.

Loiseau, C., Zoorob, R., Garnier, S., Birard, J., Federici, P., Julliard, R., & Sorci, G. (2008). Antagonistic
effects of a Mhc class I allele on malaria-infected house sparrows. *Ecology Letters*, *11*(3),
258–265.

Maizels, R. M., & Nussey, D. H. (2013). Into the wild: digging at immunology's evolutionary roots.
Nature Immunology, *14*(9), 879.

Mazerolle, M. J. (2013). AICcmodavg: Model selection and multimodel inference based on (Q) AIC (c).
R Package Version, *1*, 35.

McClelland, E. E., Penn, D. J., & Potts, W. K. (2003). Major histocompatibility complex heterozygote
superiority during coinfection. *Infection and Immunity*, *71*(4), 2079–2086.

Miller, C. M., Boulter, N. R., Ikin, R. J., & Smith, N. C. (2009). The immunobiology of the innate
response to Toxoplasma gondii. *International Journal for Parasitology*, *39*(1), 23–39.

Møller, A. P., & Saino, N. (2004). Immune response and survival. *Oikos*, *104*(2), 299–304.

Moore, A. J., & Moore, P. J. (1999). Balancing sexual selection through opposing mate choice and
male competition. *Proceedings of the Royal Society of London B: Biological Sciences*,
266(1420), 711–716.

Morellet, N., Bonenfant, C., Börger, L., Ossi, F., Cagnacci, F., Heurich, M., … Sustr, P. (2013).
Seasonality, weather and climate affect home range size in roe deer across a wide latitudinal
gradient within E urope. *Journal of Animal Ecology*, *82*(6), 1326–1339.

Mun, H.-S., Aosai, F., Norose, K., Chen, M., Piao, L.-X., Takeuchi, O., … Yano, A. (2003). TLR2 as an
essential molecule for protective immunity against Toxoplasma gondii infection.
International Immunology, *15*(9), 1081–1087.

Netea, M., Wijmenga, C., & O'Neill, L. (2012). Genetic variation in Toll-like receptors and disease
susceptibility. *Nature Immunology*, *13*, 535–542. (doi:10.1038/ni.2284).

Oliver, M. K., Telfer, S., & Piertney, S. B. (2009). Major histocompatibility complex (MHC)
heterozygote superiority to natural multi-parasite infections in the water vole (Arvicola
terrestris). *Proceedings of the Royal Society of London B: Biological Sciences*, *276*(1659),
1119–1128.

Opsteegh, M., Swart, A., Fonville, M., Dekkers, L., & Van Der Giessen, J. (2011). Age-related
Toxoplasma gondii seroprevalence in Dutch wild boar inconsistent with lifelong persistence
of antibodies. *PLoS One*, *6*(1), e16240.

Parham, P. (2003). Innate immunity: the unsung heroes. *Nature*, *423*(6935), 20–20.

Paterson, S., Wilson, K., & Pemberton, J. M. (1998). Major histocompatibility complex variation
associated with juvenile survival and parasite resistance in a large unmanaged ungulate
population (Ovis aries L.). *Proceedings of the National Academy of Sciences*, *95*(7), 3714–
3719.

Pedersen, A. B., & Greives, T. J. (2008). The interaction of parasites and resources cause crashes in a
wild mouse population. *Journal of Animal Ecology*, *77*(2), 370–377.

Phillips, K. P., Cable, J., Mohammed, R. S., Herdegen-Radwan, M., Raubic, J., Przesmycka, K. J., …
Radwan, J. (2018). Immunogenetic novelty confers a selective advantage in host–pathogen
coevolution. *Proceedings of the National Academy of Sciences*, 201708597.

Pioz, M., Loison, A., Gauthier, D., Gibert, P., Jullien, J.-M., Artois, M., & Gilot-Fromont, E. (2008).
Diseases and reproductive success in a wild mammal: example in the alpine chamois.
Oecologia, *155*(4), 691–704.

Piry, S., Guivier, E., Realini, A., & Martin, J. (2012). SESAME Barcode: NGS-oriented software for
amplicon characterization–application to species and environmental barcoding. *Molecular*
Ecology Resources, *12*(6), 1151–1157.

Quéméré, E., Galan, M., Cosson, J.-F., Klein, F., Aulagnier, S., Gilot-Fromont, E., … Charbonnel, N.
(2015). Immunogenetic heterogeneity in a widespread ungulate: the European roe deer
(Capreolus capreolus). *Molecular Ecology*. 24(15):3873-87.

Raby, A.-C., Holst, B., Davies, J., Colmont, C., Laumonnier, Y., Coles, B., … Labéta, M. O. (2011). TLR
activation enhances C5a-induced pro-inflammatory responses by negatively modulating the
second C5a receptor, C5L2. *European Journal of Immunology*, *41*(9), 2741–2752.

Roff, D. A., & Fairbairn, D. J. (2007). The evolution of trade-offs: where are we? *Journal of*
Evolutionary Biology, *20*(2), 433–447.

Romano, J. D., & Coppens, I. (2013). Host Organelle Hijackers: a similar modus operandi for
Toxoplasma gondii and Chlamydia trachomatis: co-infection model as a tool to investigate
pathogenesis. *Pathogens and Disease*, *69*(2), 72–86.

Rose, M. R. (1982). Antagonistic pleiotropy, dominance, and genetic variation. *Heredity*, *48*(1), 63.

Seabury, C. M., Bhattarai, E. K., Taylor, J. F., Viswanathan, G. G., Cooper, S. M., Davis, D. S., …
Seabury, P. M. (2011). Genome-wide polymorphism and comparative analyses in the white-
tailed deer (Odocoileus virginianus): a model for conservation genomics. *PloS One*, *6*(1),
e15811.

Sevila, J., Richomme, C., Hoste, H., Candela, M. G., Gilot-Fromont, E., Rodolakis, A., … Verheyden, H.
(2014). Does land use within the home range drive the exposure of roe deer (Capreolus
capreolus) to two abortive pathogens in a rural agro-ecosystem? *Acta Theriologica*, *59*(4),
571–581.

Siepielski, A. M., DiBattista, J. D., & Carlson, S. M. (2009). It's about time: the temporal dynamics of
phenotypic selection in the wild. *Ecology Letters*, *12*(11), 1261–1276.

Simpson, G. G. (1944). *Tempo and Mode in Evolution (Columbia Classics in Evolution Series.)*.

Sin, Y. W., Annavi, G., Dugdale, H. L., Newman, C., Burke, T., & MacDonald, D. W. (2014a). Pathogen
burden, co-infection and major histocompatibility complex variability in the E uropean
badger (M eles meles). *Molecular Ecology*, *23*(20), 5072–5088.

Sin, Y. W., Annavi, G., Dugdale, H. L., Newman, C., Burke, T., & MacDonald, D. W. (2014b). Pathogen
burden, co-infection and major histocompatibility complex variability in the European badger
(Meles meles). *Molecular Ecology*, *23*(20), 5072–5088.

Sommer, S. (2005). The importance of immune gene variability (MHC) in evolutionary ecology and
conservation. *Front Zool*, *2*(1), 16.

Spurgin, L., & Richardson, D. (2010). How pathogens drive genetic diversity: MHC, mechanisms and
misunderstandings. *Proceeding of the Royal Society B: Biological Sciences*, *277*, 979–988.

Stoffel, M. A., Nakagawa, S., & Schielzeth, H. (2017). rptR: Repeatability estimation and variance
decomposition by generalized linear mixed-effects models. *Methods in Ecology and*
Evolution, *8*(11), 1639–1644.

Takahata, N., & Nei, M. (1990). Allelic genealogy under overdominant and frequency-dependent
selection and polymorphism of major histocompatibility complex loci. *Genetics*, *124*(4), 967–
978.

Tang, J. W. (2009). The effect of environmental parameters on the survival of airborne infectious
agents. *Journal of The Royal Society Interface*, *6*(Suppl 6), S737–S746. Tenter, A. M.,
Heckeroth, A. R., & Weiss, L. M. (2000). Toxoplasma gondii: from animals to humans.
International Journal for Parasitology, *30*(12–13), 1217–1258.

Tollenaere, C., Bryja, J., Galan, M., Cadet, P., Deter, J., Chaval, Y., … Laakkonen, J. (2008). Multiple
parasites mediate balancing selection at two MHC class II genes in the fossorial water vole:
insights from multivariate analyses and population genetics. *Journal of Evolutionary Biology*,
21(5), 1307–1320.

Trowsdale, J., & Parham, P. (2004). Mini-review: defense strategies and immunity-related genes.
European Journal of Immunology, *34*(1), 7–17.

Tschirren, Barbara, Andersson, M., Scherman, K., Westerdahl, H., Mittl, P., & Raberg, L. (2013).
Polymorphisms at the innate immune receptor TLR2 are associated with Borrelia infection in
a wild rodent population. *Proceeding of the Royal Society B: Biological Sciences*, *280*,
20130364.

Tschirren, Barbara, Råberg, L., & Westerdahl, H. (2011). Signatures of selection acting on the innate
immunity gene Toll-like receptor 2 (TLR2) during the evolutionary history of rodents. *Journal*
of Evolutionary Biology, *24*(6), 1232–1240.

Turner, A. K., Begon, M., Jackson, J. A., Bradley, J. E., & Paterson, S. (2011). Genetic Diversity in
Cytokines Associated with Immune Variation and Resistance to Multiple Pathogens in a
Natural Rodent Population. *PLoS Genet*, *7*(10), e1002343. doi: 10.1371/journal.pgen.1002343

Turner, A. K., Begon, M., Jackson, J. A., & Paterson, S. (2012). Evidence for selection at cytokine loci in
a natural population of field voles (Microtus agrestis). *Molecular Ecology*, *21*(7), 1632–1646.

Wegner, K. M., Kalbe, M., Kurtz, J., Reusch, T. B., & Milinski, M. (2003). Parasite selection for
immunogenetic optimality. *Science*, *301*(5638), 1343–1343.

Wellenreuther, M. (2017). Balancing selection maintains cryptic colour morphs. *Molecular Ecology*,
26(22), 6185–6188.

Wittmann, M. J., Bergland, A. O., Feldman, M. W., Schmidt, P. S., & Petrov, D. A. (2017). Seasonally
fluctuating selection can maintain polymorphism at many loci via segregation lift.
Proceedings of the National Academy of Sciences, *114*(46), E9932–E9941.

Yang, X., & Joyee, A. (2008). Role of toll-like receptors in immune responses to chlamydial infections.
Current Pharmaceutical Design, *14*(6), 593–600.

Yarovinsky, F. (2008). Toll-like receptors and their role in host resistance to Toxoplasma gondii.
Immunology Letters, *119*(1–2), 17–21.

Zuur, A., Ieno, E. N., Walker, N., Saveliev, A. A., & Smith, G. M. (2009). *Mixed effects models and*
extensions in ecology with R. Springer Science & Business Media.

Data Accessibility:

Haplotype DNA sequences: Primers and Genbank accessions are in Table S1. SNP positions and
characteristics are in Table S2. Genotypes of all individuals at each *Tlr* and *Mhc-Drb* gene and
environmental/parasitological data can be found on Dryad: Dryad entry doi:10.5061/dryad.qt0079b.

Authors' contributions
EQ, EGM and NC designed the study and interpreted the data; AJMH, HV and YC led the data
collection process on the field; MG, JM and EQ carried out the *Tlr* and *Mhc-Drb* genotyping; EQ, PH,
NM and MF performed the statistical analyses; EQ led the writing, all authors contributed to the
manuscript and gave final approval for publication.
683

684 **Tables**

685 **Table 1 - Summary of the pathogen** data (number of samples per year in brackets). NA indicates the years for
686 which no data were available.

Pathogen	Number of samples	Number of individuals	Overall seroprevalence	Seroprevalence evaluated per year								
				2008	2009	2010	2011	2012	2013	2014	2015	2016
Toxoplasma	351	277	0.35	0.23 (40)	0.83 (42)	0.56 (43)	0.24 (50)	0.21 (42)	0.24 (38)	0.16 (31)	0.3 (49)	0.43 (16)
Chlamydia	232	196	0.16	0.00 (40)	0.14 (41)	0.37 (43)	0.28 (50)	0.05 (42)	0.00 (16)	NA	NA	NA
coïnfection	232	196	0.08	0 (40)	5 (41)	11 (43)	2 (50)	0 (42)	0 (16)	NA	NA	NA

Supporting information

Table S1. Details of primer sequences, product size and Genbank accessions.

Table S2. Summary of the 13 *Tlr* SNPs genotyped using the KASPar SNP genotyping system.

Table S3. Sample size per year and site for *Toxoplasma* and *Chlamydia*.

Table S4. Full list of models tested for genetic association with *Chlamydia* and *Toxoplasma*.

Table S5. Prevalence of Toxoplasma and Chlamydia in the seven sizes.

Figure S1. Map of the study area.

Figure S2. Model averaged parameter estimates and their 95% confidence intervals for biological and
environmental factors