Patterns of hybrid seed inviability in perennials of the *Mimulus guttatus* sp. complex reveal a potential role of parental conflict in reproductive isolation

Jenn M. Coughlan[1,2], Maya Wilson-Brown[1], John H. Willis[1]

1. *Biological Sciences, Duke University 25 Science Drive, Durham NC*

2. *Biology Department, University of North Carolina, Chapel Hill*

Author Contributions: JMC contributed substantially to the project design, data collection and analysis, and writing. MWB helped with data collection. JHW contributed substantially to the ideas presented here and the project design.

Key Words: Speciation, Endosperm, BDMIs, conflictual selection, cryptic species

Running headline: Parental conflict and hybrid seed inviability in Mimulus

Abstract

Hybrid inviability is a common reproductive barrier in nature, yet the evolutionary drivers

of post-zygotic reproductive isolation are relatively unknown. One hypothesis is that of

parental conflict, wherein differential optima for resource allocation between mothers and

fathers leads to rapid evolution of maternal and paternal resource allocation alleles. Inviable

hybrids are then the manifestation of mismatched maternal and paternal alleles, which

result in inappropriate hybrid growth and ultimately death. Here we describe a group of

relatively cryptic species nested within the *M. guttatus* species complex which show

tremendous reproductive isolation through hybrid seed inviability (HSI). We explore the

role of parental conflict in HSI between *M. guttatus* and a morphological variant, *M. decorus.*

We find that *M. decorus* is genetically distinct from *M. guttatus*, despite exceptional

phenotypic overlap. Reciprocal crosses between *M. guttatus* and *M. decorus* from across their

ranges indicate that HSI is both common and strong, but that populations of *M. decorus* vary

in their ability to cross with *M. guttatus*. Variation in HSI can be attributed in part to

variation in ploidy among populations of *M. decorus*, but we also find two distinct clades of

diploid *M. decorus* which exhibit oppositely asymmetric HSI with *M. guttatus.* Using only

diploids, we show that HSI is associated with asymmetries in reciprocal hybrid seed size,

suggesting a role of parent-of-origin effects on growth in HSI. Despite being each other's

closest relative, crosses between these two clades of *M. decorus* show complete reproductive

isolation via HSI with accompanying parent-of-origin effects on growth. Overall, these

patterns suggest the rapid evolution of parent-of-origin specific resource allocation alleles

coincident with HSI between *M. guttatus* and *M. decorus,* as well as HSI between populations

45 of *M. decorus.* Parental conflict may therefore be an important evolutionary driver of

46 reproductive isolation in this system.

Impact Statement

Despite the prevalence of intrinsic post-zygotic reproductive isolation in nature, we

generally lack an understanding of the evolutionary drivers of these incompatibilities. One

common incompatibility in plants is hybrid seed inviability. One hypothesis for the

evolution of hybrid seed inviability is that of parental conflict, which arises because of

differential optima for resource allocation to offspring between maternal and paternal

interests. In non-monogamous systems, paternal interests should select for the evolution of

greedier alleles, while maternal interests should select for growth-repressing alleles. Here,

we use a combination of common garden experiments, population genetics, and crossing

surveys to assess patterns of hybrid seed inviability in a relatively cryptic group nested

within the *M. guttatus* species complex. We find tremendous cryptic diversity in this system

which is driven, in part, by hybrid seed inviability. We use reciprocal F1 seed sizes to show

that hybrid seed inviability is associated with reciprocal F1 seed size differences, suggesting

a role of parent-of-origin effects on growth. Lastly, we show that populations of *M. decorus*

which have the most extreme values of conflict also exhibit the most reproductive isolation,

which is again paired by reciprocal F1 size differences. Our work showcases substantial

cryptic diversity within the *M. guttatus* species complex, which is driven, in part, by hybrid

seed inviability. Hybrid seed inviability in this system is associated with parent-of-origin

effects on growth, consistent with parental conflict. Lastly, we show that alleles involved in

hybrid seed inviability appear to be rapidly evolving in this group, potentially related to the

arms-race dynamics involved in parental conflict.

Introduction

The origin and maintenance of species ultimately depends on the formation of barriers to reproduction between populations. Research to date has highlighted the role of ecological and sexual selection in the formation of many pre-zygotic and extrinsic post-zygotic barriers (e.g. Bradshaw & Schemske 2003; Ramsey & Schemske 2003; reviewed in Nosil *et al.* 2005; Arngard *et al.* 2014). However, relatively less is known about the evolutionary forces driving the formation of intrinsic post-zygotic reproductive isolation.

The formation of inviable embryos is a common and powerful intrinsic reproductive barrier in nature (Tiffin *et al.* 2001; Lowry *et al.* 2008), especially in placental mammals (e.g. Vranna *et al.* 2000; Zechner *et al.* 2004; Brekke *et al.* 2014, 2016) and seed plants (e.g. Scott *et al.* 1998; Wolff *et al.* 2015; Rebernig *et al.* 2015; Oneal *et al.* 2016; Garner *et al.* 2016; Roth *et al.* 2017). One hypothesis to explain the evolution of hybrid embryo inviability is that of parental conflict (Trivers 1974; Haig and Westoby 1989). Parental conflict arises because maternal and paternal optima for resource allocation to offspring differ (Trivers 1974; Haig and Westoby 1989). Since maternity is guaranteed, selection will favor equivalent resource allocation from mothers to children. However, in non-monogamous systems, selection will favor the evolution of paternally derived alleles which increase resource allocation to offspring- potentially at the expense of the mother or other offspring (Trivers 1974; Haig and Westoby 1989). Therefore, a co-evolutionary arms race for resource-acquiring (i.e. 'greedy') paternal alleles and more resource-repressive maternal alleles can evolve. Reproductive isolation can manifest if populations have evolved at different rates and/or with different genetic responses to conflict, resulting in a mismatch between maternal and paternal alleles in hybrids.

This hypothesis can explain why embryo inviability is common in systems which

partition resources after fertilization (e.g. seed plants and placental mammals; Vranna 2007;

Lafon-Placette & Köhler 2016) relative to systems which do not (e.g. birds and frogs; Wilson

et al. 1974; Prager & Wilson 1975; Fitzpatrick 2004). It is also consistent with the observation

that many examples of hybrid embryo inviability are asymmetric (e.g. Vranna *et al.* 2000;

Rebernig *et al.* 2015). Lastly, because hybrid embryo inviability often arises from issues of

excessive or limited growth, it is consistent with the idea that these incompatibilities involve

resource allocation alleles (Vranna *et al.* 2000; Rebernig *et al.* 2015; Oneal *et al.* 2016). Yet,

hybrid inviability may also evolve as a byproduct of other selective pressures (e.g. ecological

differences), or through neutral processes (e.g. gene duplication; Lynch & Force 2000). In

addition, most studies of hybrid embryo inviability are based on a single population pair,

and as such represent a single evolutionary sample (e.g. Vranna *et al.* 2000; Rebernig *et al.*

2015; Wolff *et al.* 2015; Brekke *et al.* 2016). Thus, we lack a comparative framework for

understanding whether parental conflict can shape hybrid inviability, and what factors

affect the magnitude and direction of inviability.

In seed plants, parental conflict is thought to manifest primarily in the endosperm- a

nutritive tissue which is essential for proper embryo development (Köhler *et al.* 2010). Like

the mammalian placenta, endosperm represents a direct nutritive connection between

developing offspring and mothers. Unlike placenta, endosperm is a triploid tissue which is

comprised of 2:1 maternal:paternal genetic composition. In interploidy crosses- where

hybrid seed inviability (HSI) has been best studied- departures from this 2:1 ratio often

result in irregular endosperm growth, and seed death, even if the embryo itself is viable

('triploid block'- Comai 2005; Köhler *et al.* 2010). In *Arabidopsis* interploidy crosses, improper

dosage of normally imprinted genes in the endosperm is the cause of irregular endosperm

development and ultimately, inviable seeds (Kradolfer *et al.* 2013; Wolff *et al.* 2015). Yet,

these genome-wide differences in dosage cannot explain HSI in diploid-diploid crosses.

Indeed, the genes and evolutionary drivers responsible for HSI remain unknown in diploid

crosses.

One way to test potential drivers of reproductive isolation is to leverage natural

variation in reproductive isolation (i.e. Case *et al.* 2016). The strength of reproductive

isolation often varies between populations (Sweigart *et al.* 2007; Bomblies *et al.* 2007; Case &

Willis, 2008; Martin & Willis, 2010; Sicard *et al.* 2015; Barnard-Kubow & Galloway, 2017;

Bracewell *et al.* 2017). Assessing the covariance between reproductive isolation and potential

drivers can allow us to test what evolutionary forces cause reproductive isolation to evolve.

If parental conflict drives the evolution of HSI, we would predict that (1) reciprocal F1 seeds

should show differences in size, indicating parent-of-origin effects on growth (2) if these

parent-of-origin effects on growth cause reproductive isolation, then the degree of

asymmetry in reciprocal F1 size should correlate with inviability, and (3) *a priori*

designations of the strength of conflict should predict the degree of reproductive isolation

and seed size asymmetries in subsequent crosses.

Here we use perennial variants from the *Mimulus guttatus* species complex as a

system to study the evolution of HSI. We use a common garden experiment and population

genetics to describe perennial variants within the complex. We then complete a series of

crossing surveys to determine the occurrence and strength of HSI and assess the role of

parental conflict in HSI in this group. Overall, we suggest that parental conflict may play an

important role in the formation of reproductive barriers in this group.

Materials and Methods

Phenotypic and genetic survey

The *Mimulus guttatus* species complex is a morphologically, ecologically, and genetically diverse group (Wu *et al.* 2008). Perennial *M. guttatus* comprises several morphological variants, including two inland perennials (*M. guttatus sensu stricto,* and *M. decorus*), and a coastal perennial (*M. grandis*), that are differentiated by subtle differences in corolla length and the types, densities and locations of trichomes (Nesom 2012). We refer to these as inland *M. guttatus*, *M. decorus*, and coastal *M. guttatus*, respectively. *Mimulus tilingii* is the most divergent perennial species, and represents its own species complex (Grant 1924; Nesom 2012; Garner *et al.* 2016). Little is known about the genetic relationships and reproductive barriers between morphologically described perennials of the *M. guttatus* species complex.

To assess the degree of phenotypic and genetic divergence between perennials of the *M. guttatus* complex, we use the common garden experiment described in Coughlan & Willis (2018). Briefly, we grew 1-5 maternal families with 5 replicates per maternal family for each of six populations of annual *M. guttatus*, seven populations of coastal perennial *M. guttatus,* 16 populations of inland perennial *M. guttatus*, 18 populations of *M. decorus* and seven populations of *M. tilingii* in a common garden. Seeds were cold stratified for one week on moist Fafard 4P soil then transferred to the Duke greenhouses (18h days, 21C days/18C nights). Germinants were transplanted on the day of germination.

On the day of first flower, we measured 15 morphological traits (days from germination to first flower, node of first flower, total corolla length, corolla tube length, corolla width, stem thickness, leaf length and width, internode length between the cotyledons and first true leaf, internode length between the first and second pair of true

leaves, the number of stolons, length of the longest stolon, width of the longest stolon,

number of side branches, length of the longest side branch; Coughlan & Willis 2018; Figure

1). To determine how species varied in multivariate trait space, we completed a PCA of all

traits, with traits scaled to have unit variance. PCs 1 and 2 accounted for roughly 47.84% and

19.58% trait variance, respectively, and were the only PCs to explain more trait variance

than expected by chance using a broken stick model.

To determine the genetic relationship among these taxa, we combined a previously

published GBS dataset of *M. guttatus* (Twyford & Friedman 2015; comprising 183

individuals: 70 populations of annual and perennial *M. guttatus*; 1 *M. decorus* population

which was mis-identified as *M. guttatus* in the study, and 1 outgroup, *M. moschatus* var

moniliformis) with whole-genome re-sequenced data from *M. decorus*. We obtained whole

genomic DNA extracted from young bud tissue for 10 individuals of *M. decorus* and 1 *M.*

guttatus (each representing a unique population; outlined in Supplemental Table 1) to the

Duke sequencing facility where 150bp, paired-end read, whole genome Illumina libraries

were created and run in 1 lane on the Illumina 4000 platform (for an average of 28.9 million

reads per sample, and an average coverage of 23x). We included a previously sequenced line

of *M. decorus* from 11[th] population (*IMP*). In total, we present results from 194 individuals; 12

populations of *M. decorus*, 70 populations of annual and perennial *M. guttatus*, and 1

accession of the outgroup *M. moschatus* car *moniliformis*.

For the whole-genome sequences, we trimmed adapter and low-quality sequences

using Trim Galore! (Kreuger 2015), then aligned the trimmed sequences to the *M. guttatus*

Version 2.0 hard masked reference genome using BWA *mem* (Li & Durbin 2009;

https://phytozome.jgi.doe.gov/). We cleaned and sorted reads, and marked duplicates using

Picard tools (Li *et al.* 2009), and used GATK to call SNPs (McKenna *et al.* 2010). We processed

the raw reads from Twyford and Friedman (2015) similarly, except we split individuals by

barcode, discarded low quality reads, and trimmed barcodes using STACKs before

alignment (Catchen *et al.* 2013). We then performed *joingenocaller* in GATK with all samples.

We filtered SNPs as follows: quality score >30, minimum depth of coverage of >/=3x, a minor

allele count of >5%, and <25% missing data. Indels were removed. This resulted in 3840

SNPs across 180 accessions of *M. guttatus* (from 70 populations), 14 accessions of *M. decorus*

(from 12 populations), and one accession of *M. moschatus*. We characterized the genetic

composition of species by performing a PCA using Tassel (Bradbury *et al.* 2007). We used

the R package *ape* to build a neighbor-joining tree for *M. guttatus* and *M. decorus,* using *M.*

moniliformis as an outgroup.

Crossing survey between M. guttatus and M. decorus

To assess HSI between perennials in this group we crossed 19 populations of *M.*

decorus to an average of 4 populations of *M. guttatus* reciprocally, as well as within

population crosses, with an average of 13 replicates per cross type (totaling 1020 fruits;

Supplemental Table 1 & 2). The four populations of *M. guttatus* that were used maximize the

phenotypic and genetic diversity of the species and include one annual, two inland

perennials, and one coastal perennial population. For each fruit, we calculated the

proportion of viable seeds, assessing viability based on visible malformations. We confirmed

inviability by performing germination assays (100 seeds plated across 5 replicate plates of

0.6% agar for each cross. Plates were cold stratified for one week, then transferred to warm,

long days in the Duke growth chambers. Final germination was recorded after three weeks).

Results based on germination assays and morphologically assessed HSI agree qualitatively.

We therefore focus on morphologically determined HSI. We calculated reproductive

isolation (Ramsey & Schemske 2003):

211
$$RI = \frac{\text{Average hybrid seed viability in interspecific crosses}}{\text{Average seed viability in intraspecific crosses}}$$

213   and the symmetry of reproductive isolation, given as:

215
$$\text{symmetry of RI} = \frac{(\text{proportion viable seeds, DxG}) - (\text{proportion viable seeds, GxD})}{(\text{proportion viable seeds, DxG}) + (\text{proportion viable seeds, GxD})}$$

Where 'G' and 'D' refer to *Mimulus guttatus* and *Mimulus decorus,* respectively. These

values are bounded between -1 and 1. Values closer to -1 indicate an asymmetry where

seeds fail when *M. decorus* is the maternal donor and values closer to 1 indicate an

asymmetry where seeds fail when *M. guttatus* is the maternal donor, and values close to 0

indicate little asymmetry (but do not denote the severity of HSI).

Assessing evolutionary mechanisms for variation in hybrid seed inviability

Each population of *M. guttatus* exhibited a consistent crossing phenotype when

crossed to multiple populations of *M. deorus* (i.e. Supplemental Figure 1), but we found

substantial variation between populations of *M. decorus* in the magnitude and the direction

of HSI (Supplemental Figure 2). We sought to assess potential drivers of this variation.

1. Determining the role of genome duplication

To measure ploidy, we performed a combination of flow cytometry and chromosome

squashes. We surveyed 1-2 individuals per population of *M. decorus* to determine total

genomic content using flow cytometry (outlined in Galbraith *et al.* 1997). We chopped

freshly collected, young leaf tissue in Galbraith buffer for each individual, as well as an

internal standard, *Arabidopsis thaliana* (2n=2x, 2C=0.431pg; Schmuths *et al.* 2004). Samples in

buffer were filtered through a 40 μm, then 20 μm pore-sized nylon mesh. Shortly before

analysis, we added 0.1mg/ml of 4',6-diamidino-2-phenylindole (DAPI). Nuclei stained for at

least ten minutes before being run on a Becton Dickinson LSRII flow cytometer. Total DNA

content was calculated by the following equation:

$$\text{2C DNA content (pg DNA)} = \frac{sample\ G1\ peak\ mean}{standard\ G1\ peak\ mean} x\ standard\ 2C\ DNA\ content$$

Total genomic content ranged from 2C= 0.61- 3.06 pg (Supplemental Figure 3). To

confirm that larger genomic content corresponded with changes in ploidy, we performed

meiotic chromosomal squashes on select populations which spanned the range of genome

size (0.67pg, 1.06pg, 1.22 pg, 1.43 pg, 2 pg, and 2.96 pg). Individuals with genome sizes

<1.22pg were found to be diploid (2n=2x=14), while individuals with genomes between 1.43-

2.96pg were found to be tetraploid (2n=4x=28). We inferred ploidy for the remaining

populations based on these ranges. For the remaining analyses we use only diploid

accessions to eliminate parental conflict manifesting from differences in dosage.

2. *Assessing the role of parental conflict in hybrid seed inviability between diploids*

We tested three predictions regarding the growth of hybrid seeds if parental conflict

drives the evolution of HSI. (1) There should be parent-of-origin effects of resource

allocation to offspring, resulting in a size difference between reciprocal F1s. Specifically,

populations that have stronger conflict should produce hybrid seeds which are larger when

they are the paternal donor and smaller when they are the maternal donor when crossed to

a population with weaker conflict. Secondly, the magnitude of growth defects should

correspond to the magnitude of HSI. Lastly, *a priori* assessments of conflict should be

predictive of the outcomes of subsequent crosses.

To test these predictions, we measured seed size by photographing seeds from each

cross and measuring seed width in ImageJ (Scheider *et al.* 2012). We then assessed parent-of-

origin effects on hybrid seed size by correlating the degree of symmetry of HSI and the

degree of symmetry in reciprocal F1 seed size. Lastly, we performed subsequent crosses to

determine if our designations for the strength of conflict were predictive of reproductive

isolation.

Crossing survey within species

Given the patterns of HIS and reciprocal F1 seed sizes between diploid populations

of *M. decorus* and *M. guttatus*, we infer that some populations of *M. decorus* have weaker

conflict than *M. guttatus,* while others have stronger conflict. Parental conflict theory

predicts that populations that show the most extreme levels of conflict should display the

strongest reproductive isolation, and this reproductive isolation should be accompanied by

differences in reciprocal F1 seed size. To test this, we performed a within species crossing

survey in both *M. decorus* and *M. guttatus*. We categorized all diploid populations of *M.*

decorus as low (7 populations) or high (2 populations) conflict, then crossed all diploid

populations of *M. decorus* reciprocally to a representative low conflict and high conflict

population (*IMP* and *Odell Creek*, respectively). Crosses were processed as above to score

average seed viability based on morphology, average germination rate, and average size.

Statistical Analyses

To determine the extent of HSI and reciprocal F1 seed size, we performed two types

of analyses. First, we performed ANOVAs for each population independently with cross

type as the predictor variable (Supplemental Table 5). Based on genetic cluster, we also

performed analyses in which populations were grouped into clades. In these analyses, we

performed a linear mixed model with cross type as a fixed effect and population as a

random effect using the *lme4* and *car* packages in the statistical interface R (Bates *et al.* 2015;

Fox & Weisberg 2011). In both cases, we performed a *post hoc* pairwise T-test with Holm

correction for multiple testing to determine which crosses differed significantly.

Results

Phenotypic and genetic divergence in perennials of the Mimulus guttatus species
complex

Despite minimal phenotypic divergence between perennial variants of the *M.*

guttatus species complex (Figure 1; Coughlan & Willis 2018), *M. decorus* is genetically unique

from other perennial variants in the *M. guttatus* species complex (Figure 1). Samples of *M.*

decorus form a genetic cluster which is distinct from *M. guttatus*, while populations of

annual, inland perennial, and coastal perennial *M. guttatus* do not show any genetic

separation, consistent with previous findings (Figure 1, Supplemental Figure 4; also see

O'Neal *et al.* 2014; Twyford & Friedman 2015). *Mimulus decorus* can be categorized into three

genetic clusters: a northern-diploid group which occur primarily in the central Cascade

Mountains of Oregon, a cluster of tetraploid samples from throughout the range, and a

cluster which includes the two southern-most diploid samples from the Southern Cascade

Mountains, OR.

Figure 1: Phenotypic and genetic description of perennials in the *M. guttatus* species complex. (A) Geographic sampling of genetic samples from *M. guttatus* and *M. decorus*. Inland versus coastal perennials were not distinguished in the previous study and are thus both indicated by darker pink points in this panel. (B) PCA of GBS data for *M. guttatus* and *M. decorus*. (C) PCA of 15 morphological characteristics for perennials within the *M. guttatus* species complex, as well as perennial *M. tilingii* and annual *M. guttatus*. Trait loadings as follows: FT- days to first flower, FN- node of first flower, ST- stem thickness, CW- corolla width, TL- tube length, CL- corolla length, LL- leaf length, LW- leaf width, IL1- first internode length, IL2- second internode length, NS- number of stolons, SL- stolon length, SW- stolon width, SBL- side branch length, NSB- number of side branches. In both PCAs the percent of total variance explained by that PC is indicated in parentheses.

Divergence between northern *M. decorus* and southern *M. decorus*, as well as between

M. guttatus and each clade of *M. decorus* is ~3% (calculated as Pi for all sites using whole

genome sequences, Supplemental Table 6). In comparison, divergence at all sites between *M.*

guttatus and *M. nasutus* is ~3-4% (Brandvain *et al.* 2014) and divergence between *M. guttatus*

and *M. tilingii* is ~7% (Garner *et al.* 2016). Northern and southern clades of *M. decorus* may

thus represent a relatively recent split, and/or substantial introgression post secondary

contact.

Hybrid seed inviability between M. decorus and M. guttatus

There is significant HSI between *M. guttatus* and *M. decorus* (hybrids: mean= 57.29%

viable, se= 3.19%; *M. decorus* within population crosses: 87.47%, se=1.855%; *M. guttatus*

within population: 85.49%, se=6.52), which gives an average reproductive isolation of 0.37

(se=0.051). We find no evidence of a general reduction in seed set in interspecific crosses

relative to intraspecific crosses (i.e. although within-population *M. guttatus* crosses did

produce more seeds than all other cross types; F=4.457, p=0.004, df=1016; Supplemental

Figure 5). Thus, while there is limited reproductive isolation conferred by pollen pistil

interactions, HSI imposes a substantial reproductive barrier. We next sought to determine

whether populations of *M. decorus*, *M. guttatus,* or both varied in the level of reproductive

isolation.

Figure 2: Geographic distribution of hybrid seed inviability between *M. decorus*
and *M. guttatus*. (A) Sampling locales of 19 populations of *M. decorus* which were crossed
to four *M. guttatus* populations reciprocally, points are colored by species and genetic
clade as determined by the PCA in Fig.1 (pink= *M. guttatus*, green= tetraploid *M. decorus*,
yellow = northern *M. decorus*, and blue = southern *M. decorus*. Points which are lighter in
hue are populations that exhibit limited reproductive isolation with *M. guttatus*). (B-E)
Crossing patterns for four populations of *M. decorus*, representing the four main crossing

phenotypes observed: B) *M. guttatus* **x tetraploid** *M. decorus* **exhibits an asymmetric**
barrier, wherein seeds are largely inviable when *M. guttatus* **is the maternal donor, C) no**
hybrid seed inviability, which is exhibited by 3 populations, D) northern *M. decorus*
exhibit strong, asymmetric inviability when *M. decorus* **is the maternal parent, C)**
southern *M. decorus* **exhibit intermediate -strong asymmetric inviability when** *M.*
guttatus **is the maternal donor. F) representative viable hybrid seeds, G) representative**
inviable hybrid seeds.
344

Populations of *M. guttatus* did not drastically differ in their average HSI when

crossed with *M. decorus* (Supplemental Figure 1). However, northern, southern, and

polyploid clades of *M. decorus* differed significantly from each other in the degree and

directionality of hybrid seed lethality with *M. guttatus* (clade effect: X^2=42.1, *df*=3, *p*<0.001; as

well as a cladexcross type interaction: X^2=152.9, *df*=5, *p*<0.0001; Supplemental Figure 2).

Northern *M. decorus* populations tend to produce inviable seeds when they are the

maternal donors in crosses with *M. guttatus* (i.e. Figure 2D; although we note one population

which did not exhibit significant HSI with *M. guttatus*), while southern *M. decorus*

populations produce more inviable seeds when they are the paternal donor in crosses with

M. guttatus (i.e. Figure 2E). Tetraploids tend to produce intermediate and symmetric levels

of inviable seeds or stronger HSI when the tetraploid was the paternal donor (i.e. Figure 2B).

We note two tetraploid populations which exhibit minimal HIS with *M. guttatus* (Figure 2C,

Supplemental Table 2).

Correlates of hybrid seed inviability

To determine if HSI is associated with parent-of-origin effects on growth, we

measured reciprocal F1 seed size. Reciprocal F1 hybrid seeds often varied in size (all

populations of *M. decorus* which exhibit HSI when crossed to *M. guttatus* show significant or

nearly significant size differences in reciprocal F1s; Supplemental Table 5; Figure 3).

Northern *M. decorus* produced F1s with larger seeds when they were the maternal donor,

while southern *M. decorus* produced F1s with larger seeds when they were the paternal

donor in crosses with *M. guttatus* (Figure 3; northern clade crosses: $F=87.4$, $df=3$, $p<0.0001$;

southern clade crosses: $F=95.55$, $df=3$, $p<0.0001$). The level of asymmetry in seed size was

correlated with the level of asymmetry in viability (Figure 3; $r^2=-0.69$, $df=22$, $p<0.0001$), such

that the reciprocal F1 seed which is larger tends to be the seed which is inviable.

Figure 3: A test for conflict using reciprocal F1 seed sizes and symmetries: (A) Seed sizes for reciprocal F1s and parents for northern *M. decorus* (yellow) and southern *M. decorus* (blue) when crossed to *M. guttatus* (pink). Letters above denote significantly different groups, wherein comparable groups are denoted by color. (B) Seed viability symmetry between reciprocal F1s as a function of seed size symmetry between reciprocal F1s. Color of the points denote the line of *M. guttatus* that was used. The black line indicates the linear regression line of best fit, and grey denotes the 95% confidence intervals of the linear regression.

Based on the levels of HSI and differences in reciprocal F1 seed size, we determined

that northern populations of *M. decorus* likely have weaker conflict than *M. guttatus,* as they

are unable to prevent *M. guttatus* paternal alleles from exploiting maternal resources and

inducing growth. In contrast, southern populations of *M. decorus* have stronger conflict than

M. guttatus. We next sought to determine whether these designations of conflict were

predictive of reproductive isolation in subsequent crosses.

Hybrid seed inviability with M. decorus and M. guttatus

To determine if *a priori* designations of conflict were predictive of reproductive

isolation, we crossed all diploid accessions of *M. decorus* to two focal populations; a

northern, weaker conflict *M. decorus* population (IMP) and a southern, stronger conflict *M.*

decorus population (Odell Creek). We also crossed the four populations of *M. guttatus* used

above in a pairwise manner to determine if within-species HSI alleles segregate throughout

the range of *M. guttatus*, which exhibited much less variance in conflict.

We find nearly complete reproductive isolation between northern and southern

clades of *M. decorus,* but no reproductive isolation within groups (Figure 4; IMP crossed to

southern clade: *F*=3599, *df*=3, *p*<0.0001; Odell crossed to northern clade: *F*=864, *df*=3, *p*<0.0001;

Supplemental Figure 6). In addition, we note that geographically and phenotypically

distinct populations of *M. guttatus* do not exhibit any signs of HSI (*F*=0.772, *df*=1, *p*=0.39;

Supplemental Figure 7).

Figure 4: Seed viability and size for within species crosses. Proportion viable seed between (A) IMP (low conflict) and high conflict/southern populations, (B) Odell Creek (high conflict) and northern/low conflict populations. Seed sizes for crosses between (C) IMP (low conflict) and high conflict/southern populations, (D) Odell Creek (high conflict) and northern/low conflict populations. Crosses are denoted with the maternal parent first, S= Southern *M. decorus*, N= Northern *M. decorus*. Letters denote significantly different groups.

Reciprocal F1s between northern and southern clade *M. decorus* crosses vary in seed size, in the direction predicted by parental conflict (Figure 4; IMP crossed to southern clade: F=39.45, df=3, p<0.0001; Odell crossed to northern clade: F=129.3, df=3, p<0.0001). That is, Odell Creek produced larger seeds as the paternal donor and smaller seeds when the maternal donor when crossed to populations with less conflict, and IMP produced larger seeds when it was the maternal donor, and smaller seeds when it was the paternal donor when crossed to populations with more conflict. These differences in reciprocal F1 seed size do not occur in crosses within clades (Supplemental Figure 6). We note that, although these seeds exhibited parent of origin associated size differences, all seeds in the crosses between northern and southern clades of *M. decorus* are smaller than either of the parental seeds.

Overall, this suggests that the morphological variant *M. decorus* likely comprises at least three biological species: a northern diploid, a southern diploid, and at least one tetraploid taxon. These species 1) are phenotypically nearly indistinguishable, 2) represent each others' closest extant relatives, 3) inhabit populations which are a mere ~30 miles apart (Euclidean distance), 4) but display more reproductive isolation with one another than either does to *M. guttatus*.

Discussion

Here we explore the evolution of HSI in perennials of the *Mimulus guttatus* species complex. Despite minimal phenotypic divergence, *M. decorus,* is genetically distinct from *M. guttatus*. We crossed 19 populations of *M. decorus* to four populations of *M. guttatus* to survey for the presence of HSI, and find that HSI between these species is often quite strong, but the magnitude and direction of reproductive isolation varies throughout the range of *M. decorus.* While *M. decorus* is composed of both diploid and tetraploid populations, differences in ploidy cannot fully explain the variance in HSI between *M. guttatus* and *M. decorus.* Using only diploid accessions, we find that the strength of asymmetry in HSI is correlated with asymmetric growth defects between reciprocal F1s, suggesting a parent-of-origin effect on growth related to HSI. Populations of *M. decorus* that vary in strength of conflict were also found to be incompatible with one another, and exhibit parent-of-origin effects on growth in the direction predicted by parental conflict. Overall, we conclude that HSI in this group appears to be substantial and rapidly evolving, and the inviability of hybrid seeds is strongly linked with parent-of-origin effects on growth, consistent with parental conflict.

Phenotypic and genetic diversity within perennials of the M. guttatus species complex

Despite morphological similarity*, M. decorus* is genetically unique from and exhibits strong reproductive isolation with *M. guttatus*. Work in the *Mimulus guttatus* species complex has focused on the diversity of annual forms (e.g. Gardner & Macnair 2000; Martin & Willis 2007; Habecker 2012; Peterson *et al.* 2013; Kenney & Sweigart 2016; Ferris & Willis 2018), and differences between annuals and perennials (e.g. Hall & Willis 2006; Lowry *et al.* 2009; Peterson *et al.* 2016). Relatively less has been done to establish the presence of multiple

perennial species within the complex. Here we find substantial cryptic diversity within this

group. Not only do we find that *M. decorus* is a reproductively isolated species from *M.*

guttatus, but it likely represents three reproductively isolated taxa: a diploid northern clade

that produces inviable seeds when it is the maternal donor in crosses with *M. guttatus*, a

diploid southern clade that produces inviable seeds when it is the paternal donor in crossed

with *M. guttatus*, and at least one tetraploid taxon. This highlights a potential major

difference between annuals and perennials in the *M. guttatus* species complex: annual

species generally exhibit both ecological and phenotypic divergence, but very few exhibit

substantial post-zygotic reproductive isolation with *M. guttatus*. In contrast, *M. decorus* is

phenotypically and ecologically very similar to *M. guttatus*, but exhibits relatively strong

post-zygotic reproductive isolation. This system may thus provide an interesting system to

compare the importance of pre- vs post-zygotic reproductive isolation on rates of

introgression in nature.

HSI is a common and substantial reproductive barrier between *M. guttatus* and *M.*

decorus. This contrasts with other, more geographically restricted intrinsic postzygotic

barriers in this complex (e.g. Sweigart *et al.* 2007; Case & Willis, 2008; Case *et al.* 2016; Zuellig

& Sweigart 2018a, Zuellig & Sweigart 2018b). Because of its commonality, strength, and the

fact that it stops gene flow in the first generation of hybridization, HSI may represent an

important postzygotic barrier in nature for this group. *Mimulus decorus* exhibits striking

variation throughout the range in the extent and direction of HSI, which allows us to dissect,

at least in part, the evolutionary drivers of HSI.

Hybrid seed Inviability and parental conflict

We tested three predictions of the role of parental conflict in HSI: (1) reciprocal F1s will show differences in size, (2) size differences between F1s will correlate with the degree of reproductive isolation, and (3) *a priori* designations of parental conflict will be predictive of reproductive isolation. We find support for each of these predictions in patterns of HSI between *M. guttatus* and diploid accessions of the morphological variant *M. decorus,* highlighting the potential role of parental conflict in HSI in this group.

Evidence for parental conflict in natural systems (Willi 2013; Raunsgard *et al.* 2018), as well as its role in HSI between diploid species pairs has been found in some systems (e.g. in *Capsella* Rebernig *et al.* 2015, Lafon-Placette *et al.* 2018; *Arabidopsis* Lafon-Placette *et al.* 2017; wild tomato Roth *et al.* 2017; *peromyscus mice:* Vranna *et al.* 2000; and dwarf hampsters; Brekke *et al.* 2016). In both *Capsella* and *Mimulus* that show asymmetric inviability, a paternal-excess phenotype appears more lethal (Rebernig *et al.* 2015), which is consistent with interploidy crosses (Lafon-Placette & Köhler 2016). In systems with symmetric HSI (e.g. in *Arabidopsis lyrata* and *A. arenosa,* Lafon-Placette *et al.* 2017; wild tomato species, Roth *et al.* 2017; and the northern x southern *M. decorus* crosses here), parent-of-origin effects on growth have been shown, although both directions of the cross are often smaller than either parent, perhaps suggesting that in cases of stronger HSI, seeds are aborted earlier. In many of these systems, inviability is caused by malformation of endosperm, rather than an incompatibility manifesting in the embryo (Rebernig *et al.* 2015; Lafon-Placette *et al.* 2017, 2018), although in both wild tomatoes and *Mimulus,* HSI tends to produce both malformed endosperm and also early aborting embryos (e.g. Oneal *et al.* 2016; Roth *et al.* 2017), which may be related to differentiation in endosperm developmental programs between *Brassica* and *Solanum*/*Mimulus*.

492   The genetic mechanisms of these parent-of-origin effects on growth are hypothesized

493 to be caused by imprinted expression of nuclear genes in the endosperm of angiosperms and

494 placenta in placental mammals (Efstratiadis 1994; Haig 1997; Köhler *et al.* 2010). Two allelic

495 mechanisms have been proposed to explain the action of imprinted genes (Willi 2013;

496 Raunsgard *et al.* 2018). The 'tug of war' hypothesis posits that coevolutionary dynamics

497 caused by parental conflict will result in growth-promoting alleles to be paternally

498 expressed and maternally silenced, and growth-suppressing alleles to be maternally

499 expressed and paternally silenced. In contrast, the 'recognition avoidance' hypothesis

500 suggests that maternal alleles evolve to recognize and repress growth-promoting paternal

501 alleles. While both of these hypothesis predict that hybrid seeds should be larger than

502 parental seeds when strong-conflict fathers are crossed to weak-conflict mothers, these two

503 hypotheses differ in their predictions of how strong maternal alleles should influence size in

504 crosses with weak fathers (Raunsgard *et al.* 2018). Under the 'recognition avoidance'

505 hypothesis, maternal alleles evolve to recognize and repress paternal expression, but do not

506 have intrinsic growth-repressing properties themselves, while under the 'tug of war'

507 hypothesis, exposure of strong maternal alleles against weak paternal alleles should reveal

508 growth deficits in hybrids. In our study, we find support for the 'tug of war' hypothesis, as

509 viable hybrid seeds between *M. guttatus* and both clades of *M. decorus* show seed

510 phenotypes which are smaller than either parent. Importantly, these under-growth

511 phenotypes are not due to early abortion of seeds, as the smaller hybrid seeds are the viable

512 seeds between *M. guttatus* and both *M. decorus* clades.

513   While that patterns of HSI in the *Mimulus guttatus* species complex conform to the

514 predictions of parental conflict, it is possible that evolutionary explanations other than

515 parental conflict could explain these patterns. For example, it has been hypothesized that

mismatches in small RNAs carried by male and female gametes may result in de-repression

of TEs in developing seeds, as imprinted genes are often associated with TEs (Martienssen

2010). However, the TE and parental conflict hypotheses are not mutually exclusive; TEs

may offer a mechanistic explanation, and Parental Conflict an evolutionary explanation for

HSI (Köhler *et al.* 2010). Ultimately, a molecular genetic dissection of this incompatibility

will be needed to fully understand the proximate and ultimate causes of HSI.

Hybrid seed inviability is rapidly evolving

HSI between northern and southern clades of *M. decorus* highlights the rapidity at

which HSI has evolved in this group. These two diploid species are morphologically almost

indistinguishable, inhabit similar habitats, are geographically proximate, and are each

others' closest relative. Despite this, HSI between northern and southern *M. decorus* is

stronger than HSI between either clade of *M. decorus* and *M. guttatus*. While parental conflict

would predict rapid evolution of maternal and paternal resource allocation alleles (Trivers

1974; Haig & Westoby 1989), the variance in rates of conflict-driven evolution between

northern and southern clades of *M. decorus* is curious. Much work on the rate of evolution of

HSI due to parental conflict focuses on the role of mating system or ploidy (i.e. Brandvain

and Haig 2005; Rebernig *et al* 2015; Lafon-Placette & Köhler 2016; Lafon-Placette *et al.* 2018;

and Scott *et al.* 1998; Pennington *et al.* 2008; Lafon-Placette *et al.* 2017, respectively). Both the

southern and northern clades of *M. decorus* are diploid, and exhibit highly outcrossing floral

morphologies (i.e. large corollas and significant stigma/anther separation). One possibility is

that differences in conflict may be due to differences in clonal growth rates between the

northern and southern clades, as the rates of clonality would also affect the of variance in

paternity each clade experiences. However, this hypothesis remains uninvestigated.

Conclusions

We investigate the evolution of HSI in the *M. guttatus* species complex. We find cryptic diversity within perennials of the complex, namely, *M. decorus* which is genetically unique from and exhibits significant reproductive isolation with *M. guttatus* despite minimal phenotypic divergence. Populations of *M. decorus* vary in the magnitude and direction of HSI with *M. guttatus*. While some of the variation in HSI can be explained by variation in ploidy in *M. decorus,* we find two genetically distinct, diploid clades that exhibit oppositely asymnmetric HSI with *M. guttatus.* HSI is associated with parent-of-origin effects on size. Populations of *M. decorus* which exhibit the most extreme values of conflict exhibit the strongest HSI, and HSI is again paired with differences in reciprocal F1s size. Together, this is consistent with the idea that HSI is rapidly evolving within this group, and highlights a potential role for parent-of-origin effects on growth as the cause, consistent with parental conflict. While many early models for the evolution of intrinsic, postzygotic reproductive isolation assumed that incompatibilities evolved via the accumulation of neutral differences between species, we suggest that coevolutionary arms race dynamics caused by conflict may be important in the evolution of reproductive barriers.

Acknowledgements

We thank Shivam Dave for help with seed counting and Madison Zamora for assistance with field work. We are deeply grateful to members of the Willis and Matute lab, particularly Daniel Matute, who provided useful comments on this manuscript. This project was funded from NSF grants EF-0328636 and EF-0723814 to JHW, a DDIG (DEB- 1501758), ASN Student research award and SSE student research award to JMC. The Duke Graduate

562 School also provided funding to JMC with the Myra and William Waldo Boone fellowship,

563 and financial support was provided from the Duke Biology Department.

Data Accessibility

565 All data will be made publically available upon the acceptance of this manuscript.

References

Arnegard, M. E., McGee, M. D., Matthews, B., Marchinko, K. B., Conte, G. L., Kabir, S., …
Schluter, D. (2014). Genetics of ecological divergence during speciation. *Nature*, *511*,
307–311.

Barnard-Kubow, K. B., & Galloway, L. F. (2017). Variation in reproductive isolation across a
species range. *Ecology and Evolution*, (August), 1–11.

Bates, D., Maechler M., Bolker B., Walker S. (2015). Fitting Linear Mixed-Effects Models
Using lme4. Journal of Statistical Software, 67(1), 1-48.

Bomblies, K., & Weigel, D. (2007). Hybrid necrosis: autoimmunity as a potential gene-flow
barrier in plant species. *Nature Reviews. Genetics*, *8*(5), 382–393.

Bracewell, R. R., Bentz, B. J., Sullivan, B. T., & Good, J. M. (2017). Rapid neo-sex chromosome
evolution and incipient speciation in a major forest pest. *Nature Communications*, *8*(1),
1593.

Bradbury, P. J., Zhang, Z., Kroon, D. E., Casstevens, T. M., Ramdoss, Y., & Buckler, E. S.
(2007). TASSEL: Software for association mapping of complex traits in diverse samples.
Bioinformatics, *23*(19), 2633–2635.

Bradshaw, H., & Schemske, D. W. (2003). Allele substitution at a flower colour locus
produces a pollinator shift in monkeyflowers. *Nature*, *44*(3), 138–147.

Brandvain, Y., & Haig, D. (2005). Divergent mating systems and parental conflict as a barrier
to hybridization in flowering plants. *The American Naturalist*, *166*(3), 330–338.

Brekke, Th. D., & Good, J. M. (2014). Parent-of-origin growth effects and the evolution of
hybrid inviability in dwarf hamsters. *Evolution*, *68*(11), 3134–3148.

Brekke, T. D., Henry, L. a., & Good, J. M. (2016). Genomic imprinting, disrupted placental
expression, and speciation. *Evolution*, *3*(Haig 1996), 1–14.

Case, A. L., Finseth, F. R., Barr, C. M., & Fishman, L. (2016). Selfish evolution of cytonuclear
hybrid incompatibility in Mimulus. *Proceedings of the Royal Society B, 283*.

Case, A. L., & Willis, J. H. (2008). Hybrid male sterility in Mimulus (Phrymaceae) is
associated with a geographically restricted mitochondrial rearrangement. *Evolution,*
62(5), 1026–1039.

Catchen, J., Hohenlohe, P. a., Bassham, S., Amores, A., & Cresko, W. a. (2013). Stacks: An
analysis tool set for population genomics. *Molecular Ecology, 22*(11), 3124–3140.

Comai, L. (2005). The advantages and disadvantages of being polyploid. *Nature Reviews,*
6(11), 836–846.

Coughlan, J. M., & Willis, J. H. (2018). Dissecting the role of a large chromosomal inversion
in life history divergence throughout the Mimulus guttatus species complex. *Molecular*
Ecology, (April), 1-15.

Efstratiadis, A. (1994). Parental imprinting of autosomal mammalian genes. *Current Opinion*
in Genetics and Development, 4(2), 265–280.

Ferris, K. G., & Willis, J. H. (2018). Differential Adaptation to a Harsh Granite Outcrop
Habitat between Sympatric. *Evolution, 72*(6), 1225–1241.

Fitzpatrick, B. M. (2004). Rates of evolution of hybrid inviability in birds and mammals.
Evolution, 58(8), 1865–1870.

Fox J. & Weisberg S. (2011). An {R} Companion to Applied Regression, Second Edition.
Thousand Oaks CA: Sage.

Galbraith, D. W., Lambert, G. M., Macas, J., & Dolezel, J. (2001). Analysis of nuclear DNA
content and ploidy in higher plants. *Current Protocols in Cytometry, Chapter 7*, Unit 7.6.

Gardner, M., & Macnair, M. (2000). Factors affecting the co-existence of the serpentine
endemic Mimulus nudatus Curran and it presumed progenitor, Mimulus guttatus
Fischer ex DC. *Biological Journal of the Linnean Society, 69*, 443–459.

Garner, A. G., Kenney, A. M., Fishman, L., & Sweigart, A. L. (2016). Genetic loci with parent
of origin effects cause hybrid seed lethality between Mimulus species. *New Phytologist,*
211, 319–331.

Grant, A. L. (1924). A monograph of the genus Mimulus. *Annals of the Missouri Botanical*
Garden, 11(2), 99–388.

Habecker, N. M. (2012). *Ecogeographical and intrinsic postzygotic isolation between Mimulus*
glaucescens and Mimulus guttatus (Masters thesis). Retrieved from California State
University Chico repository.

Haig, D. (1997). Parental antagonism, relatedness asymmetries, and genomic imprinting.
Proceedings of the Royal Society B: Biological Sciences, 264(1388), 1657–1662.

Haig, D., & Mark Westoby. (1989). Parent-specific gene expression and the triploid
endosperm. *American Naturalist, 134*(1), 147–155.

Hall, M. C., & Willis, J. H. (2006). Divergent selection on flowering time contributes to local
adaptation in Mimulus guttatus populations. *Evolution, 60*(12), 2466–2477.

Kenney, A. M., Sweigart, A. L., & Kenney, A. M. (2016). Reproductive isolation and
introgression between sympatric Mimulus species. *Molecular Ecology, 25*, 2499–2517.

Ko, C., Köhler, C., & Weinhofer-Molisch, I. (2010). Mechanisms and evolution of genomic
imprinting in plants. *Heredity, 105*(1), 57–63.

Köhler, C., Mittelsten Scheid, O., & Erilova, A. (2010). The impact of the triploid block on the
origin and evolution of polyploid plants. *Trends in Genetics, 26*(3), 142–148.

Kradolfer, D., Wolff, P., Jiang, H., Siretskiy, A., & Köhler, C. (2013). An imprinted gene
underlies postzygotic reproductive isolation in arabidopsis thaliana. *Developmental Cell,*
26(5), 525–535.

Kreuger, F. Trim Galore! 0.3. 7 In.

Lafon-Placette, C., Hatorangan, M. R., Steige, K. A., Cornille, A., Lascoux, M., Slotte, T., &
Köhler, C. (2018). Paternally expressed imprinted genes associate with hybridization
barriers in Capsella. *Nature Plants*.

Lafon-Placette, C., Johannessen, I. M., Hornslien, K. S., Ali, M. F., Bjerkan, K. N., Bramsiepe,
J., … Köhler, C. (2017). Endosperm-based hybridization barriers explain the pattern of
gene flow between *Arabidopsis lyrata* and *Arabidopsis arenosa* in Central Europe.
Proceedings of the National Academy of Sciences, 201615123.

Lafon-Placette, C., & Köhler, C. (2016). Endosperm-based postzygotic hybridization barriers:
Developmental mechanisms and evolutionary drivers. *Molecular Ecology*, 2620–2629.

Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics, 25*(14), 1754–1760.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,
Durbin, R. (2009). The Sequence Alignment/Map format and SAMtools. *Bioinformatics,*
25(16), 2078–2079.

Lowry, D. B., Hall, M. C., Salt, D. E., & Willis, J. H. (2009). Genetic and physiological basis of
adaptive salt tolerance divergence between coastal and inland Mimulus guttatus. *The*
New Phytologist, 183(3), 776–788.

Lowry, D. B., Modliszewski, J. L., Wright, K. M., Wu, C. A., & Willis, J. H. (2008). The strength and genetic basis of reproductive isolating barriers in flowering plants. *Philosophical Transactions of the Royal Society B*, *363*(1506), 3009–3021.

Lynch, M., & Force, A. G. (2000). The Origin of Interspecific Genomic Incompatibility via Gene Duplication. *The American Naturalist*, *156*(6), 590–605.

Martienssen, R. A. (2010). Heterochromatin, small RNA and post-fertilization dysgenesis in allopolyploid and interploid hybrids of Arabidopsis. *New Phytologist*, *186*(1), 46–53.

Martin, N. H., & Willis, J. H. (2010). Geographical variation in postzygotic isolation and its genetic basis within and between two Mimulus species. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *365*(1555), 3261–3261.

Martin, N. H., & Willis, J. H. (2007). Ecological divergence associated with mating system causes nearly complete reproductive isolation between sympatric Mimulus species. *Evolution; International Journal of Organic Evolution*, *61*(1), 68–82.

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kerntsky, A., … DePristo, M. A. (2010). The Genome Analysis Toolkit: A MapReduce framework for analysing next-generation DNA sequencing data. *Genome Research*, *20*, 1297–1303.

Nesom, G. L. (2012). Taxonomy of erythranthe sect. simiolus (phrymaceae) in the usa and mexico. *Phytoneuron*, (1978).

Nosil, P., Vines, T. H., & Funk, D. J. (2005). Reproductive isolation casued by natural selection against immigrants from divergent habitats. *Evolution*, *59*, 705–719.

Oneal, E., Willis, J. H., & Franks, R. G. (2016). Disruption of endosperm development is a major cause of hybrid seed inviability between Mimulus guttatus and Mimulus nudatus. *New Phytologist*, *210*, 1107–1120.

Pennington, P. D., Costa, L. M., Gutierrez-Marcos, J. F., Greenland, A. J., & Dickinson, H. G. (2008). When genomes collide: Aberrant seed development following maize interploidy crosses. *Annals of Botany*, *101*(6), 833–843.

Peterson, M. L., Kay, K. M., & Angert, A. L. (2016). The scale of local adaptation in Mimulus guttatus : comparing life history races , ecotypes , and populations. *New Phytologist*, *211*, 345–356.

Peterson, M. L., Rice, K. J., & Sexton, J. P. (2013). Niche partitioning between close relatives suggests trade-offs between adaptation to local environments and competition. *Ecology and Evolution*, *3*(3), 512–522.

Prager, E. M., & Wilson, a. C. (1975). Slow evolutionary loss of the potential for interspecific hybridization in birds: a manifestation of slow regulatory evolution. *Proceedings of the National Academy of Sciences*, *72*(1), 200–204.

Ramsey, J., Bradshaw, H. D., & Schemske, D. W. (2003). Components of reproductive
isolation between the monkeyflowers Mimulus lewisii and M. cardinalis (Phrymaceae).
Evolution, *57*(7), 1520–1534.

Raunsgard, A., Opedal, Ø. H., Ekrem, R. K., Wright, J., Bolstad, G. H., Armbruster, W. S., &
Pélabon, C. (2018). Intersexual conflict over seed size is stronger in more outcrossed
populations of a mixed-mating plant. *Proceedings of the National Academy of Sciences*,
201810979.

Rebernig, C. a., Lafon-Placette, C., Hatorangan, M. R., Slotte, T., & Köhler, C. (2015). Non-
reciprocal Interspecies Hybridization Barriers in the Capsella Genus Are Established in
the Endosperm. *PLOS Genetics*, *11*(6), e1005295.

Roth, M., Florez-Rueda, A. M., Griesser, S., Paris, M., & Städler, T. (2017). Incidence and
developmental timing of endosperm failure in postzygotic isolation between wild
tomato lineages. *Annals of Botany*, (January), 1–12.

Schneider, C. a., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of
image analysis. *Nature Methods*, *9*(7), 671–675.

Scott, R. J., Spielman, M., Bailey, J., & Dickinson, H. G. (1998). Parent-of-origin effects on
seed development in Arabidopsis thaliana. *Development (Cambridge, England)*,
125(October 2014), 3329–3341.

Sicard, A., Kappel, C., Josephs, E. B., Lee, Y. W., Marona, C., Stinchcombe, J. R., … Lenhard,
M. (2015). Divergent sorting of a balanced ancestral polymorphism underlies the
establishment of gene-flow barriers in Capsella. *Nature Communications*, *6*, 7960.

Sweigart, A. L., Mason, A. R., & Willis, J. H. (2007). Natural variation for a hybrid
incompatibility between two species of mimulus. *Evolution*, *61*(1), 141–151.

Tiffin, P., Olson, M. S., & Moyle, L. C. (2001). Asymmetrical crossing barriers in
angiosperms. *Proceedings. Biological Sciences / The Royal Society*, *268*(1469), 861–867.

Trivers, R. L. (1974). Parent-Offspring Conflict. *American Zoology*, *14*(1), 249–264.

Twyford, A. D., & Friedman, J. (2015). Adaptive divergence in the monkey flower Mimulus
guttatus is maintained by a chromosomal inversion. *Evolution*, *69*(9), 1476–1486.

Vrana, P. B. (2007). Genomic Imprinting As a Mechanism of Reproductive Isolation in
Mammals, *88*(1), 5–23.

Vrana, P. B., Fossella, J. a, Matteson, P., del Rio, T., O'Neill, M. J., & Tilghman, S. M. (2000).
Genetic and epigenetic incompatibilities underlie hybrid dysgenesis in Peromyscus.
Nature Genetics, *25*(1), 120–124.

725 Willi, Y. (2013). The Battle of the Sexes over Seed Size: Support for Both Kinship Genomic
726   Imprinting and Interlocus Contest Evolution. *The American Naturalist*, *181*(6), 787–798.

727 Wilson, a. C., Maxson, L. R., & Sarich, V. M. (1974). Two types of molecular evolution:
728   evidence from studies of interspecific hybridization. *Proceedings of the National Academy*
729   *of Sciences of the United States of America*, *71*(7), 2843–2847.

730 Wolff, P., Jiang, H., Wang, G., Santos-González, J., & Köhler, C. (2015). Paternally expressed
731   imprinted genes establish postzygotic hybridization barriers in *Arabidopsis thaliana*.
732   *ELife*, *4*, 1–14.

733 Wu, C. A., Lowry, D. B., Cooley, a M., Wright, K. M., Lee, Y. W., & Willis, J. H. (2008).
734   Mimulus is an emerging model system for the integration of ecological and genomic
735   studies. *Heredity*, *100*(2), 220–230.

736 Zechner, U., Shi, W., Hemberger, M., Himmelbauer, H., Otto, S., Orth, a., … Fundele, R.
737   (2004). Divergent genetic and epigenetic post-zygotic isolation mechanisms in Mus and
738   Peromyscus. *Journal of Evolutionary Biology*, *17*(2), 453–460.

739 Zuellig, M. P., & Sweigart, A. L. (2018)a. A two-locus hybrid incompatibility is widespread,
740   polymorphic, and active in natural populations of Mimulus. *Evolution*, 1–12.

741 Zuellig, M. P., & Sweigart, A. L. (2018)b. Gene duplicates cause hybrid lethality between
742   sympatric species of Mimulus. *PLoS*, *14*(4), e1007130.