Revealing metabolic flexibility of *Candidatus* Accumulibacter phosphatis through redox cofactor analysis and metabolic network modeling

Leonor Guedes da Silva#, Karel Olavarria Gamez, Joana Castro Gomes, Kasper Akkermans, Laurens Welles, Ben Abbas, Mark C.M. van Loosdrecht, Sebastian Aljoscha Wahl

Department of Biotechnology, Delft University of Technology, The Netherlands

#Address correspondence to Leonor Guedes da Silva, LeonorGuedesdaSilva@gmail.com

These investigations were supported by the SIAM Gravitation Grant 024.002.002, the Netherlands Organization for Scientific Research (NWO). The authors declare no conflict of interest.

KEYWORDS

Candidatus Accumulibacter phosphatis, Polyphosphate Accumulating Organisms (PAO), Enhanced Biological Phosphate Removal (EBPR), Central carbon metabolism, Redox cofactors, Flux Balance Analysis, Enzymatic assays

ABSTRACT

Environmental fluctuations in the availability of nutrients lead to intricate metabolic strategies. *Candidatus* Accumulibacter phosphatis, a polyphosphate accumulating organism (PAO) responsible for enhanced biological phosphorus removal (EBPR) from wastewater treatment systems, is prevalent in aerobic/anaerobic environments. While the overall metabolic traits of these bacteria are well described, the inexistence of isolates has led to controversial conclusions on the metabolic pathways used.

Here, we experimentally determined the redox cofactor preference of different oxidoreductases in the central carbon metabolism of a highly enriched *Ca.* A. phosphatis culture. Remarkably, we observed that the acetoacetyl-CoA reductase engaged in polyhydroxyalkanoates (PHA) synthesis is NADH-preferring instead of the generally assumed NADPH dependency. Based on previously published meta-omics data and the results of enzymatic assays, a reduced central carbon metabolic network was constructed and used for simulating different metabolic operating modes. In particular, scenarios with different acetate-to-glycogen consumption ratios were simulated. For a high ratio (i.e. more acetate), a polyphosphate-based metabolism arises as optimal with a metabolic flux through the glyoxylate shunt. In case of a low acetate-to-glycogen ratio, glycolysis is used in combination with reductive branch of the TCA cycle. Thus, optimal metabolic flux strategies will depend on the environment (acetate uptake) and on intracellular storage compounds availability (polyphosphate/glycogen).

This metabolic flexibility is enabled by the NADH-driven PHA synthesis. It allows for maintaining metabolic activity under varying environmental substrate

conditions, with high carbon conservation and lower energetic costs compared

to NADPH dependent PHA synthesis. Such (flexible) metabolic redox coupling

can explain PAOs' competitiveness under oxygen-fluctuating environments.

IMPORTANCE

Here we demonstrate how microbial metabolism can adjust to a wide range

of environmental conditions. Such flexibility generates a selective advantage

under fluctuating environmental conditions. It can also explain the different

observations reported in PAO literature, including the capacity of *Ca.*

Accumulibacter phosphatis to act like glycogen accumulating organisms

(GAO). These observations stem from slightly different experimental conditions

and controversy only arises when one assumes metabolism can only operate

in one single mode. Furthermore, we also show how the study of metabolic

strategies is possible when combining -omics data with functional assays and

modeling. Genomic information can only provide the potential of a

microorganism. The environmental context and other complementary

approaches are still needed to study and predict the functional application of

such metabolic potential.

INTRODUCTION

Natural habitats of microorganisms are dynamic environments with intermittent supply of nutrients. Under these dynamic conditions, organisms are selected that can accumulate growth substrates when these are abundant to compensate for periods when these are exhausted (1).

Figure 1 – Schematic diagram of an EBPR process with nutrient/polymer profiles (A) and corresponding metabolic strategy (B). Recirculated activated sludge containing PAOs is mixed with influent wastewater in an anaerobic reactor. To compensate for the absence of an external electron acceptor, Accumulibacter use their polyphosphate and glycogen reserves to take up organic carbon sources (e.g. acetate) and accumulate them in the form of polyhydroxyalkanoates (PHA). Phosphate is released at this stage. The TCA cycle is represented by a dashed line has it can have different operating modes. When oxygen becomes available, Accumulibacter makes use of their PHA storage to grow, to replenish glycogen and polyphosphate. The re-accumulation of polyphosphate in growing biomass and subsequent purge of this biomass leads to the net removal of phosphate from wastewater. Adapted from (2, 3). PHA, polyhydroxyalkanoates; PolyP, polyphosphate; PO_4, phosphate; TCA, tricarboxylic acid cycle.

Enhanced biological phosphate removal (EBPR) from wastewater is

designed to make use of such physiological feature by circulating activated

sludge through alternating zones with or without an external electron acceptor

here respectively defined as aerobic/anaerobic (see Figure 1A) (4, 5). This

environment selects for polyphosphate accumulating organisms (PAOs) like

Candidatus Accumulibacter phosphatis (hereafter referred to as

Accumulibacter). These bacteria thrive under these dynamic conditions thanks

to a complex metabolic strategy encompassing the cycling of three common

storage polymers: polyphosphate, glycogen and polyhydroxyalkanoates (PHA).

Of these, polyphosphate stands out for allowing for fast and competitive

harvesting of organic matter in the absence of an external electron acceptor

(see Figure 1B).

In the past decades, a number of researchers have derived hypotheses

about Accumulibacter's (anaerobic) physiology (4, 6–8). One of the most

important inconclusive discussions is the source of reducing power for the

anaerobic accumulation of polyhydroxyalkanoates (PHA) from volatile fatty

acids (e.g. acetate, propionate), as reviewed by Zhou *et al* (9). Most

experimental approaches were adequate to study the general physiology of

Accumulibacter species, however the inherent population heterogeneity has

not yet been sufficiently addressed; for example, only recently researchers

have reported different Accumulibacter clades showing different kinetic

characteristics leading to different metabolic operations (10). Furthermore,

novel meta-omics approaches were used, which allowed for Accumulibacter-

targeted (culture-independent) analysis without the interference of other, non-

Accumulibacter, sub-populations. A comprehensive overview of the

physiological studies published on Accumulibacter can be found in the

Supplementary Document S1. In one of the metatranscriptomic studies (11),

Oyserman and colleagues highlighted the need for validating assumptions

often made in metabolic models of Accumulibacter. In particular, they refined

the discussion on reducing power sources and requirements by making a

distinction between the different redox cofactors, NADH and NADPH. This

distinction imposes a constraint between sources and sinks for each redox

cofactor, which can only be alleviated by energy-consuming

transhydrogenase(-like) mechanisms or using electrons for hydrogen

production. However, in their analysis, the specificity of the involved

oxidoreductases was derived from analogy with other organisms.

In Accumulibacter, reducing power is required for the conversion of acetate

to 3-hydroxybutyrate (3HB), the monomer of the reserve polymer poly-3-

hydroxybutyrate (PHB). The intermediate reduction step is catalyzed by the

enzyme acetoacetyl-CoA reductase (AAR). In PHA accumulating bacteria such

as Cupriavidus necator and Zoogloea ramigera, the preferred electron donor

cofactor of the AAR is NADPH (12), which was the cofactor assumed by

analogy for Accumulibacter (11). However, AARs accepting both NADH and

NADPH have been reported for Azotobacter beijerinckii (13, 14) and

Allochromatium vinosum (15). A brief thermodynamic feasibility analysis of the

coupling between glycolysis and either NADPH- or NADH-preferring AAR is

present in the Supplementary Document S2.

So far and to our knowledge, the nature of the cofactor accepted by

Accumulibacter's acetoacetyl-CoA reductase has not yet been experimentally

established. In this study, we addressed this knowledge gap by measuring the

NADH- and NADPH-dependent acetoacetyl-CoA reductase activities in a cell

free extract from a highly enriched culture of Accumulibacter. We further

extended the analysis to other key oxidoreductases involved in the anaerobic

metabolism of Accumulibacter. These results were then used to update

Accumulibacter's biochemical model and as constraints in a flux balance

analysis (FBA) framework. Our simulations show how the use of different

pathways and storage compounds lead to metabolic flexibility. Without this

flexibility, Accumulibacter's metabolism would become very restrictive and,

under unpredictable anaerobic environments, it could lead to situations were

only a limited fraction of the available substrate could be consumed.

MATERIALS AND METHODS

Reactor operation

Two independent Accumulibacter enrichments were obtained in sequencing batch reactors (SBR). The main operational conditions are described in Table 2 and were adapted from the SBR-S as described in Welles *et al* (10).

Cell free extracts preparation

Broth samples (10 mL) collected from the bioreactor during both anaerobic and aerobic phases were centrifuged (2500g, 10 min, 4°C) and the pellet was washed using 10 mL buffer (hereafter named Buffer 1X) containing 50 mM Tris at pH 8, 5 mM $MgCl_2$, 5 mM NaCl and 5% (v/v) glycerol, and the obtained suspension was centrifuged again (2500g, 10 min, 4°C). After centrifugation, pellets were kept at -20°C for no longer than four days until further analysis. For the enzymatic assay, cellular pellets were suspended in 10 mL of Buffer 1X supplemented with 2 mM (L+D) 1,4-Dithiothreitol (DTT) and cOmplete™ mini protease inhibitor cocktail (Roche). To avoid overheating and protein denaturation, cells were kept on ice while being sonicated until the cell suspension was homogenized (i.e. no granules visible). The resulting suspension was centrifuged (15000g for at least 45 min, 4°C). The obtained supernatant, a cell free extract (CFE), was used for the enzymatic assays.

Enzymatic assays

The reaction mixtures used for each enzymatic assay are described in Table 1. Activities were calculated using the initial rate of reduction of $NAD(P)^+$ or oxidation of NAD(P)H, which was obtained by following the changes in the absorbance at 340 nm, at 30°C. To control the contribution of putative background reactions, reaction mixtures were monitored that contained all the

components except one substrate or without addition of CFE (see "Controls" in

Supplementary Document S4). The measured rates were normalized to the

total protein concentration in the respective CFE. The protein concentration was

measured using the Bradford dye-binding protein assay (Bio-Rad), using

known concentrations of bovine serum albumin as external standards.

Table 1 – Origin of the cell free extract (CFE), buffer, substrates, controls and references
(16–24) used for each enzymatic assay. Controls used can be found in Supplementary
Document S4.

CFE	Tested enzyme	Buffer	Substrates	Reference
SBR-1	acetoacetyl-CoA reductase (AAR)	1X, supplemented	200 µM acetoacetyl-CoA, 200 µM NADH or 200 µM NADPH	(16)
SBR-2	acetoacetyl-CoA reductase (AAR)	1X, supplemented	40 µM acetoacetyl-CoA, 100 µM NADH or 200 µM NADPH	(17)
SBR-2	3-hydroxybutyryl-CoA dehydrogenase (3HBDH)	1X, supplemented	100 µM 3-hydroxybutyryl-CoA 1000 µM NAD(P)$^+$	(17)
SBR-1	glucose-6-phosphate dehydrogenase (G6PDH)	1X, supplemented	2 mM glucose-6-phosphate; 2000 µM NAD$^+$ or 200 µM NADP$^+$	(18)
SBR-2	glucose-6-phosphate dehydrogenase (G6PDH)	1X, supplemented	4 mM glucose-6-phosphate; 400 µM NAD(P)$^+$	
SBR-2	isocitrate dehydrogenase (ICDH)	1X, supplemented	400 µM NAD(P)$^+$ 400 µM isocitrate	(19)
	malate dehydrogenase (MDH)	1X, supplemented	200 µM NAD(P)H 200 µM oxaloacetate	(20)
	malic enzyme (ME)	1X, supplemented	200 µM NADP 200 µM L-malate	(21)
	isocitrate lyase (ICL)	40 mM HEPES, pH 7, 6 mM MgCl$_2$	4 mM isocitrate, 280 µM NADH 45 units lactate dehydrogenase from rabbit muscle (Roche)	(22)
	fumarate reductase (FR)	50 mM HEPES, pH 7	15 mM fumarate 250 µM NADH	(23)
	α-ketoglutarate dehydrogenase (AKGDH)	50 mM HEPES, pH 7, 1mM MgCl$_2$, 1mM Thiamine pyrophosphate, 2.5 mM DTT	2 mM α-ketoglutaric acid 100 µM Coenzyme A 2.5 mM NAD(P)$^+$	(24)

Microbial community characterization

The microbial community present in our enrichments has been characterized

by three orthogonal approaches as described below and the detailed results

can be found in Supplementary Document S3.

Fluorescence *in situ* hybridization (FISH) was used to qualitatively assess

the presence of Accumulibacter in the enrichment cultures. All bacteria were

targeted by the EUB338 mix (general bacteria probe) (25–27). Accumulibacter

clade I and II were targeted by the probes Acc-1-444 and Acc-2-444 (28),

respectively. Hybridized samples were examined with a Zeiss Axioplan-2

epifluorescence microscope.

To further confirm the specific Accumulibacter clade, the presence of the

gene encoding for the polyphosphate kinase I (*ppk1*) present in Accumulibacter

was tested as described in (29, 30) using the primers ACCppk1-254F and

ACCppk1-1376R targeting the *ppk1* gene from Accumulibacter-like bacteria

(31). The *ppk1* gene sequences obtained in this study have been deposited in

the GenBank database under accession numbers MH899084-MH899086.

To identify putative side-populations, 16S-rRNA gene amplicon sequencing

was applied. DNA samples from cell pellets were extracted using the DNeasy

UltraClean Microbial Kit (Qiagen, The Netherlands). Approximately 250 mg wet

biomass was treated according to the standard protocol except an alternative

lysis was implemented. This included a combination of 5 min of heat (65^{o}C)

followed by 5 min of bead-beating for cell disruption on a Mini-Beadbeater-24

(Biospec, U.S.A.). After extraction the DNA was checked for quality by gel

electrophoresis and quantified using a Qubit 4 (Thermo Fisher Scientific,

U.S.A.).

202 After quality control, samples were sent to Novogene Ltd. (Hongkong, China)

203 for amplicon sequencing of the V3-4 region of the 16S-rRNA gene (position

204 341-806) on an Illumina paired-end platform. After sequencing, the raw reads

205 were quality filtered, chimeric sequences were removed and OTUs were

206 generated on the base of ≥ 97% identity. Subsequently, microbial community

207 analysis was performed by Novogene using Mothur & Qiime software(V1.7.0)

208 (32, 33). For phylogenetical determination, the most recent SSURef database

209 from SILVA (34) was used. The microbial communities in each enrichment were

210 compared based on the 10 most abundant OTUs with a distinctive genus (i.e.

211 with most reads assigned to it). The 16S-rRNA gene amplicon data have been

212 deposited in GenBank under Bioproject PRJNA490689.

213 **Anaerobic biochemical model of Accumulibacter**

214 The metabolic network shown in Figure 2 is based on the ancestral genome

215 reconstruction of Accumulibacter proposed by (2) as well as experimental

216 observations obtained here. Additionally, an extensive literature overview

217 supporting this biochemical model of Accumulibacter can be found in

218 Supplementary Document S1.

219 **Flux balance analysis (stoichiometric modeling)**

220 To estimate the possible metabolic flux distributions Flux Balance Analysis

221 (FBA) has been applied (35). The flux distribution can be obtained by

222 optimization:

223
$$v_{opt} = \arg \min_{v} v_{CO2_{prod}} \text{ subject to } \begin{cases} Nv = 0 \\ v_{irr} \geq 0 \\ v_{Glyc_{deg}} = 1 - f \\ v_{Ac_{upt}} = f \end{cases}$$

224
$$\text{with } f = 0 \dots 1 \frac{Cmol}{Cmol \; C_{consumed}}$$

where N is the stoichiometry matrix containing the reactions later shown in

Table 3 and v is the vector containing all reaction fluxes (see also

Supplementary Document S6). The simulation assumes that none of the

balanced intermediates is accumulating inside the cell (steady-state

assumption). This is acceptable given the long simulation period compared to

the turnover time of these intermediates. Inequality constraints were introduced

for physiologically irreversible fluxes (i.e. these should always be positive). The

consumption of acetate and glycogen is varied between only acetate ($f = 1$) to

only glycogen ($f = 0$). The respective experimental data is normalized to a

summed consumption of 1 Cmol (see Supplementary Document S5).

Commonly, maximization of biomass synthesis is used in FBA as an

optimization objective. However, Accumulibacter only grows aerobically and on

the intracellular carbon reserves (PHAs) accumulated during substrate uptake

in the anaerobic period. Biomass synthesis was thus assumed proportional to

the carbon stored as PHA. Consequently, maximal carbon conservation in

PHA, resp. minimal CO_2 production, is set as the cellular objective for the

anaerobic phase.

To prevent adding a stoichiometric constraint due to an assumption on the

proportion of the different PHA polymers possible, i.e. poly-3-hydroxybutyrate

(PHB), poly-3-hydroxyvalerate (PHV), poly-3-hydroxy-2-methylbutyrate

(PH2MB), or poly-3-hydroxy-2-methylvalerate (PH2MV), the respective

monomer amount is introduced, i.e. the reduced precursors Ac-CoA* and Pr-

CoA* such that:

$$2 \text{ Ac-CoA}^* \rightarrow 3\text{HB};$$

$$\text{Ac-CoA}^* + \text{Pr-CoA}^* \rightarrow 3\text{HV (or 3H2MB)};$$

$$2 \text{ Pr-CoA}^* \rightarrow 3\text{H2MV};$$

Compilation and normalization of stoichiometric data

The results of the aforementioned simulations were compared to

stoichiometric data reported in literature for several different Accumulibacter

enrichments. Anaerobic-feast yields were used when available, otherwise the

yields of the whole anaerobic phase were used. In case of missing compound

rates (usually CO_2, PH2MV), these were estimated using electron and carbon

balancing (see Supplementary Document S5). For datasets with redundant

measurements, a data reconciliation method was applied (36). For all

calculations, we assumed acetate and glycogen monomers were the only

substrates and CO_2, PHB, PHV and PH2MV were the only products. In case

yields were reported without the respective error, the error was calculated using

error propagation. Here, the relative errors were assumed as follows: 5% for

acetate and PHB measurements, and 10% for PHV, PH2MV and glycogen

measurements. From the (reconciled) rates for PHB, PHV and PH2MV, the

respective Ac-CoA* and Pr-CoA* rates were determined and normalized to the

amount of consumed substrates, in Cmol.

RESULTS

Characterization of Accumulibacter enrichments

RESULTS

Characterization of Accumulibacter enrichments

This study was carried out using two independent Accumulibacter

enrichment cultures. SBR-1 contained the highest enrichment of

Accumulibacter observed in our lab. However, this cultivation was operating

close to a critical dilution rate which, unfortunately, resulted in washout (i.e.

enrichment deterioration) after sampling. In SBR-2, conditions were adjusted to

reduce the risk of washout, i.e. higher COD load and higher SRT were used.

Table 2 – Process parameters and key performance indicators of the two independent
enrichments of Accumulibacter.

	SBR-1	SBR-2	
pH	7.6	7.6	
T	20ºC	20ºC	
Carbon source	210 mg COD/L (63:37 Acetate:Propionate)	400 mg COD/L (75:25 Acetate:Propionate)	
Phosphate load per C	0.1 Pmol/Cmol	0.1 Pmol/Cmol	
Cycle time	6 h Settling – 76 min; Anaerobic – 135 min; Aerobic – 135 min.	6 h Settling – 30 min; Anaerobic – 112 min; Aerobic – 200 min.	
HRT	12 h	12 h	
SRT	4 days	5.6 days	
Aerobic SRT	1.5 days	3.1 days	
Anaerobic P-release per C-fed	0.58 Pmol/Cmol	~0.75 Pmol/Cmol	
Anaerobic-feast length	~41 min	~42 min	
FISH	PAO Acc I	PAO Acc I	
16S rRNA gene amplicon sequencing (dominant OTUs)	Ca_Accumulibacter Rhodopseudomonas Blastochloris	Start: Dechloromonas Ca_Accumulibacter genus from f__Hyphomonadaceae	End: Ca_Accumulibacter Chryseobacterium genus from o__Sphingobacteriales
***ppk1* analysis**	Dominant: clade IC	Dominant: clade IA	

For both cultures, anaerobic P-release per C-fed indicated that the PAO

present in the sludge are likely saturated with polyphosphate and that glycogen

accumulating organisms (GAO) are not present (37). Furthermore, microbial

characterization by FISH showed Accumulibacter from clade I were the majority

of the microorganisms across all biomass samples as it can be seen by the high

overlap between the Accumulibacter-specific probe and the general Bacteria

probe (micrographs available in Supplementary Document S3). The *ppk1* gene

analysis further specified which subclade was dominant and the 16S-rRNA

gene analysis provided information on the genus of the most abundant sub-

populations next to Accumulibacter (see Table 2 and Supplementary Document

S3).

Redox cofactor preferences of key oxidoreductases

For the simulation and interpretation of the metabolic network function it was

essential to identify the cofactor specificity of relevant reactions. Enzymatic

assays were performed using cell free extracts. While these assays cannot

discriminate which organism has the respective activity, the performed assays

still answer: 1) if an activity is not observed for the whole community, it is also

not present in Accumulibacter, 2) if the cell extract shows a clear preference for

NAD(H) or NADP(H) this is most probably also the preferred cofactor of

Accumulibacter which is highly enriched in the culture.

The enzymatic activity assay results display a clear preference which is

sufficient to annotate a specific cofactor. These are shown in blue in the

metabolic network in Figure 2. Further details can be found in Supplementary

Document S4. The different assay results will be presented from the sinks

(PHA) to the putative sources of reducing power:

(1) Redox cofactor preference (NADH or NADPH) of the PHA synthesis. The CFE from SBR-1 had a NADH preferring acetoacetyl-CoA reductase. The activity was 10x higher with NADH compared to NADPH as substrate. To further confirm this cofactor preference, the acetoacetyl-CoA reductase activity was assayed using CFEs from a second enrichment (SBR-2) and both directions of the reaction were monitored (i.e. NAD(P)H + acetoacetyl-CoA $\leftrightarrow$ NAD(P)$^+$ + 3-hydroxybutyryl-CoA).

(2) Stoichiometry of glycolysis – Embden-Meyerhof-Parnas (EMP) or Entner-Doudoroff (ED)? To discriminate between these pathways, the presence of the glucose-6-phosphate dehydrogenase, the enzyme that catalyzes the first step of ED was measured. This reaction is also common to the oxidative branch of the pentose phosphate pathway (oxPP). No activity was found in the CFEs from both SBRs. The biological positive control, a CFE from *P. putida* KT2440, showed activity with both NAD$^+$ and NADP$^+$, as expected (38). Therefore, Accumulibacter nor the community have an active ED or oxPP pathway leaving EMP as glycolytic route.

(3) Alternative NADPH sources. For many organisms oxPP is an important source of NADPH for growth (20). This activity was not found in Accumulibacter, raising the question if NADPH could be provided by other reaction like isocitrate dehydrogenase. The oxidation of isocitrate was tested using either NAD$^+$ or NADP$^+$. The activity with NADP$^+$ was more than 70x higher, indicating that isocitrate dehydrogenase is a relevant source of NADPH in Accumulibacter.

Additionally, the activity of other oxidoreductases and anaplerotic routes was measured: i) the reduction of oxaloacetate to malate (catalyzed by malate dehydrogenase) had, at least, 55x higher activity when using NADH

than NADPH. Because of the small activity observed when using NADPH,

it was not possible to independently study the cofactor preference of the

malic enzyme (oxidation of malate to pyruvate); ii) the glyoxylate shunt was

also found to be active and to have an activity comparable to the positive

biological control *E. coli* grown in acetate (well known to make use of the

glyoxylate shunt (39)); and lastly, iii) fumarate reductase and α-ketoglutarate

dehydrogenase were also tested but their activities were very low compared

to all other enzymes tested in our study.

Figure 2 – Central carbon metabolic network of Accumulibacter including redox cofactor
preference of oxidoreductases assayed in this study. A simplified version of this network (Table
3) was used for the simulations. The values shown corresponds to the factor difference between
the preferred cofactor and the alternative one. This metabolic network is based on the ancestral
genome reconstruction done by (2). The arrows show the expected flux direction under
anaerobic conditions. The untested cofactor preferences (in gray) remain as assumed by
analogy with other microorganisms. Despite not being annotated in the genome, the ED
pathway has been suggested as the route for glycogen degradation (40, 41). A more extensive

Anaerobic stoichiometric model construction

Based on the defined reaction network (Figure 2 and Table 3) the balance of reducing equivalents by Accumulibacter during anaerobic acetate conversion to PHAs was studied quantitatively. Note that the network from Figure 2 was further simplified (Table 3) based on the presented evidence that PHA accumulation in Accumulibacter is NADH-consuming rather than previously assumed NADPH preference based on genome annotations. The cofactor NADH can be regenerated from other electron carriers like NADPH or ferredoxin using transhydrogenases without any energetic cost for the cell (20).

This is not the case for FADH, which would require an input of energy (ATP) to be re-oxidized using NAD^+. Currently, there is still no experimentally validated mechanism on how Accumulibacter could re-oxidize FADH in the absence of an external electron acceptor (e.g. oxygen or nitrate). Therefore, for this simulation, we blocked this FADH producing step in the TCA cycle (i.e. Succ-CoA to Fum) and for the remaining reactions we neglected the different types of redox cofactors by simply balancing "electrons" (note that each redox cofactor carries 2 electrons and, in some publications, these are simply referred to as [H]). Since the FADH producing step in the TCA cycle is blocked, only the oxidative branch (OAA to Succ-CoA via ICDH and AKGDH) and/or the reductive branch (OAA to Succ-CoA via MDH and FR) are possible.

Pathway	Stoichiometry	Enzyme assayed in this study
Glycolysis (EMP)	$(Glycogen)_1 \rightarrow 2\ Pyr + 4\ electrons$	G6PDH
Pyruvate to Acetyl-CoA	$Pyr \rightarrow Ac\text{-}CoA + 2\ electrons + CO_2$	
Pyruvate to Oxaloacetate	$Pyr + CO_2 \leftrightarrow OAA$	
Oxidative TCA branch	$OAA + Ac\text{-}CoA \rightarrow Succ\text{-}CoA + 2\ CO_2 + 4\ electrons$	ICDH AKGDH
Reductive TCA branch (reverse TCA)	$OAA + 4\ electrons \rightarrow Succ\text{-}CoA$	MDH FR
Glyoxylate shunt	$2\ Ac\text{-}CoA \rightarrow Succ\text{-}CoA + 2\ electrons$	ICL
Succinate-propionate shunt	$Succ\text{-}CoA \leftrightarrow Pr\text{-}CoA + CO_2$	
Ac-CoA* production (precursor for 3HB, 3HV and 3HMB)	$Ac\text{-}CoA + electron \rightarrow Ac\text{-}CoA*$ (* means "+1electron")	AcAc-CoA red
Pr-CoA* production (precursor for 3H2MV, 3HV and 3HMB)	$Pr\text{-}CoA + electron \rightarrow Pr\text{-}CoA*$ (* means "+1electron")	AcAc-CoA red

In contrast to electron balances, the ATP balance cannot be used as there are still too many unknowns: 1) how much ATP can be generated from efflux of ions (potassium, magnesium and phosphate) associated with polyphosphate hydrolysis; 2) how much ATP is required for acetate uptake; 3) how much is

needed to upgrade redox cofactors (i.e. transfer electrons from low to high

potential cofactors); 4) and how much ATP is used for cellular maintenance.

Therefore, ATP is not balanced in this simulation. Nevertheless, it is important

to note that the different pathways used for redox balancing will lead to different

levels of ATP generation. Thus, in cases of polyphosphate limitation, the

metabolism could prefer pathways with higher ATP-yield.

Experiments show that the ratio between glycogen and acetate consumption

by PAOs is variable, e.g. between high/low temperature (42) or

polyphosphate/glycogen availability (37, 43, 44). To reflect this variability,

simulations were performed for a range of acetate and glycogen mixtures

ranging from only acetate to only glycogen and results were compared to

experimental data reported in literature.

For the anaerobic sequestration of acetate as PHA, acetate needs to be

reduced (Ac-CoA*). The Ac-CoA* optimum in Figure 3 corresponds to the

situation in which most carbon is kept inside the cell. The required reducing

power can come from either glycolysis (glycogen breakdown) or glyoxylate

shunt (acetate oxidation) operation. The different modes are presented in

Figure 4. These modes respectively form the extreme left and right values of

the optimum line in Figure 3. If glycogen degradation can supply exactly the

amount of electrons needed for the uptake and conversion of acetate, then only

Ac-CoA* (as PHB) is produced and there is no need to use any part of the TCA

cycle (Figure 4B, and highest PHB yield in Figure 3).

Figure 3 – The amount of Ac-CoA* accumulated (PHB and PHV precursor) depends on the different proportion of glycogen to acetate consumed and on the available pathways. The three optimum redox balancing strategies are shown in Figure 4 ABC. From 0 to approx. 2/6 glycogen Cmol, it can be considered that bacteria are performing a polyphosphate-based metabolism (PAM, optimum line in red) and from approx. 3/6 to 1 glycogen Cmol, a glycogen-based metabolism (GAM, optimum line in green) is used. The "optimum" and "minimum" lines were obtained by minimizing and maximizing CO_2 production, respectively. The feasible solution space (in yellow) lies in between these two cellular objectives, in which a mixture of different redox balancing strategies is used. The minimum carbon conserving strategy is the one that only uses the oxidative TCA branch. Experimental datasets were retrieved from (10, 37, 50–59, 41, 42, 44–49) and normalized to respect carbon and energy conservation principles (see Supplementary Document S5). Similar plots for Pr-CoA*, CO_2, alternative simulations and including error bars on the experimental data points can be found in Supplementary Document S6.

The optimum can probably only be achieved with polyphosphate as ATP

source. In a typical GAO-like metabolism, an excess of glycogen is degraded

compared to the acetate consumed to supply all ATP needed. This leads to the

generation of an excess of reducing power, which is used to produce Pr-CoA*

leading to PHAs more reduced than PHB (e.g. PHV or PH2MV), using the

reductive TCA branch (Figure 4C). On the other hand, in a PAO-like metabolism

the amount of reducing equivalents generated by glycolysis is usually lower

than required to reduce all consumed acetate. In this situation, part of the

acetate is oxidized to produce reducing equivalents. Such oxidation is possible

via an active glyoxylate shunt which was also found from the enzymatic assays.

The simulations show this as the optimal pathway to produce PHAs when

glycogen is limiting (Figure 4A), i.e. it allows for higher carbon conservation (i.e.

higher PHA yield) than the "horseshoe" TCA operation.

Figure 4 – Different optimum redox balancing strategies for Accumulibacter under anaerobic conditions. A) When little glycogen is degraded and is not enough

to reduce all acetate to PHB, the glyoxylate shunt as recommended by (55) is the most optimal way to provide these electrons. B) When the stoichiometric amount

of glycogen is degraded for acetate reduction to PHB, there is no real need to operate any part of the TCA cycle to balance redox and only PHB is then produced,

as proposed in (60). C) When more glycogen than needed for acetate reduction is degraded, the reductive TCA branch can be used as proposed by (55) for GAOs

to sink electrons in more reduced PHAs (e.g. PHV and PH2MV).

DISCUSSION

Characterization of Accumulibacter enrichments

The discussion will be based on the observed high PAO enrichment of mainly *Ca.* Accumulibacter phosphatis clade I. Consequently, extrapolations of these results to Accumulibacter clade II or GAOs are speculative until further validation using respective environments for enrichments (10, 47, 61–64).

Updated biochemical (stoichiometric) model

The presented study shows that the metabolic traits of Accumulibacter enable a flexible metabolic operation under anaerobic conditions. This flexibility is made possible by their energy and reducing power storage, enabling the observed phenotypes: fast anaerobic acetate uptake and anaerobic PHB synthesis (decoupled from growth). The flexible reducing power balancing by Accumulibacter depends on the nature of the redox cofactors used – here, enzymatic assays were performed to define the redox cofactor preferences of the main oxidoreductases in the central metabolic pathways of Accumulibacter.

The key finding from these assays is the NADH-preferring PHA accumulation in Accumulibacter. This NADH-preference allows for a direct consumption of the NADH produced in most of Accumulibacter's reducing power sources. This also eliminates the need for NADH into NADPH conversion, which was suggested earlier by (11) using the membrane-bound transhydrogenase (PntAB) driven by proton motive force. Although there are previous reports showing NADH-driven PHB accumulation (13, 65–68), the level of NADH preference of the acetoacetyl-CoA reductase from Accumulibacter is striking. Further characterization of this enzyme was undertaken by Olavarria and

colleagues (in preparation). This observation allows to re-think the role of PHAs:

we hypothesize that depending on the environment where microorganisms

thrive, PHA accumulation will play a role as carbon reservoir during metabolic

over-flow (NADPH-driven accumulation) or as an electron reservoir during

scarcity of external electron acceptors (NADH-driven accumulation). Thus, for

Accumulibacter, PHA is essentially a fermentation product.

A related finding was the absence of activity of the first step of the

sometimes-implicated ED glycolytic pathway. These results match those

observed by (42) and in (69) for GAOs, where no $NADP^+$ dependent glucose-

6-phosphate dehydrogenase activity was found. The experimental findings are

in line with the absence of key ED genes in the genome annotations of

Accumulibacter and closest relatives *Dechloromonas aromatica* and *Azoarcus*

sp. EbN1 (64, 70). Furthermore, no enzymatic activity of the ED pathway with

NAD^+ as electron acceptor (38) was observed.

These findings are in contradiction with earlier [13]C NMR studies (40, 41),

which indicated that the ED pathway was more likely the route for glycolysis

than EMP. Nevertheless, it has to be noted that these early studies comprise

interpretations of [13]C patterns using simplified metabolic models, with limited

information on the reversibility of each reaction and potentially with reactions

missing; these are common pitfalls of the [13]C-labelling method (71). Also note

that these [13]C NMR studies were performed before the first draft genome of

Accumulibacter was available (70), and only the [13]C pattern in the different

PHAs was measured and not in the metabolic intermediates of each pathway

used.

While NADPH is not required anaerobically by Accumulibacter, this reducing

cofactor has to be produced aerobically to drive biomass synthesis. Here,

isocitrate dehydrogenase was found $NADP^+$ dependent. This is consistent with

the observation that most acetate consumers will use this conversion to

produce NADPH for their anabolism (72) and confirms the protein annotation

found in (64). Since PHA accumulation is now known to be NADH-preferring

and without another sink of NADPH, the latter might accumulate anaerobically

and thereby inhibit the isocitrate dehydrogenase reaction. Alternatively, a

soluble transhydrogenase could convert this NADPH into NADH and allow the

oxidative branch of the TCA to be operational under anaerobic conditions.

Regarding the activity of other TCA oxidoreductases and anaplerotic routes:

i) the oxidation of malate using NAD^+ was also found in the studies of (42); ii)

the glyoxylate shunt was also found active as observed in assays done by (41,

42, 73) and metatranscriptomics/proteomics studies by (2, 63, 74–76), which is

expected as this is the anaplerotic route that allows for microorganisms to

convert C2 sources like acetate into C4 building blocks for anabolism (72); and

iii) fumarate reductase and α-ketoglutarate dehydrogenase activities were very

low compared to all other enzymes tested in our study and alike in the studies

of (42). An α-ketoglutarate:ferredoxin oxidoreductase has been identified in

Accumulibacter's genome and proteome (64). This has not been assayed here

as sufficient evidence was collected suggesting an anaerobic operation mode

via the glyoxylate shunt rather than a full or partial oxidative TCA cycle.

Flexible anaerobic metabolism – Adjustments depending on the environment and intracellular storage compounds

As (9) suggested in their review, the flexibility of Accumulibacter's metabolism is likely the major reason why there is controversy in literature regarding how reducing power is balanced under anaerobic conditions. Long-term exposure to set conditions (e.g. pH, temperature, oxidation level of the substrate, nutrient availability, counter-ions, SRT, settling time) will select for the best strategy for those conditions, but short-term perturbations of those set conditions have shown that Accumulibacter seems to still be able to solve the redox balancing problem even if sub-optimally regarding carbon conservation (10, 43, 44, 77). Thus, based on our analysis, we observe that there is not one fixed stoichiometry, but a range of possible stoichiometries that in the end are defined by the relative proportions of each of the intervening substrates, supplied to the system or produced by a side-population (e.g. acetate, glycogen, polyphosphate, other VFAs, oxygen, nitrate, hydrogen).

The simulations showed Accumulibacter can operate in three distinct modes (and combinations in between): A) when reducing equivalents from glycolysis are limiting compared to the acetate imported (Yagci's model (55)); B) when glycolysis supplies exactly enough reducing equivalents to convert all imported acetate into PHB (Mino's model (60)); and lastly C) when glycolysis is producing an excess of reducing equivalents (alike Yagci's model for GAOs (55)).

The anaerobic use of the glyoxylate shunt in a PAO-like metabolism to provide for the reducing equivalents needed for PHA accumulation is supported by the studies of (55, 73); However, as seen in Figure 3, it does not explain the higher levels of Ac-CoA* found (points above the red optimum line) when a shift

to more reduced PHAs (Pr-CoA*) was expected. This indicates that there might

be an alternative process that allows the cell to conserve extra carbon, in other

words, accumulate more Ac-CoA* and less Pr-CoA*. It could be that 1) fully

anaerobic conditions were not attained when performing the experiments, or

that 2) another side population is providing electrons in the form of a more

reduced organic substrate or even hydrogen gas (78) or 3) Accumulibacter has

yet another, alternative way of balancing redox which has yet to be described

and demonstrated.

Also for the scenario where glycolysis produces an excess of reduced

cofactors as in a GAO-like metabolism, a few experimental data points fall

outside the feasible solution space (points above the green optimum line in

Figure 3); these could be explained in the case that Accumulibacter produces

H_2 to solve an excess of NADH as described by (11).

The multilayered complexity of a physiological analysis of

Accumulibacter

The experimental data gathered are likely also influenced by the intrinsic

heterogeneity of Accumulibacter enrichments; clade differences, population

heterogeneity or even different environment (e.g. depending on the position of

the cell in a floc/granule). Therefore, experimental data points under the

optimum line in Figure 3 likely represent mixtures of cells, each with a different

metabolic mode (Figure 4 A, B or C). Furthermore, any kinetic limitation (i.e.

pathway capacity) may also explain a sub-optimal, mixed phenotype.

Additionally, if all enzymatic machinery is available, it seems possible that

the same cell might change modes depending on the environment and its

intracellular storage dynamics. Particularly, availability of acetate (e.g.

transition feast/famine) and/or polyphosphate storage as well as glycogen can

trigger shifts in metabolic mode.

Despite the heterogeneity and noise, the experimental data outside the

solution space suggests this stoichiometric model is still incomplete to

represent all experimental conditions. The solution space expands for example

when (1) allowing for hydrogen production in excess glycogen-to-acetate

conditions or (2) the oxidation of acetate in a fully operational TCA cycle in a

glycolysis-limiting scenario. With these reactions, all experimental data can be

explained (see Supplementary Document S6). The first is supported by the

detection of hydrogen gas produced by an Accumulibacter enrichment (11). For

the latter, a full TCA cycle has been previously suggested based on ^{13}C NMR

observations (58) and stoichiometric analyses by (43). Additionally, a novel

protein has been proposed based on metagenomic analysis that could allow for

full TCA operation under anaerobic conditions (70); however, there is still no

direct biochemical experimental evidence that indeed validates a full anaerobic

TCA operation. Some indirect observation is presented in (79). Here, labelled

propionate was used as substrate and ^{13}C enrichment was found in the PHV

fragments that come from acetyl-CoA. This labeling pattern can be explained

by a full TCA, but this is not the only metabolic route that could explain this

observation.

Alternatively, it should also be considered that, alike many (strict) anaerobes,

Accumulibacter may use electron bifurcation mechanisms (80) via ferredoxin

oxidoreductases to allow for alternative pathways in the central carbon

metabolism.

CONCLUSION

In this study a network-based modeling approach was used to unravel the metabolic flexibility of Accumulibacter under dynamic conditions. The approach integrates findings and hypotheses derived from previous meta-omics studies as well as different physiological datasets and biochemical assays.

The developed metabolic model demonstrates the flexibility in metabolic function and could explain previous controversy in PAO literature. The NADH dependent PHA synthesis is both an efficient link to catabolic pathways as well as key for enabling metabolic flexibility. Depending on the exact history and cultivation conditions, Accumulibacter can exhibit different metabolic phenotypes; the metabolic network can handle different combinations of carbon sources (i.e. acetate and glycogen) adjusting the use of glycolysis (EMP), different branches of the TCA cycle (incl. glyoxylate shunt) and potentially other pathways not yet considered. This poses a challenge to predictive EBPR modeling, as this implies that stoichiometry is not fixed, but variable, spanning continuously from polyphosphate to glycogen-based phenotypes.

This metabolic versatility is likely what allows Accumulibacter to be, in most situations, very close to a carbon conservation optimum, which is key to ensure their competitiveness in the dynamic environments of EBPR systems.

ACKNOWLEDGEMENTS

These investigations were supported by the SIAM Gravitation Grant 024.002.002, the Netherlands Organization for Scientific Research (NWO). The authors would like to thank Roel van de Wijgaart, Alexandre Carnet, Hein van der Wall, Koen Verhagen, David Weissbrodt, Alex Salazar, and Thomas Abeel for their collaboration in this project. The authors would also like to thank to Sergio Tomás Martínez and Eleni Vasilakou for proofreading and Ben Oyserman for his invaluable advice on this manuscript.

COMPETING INTERESTS

The authors declare no conflict of interest.

REFERENCES

1. Van Loosdrecht MCM, Pot M a., Heijnen JJ. 1997. Importance of bacterial storage polymers in bioprocesses. Water Sci Technol 35:41–47.

2. Oyserman BO, Moya F, Lawson CE, Garcia AL, Vogt M, Heffernen M, Noguera DR, McMahon KD. 2016. Ancestral genome reconstruction identifies the evolutionary basis for trait acquisition in polyphosphate accumulating bacteria. ISME J 10:2931–2945.

3. McMahon KD, Read EK. 2013. Microbial contributions to phosphorus cycling in eutrophic lakes and wastewater. Annu Rev Microbiol 67:199–219.

4. Seviour RJ, Mino T, Onuki M. 2003. The microbiology of biological phosphorus removal in activated sludge systems. FEMS Microbiol Rev 27:99–127.

5. Barnard JL. 1976. A review of Biological Phosphorous Removal in the Activated Sludge Process. Water Sa 2:136–144.

6. Seviour RJ, McIlroy S. 2008. The microbiology of phosphorus removal in activated sludge processes-the current state of play. J Microbiol 46:115–124.

7. Mino T, van Loosdrecht MCM, Heijnen JJ. 1998. Microbiology and biochemistry of the enhanced biological phosphate removal process. Water Res 32:3193–3207.

8. Oehmen A, Lemos PC, Carvalho G, Yuan Z, Keller J, Blackall LL, Reis M a M. 2007. Advances in enhanced biological phosphorus removal: From micro to macro scale. Water Res 41:2271–2300.

9. Zhou Y, Pijuan M, Oehmen A, Yuan Z. 2010. The source of reducing power in the anaerobic metabolism of polyphosphate accumulating organisms (PAOs) - A mini-review. Water Sci Technol 61:1653–1662.

10. Welles L, Tian WD, Saad S, Abbas B, Lopez-Vazquez CM, Hooijmans CM, van Loosdrecht MCM, Brdjanovic D. 2015. Accumulibacter clades Type I and II performing kinetically different glycogen-accumulating organisms metabolisms for anaerobic substrate uptake. Water Res 83:354–366.

11. Oyserman BO, Noguera DR, del Rio TG, Tringe SG, McMahon KD. 2016. Metatranscriptomic insights on gene expression and regulatory controls in Candidatus Accumulibacter phosphatis. ISME J 10:810–22.

12. Madison LL, Huisman GW. 1999. Metabolic Engineering of Poly (3-Hydroxyalkanoates): From DNA to Plastic. Microbiol Mol Biol Rev 63:21–53.

13. Ritchie GAF, Senior PJ, Dawes EA. 1971. The Purification and Characterization of Acetoacetyl-Coenzyme A Reductase from Azotobacter beijerinckii. Biochem J 121:309–316.

14. Senior PJ, Beech GA, Ritchie GA, Dawes EA. 1972. The role of oxygen limitation in the formation of poly- -hydroxybutyrate during batch and continuous culture of Azotobacter beijerinckii. Biochem J 128:1193–201.

15. Liebergesell M, Steinbüchel A. 1992. Cloning and nucleotide sequences of genes relevant for biosynthesis of poly(3-hydroxybutyric acid) in Chromatium vinosum strain D. Eur J Biochem 209:135–50.

16. Satoh Y, Tajima K, Tannai H, Munekata M. 2003. Enzyme-catalyzed poly(3-hydroxybutyrate) synthesis from acetate with CoA recycling and NADPH regeneration in Vitro. J Biosci Bioeng 95:335–341.

17. Chohan, Copeland. 1998. Acetoacetyl coenzyme A reductase and polyhydroxybutyrate synthesis in rhizobium (Cicer) sp. Strain CC 1192. Appl Environ Microbiol 64:2859–63.

18. Olavarria K, Marone MP, da Costa Oliveira H, Roncallo JC, da Costa Vasconcelos FN, da Silva LF, Gomez JGC. 2015. Quantifying NAD(P)H production in the upper

Entner-Doudoroff pathway from Pseudomonas putida KT2440. FEBS Open Bio
5:908–15.

19. Dean AM, Golding GB. 1997. Protein engineering reveals ancient adaptive
replacements in isocitrate dehydrogenase. Proc Natl Acad Sci U S A 94:3104–9.

20. Fuhrer T, Sauer U. 2009. Different biochemical mechanisms ensure network-wide
balancing of reducing equivalents in microbial metabolism. J Bacteriol 191:2112–
2121.

21. Bologna FP, Andreo CS, Drincovich MF. 2007. Escherichia coli malic enzymes: two
isoforms with substantial differences in kinetic properties, metabolic regulation, and
structure. J Bacteriol 189:5937–46.

22. Giachetti E, Pinzauti G, Vanni P. 1984. A new continuous optical assay for isocitrate
lyase. Experientia 40:227–228.

23. Hillier AJ, Jericho RE, Green SM, Jago GR. 1979. Properties and function of
fumarate reductase (NADH) in Streptococcus lactis. Aust J Biol Sci 32:625–35.

24. Bunik VI, Denton TT, Xu H, Thompson CM, Cooper AJL, Gibson GE. 2005.
Phosphonate Analogues of α-Ketoglutarate Inhibit the Activity of the α-Ketoglutarate
Dehydrogenase Complex Isolated from Brain and in Cultured Cells †. Biochemistry
44:10552–10561.

25. Daims H, Brühl A, Amann R, Schleifer KH, Wagner M. 1999. The domain-specific
probe EUB338 is insufficient for the detection of all Bacteria: development and
evaluation of a more comprehensive probe set. Syst Appl Microbiol 22:434–44.

26. Amann RI. 1995. In situ identification of micro-organisms by whole cell hybridization
with rRNA-targeted nucleic acid probes, p. 331–345. In Akkermans, ADL, Van
Elsas, JD, De Bruijn, FJ (eds.), Molecular Microbial Ecology Manual. Springer
Netherlands, Dordrecht.

27. Amann RI, Binder BJ, Olson RJ, Chisholm SW, Devereux R, Stahl DA. 1990.
Combination of 16S rRNA-targeted oligonucleotide probes with flow cytometry for
analyzing mixed microbial populations. Appl Environ Microbiol 56:1919–25.

28. Flowers JJ, He S, Yilmaz S, Noguera DR, McMahon KD. 2009. Denitrification
capabilities of two biological phosphorus removal sludges dominated by different
"Candidatus Accumulibacter" clades. Environ Microbiol Rep 1:583–588.

29. Rubio-Rincón FJ, Welles L, Lopez-Vazquez CM, Nierychlo M, Abbas B, Geleijnse
M, Nielsen PH, van Loosdrecht MCM, Brdjanovic D. 2017. Long-term effects of
sulphide on the enhanced biological removal of phosphorus: The symbiotic role of
Thiothrix caldifontis. Water Res 116:53–64.

30. Saad SA, Welles L, Abbas B, Lopez-Vazquez CM, van Loosdrecht MCM,
Brdjanovic D. 2016. Denitrification of nitrate and nitrite by 'Candidatus
Accumulibacter phosphatis' clade IC. Water Res 105:97–109.

31. McMahon KD, Yilmaz S, He S, Gall DL, Jenkins D, Keasling JD. 2007.
Polyphosphate kinase genes from full-scale activated sludge plants. Appl Microbiol
Biotechnol 77:167–173.

32. Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski
RA, Oakley BB, Parks DH, Robinson CJ, Sahl JW, Stres B, Thallinger GG, Horn DJ
Van, Weber CF. 2009. Introducing mothur: Open-Source, Platform-Independent,
Community-Supported Software for Describing and Comparing Microbial
Communities. Appl Environ Microbiol 75:7537–7541.

33. Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK,
Fierer N, Peña AG, Goodrich JK, Gordon JI, Huttley G a, Kelley ST, Knights D,
Koenig JE, Ley RE, Lozupone C a, McDonald D, Muegge BD, Pirrung M, Reeder J,
Sevinsky JR, Turnbaugh PJ, Walters W a, Widmann J, Yatsunenko T, Zaneveld J,
Knight R. 2010. QIIME allows analysis of high-throughput community sequencing

data. Nat Methods 7:335–6.

34.   Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner
711   FO. 2013. The SILVA ribosomal RNA gene database project: improved data
712   processing and web-based tools. Nucleic Acids Res 41:D590-6.

713   35.   Orth JD, Thiele I, Palsson BØ. 2010. What is flux balance analysis? Nat Biotechnol
714   28:245–8.

715   36.   van der Heijden RTJM, Heijnen JJ, Hellinga C, Romein B, Luyben KCAM. 1994.
716   Linear constraint relations in biochemical reaction systems: I. Classification of the
717   calculability and the balanceability of conversion rates. Biotechnol Bioeng 43:3–10.

718   37.   Welles L, Abbas B, Sorokin DY, Lopez-Vazquez CM, Hooijmans CM, van
719   Loosdrecht MCM, Brdjanovic D. 2017. Metabolic response of "candidatus
720   accumulibacter phosphatis" clade II C to changes in influent P/C ratio. Front
721   Microbiol 7.

722   38.   Olavarría K, Valdés D, Cabrera R. 2012. The cofactor preference of glucose-6-
723   phosphate dehydrogenase from Escherichia coli--modeling the physiological
724   production of reduced cofactors. FEBS J 279:2296–309.

725   39.   Gottschalk G. 1986. Bacterial Metabolism. Springer New York, New York, NY.

726   40.   Maurer M, Gujer W, Hany R, Bachmann S. 1997. Intracellular carbon flow in
727   phosphorus accumulating organisms from activated sludge systems. Water Res
728   31:907–917.

729   41.   Hesselmann RPX, Von Rummell R, Resnick SM, Hany R, Zehnder a. JB. 2000.
730   Anaerobic metabolism of bacteria performing enhanced biological phosphate
731   removal. Water Res 34:3487–3494.

732   42.   Erdal ZK, Erdal UG, Randall CW. 2005. Biochemistry of enhanced biological
733   phosphorus removal and anaerobic COD stabilization. Water Sci Technol 52:557–
734   567.

735   43.   Zhou Y, Pijuan M, Zeng RJ, Yuan Z. 2009. Involvement of the TCA cycle in the
736   anaerobic metabolism of polyphosphate accumulating organisms (PAOs). Water
737   Res 43:1330–1340.

738   44.   Zhou Y, Pijuan M, Zeng RJ, Lu H, Yuan Z. 2008. Could polyphosphate-
739   accumulating organisms (PAOs) be glycogen-accumulating organisms (GAOs)?
740   Water Res 42:2361–2368.

741   45.   Brdjanovic D, Loosdrecht MCM van, Hooijmans CM, Alaerts GJ, Heijnen JJ. 1997.
742   Temperature Effects on Physiology of Biological Phosphorus Removal. J Environ
743   Eng 123:144–153.

744   46.   Brdjanovic D, Van Loosdrecht MCM, Hooijmans CM, Mino T, Alaerts GJ, Heijnen
745   JJ. 1998. Effect of polyphospbate limitation on the anaerobic metabolism of
746   phosphorus-accumulating microorganisms. Appl Microbiol Biotechnol 50:273–276.

747   47.   Acevedo B, Oehmen A, Carvalho G, Seco a., Borrás L, Barat R. 2012. Metabolic
748   shift of polyphosphate-accumulating organisms with different levels of
749   polyphosphate storage. Water Res 46:1889–1900.

750   48.   Smolders GJF, van der Meij J, van Loosdrecht MCM, Heijnen JJ. 1994.
751   Stoichiometric model of the aerobic metabolism of the biological phosphorus
752   removal process. Biotechnol Bioeng 44:837–848.

753   49.   Smolders GJF, Klop JM, van Loosdrecht MCM, Heijnen JJ. 1995. A metabolic
754   model of the biological phosphorus removal process: I. Effect of the sludge retention
755   time. Biotechnol Bioeng 48:222–233.

756   50.   Tian W-D, Lopez-Vazquez CM, Li W-G, Brdjanovic D, van Loosdrecht MCM. 2013.
757   Occurrence of PAOI in a low temperature EBPR system. Chemosphere 92:1314–
758   20.

51.   Zeng RJ, Yuan Z, Keller J. 2003. Model-based analysis of anaerobic acetate uptake
by a mixed culture of polyphosphate-accumulating and glycogen-accumulating
organisms. Biotechnol Bioeng 83:293–302.

52.   López-Vázquez CM, Hooijmans CM, Brdjanovic D, Gijzen HJ, van Loosdrecht
MCM. 2007. A practical method for quantification of phosphorus- and glycogen-
accumulating organism populations in activated sludge systems. Water Environ Res
79:2487–2498.

53.   Lanham AB. 2012. Full-scale biological phosphorus removal: quantification of
storage polymers, microbial performance and metabolic modelling. Universidade
Nova de Lisboa.

54.   Pijuan M, Saunders AM, Guisasola A, Baeza JA, Casas C, Blackall LL. 2004.
Enhanced biological phosphorus removal in a sequencing batch reactor using
propionate as the sole carbon source. Biotechnol Bioeng 85:56–67.

55.   Yagci N, Artan N, Çokgör EU, Randall CW, Orhon D. 2003. Metabolic model for
acetate uptake by a mixed culture of phosphate- and glycogen-accumulating
organisms under anaerobic conditions. Biotechnol Bioeng 84:359–373.

56.   Filipe CDM, Daigger GT, Grady CPL. 2001. A metabolic model for acetate uptake
under anaerobic conditions by glycogen accumulating organisms: Stoichiometry,
kinetics, and the effect of pH. Biotechnol Bioeng 76:17–31.

57.   Zeng R, Yuan Z, van Loosdrecht MCM, Keller J. 2002. Proposed modifications to
metabolic model for glycogen-accumulating organisms under anaerobic conditions.
Biotechnol Bioeng 80:277–279.

58.   Pereira H, Lemos PC, Reis MAM, Crespo JPSG, Carrondo MJT, Santos H. 1996.
Model for carbon metabolism in biological phosphorus removal processes based on
in vivo 13C-NMR labelling experiments. Water Res 30:2128–2138.

59.   Carvalheira M. 2014. The effect of key process operational conditions on enhanced
biological phosphorus removal from wastewater. Universidade Nova de Lisboa.

60.   Mino T, Tsuzuki Y, Matsuo T. 1987. Effect of phosphorus accumulation on acetate
metabolism in the biological phosphorus removal process, p. 27–38. *In* Ramadori,
R (ed.), Biological phosphate removal from wastewaters: proceedings from the
International Conference of Advanced Water Pollution Control (IAWPRC).
Pergamon Press, Oxford, England.

61.   Flowers JJ, He S, Malfatti S, del Rio TG, Tringe SG, Hugenholtz P, McMahon KD.
2013. Comparative genomics of two "Candidatus Accumulibacter" clades
performing biological phosphorus removal. ISME J 7:2301–14.

62.   Welles L, Lopez-Vazquez CM, Hooijmans CM, van Loosdrecht MCM, Brdjanovic D.
2016. Prevalence of 'Candidatus Accumulibacter phosphatis' type II under
phosphate limiting conditions. AMB Express 6.

63.   Wexler M, Richardson DJ, Bond PL. 2009. Radiolabelled proteomics to determine
differential functioning of Accumulibacter during the anaerobic and aerobic phases
of a bioreactor operating for enhanced biological phosphorus removal. Environ
Microbiol 11:3029–3044.

64.   Barr JJ, Dutilh BE, Skennerton CT, Fukushima T, Hastie ML, Gorman JJ, Tyson
GW, Bond PL. 2016. Metagenomic and metaproteomic analyses of Accumulibacter
phosphatis-enriched floccular and granular biofilm. Environ Microbiol 18:273–87.

65.   Kim J, Chang JH, Kim K-J. 2014. Crystal structure and biochemical properties of
the (S)-3- hydroxybutyryl-CoA dehydrogenase PaaH1 from Ralstonia eutropha.
Biochem Biophys Res Commun 448:163–168.

66.   Barycki JJ, O LK, Strauss AW, Banaszak LJ, Author C, Banaszak Dietrich Professor
LJ. 2000. Sequestration of the Active Site by Interdomain Shifting: Crystallographic
and Spectroscopic Evidence for Distinct Conformations of L-3-Hydroxyacyl-CoA

Dehydrogenase Running Title: Human L-3-Hydroxyacyl-CoA Dehydrogenase. JBC
Pap Press Publ.

67.   Haywood GW, Anderson AJ, Chu L, Dawes EA. 1988. The role of NADH- and
NADPH-linked acetoacetyl-CoA reductases in the poly-3-hydroxybutyrate
synthesizing organism Alcaligenes eutrophus. FEMS Microbiol Lett 52:264–259.

68.   Ling C, Qiao G-Q, Shuai B-W, Olavarria K, Yin J, Xiang R-J, Song K-N, Shen Y-H,
Guo Y, Chen G-Q. 2018. Engineering NADH/NAD+ ratio in Halomonas
bluephagenesis for enhanced production of polyhydroxyalkanoates (PHA). Metab
Eng 49:275–286.

69.   Filipe CDM, Daigger GT, Grady CPL. 2001. A metabolic model for acetate uptake
under anaerobic conditions by glycogen accumulating organisms: Stoichiometry,
kinetics, and the effect of pH. Biotechnol Bioeng 76:17–31.

70.   García Martín H, Ivanova N, Kunin V, Warnecke F, Barry KW, McHardy AC, Yeates
C, He S, Salamov A a, Szeto E, Dalin E, Putnam NH, Shapiro HJ, Pangilinan JL,
Rigoutsos I, Kyrpides NC, Blackall LL, McMahon KD, Hugenholtz P. 2006.
Metagenomic analysis of two enhanced biological phosphorus removal (EBPR)
sludge communities. Nat Biotechnol 24:1263–1269.

71.   Van Winden W, Verheijen P, Heijnen S. 2001. Possible pitfalls of flux calculations
based on13C-labeling. Metab Eng 3:151–162.

72.   Gottschalk G. 1986. Bacterial Metabolism. Springer New York, New York, NY.

73.   Burow LC, Mabbett AN, Blackall LL. 2008. Anaerobic glyoxylate cycle activity during
simultaneous utilization of glycogen and acetate in uncultured Accumulibacter
enriched in enhanced biological phosphorus removal communities. ISME J 2:1040–
51.

74.   Skennerton CT, Barr JJ, Slater FR, Bond PL, Tyson GW. 2015. Expanding our view
of genomic diversity in Candidatus Accumulibacter clades. Environ Microbiol
17:1574–1585.

75.   Wilmes P, Andersson AF, Lefsrud MG, Wexler M, Shah M, Zhang B, Hettich RL,
Bond PL, VerBerkmoes NC, Banfield JF. 2008. Community proteogenomics
highlights microbial strain-variant protein expression within activated sludge
performing enhanced biological phosphorus removal. ISME J 2:853–864.

76.   He S, McMahon KD. 2011. "Candidatus Accumulibacter" gene expression in
response to dynamic EBPR conditions. ISME J 5:329–340.

77.   Erdal UG, Erdal ZK, Daigger GT, Randall CW. 2008. Is it PAO-GAO competition or
metabolic shift in EBPR system? Evidence from an experimental study. Water Sci
Technol 58:1329–34.

78.   Lawson CE, Strachan BJ, Hanson NW, Hahn AS, Hall ER, Rabinowitz B, Mavinic
DS, Ramey WD, Hallam SJ. 2015. Rare taxa have potential to make metabolic
contributions in enhanced biological phosphorus removal ecosystems. Environ
Microbiol 17:4979–4993.

79.   Lemos PC, Serafim LS, Santos MM, Reis M a M, Santos H. 2003. Metabolic
Pathway for Propionate Utilization by Phosphorus-Accumulating Organisms in
Activated Sludge: 13C Labeling and In Vivo Nuclear Magnetic Resonance. Appl
Environ Microbiol 69:241–251.

80.   Buckel W, Thauer RK. 2013. Energy conservation via electron bifurcating ferredoxin
reduction and proton/Na+ translocating ferredoxin oxidation. Biochim Biophys Acta
- Bioenerg 1827:94–113.