# Revealing metabolic flexibility of *Candidatus* Accumulibacter
# phosphatis through redox cofactor analysis and metabolic
# network modeling

**Running title: Metabolic flexibility of Accumulibacter**

Leonor Guedes da Silva, Karel Olavarria Gamez, Joana Castro Gomes, Kasper Akkermans,
Laurens Welles, Ben Abbas, Mark C.M. van Loosdrecht, Sebastian Aljoscha Wahl

Department of Biotechnology, Delft University of Technology, The Netherlands

Corresponding Author: Leonor Guedes da Silva
Address: Van der Maasweg 9, 2629HZ Delft, The Netherlands
Telephone number: +31 15 278 5307
E-Mail: LeonorGuedesdaSilva@gmail.com

22   These investigations were supported by the SIAM Gravitation Grant 024.002.002, the Netherlands
23   Organization for Scientific Research (NWO).

27   The authors declare no conflict of interest.

## ABSTRACT

*Candidatus* Accumulibacter phosphatis, a polyphosphate accumulating organism (PAO) growing under alternating anaerobic/aerobic conditions in enhanced biological phosphorus removal (EBPR) systems, displays an intricate metabolic strategy. The organism relies on combining features from either anaerobic- or aerobic-only metabolisms and connecting phases using different reserve polymers. Analysis of the metabolic functions is not trivial, specially without isolates of this microorganism available and without employing advanced metabolic modelling techniques. So far, there is no consensus regarding the type of glycolysis nor the operating mode of the tricarboxylic acid cycle anaerobically. Here, we simulated different conditions using flux balance analyses and a reduced central carbon metabolic network based on previously published meta-omics data and validated with enzyme activity assays. Interestingly, we observed that the polyhydroxyalkanoates (PHA) production was NADH preferring instead of the commonly reported NADPH dependency. This feature allows for higher metabolic flexibility while maintaining a near-optimal carbon conservation and without extra energetic costs, which explains PAOs' competitiveness under anaerobic/aerobic shifting environments.

## KEYWORDS

*Candidatus* Accumulibacter phosphatis, Polyphosphate Accumulating Organisms (PAO), Enhanced Biological Phosphate Removal (EBPR), Anaerobic, Central carbon metabolism, Redox cofactors, Flux Balance Analysis, Enzymatic assays

## INTRODUCTION

Enhanced biological phosphate removal (EBPR) from wastewater is enabled by circulating activated sludge through alternating anaerobic/aerobic and substrate feast/famine conditions. Under these dynamic conditions, organisms are selected that can accumulate growth substrates when these are abundant to compensate for moments when these are exhausted (1). *Candidatus* Accumulibacter phosphatis (hereafter referred to as Accumulibacter) makes use of such strategy for multiple types of nutrients; particularly, by aerobic uptake of phosphate (removal from the wastewater) and intracellular accumulation as polyphosphate that can be later used as anaerobic energy source. For this feature Accumulibacter is known as polyphosphate accumulating organism (PAO) as it allows for fast and competitive harvesting of organic matter in the absence of an external electron acceptor (i.e. anaerobic conditions).

In the past decades, a number of researchers have derived hypotheses about Accumulibacter's physiology (2–5). One of the most significant inconclusive discussions is the source of reducing power for the anaerobic accumulation of polyhydroxyalkanoates (PHA) from volatile fatty acids (e.g. acetate, propionate), as reviewed by (6). While most experimental approaches were adequate to study the population's physiology, the inherent heterogeneity of the system has not yet sufficiently been addressed; for example, in the last decade, researchers have reported different Accumulibacter clades can each have different kinetic characteristics (7). Furthermore, novel meta-omics approaches were used, which allowed for Accumulibacter-targeted analysis without the interference of side-populations. One of the metatranscriptomic studies, (8) highlighted the need of validating assumptions often made in metabolic models of Accumulibacter. In particular, they refined the discussion on reducing power sources and requirements by making a distinction between the different redox cofactors, NADH and NADPH. This distinction imposes a constraint between sources and sinks for each redox cofactor, which can only be alleviated by energy-consuming transhydrogenase(-like) mechanisms or using

electrons for hydrogen production. However, in their analysis, the specificity of the

involved oxidoreductases was derived from analogy with other organisms.

In Accumulibacter, the reducing power is required for the conversion of acetate to 3-

hydroxybutyrate (3HB), the monomer of the reserve polymer poly-3-hydroxybutyrate

(PHB). The intermediate reduction step is catalyzed by the enzyme acetoacetyl-CoA

reductase (AAR). In common PHA accumulating bacteria such as *Cupriavidus necator* and

*Zoogloea ramigera*, the preferred electron donor cofactor of the acetoacetyl-CoA

reductase is NADPH (9), which was then assumed by analogy for Accumulibacter.

However, dual preference (accepting both NADH and NADPH) has been reported for

*Azotobacter beijerinckii* (10,11) and *Allochromatium vinosum* (12).

So far and to our knowledge, the nature of the cofactor accepted by Accumulibacter's

acetoacetyl-CoA reductase has not yet been experimentally established. In this study, we

addressed this knowledge gap by measuring the NADH- and NADPH-dependent

acetoacetyl-CoA reductase activities in a cell free extract from a highly enriched culture

of Accumulibacter. We further extended the analysis to the other key oxidoreductases

involved in the anaerobic metabolism of Accumulibacter. These results were then used to

update Accumulibacter's biochemical model and as constraints in a flux balance analysis

(FBA) framework. Our simulations demonstrate how flexible Accumulibacter's

metabolism can be by making use of different pathways as reducing power sources to

balance their redox state under unpredictable anaerobic environments.

## MATERIALS AND METHODS

### Reactor operation

Two independent Accumulibacter enrichments were obtained in sequencing batch reactors (SBR). The main operational conditions are described in Table 1 and were adapted from the SBR-S as described in (7).

### Cell free extracts preparation

Broth samples (10 mL) collected from the bioreactor during both anaerobic and aerobic phases were centrifuged (2500 g, 10 min, 4°C) and the pellet was washed using 10 mL buffer (hereafter named Buffer 1X) containing 50 mM Tris at pH 8, 5 mM $MgCl_2$, 5 mM NaCl and 5% (v/v) glycerol, and the obtained suspension was centrifuged again (2500 g, 10 min, 4°C). After centrifugation, pellets were kept at -20°C for no longer than four days until further analysis. For the enzymatic assay, cellular pellets were suspended in 10 mL of Buffer 1X supplemented with 2 mM (L+D) 1,4-Dithiothreitol (DTT) and protease inhibitor cocktail (Roche). The cells were disrupted using a sonicator (Sonifier 250, Branson) setting the output power to the level 3, 30% duty cycle, during 3 min. The cellular suspension was kept cold during the disruption using ice. The resulting suspension was centrifuged (4°C, 15000 g for at least 45 min). The obtained supernatant, a cell free extract (CFE), was used for the enzymatic assays.

### Enzymatic assays

The reaction mixtures used for each enzymatic assay are described in

Table 1. Activities were calculated using the initial rate of reduction of $NAD(P)^+$ or oxidation of NAD(P)H, which was obtained by following the changes in the absorbance at 340 nm, at 30°C. To control the contribution of putative background reactions, reaction mixtures were monitored that contained all the components except one substrate or without addition of cell free extract (CFE, see "Controls" in Supplementary Document 4).

The measured rates were normalized to the total protein concentration in the

respective CFE. The protein concentration was measured using the Bradford dye-binding

protein assay (Bio-Rad), using known concentrations of bovine serum albumin as

external standards.

Table 1 – Origin of CFE, buffer, substrates, controls and references (26,53–60) used for each
enzymatic assay. Controls used can be found in Supplementary Document 4.

| CFE | Tested enzyme | Buffer | Substrates | Reference |
|---|---|---|---|---|
| SBR-1 | acetoacetyl-CoA reductase | 1X, supplemented | 200 µM acetoacetyl-CoA, 200 µM NADH or 200 µM NADPH | (53) |
| SBR-2 | acetoacetyl-CoA reductase | 1X, supplemented | 40 µM acetoacetyl-CoA, 100 µM NADH or 200 µM NADPH | (54) |
| SBR-2 | 3-hydroxybutyryl-CoA dehydrogenase | 1X, supplemented | 100 µM 3-hydroxybutyryl-CoA 1000 µM NAD(P)$^+$ | (54) |
| SBR-1 | glucose-6-phosphate dehydrogenase | 1X, supplemented | 2 mM glucose-6-phosphate; 2000 µM NAD$^+$ or 200 µM NADP$^+$ | (55) |
| SBR-2 | glucose-6-phosphate dehydrogenase | 1X, supplemented | 4 mM glucose-6-phosphate; 400 µM NAD(P)$^+$ | |
| SBR-2 | isocitrate dehydrogenase | 1X, supplemented | 400 µM NAD(P)$^+$ 400 µM isocitrate | (56) |
| | malate dehydrogenase | 1X, supplemented | 200 µM NAD(P)H 200 µM oxaloacetate | (26) |
| | malic enzyme | 1X, supplemented | 200 µM NADP 200 µM L-malate | (57) |
| | isocitrate lyase | 40 mM HEPES, pH 7, 6 mM MgCl$_2$ | 4 mM isocitrate, 280 µM NADH 45 units lactate dehydrogenase from rabbit muscle (Roche) | (58) |
| | fumarate reductase | 50 mM HEPES, pH 7 | 15 mM fumarate 250 µM NADH | (59) |
| | α-ketoglutarate dehydrogenase | 50 mM HEPES, pH 7, 1mM MgCl$_2$, 1mM Thiamine pyrophosphate, 2.5 mM DTT | 2 mM α-ketoglutaric acid 100 µM Coenzyme A 2.5 mM NAD(P)$^+$ | (60) |

130 **Microbial community characterization**

**FISH**

Fluorescence *in situ* hybridization (FISH) was used to qualitatively assess the presence of Accumulibacter in our enrichments. All bacteria were targeted by the EUB338 mix (general bacteria probe) (13–15). Accumulibacter clade I and II were targeted by the probes Acc-1-444 and Acc-2-444 (16), respectively. Hybridized samples were examined with a Zeiss Axioplan-2 epifluorescence microscope.

**16S-rRNA amplicon sequencing**

To identify typical side-populations present in our enrichments, 16S-rRNA amplicon sequencing was applied. DNA samples from cell pellets of our enrichments were extracted using the DNeasy UltraClean Microbial Kit (Qiagen, The Netherlands). Approximately 250 mg wet biomass was treated according to the standard protocol except an alternative lysis was implemented. This included a combination of 5 min of heat (65°C) followed by 5 min of bead-beating for cell disruption on a Mini-Beadbeater-24 (Biospec, U.S.A.). After extraction the DNA was checked for quality by gel electrophoresis and quantified using a Qubit 4 (Thermo Fisher Scientific, U.S.A.).

After quality control, samples were sent to Novogene Ltd. (Hongkong, China) for amplicon sequencing of the V3-4 region of the 16S-rRNA gene (position 341-806) on a Illumina paired-end platform. After sequencing, the raw reads were quality filtered, chimeric sequences were removed and OTUs were generated on the base of ≥ 97% identity. Subsequently microbial community analysis was performed by Novogene using Mothur & Qiime software(V1.7.0). For phylogenetical determination, the most recent SSURef database from SILVA (http://www.arb-silva.de/) was used. The microbial communities were compared by generating a top 10 genera graph. The 16S-rRNA gene amplicon libraries have been deposited under project PRJNA490689.

**PPK analysis**

To further confirm the specific PAO clade, the presence of the gene encoding for the polyphosphate kinase I (*ppk1*) present in Accumulibacter was tested as described in

(17,18) and using the primers ACCppk1-254F and ACCppk1-1376R targeting the *ppk1*

gene from Accumulibacter sp. like bacteria (19). The *ppk1* gene sequences obtained in this

study have been deposited in the GenBank database under accession numbers

MH899084-MH899086.

**Flux balance analysis (stoichiometric modelling)**

The metabolic network shown in Figure 1 is based on the ancestral genome

reconstruction done by (20). Despite not being annotated in the genome, the ED pathway

has been suggested as the route for glycogen degradation (21,22). A more extensive

overview of evidence for each pathway can be found in Supplementary Document 1. The

stoichiometric model (see Table 2) used in the simulations was built based on the

metabolic network shown in Figure 1, including the experimental observations obtained

in this present study.

To estimate the possible metabolic flux distributions through this underdetermined

system, Flux Balance Analysis (FBA) can be applied (23). Inequality constraints were

introduced for physiologically irreversible fluxes (i.e. these should always be positive). All

fluxes were normalized to the sum of acetate and glycogen consumption rates (hereafter,

C consumed). To simulate the different ratios of glycogen over C consumed, glycogen

dissimilation ratio was varied between 0 to 1 Cmol glucose per Cmol consumed

(inversely, acetate varied from 1 to 0). This simulation assumes that none of the balanced

intermediates is accumulated inside the cell (steady-state assumption).

Commonly, maximization of biomass synthesis is used in FBA as optimization

objective. However, Accumulibacter only grow later when oxygen is available and from

the intracellular carbon reserves (PHAs) accumulated during anaerobiosis. Therefore, we

assume maximal carbon conservation, resp. minimal $CO_2$ production, as the cellular

objective to simulate Accumulibacter's metabolism anaerobically.

$$v_{opt} = \arg\min v_{CO2_{prod}} \quad \text{subject to} \begin{cases} Nv = 0 \\ v_{Ac_{upt}} = 1 : 0 \; Cmol/Cmol \; C_{consumed} \\ v_{Glyc_{deg}} = 0 : 1 \; Cmol/Cmol \; C_{consumed} \\ v_{irr} \geq 0 \end{cases}$$

For this optimization problem, MATLAB's function *linprog* was used.

To prevent stoichiometric constraints from specific assumption on the polymer

composition (PHB, PHV, PH2MV, etc), we introduce the polymer amount as reduced

precursors Ac-CoA* and Pr-CoA* such that:

188                    2Ac-CoA* → 3HB;

189              Ac-CoA* + Pr-CoA* → 3HV (or 3H2MB);

190                    2Pr-CoA* → 3H2MV;

**Experimental data reconciliation**

The results of the aforementioned simulations were compared to experimental data

taken from literature. Anaerobic-feast yields were used when available, otherwise the

yields of the whole anaerobic phase were used. In case of missing compound rates

(usually $CO_2$, $PH_2MV$), these were estimated using electron and carbon balancing (see

Supplementary Document 5). For datasets with redundant measurements, data

reconciliation was applied (24). For all calculations, we assumed acetate and glycogen

monomers were the only substrates and $CO_2$, PHB, PHV and $PH_2MV$ were the only

products. For yields reported without the respective error, we estimated their error by

propagating the assumed relative errors of 5% for acetate and PHB measurements and

10% for PHV, $PH_2MV$ and glycogen measurements. From the (reconciled) rates for PHB,

PHV and $PH_2MV$, the respective Ac-CoA* and Pr-CoA* rates were determined and

normalized to the amount of consumed substrates, in C-mol.

## RESULTS

### Characterization of Accumulibacter enrichments

This study was carried out using two independent Accumulibacter enrichment cultures. Performance data indicated a high level of PAO activity and microbial characterization confirmed by FISH, 16S-rRNA and *ppk1* gene analysis that the biomass was highly dominated by Accumulibacter clade I (see Table 1 and Supplementary Document 3).

Table 1 – Process parameters and key performance indicators of the two independent enrichments of Accumulibacter.

| | SBR-1 | SBR-2 | |
|---|---|---|---|
| pH | 7.6 | 7.6 | |
| T | 20ºC | 20ºC | |
| Carbon source | 210 mg COD/L (63:37 Acetate:Propionate) | 400 mg COD/L (75:25 Acetate:Propionate) | |
| Phosphate load per C | 0.1 Pmol/Cmol | 0.1 Pmol/Cmol | |
| | | | |
| Cycle time | 6 h Settling – 76 min; Anaerobic – 135 min; Aerobic – 135 min. | 6 h Settling – 30 min; Anaerobic – 112 min; Aerobic – 200 min. | |
| HRT | 12 h | 12 h | |
| SRT | 4 days | 5.6 days | |
| Aerobic SRT | 1.5 days | 3.1 days | |
| | | | |
| Anaerobic P-release per C-fed | 0.58 Pmol/Cmol | ~0.75 Pmol/Cmol | |
| Anaerobic-feast length | ~41 min | ~42 min | |
| | | | |
| FISH | PAO Acc I | PAO Acc I | |
| 16S amplicon sequencing (dominant OTUs) | Ca_Accumulibacter Rhodopseudomonas Blastochloris | Start: Dechloromonas Ca_Accumulibacter genus from f__Hyphomonadaceae | End: Ca_Accumulibacter Chryseobacterium genus from o__Sphingobacteriales |
| PPK analysis | Dominant: clade IC | Dominant: clade IA | |

### Redox cofactor preferences of key oxidoreductases

Enzymatic assays can be a useful tool to investigate the presence and potential extension of an enzymatically catalyzed conversion in a microbial community. However,

employed at that level, these assays will not specify "who" in the community is performing

the assessed conversion. Considering this limitation, we can still say 1) Accumulibacter

seems to not have a given function when such activity is not detected during the assay for

the whole community, 2) an enzyme from Accumulibacter is NAD(H) or NADP(H)

preferring when the community also shows the same clear preference. To further

characterize Accumulibacter's oxidoreductases (including their exact relative cofactor

preference), specific assays with purified enzymes can be performed as described in (25),

however this goes beyond the scope of our current analysis. The enzymatic activity assay

results are shown in blue in the metabolic network in Figure 1 and further details can be

found in Supplementary Document 4.

Figure 1 – Central carbon metabolic network of Accumulibacter including redox cofactor
preference of oxidoreductases assayed in this study. The value shown corresponds to the factor
difference between the preferred cofactor and the alternative one. This metabolic network is based
on the ancestral genome reconstruction done by (20). The arrows show the expected flux direction
under anaerobic conditions. The remaining cofactor preferences (in gray) remain as assumed by
analogy with other microorganisms. Despite not being annotated in the genome, the ED pathway
has been suggested as the route for glycogen degradation (21,22). A more extensive overview of
evidence for each pathway can be found in Supplementary Document 1.

The first question addressed was whether PHA accumulation is NADPH or NADH

dependent; it is clear from these results that the microbial community under study has a

NADH preferring accumulation of PHA with, at least, 10x more activity with NADH than

with NADPH. To further confirm this cofactor preference, the acetoacetyl-CoA reductase

activity was assayed using cell-free extracts from a second enrichment and both

directions of the reaction were monitored (i.e. $NAD(P)H$ + acetoacetyl-CoA $\leftrightarrow$ $NAD(P)^+$ +

3-hydroxybutyryl-CoA). In addition, a purified acetoacetyl-CoA reductase, encoded by a

gene highly similar to Accumulibacter's *phaB* (see Supplementary Document 4), was

employed as positive control as it is NAD(H)-preferring.

The second question tackled was related to the long-standing debate on glycolysis

proceeding via EMP or ED. To clarify this, we looked for the presence of the glucose-6-

phosphate dehydrogenase, the enzyme that catalyzes the first step of ED, and which is

also common to the oxidative branch of the pentose phosphate pathway (oxPP). No

activity was found in the cell-free extracts obtained from our enrichments, while the

biological positive control used, a cell-free extract from *P. putida* KT2440, showed clear

activity both with $NAD^+$ and $NADP^+$. Therefore, Accumulibacter is likely to use EMP as

glycolytic route and to produce the NADH required for PHA synthesis.

Since for many organisms oxPP is an important source of NADPH for growth (26) and

it is not used by Accumulibacter, the next question was if NADPH could be provided by

the isocitrate dehydrogenase (a TCA cycle enzyme). The oxidation of isocitrate was tested

using either $NAD^+$ or $NADP^+$ and the activity with $NADP^+$ was, at least, 70x higher, thus

isocitrate dehydrogenase is indeed a candidate supplier of NADPH in Accumulibacter.

Additionally, we tested the activity of other TCA oxidoreductases and anaplerotic

routes: 1) the reduction of oxaloacetate to malate (catalyzed by malate dehydrogenase)

had, at least, 55x higher activity when using NADH than NADPH. Unfortunately, the small

activity when using NADPH impeded the study of the cofactor preference of the malic

enzyme (oxidation of malate to pyruvate) using our assay; 2) the glyoxylate shunt was

also found active and with an activity comparable to the positive biological control *E. coli*

grown in acetate; and lastly, 3) fumarate reductase and α-ketoglutarate dehydrogenase

were also tested but their activity was very low compared to all other enzymes tested in

our study.

Next to assaying the cofactor preference of the aforementioned TCA oxidoreductases,

we also evaluated the differences in enzymatic activity between the anaerobic and aerobic

phases. The enzyme assays gave similar activities for anaerobic and aerobically harvested

biomass and thus do not allow for concluding on the anaerobic operation mode of the

TCA. These results are in line with the previous observation of proteomics studies where

relative protein abundances do not change notably across an anaerobic/aerobic cycle

(27,28). Nevertheless, the collected information on cofactor preferences and glycolysis

type (summarized in Figure 1) allowed for setting up an intracellular carbon flux balance

analysis (FBA) simulator to test different hypotheses for the anaerobic metabolism of

Accumulibacter against published experimental data (Figure 2).

**Lumped stoichiometric model of Accumulibacter for Flux Balance Analysis**

The goal of the flux balance analysis was to assess the balance of reducing equivalents

by Accumulibacter during anaerobic acetate conversion to PHAs. To this end, we used the

stoichiometric model of Accumulibacter's anaerobic metabolism presented in Table 2,

which is a lumped version of the central carbon metabolic network shown in Figure 1.

Given the presented evidence that PHA accumulation in Accumulibacter is NADH-

consuming (instead of the previously assumed NADPH preference), it follows that the

previously-made assumption in many Accumulibacter models that the type of redox

cofactor used does not affect the stoichiometry is valid: If the reducing reaction is feasible

using NADH, which is usually a worse electron donor than reduced ferredoxin or NADPH,

then the latter two can always be converted to NADH using transhydrogenases and

without any energetic cost for the cell (26). The same does not hold for FADH, which

would require an input of energy (ATP) to be re-oxidized using $NAD^+$. Moreover, there is

still no experimentally validated mechanism on how Accumulibacter could re-oxidize

FADH in the absence of oxygen or nitrate. Therefore, for this simulation, we blocked this

FADH producing step in the TCA cycle (i.e. Succ-CoA to Fum) and for the remaining

reactions we neglected the different types of redox cofactors by simply balancing

"electrons" (note that each redox cofactor carries 2 electrons and in some publications

these are referred to as [H]). Since the FADH producing step in the TCA cycle is blocked,

only the oxidative branch (OAA to Succ-CoA via ICDH and AKGDH) and/or the reductive

branch (OAA to Succ-CoA via MDH and FR) are possible.

Table 2 – Stoichiometry used for flux balance analysis (in mol basis) to simulate
Accumulibacter's <u>anaerobic</u> metabolism. Reversible reaction: <->, irreversible reaction: ->. Note
that these are lumped reactions and each may involve several enzymatic steps.

| Pathway | Stoichiometry | Enzyme assayed in this study |
|---|---|---|
| Glycolysis (EMP) | $(Glycogen)_1$ -> $2Pyr$ + 4electrons | G6PDH 6PGDH GAPDH |
| Pyruvate to Acetyl-CoA | Pyr -> Ac-CoA + 2electrons + $CO_2$ | |
| Pyruvate to Oxaloacetate | Pyr + $CO_2$ <-> OAA | |
| Oxidative TCA branch | OAA + Ac-CoA -> Succ-CoA + $2CO_2$ + 4electrons | ICDH AKGDH |
| Reductive TCA branch (reverse TCA) | OAA + 4electrons -> Succ-CoA | MDH FR |
| Glyoxylate shunt | 2Ac-CoA -> Succ-CoA + 2electrons | ICL |
| Succinate-propionate shunt | Succ-CoA <-> Pr-CoA + $CO_2$ | |
| Ac-CoA* production (precursor for 3HB, 3HV and 3HMB) | Ac-CoA + electron -> Ac-CoA* (* means "+1electron") | AcAc-CoA red |
| Pr-CoA* production (precursor for 3H2MV, 3HV and 3HMB) | Pr-CoA + electron -> Pr-CoA* (* means "+1electron") | AcAc-CoA red |

301 Regarding the ATP generation, there are still, unfortunately, too many unknowns: 1)

302 how much ATP can be generated from efflux of ions (potassium, magnesium and

303 phosphate) associated with polyphosphate hydrolysis; 2) how much is required for

acetate uptake; 3) how much is needed to upgrade redox cofactors; 4) and how much ATP

is actually used for cellular maintenance. Therefore, we will not balance ATP in this

simulation. Still, it is important to note that the different pathways used for redox

balancing lead to different levels of ATP generation via direct substrate-level

phosphorylation, and that, in cases of polyphosphate limitation, more ATP-generating

pathways will be preferred over less generating ones.

Experiments show that the ratio between glycogen and acetate consumption by PAO's

is variable, e.g. between high/low temperature (29) or polyphosphate/glycogen

availability (30–32). Therefore, in our simulations, we varied this ratio between the

glycogen and acetate fluxes and we compared the results with experimental data reported

in literature.

Figure 2 – The amount of Ac-CoA* accumulated (PHB and PHV precursor) depends on the different proportion of glycogen to acetate consumed and on the available pathways. The three optimum redox balancing strategies are shown in Figure 3 ABC. From 0 to approx. 2/6 glycogen Cmol, it can be considered that bacteria are performing a polyphosphate-based metabolism (PAM) and from approx. 3/6 to 1 glycogen Cmol, a Glycogen-based metabolism (GAM) is used. The "optimum" and "minimum" lines were obtained by minimizing and maximizing $CO_2$ production, respectively. The feasible solution space (in yellow) lies in between these two cellular objectives, in which a mixture of different redox balancing strategies is used. The minimum carbon conserving strategy is that which only uses the oxidative TCA branch. Experimental data was retrieved from (7,22,29,31,32,35,47,50,61–72) and reconciled to respect carbon and energy conservation principles. Similar plots for Pr-CoA*, $CO_2$, alternative simulations and including error bars on the experimental data points can be found in Supplementary Document 6.

For the anaerobic sequestration of acetate as PHA, acetate needs to be reduced to form

PHA. The Ac-CoA* optimum in Figure 2 corresponds to the situation in which most carbon

is kept inside the cell, most likely in the form of PHB. The source of reducing power can

come from either glycolysis or partial TCA cycle operation and the corners of the optimum

line in Figure 2 represent the extreme cases, which are presented in Figure 3. If the

amount of glycogen degraded can supply exactly the amount of electrons needed for the

uptake and conversion of acetate, then only Ac-CoA* (as PHB) is expected and there is no

need to use any part of the TCA cycle (Figure 3B). However, in a typical GAO-like

metabolism, an excess of glycogen is degraded compared to the acetate consumed to

supply all ATP needed. This leads to the generation of an excess of reducing power, which

is used to produce Pr-CoA* leading to PHAs more reduced than PHB (e.g. PHV or even

$PH_2MV$), using the reductive TCA branch (Figure 3C). On the other hand, in a PAO-like

metabolism when most ATP is provided by polyphosphate metabolism, the amount of

reducing equivalents generated by glycolysis is usually limited when compared to the

amount of acetate consumed. In this situation, part of the acetate also needs to be oxidized

to produce reducing equivalents. With the enzymatic activity observation of an active

glyoxylate shunt, our simulations show this as the optimal pathway to produce PHAs

when glycogen is limiting (Figure 3A); it allows for higher carbon conservation (i.e. higher

PHA yield) than the "horseshoe" TCA operation and also avoids the problem of

regenerating FADH produced in a full TCA cycle run.

349   Figure 3 – Different optimum redox balancing strategies for Accumulibacter under anaerobic conditions. A) When little glycogen is degraded and is not enough to
350 reduce all acetate to PHB, the glyoxylate shunt as recommended by (47) is the most optimal way to provide these electrons. B) When the stoichiometric amount of
351 glycogen is degraded for acetate reduction to PHB, there is no real need to operate any part of the TCA cycle to balance redox and only PHB is then produced, as proposed
352 in (48). C) When more glycogen than needed for acetate reduction is degraded, the reductive TCA branch can be used as proposed by (47) for GAOs to sink electrons in
353 more reduced PHAs (e.g. PHV and PH2MV).

## DISCUSSION

**Characterization of Accumulibacter enrichments**

Firstly, it is important to note that all presented results and following interpretations hold for the microbial community present in our PAO enrichments (mainly *Ca.* Accumulibacter phosphatis clade I). Any extrapolation of these results to Accumulibacter clade II or GAOs must be considered an assumption until further validation. This is due to the well-known fact that, depending on their environment, different types of Accumulibacter clades can be found, and each may have a slightly different function (7,28,33–36).

**Updated biochemical (stoichiometric) model**

Accumulibacter and their flexible metabolism have shown to defy many of the traditional characteristics of anaerobic metabolism due to their energy and reducing power storage, for example: anaerobic PHB synthesis decoupled from growth or the fast anaerobic acetate uptake are thanks to ATP provided by their polyphosphate storage. Consequently, general biochemical knowledge may only help guiding thoughts and analyses on Accumulibacter's unique metabolism, and the actual characteristics of their biochemistry should still be validated. Here in this study, enzymatic assays were performed to define what are the redox cofactor preferences of the main oxidoreductases in the central metabolic pathways of Accumulibacter. The key finding in this study is the NADH-preferring PHA accumulation in Accumulibacter. Unlike for NAPDH-PHA producers, this NADH-preference allows for a direct consumption of the NADH produced in most of Accumulibacter's reducing power sources. This prevents the need for an additional conversion of NADH into NADPH at the cost of proton motive force via the membrane-bound transhydrogenase (PntAB) as suggested by (8). Although previous reports of the use of NADH to drive PHB accumulation are found in literature (10,37–40), the level of NADH preference of the acetoacetyl-CoA reductase from Accumulibacter is the

largest so far observed with virtually no NADPH-activity (see Supplementary Document

S4). This observation allows to re-think the role of PHAs: we hypothesize that depending

on the environment where microorganisms thrive, PHA accumulation will play a role as

carbon reservoir during metabolic over-flow (NADPH-driven accumulation) or as an

electron reservoir during scarcity of external electron acceptors (NADH-driven

accumulation). Thus, for Accumulibacter, PHA is essentially a fermentation product.

Another important finding was the absence of activity of the first step of the

sometimes-implicated ED glycolytic pathway. These results match those observed by (29)

and in (41) for GAOs, where no glucose-6-phosphate dehydrogenase activity was found,

using $NADP^+$ only. The experimental findings are in line with the absence of key ED genes

in the genome annotations of Accumulibacter and closest relatives *Dechloromonas*

*aromatica* and *Azoarcus sp. EbN1* (36,42). Furthermore, we also verified the absence of

the ED pathway with $NAD^+$ as electron acceptor since there are microorganisms known

to use both cofactors (25). On the other hand, these findings are in contradiction with

earlier [13]C NMR studies (21,22), which indicated that the ED pathway was more likely the

route for glycolysis than EMP. Nevertheless, it has to be noted that these early studies

comprise interpretations of [13]C patterns using simplified metabolic models, with limited

information on the reversibility of each reaction and potentially with reactions missing;

these are common pitfalls of the [13]C-labelling method (43). Furthermore, these [13]C NMR

studies were performed before the first draft genome of Accumulibacter was available

(42), and only the [13]C pattern in the different PHAs was measured and not in the metabolic

intermediates of each pathway used.

While NADPH is not required anaerobically, this reducing cofactor still has to be

produced aerobically for growth. Given the observation that NADPH is not produced by

the oxPP pathway, the next logical candidate is the isocitrate dehydrogenase in the TCA

cycle. This is indeed the case, which is not unexpected as most acetate consumers will use

this conversion to produce NADPH for their anabolism (44) and confirms the protein

annotation done by (36). Since PHA accumulation is now known to be NADH-preferring

and without another sink of NADPH, the latter might accumulate anaerobically and

thereby inhibit the isocitrate dehydrogenase reaction. Alternatively, a soluble

transhydrogenase can convert this NADPH into NADH and allow the oxidative branch of

the TCA to be operational under anaerobic conditions.

Regarding the activity of other TCA oxidoreductases and anaplerotic routes: 1) the

oxidation of malate using $NAD^+$ was also found in the studies of (29); 2) the glyoxylate

shunt was also found active as observed by (22,29,45), which is expected as this is the

anaplerotic route that allows for microorganisms to convert C2 sources like acetate into

C4 building blocks for anabolism (44); and 3) fumarate reductase and α-ketoglutarate

dehydrogenase activities were very low compared to all other enzymes tested in our

study and alike in the studies of (29). Since we did not include biological controls, we

cannot interpret these low activities. We recommend that these enzymes are assayed

again next to biological controls to at least show that the conditions used in the assay were

appropriate. We also propose to assay the oxidation of α-ketoglutarate using ferredoxin

as electron acceptor as the α-ketoglutarate:ferredoxin oxidoreductase has been identified

in Accumulibacter's genome and proteome (36).

All the aforementioned redox cofactor specificities were used to update

Accumulibacter's central carbon metabolic network. With this biochemical model, we

performed a Flux Balance Analysis to study Accumulibacter's main activity under

anaerobic conditions. Since growth is negligible, the main anaerobic metabolic activity is

restricted to the central pathways, hence why we consider a reduced metabolic network

sufficient for this analysis. Furthermore, the inherent FBA assumption of no intermediate

accumulation is acceptable while the flux going through each intermediate is much larger

than its own concentration. This is the case during acetate uptake when it is present at

high concentrations (i.e. uptake rate is maximal and steady). Inversely, it might not be

valid during the anaerobic famine phase when no acetate is present since it will depend

on how large the glycogen degradation flux is compared to the intracellular concentration

of intermediates.

**Flexible anaerobic metabolism – Adjustments depending on available compounds**

**accumulated and on the environment**

As (6) suggested in their review, the flexibility of Accumulibacter's metabolism is likely

the major reason why there is controversy in literature regarding how reducing power is

balanced under anaerobic conditions. Long-term exposure to set conditions (e.g. pH,

temperature, oxidation level of the substrate, nutrient availability, counter-ions, SRT,

settling time) will select for the best strategy for those conditions, but short-term

perturbations of those set conditions have shown that Accumulibacter seems to still be

able to solve the redox balancing problem even if sub-optimally regarding carbon

conservation (7,30,31,46). Thus, based on our analysis, we believe that there is not one

fixed stoichiometry, but a range of possible stoichiometries that in the end are defined by

the relative proportions of each of the intervening substrates, supplied to the system or

produced by a side-population (e.g. acetate, glycogen, polyphosphate, other VFAs, oxygen,

nitrate, hydrogen).

The simulations showed Accumulibacter can face three distinct situations: A) when

reducing equivalents from glycolysis are limiting compared to the acetate imported

(Yagci's model (47)); B) when glycolysis supplies exactly enough reducing equivalents to

convert all imported acetate into PHB (Mino's model (48)); and lastly C) when glycolysis

is producing an excess of reducing equivalents (alike Yagci's model for GAOs (47)).

The anaerobic use of the glyoxylate shunt in a PAO-like metabolism to provide for the

reducing equivalents needed for PHA accumulation is supported by the studies of (45,47);

However, as seen in Figure 2, it does not explain the higher levels of Ac-CoA* found (points

above the optimum line) when more reduced PHAs are expected. This indicates that there

might be an alternative process that allows the cell to conserve extra carbon, in other

words, more Ac-CoA* and less Pr-CoA*. It could be that 1) fully anaerobic conditions were

not attained when performing the experiments, or that 2) another side population is

providing electrons in the form of a more reduced organic substrate or even hydrogen gas

(49) or 3) Accumulibacter has yet another, alternative way of balancing redox which has

yet to be described/demonstrated.

Also for the scenario where glycolysis produces an excess of reduced cofactors as in a

GAO-like metabolism, a few experimental data points fall outside of the feasible solution

space of our simulations (points above the optimum line in Figure 2); these could be

explained in the case that Accumulibacter produces $H_2$ to solve an excess of NADH as

described by (8).

**The multilayered complexity of a physiological analysis of Accumulibacter**

The experimental data gathered are likely also noisy due to the intrinsic heterogeneity

of Accumulibacter enrichments; clade differences, population heterogeneity or even

different environment (e.g. depending on the position of the cell in a floc/granule).

Therefore, experimental data points under the optimum line in Figure 2 likely represent

mixtures of cells, each with a different mode of operation (Figure 3 A, B or C).

Furthermore, any kinetic limitation (i.e. pathway capacity) may also explain a sub-

optimal, mixed phenotype. And from this analysis, when all enzymatic machinery is

available anyway, it seems possible that the same cell might be able to go from A to B to C

depending on the availability of acetate (e.g. transition feast/famine) or if polyphosphate

becomes limiting, or even from C to A when glycogen becomes limiting.

Nevertheless, despite the noise, the experimental data outside the solution space

suggests this stoichiometric model is still incomplete. For example, when allowing for

hydrogen production in excess glycogen-to-acetate conditions or the oxidation of acetate

in a fully operational TCA cycle in a glycolysis-limiting scenario, the observed data can be

explained (see Supplementary Document 6). The first is supported by the detection of

hydrogen gas produced by an Accumulibacter enrichment (8), while the latter has been

previously suggested based on [13]C NMR observations (50) and stoichiometric analyses

such as done by (30). Furthermore, a novel protein has been proposed based on

metagenomic analysis to allow for full TCA operation under anaerobic conditions (42);

however, there is still no experimental evidence that indeed validates this peculiar

anaerobic TCA operation. Moreover, to confirm full TCA cycle activity anaerobically, we

suggest the use of $^{13}$C labelled acetate together with MS techniques, which allow for

tracking $^{13}$C patterns in the different pathway intermediates. In (51), labelled propionate

has been used and $^{13}$C was found back in the PHV fragments that come from acetyl-CoA,

thus supporting the full TCA should be active although alternative metabolic routes could

also explain this observation. Furthermore, it may also be considered that Accumulibacter

can also make use of energy conservation features employed by many anaerobes such as

electron bifurcation (52), which employs ferredoxin oxidoreductases to allow for

alternative pathways in the central carbon metabolism.

## CONCLUSION

In this study an interdisciplinary approach was used, which integrated findings from genomics, transcriptomics and different physiological datasets as well as biochemical assays, to unravel the metabolic flexibility of Accumulibacter under dynamic conditions.

This flexible metabolic function can explain previous controversy in PAO literature. Depending on the exact history and cultivation conditions, Accumulibacter can exhibit different metabolic phenotypes. The NADH dependent PHA synthesis generates a direct link to central pathways used by Accumulibacter without the previously assumed requirement to generate NADPH for PHA synthesis through an oxidative PPP or ED pathways, or an energy-consuming transhydrogenase. Thus, under anaerobic conditions, Accumulibacter can make combinations of glycolysis (EMP), different branches of the TCA cycle (incl. glyoxylate shunt) and potentially other pathways not yet considered. This poses a challenge to predictive EBPR modelling, as it implies that stoichiometry is not fixed, but variable, spanning continuously from polyphosphate to glycogen-based phenotypes.

This metabolic versatility is likely what allows Accumulibacter to be, in most situations, very close to a carbon conservation optimum, which is key to ensure their competitiveness in the dynamic environments of EBPR systems.

## ACKNOWLEDGEMENTS

These investigations were supported by the SIAM Gravitation Grant 024.002.002, the Netherlands Organization for Scientific Research (NWO). The authors would like to thank Roel van de Wijgaart, Alexandre Carnet, Hein van der Wall, Koen Verhagen, David Weissbrodt, Alex Salazar, and Thomas Abeel for their collaboration in this project. The authors would also like to thank Sergio Tomas Martinez and Eleni Vasilakou for proofreading this manuscript.

## COMPETING INTERESTS

The authors declare no conflict of interest.

## REFERENCES

1. Van Loosdrecht MCM, Pot M a., Heijnen JJ. Importance of bacterial storage polymers in bioprocesses. Water Sci Technol. 1997;35(1):41–7.

2. Seviour RJ, Mino T, Onuki M. The microbiology of biological phosphorus removal in activated sludge systems. FEMS Microbiol Rev. 2003;27(1):99–127.

3. Seviour RJ, McIlroy S. The microbiology of phosphorus removal in activated sludge processes-the current state of play. J Microbiol. 2008;46(2):115–24.

4. Mino T, van Loosdrecht MCM, Heijnen JJ. Microbiology and biochemistry of the enhanced biological phosphate removal process. Water Res. 1998 Nov;32(11):3193–207.

5. Oehmen A, Lemos PC, Carvalho G, Yuan Z, Keller J, Blackall LL, et al. Advances in enhanced biological phosphorus removal: From micro to macro scale. Water Res. 2007;41:2271–300.

6. Zhou Y, Pijuan M, Oehmen A, Yuan Z. The source of reducing power in the anaerobic metabolism of polyphosphate accumulating organisms (PAOs) - A mini-review. Water Sci Technol. 2010 Apr;61(7):1653–62.

7. Welles L, Tian WD, Saad S, Abbas B, Lopez-Vazquez CM, Hooijmans CM, et al. Accumulibacter clades Type I and II performing kinetically different glycogen-accumulating organisms metabolisms for anaerobic substrate uptake. Water Res. Elsevier Ltd; 2015 Jul;83:354–66.

8. Oyserman BO, Noguera DR, del Rio TG, Tringe SG, McMahon KD. Metatranscriptomic insights on gene expression and regulatory controls in Candidatus Accumulibacter phosphatis. ISME J. 2016 Apr 10;10(4):810–22.

9. Madison LL, Huisman GW. Metabolic Engineering of Poly ( 3-Hydroxyalkanoates ): From DNA to Plastic. Microbiol Mol Biol Rev. 1999;63(1):21–53.

10. Ritchie GAF, Senior PJ, Dawes EA. The Purification and Characterization of Acetoacetyl-Coenzyme A Reductase from Azotobacter beijerinckii. Biochem J. 1971;121:309–16.

11. Senior PJ, Beech GA, Ritchie GA, Dawes EA. The role of oxygen limitation in the formation of poly- -hydroxybutyrate during batch and continuous culture of Azotobacter beijerinckii. Biochem J. 1972;128(5):1193–201.

12. Liebergesell M, Steinbüchel A. Cloning and nucleotide sequences of genes relevant for biosynthesis of poly(3-hydroxybutyric acid) in Chromatium vinosum strain D. Eur J Biochem. 1992 Oct 1;209(1):135–50.

13. Daims H, Brühl A, Amann R, Schleifer KH, Wagner M. The domain-specific probe EUB338 is insufficient for the detection of all Bacteria: development and evaluation of a more comprehensive probe set. Syst Appl Microbiol. 1999 Sep;22(3):434–44.

14. Amann RI. In situ identification of micro-organisms by whole cell hybridization with rRNA-targeted nucleic acid probes. In: Akkermans ADL, Van Elsas JD, De Bruijn FJ, editors. Molecular Microbial Ecology Manual. Dordrecht: Springer Netherlands; 1995. p. 331–45.

15. Amann RI, Binder BJ, Olson RJ, Chisholm SW, Devereux R, Stahl DA. Combination of 16S rRNA-targeted oligonucleotide probes with flow cytometry for analyzing mixed microbial populations. Appl Environ Microbiol. 1990 Jun;56(6):1919–25.

16. Flowers JJ, He S, Yilmaz S, Noguera DR, McMahon KD. Denitrification capabilities of two biological phosphorus removal sludges dominated by different "Candidatus

573    Accumulibacter" clades. Environ Microbiol Rep. 2009;1:583–8.

574    17.    Rubio-Rincón FJ, Welles L, Lopez-Vazquez CM, Nierychlo M, Abbas B, Geleijnse M,
575           et al. Long-term effects of sulphide on the enhanced biological removal of
576           phosphorus: The symbiotic role of Thiothrix caldifontis. Water Res. 2017;116:53–
577           64.

578    18.    Saad SA, Welles L, Abbas B, Lopez-Vazquez CM, van Loosdrecht MCM, Brdjanovic
579           D. Denitrification of nitrate and nitrite by 'Candidatus Accumulibacter phosphatis'
580           clade IC. Water Res. 2016 Nov;105:97–109.

581    19.    McMahon KD, Yilmaz S, He S, Gall DL, Jenkins D, Keasling JD. Polyphosphate kinase
582           genes from full-scale activated sludge plants. Appl Microbiol Biotechnol.
583           2007;77:167–73.

584    20.    Oyserman BO, Moya F, Lawson CE, Garcia AL, Vogt M, Heffernen M, et al. Ancestral
585           genome reconstruction identifies the evolutionary basis for trait acquisition in
586           polyphosphate accumulating bacteria. ISME J. 2016 Dec 29;10(12):2931–45.

587    21.    Maurer M, Gujer W, Hany R, Bachmann S. Intracellular carbon flow in phosphorus
588           accumulating    organisms    from    activated    sludge    systems.    Water    Res.
589           1997;31(4):907–17.

590    22.    Hesselmann RPX, Von Rummell R, Resnick SM, Hany R, Zehnder  a. JB. Anaerobic
591           metabolism of bacteria performing enhanced biological phosphate removal. Water
592           Res. 2000;34(14):3487–94.

593    23.    Orth JD, Thiele I, Palsson BØ. What is flux balance analysis? Nat Biotechnol. Nature
594           Publishing Group; 2010 Mar 1;28(3):245–8.

595    24.    van der Heijden RTJM, Heijnen JJ, Hellinga C, Romein B, Luyben KCAM. Linear
596           constraint relations in biochemical reaction systems: I. Classification of the
597           calculability and the balanceability of conversion rates. Biotechnol Bioeng. 1994
598           Jan 5;43(1):3–10.

599    25.    Olavarría K, Valdés D, Cabrera R. The cofactor preference of glucose-6-phosphate
600           dehydrogenase from Escherichia coli--modeling the physiological production of
601           reduced cofactors. FEBS J. 2012 Jul;279(13):2296–309.

602    26.    Fuhrer T, Sauer U. Different biochemical mechanisms ensure network-wide
603           balancing of reducing equivalents in microbial metabolism. J Bacteriol. American
604           Society for Microbiology (ASM); 2009 Apr;191(7):2112–21.

605    27.    Wilmes P, Andersson AF, Lefsrud MG, Wexler M, Shah M, Zhang B, et al. Community
606           proteogenomics highlights microbial strain-variant protein expression within
607           activated sludge performing enhanced biological phosphorus removal. ISME J.
608           2008;2:853–64.

609    28.    Wexler M, Richardson DJ, Bond PL. Radiolabelled proteomics to determine
610           differential functioning of Accumulibacter during the anaerobic and aerobic
611           phases of a bioreactor operating for enhanced biological phosphorus removal.
612           Environ Microbiol. 2009;11:3029–44.

613    29.    Erdal ZK, Erdal UG, Randall CW. Biochemistry of enhanced biological phosphorus
614           removal and anaerobic COD stabilization. Water Sci Technol. 2005;52(10–
615           11):557–67.

616    30.    Zhou Y, Pijuan M, Zeng RJ, Yuan Z. Involvement of the TCA cycle in the anaerobic
617           metabolism of polyphosphate accumulating organisms (PAOs). Water Res. Elsevier
618           Ltd; 2009;43(5):1330–40.

619    31.    Zhou Y, Pijuan M, Zeng RJ, Lu H, Yuan Z. Could polyphosphate-accumulating
620           organisms (PAOs) be glycogen-accumulating organisms (GAOs)? Water Res.

621    2008;42:2361–8.

622    32.    Welles L, Abbas B, Sorokin DY, Lopez-Vazquez CM, Hooijmans CM, van Loosdrecht
623           MCM, et al. Metabolic response of "candidatus accumulibacter phosphatis" clade II
624           C to changes in influent P/C ratio. Front Microbiol. 2017 Jan 5;7(JAN).

625    33.    Flowers JJ, He S, Malfatti S, del Rio TG, Tringe SG, Hugenholtz P, et al. Comparative
626           genomics of two "Candidatus Accumulibacter" clades performing biological
627           phosphorus removal. ISME J. Nature Publishing Group; 2013;7(12):2301–14.

628    34.    Welles L, Lopez-Vazquez CM, Hooijmans CM, van Loosdrecht MCM, Brdjanovic D.
629           Prevalence of 'Candidatus Accumulibacter phosphatis' type II under phosphate
630           limiting conditions. AMB Express. 2016;6(1).

631    35.    Acevedo B, Oehmen  a., Carvalho G, Seco  a., Borrás L, Barat R. Metabolic shift of
632           polyphosphate-accumulating organisms with different levels of polyphosphate
633           storage. Water Res. 2012;46:1889–900.

634    36.    Barr JJ, Dutilh BE, Skennerton CT, Fukushima T, Hastie ML, Gorman JJ, et al.
635           Metagenomic and metaproteomic analyses of Accumulibacter phosphatis-
636           enriched floccular and granular biofilm. Environ Microbiol. 2016 Jan;18(1):273–
637           87.

638    37.    Kim J, Chang JH, Kim K-J. Crystal structure and biochemical properties of the (S)-3-
639           hydroxybutyryl-CoA dehydrogenase PaaH1 from Ralstonia eutropha. Biochem
640           Biophys Res Commun. 2014;448:163–8.

641    38.    Barycki JJ, O LK, Strauss AW, Banaszak LJ, Author C, Banaszak Dietrich Professor
642           LJ. Sequestration of the Active Site by Interdomain Shifting: Crystallographic and
643           Spectroscopic Evidence for Distinct Conformations of L-3-Hydroxyacyl-CoA
644           Dehydrogenase Running Title: Human L-3-Hydroxyacyl-CoA Dehydrogenase. JBC
645           Pap Press Publ. 2000;

646    39.    Haywood GW, Anderson AJ, Chu L, Dawes EA. The role of NADH- and NADPH-
647           linked acetoacetyl-CoA reductases in the poly-3-hydroxybutyrate synthesizing
648           organism Alcaligenes eutrophus. FEMS Microbiol Lett. 1988 Aug;52(3):264–259.

649    40.    Ling C, Qiao G-Q, Shuai B-W, Olavarria K, Yin J, Xiang R-J, et al. Engineering
650           NADH/NAD+ ratio in Halomonas bluephagenesis for enhanced production of
651           polyhydroxyalkanoates    (PHA).    Metab    Eng.    Elsevier    Inc.;    2018
652           Sep;49(September):275–86.

653    41.    Filipe CDM, Daigger GT, Grady CPL. A metabolic model for acetate uptake under
654           anaerobic conditions by glycogen accumulating organisms: Stoichiometry,
655           kinetics, and the effect of pH. Biotechnol Bioeng. 2001;76(1):17–31.

656    42.    García Martín H, Ivanova N, Kunin V, Warnecke F, Barry KW, McHardy AC, et al.
657           Metagenomic analysis of two enhanced biological phosphorus removal (EBPR)
658           sludge communities. Nat Biotechnol. 2006;24(10):1263–9.

659    43.    Van Winden W, Verheijen P, Heijnen S. Possible pitfalls of flux calculations based
660           on13C-labeling. Metab Eng. 2001;3(2):151–62.

661    44.    Gottschalk G. Bacterial Metabolism. Gottschalk G, editor. New York, NY: Springer
662           New York; 1986. 359 p.

663    45.    Burow LC, Mabbett AN, Blackall LL. Anaerobic glyoxylate cycle activity during
664           simultaneous utilization of glycogen and acetate in uncultured Accumulibacter
665           enriched in enhanced biological phosphorus removal communities. ISME J. 2008
666           Oct;2(10):1040–51.

667    46.    Erdal UG, Erdal ZK, Daigger GT, Randall CW. Is it PAO-GAO competition or
668           metabolic shift in EBPR system? Evidence from an experimental study. Water Sci

669   Technol. 2008 Oct;58(6):1329–34.

670 47. Yagci N, Artan N, Çokgör EU, Randall CW, Orhon D. Metabolic model for acetate
671   uptake by a mixed culture of phosphate- and glycogen-accumulating organisms
672   under anaerobic conditions. Biotechnol Bioeng. 2003;84:359–73.

673 48. Mino T, Tsuzuki Y, Matsuo T. Effect of phosphorus accumulation on acetate
674   metabolism in the biological phosphorus removal process. In: Ramadori R, editor.
675   Biological phosphate removal from wastewaters: proceedings from the
676   International Conference of Advanced Water Pollution Control (IAWPRC). Oxford,
677   England: Pergamon Press; 1987. p. 27–38.

678 49. Lawson CE, Strachan BJ, Hanson NW, Hahn AS, Hall ER, Rabinowitz B, et al. Rare
679   taxa have potential to make metabolic contributions in enhanced biological
680   phosphorus removal ecosystems. Environ Microbiol. 2015 Dec;17(12):4979–93.

681 50. Pereira H, Lemos PC, Reis MAM, Crespo JPSG, Carrondo MJT, Santos H. Model for
682   carbon metabolism in biological phosphorus removal processes based on in vivo
683   13C-NMR labelling experiments. Water Res. 1996 Sep;30(9):2128–38.

684 51. Lemos PC, Serafim LS, Santos MM, Reis M a M, Santos H. Metabolic Pathway for
685   Propionate Utilization by Phosphorus-Accumulating Organisms in Activated
686   Sludge: 13C Labeling and In Vivo Nuclear Magnetic Resonance. Appl Environ
687   Microbiol. 2003 Jan 1;69(1):241–51.

688 52. Buckel W, Thauer RK. Energy conservation via electron bifurcating ferredoxin
689   reduction and proton/Na+ translocating ferredoxin oxidation. Biochim Biophys
690   Acta - Bioenerg. 2013 Feb;1827(2):94–113.

691 53. Satoh Y, Tajima K, Tannai H, Munekata M. Enzyme-catalyzed poly(3-
692   hydroxybutyrate) synthesis from acetate with CoA recycling and NADPH
693   regeneration in Vitro. J Biosci Bioeng. 2003 Jan;95(4):335–41.

694 54. Chohan, Copeland. Acetoacetyl coenzyme A reductase and polyhydroxybutyrate
695   synthesis in rhizobium (Cicer) sp. Strain CC 1192. Appl Environ Microbiol.
696   American Society for Microbiology; 1998 Aug;64(8):2859–63.

697 55. Olavarria K, Marone MP, da Costa Oliveira H, Roncallo JC, da Costa Vasconcelos FN,
698   da Silva LF, et al. Quantifying NAD(P)H production in the upper Entner-Doudoroff
699   pathway from Pseudomonas putida KT2440. FEBS Open Bio. 2015 Jan 1;5(1):908–
700   15.

701 56. Dean AM, Golding GB. Protein engineering reveals ancient adaptive replacements
702   in isocitrate dehydrogenase. Proc Natl Acad Sci U S A. 1997 Apr 1;94(7):3104–9.

703 57. Bologna FP, Andreo CS, Drincovich MF. Escherichia coli malic enzymes: two
704   isoforms with substantial differences in kinetic properties, metabolic regulation,
705   and structure. J Bacteriol. 2007 Aug 15;189(16):5937–46.

706 58. Giachetti E, Pinzauti G, Vanni P. A new continuous optical assay for isocitrate lyase.
707   Experientia. 1984 Feb;40(2):227–8.

708 59. Hillier AJ, Jericho RE, Green SM, Jago GR. Properties and function of fumarate
709   reductase (NADH) in Streptococcus lactis. Aust J Biol Sci. 1979 Dec;32(6):625–35.

710 60. Bunik VI, Denton TT, Xu H, Thompson CM, Cooper AJL, Gibson GE. Phosphonate
711   Analogues of α-Ketoglutarate Inhibit the Activity of the α-Ketoglutarate
712   Dehydrogenase Complex Isolated from Brain and in Cultured Cells †. Biochemistry.
713   2005 Aug;44(31):10552–61.

714 61. Smolders GJF, van der Meij J, van Loosdrecht MCM, Heijnen JJ. Stoichiometric model
715   of the aerobic metabolism of the biological phosphorus removal process.
716   Biotechnol Bioeng. Wiley Subscription Services, Inc., A Wiley Company; 1994 Sep

20;44(7):837–48.

62.    Smolders GJF, Klop JM, van Loosdrecht MCM, Heijnen JJ. A metabolic model of the
biological phosphorus removal process: I. Effect of the sludge retention time.
Biotechnol Bioeng. Wiley Subscription Services, Inc., A Wiley Company; 1995 Nov
5;48(3):222–33.

63.    Tian W-D, Lopez-Vazquez CM, Li W-G, Brdjanovic D, van Loosdrecht MCM.
Occurrence of PAOI in a low temperature EBPR system. Chemosphere. 2013
Aug;92(10):1314–20.

64.    Zeng RJ, Yuan Z, Keller J. Model-based analysis of anaerobic acetate uptake by a
mixed culture of polyphosphate-accumulating and glycogen-accumulating
organisms. Biotechnol Bioeng. 2003;83:293–302.

65.    López-Vázquez CM, Hooijmans CM, Brdjanovic D, Gijzen HJ, van Loosdrecht MCM.
A practical method for quantification of phosphorus- and glycogen-accumulating
organism populations in activated sludge systems. Water Environ Res. 2007
Dec;79(13):2487–98.

66.    Lanham AB. Full-scale biological phosphorus removal: quantification of storage
polymers, microbial performance and metabolic modelling. Universidade Nova de
Lisboa; 2012.

67.    Pijuan M, Saunders AM, Guisasola A, Baeza JA, Casas C, Blackall LL. Enhanced
biological phosphorus removal in a sequencing batch reactor using propionate as
the sole carbon source. Biotechnol Bioeng. 2004 Jan 5;85(1):56–67.

68.    Filipe CDM, Daigger GT, Grady CPL. A metabolic model for acetate uptake under
anaerobic conditions by glycogen accumulating organisms: Stoichiometry,
kinetics, and the effect of pH. Biotechnol Bioeng. 2001;76:17–31.

69.    Zeng R, Yuan Z, van Loosdrecht MCM, Keller J. Proposed modifications to metabolic
model for glycogen-accumulating organisms under anaerobic conditions.
Biotechnol Bioeng. Wiley Subscription Services, Inc., A Wiley Company; 2002 Nov
5;80(3):277–9.

70.    Carvalheira M. The effect of key process operational conditions on enhanced
biological phosphorus removal from wastewater. Universidade Nova de Lisboa;
2014.

71.    Brdjanovic D, Loosdrecht MCM van, Hooijmans CM, Alaerts GJ, Heijnen JJ.
Temperature Effects on Physiology of Biological Phosphorus Removal. J Environ
Eng. American Society of Civil Engineers; 1997 Feb 1;123(2):144–53.

72.    Brdjanovic D, Van Loosdrecht MCM, Hooijmans CM, Mino T, Alaerts GJ, Heijnen JJ.
Effect of polyphospbate limitation on the anaerobic metabolism of phosphorus-
accumulating microorganisms. Appl Microbiol Biotechnol. 1998 Aug
27;50(2):273–6.