AMPK-Fyn signaling promotes Notch1 stability to potentiate hypoxia-induced breast cancer stemness and drug resistance

Mohini Lahiry[1], Saurav Kumar[1], Kishore Hari[2], Adithya Chedere[1], Mohit Kumar Jolly[2], and Annapoorni Rangarajan[1]*

[1]Department of Molecular Reproduction, Development and Genetics, Indian Institute of Science, Bangalore-560012, India

[2]Centre for BioSystems Science and Engineering, Indian Institute of Science, Bangalore-560012, India

***Name and address for correspondence:** Dr. Annapoorni Rangarajan, Department of Molecular Reproduction, Development and Genetics, Indian Institute of Science, Bangalore 560012, Karnataka, India, Phone: 91-80-22933263; Fax: 91-80-23600999

e-mail: anu@iisc.ac.in

Summary:

Hypoxia is a hall mark of solid tumor microenvironment and contributes to tumor progression and therapy failure. The developmentally important Notch pathway is implicated in cellular response of cancer cells to hypoxia. Yet, the mechanisms that potentiate Notch signaling under hypoxia are not fully understood. Hypoxia is also a stimulus for AMP-activated protein kinase (AMPK), a major cellular energy sensor. In this study, we investigated if AMPK interacts with the Notch pathway and influences the hypoxia-response of breast cancer cells. Activating AMPK with pharmacological agent or genetic approaches led to an increase in the levels of cleaved Notch1 and elevated Notch signaling in invasive breast cancer cell lines. In contrast, inhibition or depletion of AMPK reduced the amount of cleaved Notch1. Significantly, we show that the hypoxia-induced increase in cleaved Notch1 levels requires AMPK activation. Probing into the mechanism, we demonstrate that AMPK activation impairs the interaction between cleaved Notch1 and its ubiquitin ligase, Itch/AIP4 through the tyrosine kinase Fyn. Under hypoxia, the AMPK-Fyn axis promotes inhibitory phosphorylation of Itch which abrogates its interaction with substrates, thus stabilizing cleaved Notch1 by reducing its ubiquitination and degradation. We further show that inhibition of AMPK alleviates the hypoxia-triggered, Notch-mediated stemness and drug resistance phenotype. Breast cancer patient samples also showed co-expression of hypoxia/AMPK/Notch gene signature. Our work thus establishes AMPK as a key component in the adaptation of breast cancer cells to hypoxia, and proposes therapeutic inhibition of AMPK to mitigate the hypoxia-triggered aggressiveness.

Keywords: Hypoxia, AMP-activated protein kinase (AMPK), Itch/AIP4, cleaved Notch1, breast cancer, cancer stem cell

Introduction:

Developmental pathways such as Notch are known to regulate self-renewal and cell fate decisions in embryonic and tissue-specific stem cells (Artavanis-Tsakonas et al., 1999). In contrast, deregulation of Notch signaling is associated with malignant transformation (Radtke and Raj, 2003). Chromosomal translocation and activating mutations in Notch1 promote T-acute lymphoblastic leukemia (Aster et al., 2008). Aberrant Notch signaling has also been observed in several solid tumors including breast, cervical and prostate cancer (Allenspach et al., 2002). In breast cancer, we and others have reported an accumulation of intracellular domain of Notch1, as well as elevated expression of Notch ligand Jagged1 and its correlation with poor patient survival (Mittal et al., 2009; Reedijk et al., 2005; Stylianou et al., 2006). In contrast, there is loss of expression of negative regulators of Notch like Numb (Pece et al., 2004). Aberrant Notch signaling promotes cancer cell growth and survival, and alters metabolism. Recent studies have additionally implicated a critical involvement of Notch pathway in breast cancer stem cell self-renewal (D'Angelo et al., 2015) and therapy resistance (Wang et al., 2010).

Notch belongs to a family of four transmembrane receptors, named Notch 1-4 (Gordon et al., 2008). Ligand binding results in multiple proteolytic cleavages leading to the generation of cleaved Notch1 proteins, and finally the release of the transcriptionally active Notch intracellular domain (NICD) from the plasma membrane. NICD translocates to the nucleus where it forms a DNA-binding complex with other co-activators like MAML, CSL and p300, and activates expression of target genes like HES and HEY (Jarriault et al., 1995; Schroeter et al., 1998). Notch signaling is regulated at multiple levels, including presentation & availability of ligand, endocytosis & trafficking of processed receptors, and degradation by recognition of C-terminal PEST domain (Bray, 2006). In addition, post-translational

modifications including phosphorylation, ubiquitination, hydroxylation and acetylation affect

the stability, transcriptional activity and localization of NICD (Andersson et al., 2011).

Hypoxia develops in solid tumor environment due to rapid proliferation of cancer cells and an

inadequate supply of oxygen to meet the demand. Under most scenarios, this hypoxia is

likely to be cyclic in nature (Saxena and Jolly, 2019). Adaptation of tumor cells to hypoxia

contributes to more aggressive and therapy-resistant phenotypes (Kim and Lee, 2017; Kunz

and Ibrahim, 2003). Cellular adaptation to hypoxia is mainly orchestrated by the transcription

factor HIF1-α (Wenger and Gassmann, 1997). The Hypoxia-HIF1 axis regulates cancer

stemness by activating several pathways like TAZ (Xiang et al., 2014), CD47 (Zhang et al.,

2015), PHGDH (Samanta et al., 2016), and ALKBH5 (Zhang et al., 2016). It also aids breast

cancer cell migration and invasion through the Notch pathway (Sahlgren et al., 2008).

Hypoxia potentiates Notch signaling by multiple mechanisms. It induces the expression of

Notch ligands like Jagged2 and Delta-like ligand 4 (DLL4) (Jubb et al., 2009; Xing et al.,

2011). Direct binding of HIF1-α to NICD leads to the upregulation of Notch-downstream

genes (Gustafsson et al., 2005). In addition, the hypoxia-HIF1α axis also stabilizes NICD

(Gustafsson et al., 2005); however, the downstream molecular mechanisms that lead to Notch

stabilization remains poorly understood.

Yet another protein activated by hypoxia is the AMP-activated protein kinase, AMPK

(Gusarova et al., 2011; Mungai et al., 2011). It is a heterotrimeric protein composed of

catalytic α subunits, and regulatory β and γ subunits. AMPK is activated by stimuli that

trigger an increase in AMP/ATP ratio, such as, glucose deprivation and chronic hypoxia

(Hardie et al., 2012). Hypoxia is also known to activate AMPK independent of any changes

in AMP/ATP ratio, through oxidant signaling (Emerling et al., 2009; Mungai et al., 2011).

Upon activation, AMPK inhibits anabolic pathways while activating catabolic pathways, thus

bringing about energy homeostasis (Hardie et al., 2012). In addition to its central role in

metabolism, recent studies have highlighted AMPK functions in regulating several

physiological processes such as cell growth, polarity, apoptosis, autophagy and cell fate

(Mihaylova and Shaw, 2011).

Owing largely to its growth suppressive effects, AMPK is mainly recognized for its tumor

suppressive properties (Hardie and Alessi, 2013), and consistent with this notion, AMPK

activating agents have been proposed for cancer treatment (Hardie, 2013). However,

accumulating evidence points to a contextual oncogenic role of AMPK by enabling cancer

cell survival under challenges faced in the tumor microenvironment, such as, glucose,

oxygen, and matrix deprivation (Hindupur et al., 2014; Jeon et al., 2012; Jeon and Hay, 2012,

2015; Liang and Mills, 2013). In breast cancer, we and others have reported an increasing,

grade-specific activation of AMPK in patient samples (Hart et al., 2015b; Sundararaman et

al., 2016). We have also highlighted a novel role for AMPK in inhibiting apoptosis by

phosphorylating PEA15 (Hindupur et al., 2014) and in mitigating the anchorage-deprivation

stress by activating autophagy in breast cancer cells (Hindupur et al., 2014; Jeon et al., 2012;

Saha et al., 2018). Hence, AMPK is increasingly recognized to mediate several non-

metabolic functions by phosphorylating a plethora of proteins that participate in diverse

cellular signaling pathways (Schaffer et al., 2015) which may be beneficial for tumor

progression.

Thus, independent studies have demonstrated hypoxia-triggered activation of AMPK and

Notch pathway; yet, a cross talk between these two pathways remains unexplored. In this

study, we investigated if AMPK interacts with the Notch pathway and influences the

response of breast cancer cells to hypoxia. We show that AMPK activation potentiates Notch

signaling by stabilizing cleaved Notch1 through Fyn tyrosine kinase. We further show that

inhibition or depletion of AMPK mitigates the hypoxia-driven, Notch1-mediated cancer stem

cell and drug resistance phenotype. Our study thus highlights AMPK as a crucial component

of the hypoxia-Notch1 signaling in promoting breast cancer aggressiveness.

Results:

AMPK regulates cleaved Notch1 levels in breast cancer cells

To test if AMPK is involved in regulating Notch signaling, we first gauged the effect of

pharmacologic and genetic activation of AMPK on Notch1 expression in breast cancer cell

lines. To measure the Notch1 protein levels, we performed immunoblotting using an antibody

specific to the C-terminus of Notch1 which recognizes both the full length (FL-Notch1; 300

kD) and proteolytically cleaved (cl-Notch1; ~120 kD) forms of Notch1. We used A769662 as

a pharmacological activator of AMPK (Cool et al., 2006). Detection of an increase in

phosphorylation of ACC, a bonafide substrate of AMPK (Winder and Hardie, 1996),

confirmed AMPK activation upon treatment with A769662 **(Figure 1A)**. AMPK activation

did not alter the transcript level of Notch1 **(Figure S1A)** or the full length Notch1 protein

levels in MDA-MB-231 invasive breast cancer cell line **(Figure 1A)**; however, we observed

elevated levels of cleaved Notch1 **(Figure 1A)**, suggesting possible activation of Notch

signaling by AMPK.

We next gauged the effect of AMPK activation on cleaved Notch1 levels in a variety of

invasive, non-invasive and immortalized breast cells. We found elevated levels of cleaved

Notch1 in invasive BT 474 and HCC 1806 breast cancer cells when treated with A769662

(Figure S1B). However, in non-invasive MCF-7 and immortalized HMLE breast cells

(Elenbaas et al., 2001), cleaved Notch1 levels were not modulated by AMPK activator

(Figure S1B), suggesting that the regulation of cleaved Notch1 protein levels by AMPK is

likely to be cell-type and context-specific.

Since pharmacological agents can have non-specific effects, we additionally used genetic

approaches to activate AMPK and checked for cleaved Notch1 levels. Over expression of

constitutively activated AMPK with γR70Q mutation (Hamilton et al., 2001) led to increased

AMPK activity, as gauged by increase in pACC levels, as well as higher levels of cleaved

Notch1 **(Figure 1B)**. Similarly, overexpression of a constitutively active form of an AMPK

upstream kinase CaMKKβ also resulted in elevated cleaved Notch1 levels **(Figure S1C)**.

Thus, increasing AMPK activity led to elevated cleaved Notch1 levels in invasive breast

cancer cells.

To further address the role of AMPK in regulating cleaved Notch1 levels, we investigated the

effects of AMPK inhibition and depletion using pharmacologic and genetic approaches. Upon

treatment with a pharmacological inhibitor of AMPK, Compound C (Zhou et al., 2001), we

observed reduced phosphorylated ACC levels (confirming the efficacy of the inhibitor) along

with reduced levels of cleaved Notch1 **(Figure 1C)**. To further confirm, this effect using a

genetic approach, we employed doxycycline-inducible knockdown strategy targeting the

catalytic α subunit of AMPK. In keeping with the higher expression and activity of AMPKα2

isoform, compared to AMPKα1, in breast cancer cell lines (Hindupur et al., 2014), we

gauged the effect of AMPKα2 knockdown on cleaved Notch1 levels. Addition of

doxycycline to MDA-MB-231 cells stably expressing the inducible shAMPKα2 (clone#4)

construct led to a reduction in the levels of total AMPK α2, as well as a reduction in the

levels of cleaved Notch1, compared to un-induced control cells **(Figure 1D)**. AMPK

depletion using an independent, doxycycline-inducible, shRNA construct (clone#1) against

AMPKα2 also led to a reduction in cleaved Notch1 levels **(Figure S1D)**. To further confirm

the role of AMPK, we used AMPK α1/α2 double knock out (AMPK-DKO) MEFs.

Compared to wild type (WT) MEFs, AMPK-DKO MEFs showed reduced levels of cleaved

Notch1 **(Figure 1E)**. Restoration of AMPK expression in DKO MEFs by exogenous

expression of catalytic subunits of AMPK led to increase in cleaved Notch1 levels **(Figure**

S1E), further confirming a direct role for AMPK in the positive regulation of cleaved Notch1

levels.

AMPK activation promotes Notch signaling

We next investigated whether an increase in cleaved Notch1 levels upon AMPK activation

led to higher Notch downstream signaling. To do so, we first checked if AMPK activation

leads to elevated levels of the active form of cleaved Notch1. The C-terminal specific

antibody that we have used is likely to recognize both cleaved but membrane tethered form of

Notch1, as well as the active intracellular domain (ICD) of Notch1 that translocates to the

nucleus (Brou et al., 2000). Cleavage by gamma-secretase at Valine 1744 results in the

generation of active form of Notch1 (NICD) which enhances transcription of downstream

genes. We therefore checked the effect of AMPK activation in the generation of NICD using

an antibody that specifically recognizes the gamma secretase-specific cleaved Notch1 (Valine

1744). Immunoblotting revealed elevated levels of NICD in the presence of AMPK activator

A769662 **(Figure 1F)**. To further confirm a role for AMPK in modulating NICD levels, we

tested the effect of AMPK knockdown in cells ectopically expressing NICD. We chose

HEK293T cells for this experiment as they show very minimal basal levels of endogenous

Notch1 expression. AMPK depletion led to reduction in the levels of exogenously expressed

NICD **(Figure S1F)**, additionally suggesting that the effect of AMPK on accumulation of cleaved Notch1 may be downstream of gamma secretase cleavage, perhaps by altering NICD stability.

In light of our data revealing an increase in active, cleaved Notch1 levels upon AMPK activation, we next tested if AMPK modulates Notch signaling. To do so, we used the Notch-responsive 12xCSL-luciferase reporter assay (Blokzijl et al., 2003). Treatment with AMPK activator A769662 led to a 2.6-fold increase in 12xCSL-luciferase reporter activity **(Figure 1G)**, indicative of increase in canonical Notch signaling upon AMPK activation. To further gauge the effect of AMPK activation on Notch signaling, we measured the endogenous levels of Notch target genes. Treatment with AMPK activator also led to an increase in the levels of two Notch pathway downstream genes Hes1 **(Figure 1H)** and Hes5 **(Figure S1G)**, as gauged by RT-PCR and immunocytochemistry, respectively. In addition, a microarray-based data of A769662-treated MDA-MB-231 cells revealed upregulation of several Notch pathway-regulated genes including HES5, HEY2, SNAI2, HEYL, IFNG and CDH6 **(Figure S1H)**. Taken together, these observations reveal that elevated cleaved Notch1 levels downstream to AMPK activation leads to enhanced Notch signaling.

AMPK inhibition or depletion impairs hypoxia-induced NICD generation

In response to hypoxia, independent reports have shown elevated Notch signaling in breast cancer (Sahlgren et al., 2008) and AMPK activation in other cell types (Emerling et al., 2009; Mungai et al., 2011). In light of our data revealing elevated Notch signaling upon AMPK activation in breast cancer cells, we investigated if AMPK is involved in hypoxia-induced Notch signaling. To address this, we treated MDA-MB-231 cells with a hypoxia-mimetic agent $CoCl_2$, which simulates hypoxic effects by stabilizing HIF-1α (Xi et al., 2004). As

shown before (Chen et al., 2017), treatment with $CoCl_2$ led to an increase in phosphorylated

AMPK levels in MDA MB 231 cells **(Figure 2A)**. It also led to an increase in cleaved

Notch1 levels **(Figure 2A)**. Treatment of yet another invasive breast cancer cell line BT474

with $CoCl_2$ led to increase in carbonic anhydrase IX (CA9), which is a direct readout of HIF-

1α, and pACC levels, which is an indicator of AMPK activity **(Figure S2A)**. Consistent with

our previous results, we observed elevated levels of cleaved Notch1 **(Figure S2A)**. Having

confirmed both AMPK activation and elevated cleaved Notch1 levels in the presence of

$CoCl_2$, we investigated if AMPK is involved in hypoxia-triggered elevated cleaved Notch1

levels. To do so, we tested the effect of AMPK inhibition in this process under hypoxia.

$CoCl_2$-induced elevated cleaved Notch1 was impaired in the presence of AMPK inhibitor,

Compound C **(Figure 2B)**. Similar results were observed in BT-474 cells also **(Figure S2A)**.

To further confirm this, we measured the effect of AMPK depletion in the $CoCl_2$-induced

cleaved Notch 1 levels. Cells expressing the control pTRIPZ construct showed expected

increase in cleaved Notch1 levels in the presence of $CoCl_2$ when treated with doxycycline,

whereas those expressing inducible $AMPK\alpha2$ shRNA construct (clone#1) failed to do so

(Figure 2C). We obtained similar results with an independent inducible $AMPK\alpha2$ (clone#4)

shRNA construct **(Figure S2B)**. Together, these data indicated a role for AMPK in the

$CoCl_2$-induced increase in cleaved Notch1 levels.

To better study the AMPK-Notch crosstalk under hypoxia relevant to physiological and

patho-physiological condition, we used oxygen-regulatable trigas incubator to create hypoxic

3% oxygen condition in contrast to ambient 21% oxygen in regular incubators. Cells cultured

in hypoxia showed increased levels of HIF-1α and CA 9, indicative of successful generation

of hypoxic condition **(Figure 2D)**. As expected, we also observed elevated levels of cleaved

Notch1 and pAMPK under hypoxia compared to normoxia **(Figure 2D)**. However, AMPK

inhibition with Compound C or depletion of AMPKα2 impaired the hypoxia-induced

increase in cleaved Notch1 levels **(Figure 2E-F)**, thus confirming a role for AMPK in the

hypoxia-induced elevated cleaved Notch1 levels.

To further probe if AMPK plays a role in the generation of the active form of cleaved Notch1

under hypoxia, we used the gamma secretase cleavage specific Notch1 (Valine 1744)

antibody. We observed elevated NICD levels under hypoxia by immunoblotting **(Figure 2D)**.

In addition, we performed immunocytochemistry to gauge NICD levels. As expected, we

observed an increase in pAMPK staining intensity under hypoxia, confirming the generation

of hypoxic condition. We also observed an increase in NICD staining intensity under hypoxia

compared to cells grown in normoxia **(Figure S2C)**. Further, AMPK inhibition reduced the

hypoxia-triggered NICD staining intensity **(Figure 2G)**, suggesting a role for AMPK in the

generation of active, cleaved form of Notch1 in hypoxia.

We next asked if the positive regulation of cleaved Notch1 levels by AMPK in response to

hypoxic condition is also reflected in Notch signaling. To test this, we probed the activity of

the Notch-responsive 12XCSL-reporter construct in the presence of $CoCl_2$ to mimic hypoxic

condition, and tested the role of AMPK. As expected, treatment with $CoCl_2$ increased the

activity of the 12XCSL-reporter; however, inhibition of AMPK impaired this **(Figure 2H)**.

Taken together, these results reveal a key role for AMPK in the accumulation of active,

cleaved Notch 1 and positive regulation of the Notch signaling pathway in response to

hypoxic stress.

AMPK inhibition affects stability of cleaved Notch1 under hypoxia

In light of our observation that AMPK activation increases the amounts of cleaved Notch1,

but not full-length receptor, we next explored if AMPK enhances the stability of cleaved

Notch1. To do so, we performed a cycloheximide chase assay in MDA-MB-231 cells

harboring doxycycline-inducible AMPKα2 shRNA. Addition of cycloheximide revealed a

faster reduction in the amount of cleaved Notch1 under AMPK depleted conditions **(Figure**

3A), suggesting a possible role for AMPK in regulating cleaved Notch1 stability. To further

confirm this, we subjected AMPK DKO MEFs to cycloheximide treatment. Treatment with

cycloheximide revealed a higher reduction in cleaved Notch1 levels in AMPK DKO MEFs

compared to WT MEFs **(Figure S3A)**. Together, these data showed that the absence of

AMPK enhances the rate of Notch1 degradation, suggesting a role for AMPK in stabilization

of cleaved Notch1.

We next investigated if AMPK is involved in the hypoxia-induced stabilization of cleaved

Notch1 as shown earlier **(Figure 2)**, and as reported previously in P19 cells (Gustafsson et

al., 2005). To do so, we grew MDA-MB-231 cells harbouring doxycycline-inducible

AMPKα2 shRNA constructs in trigas incubator with 3% oxygen and subsequently treated

with cycloheximide. Immunoblotting revealed heightened reduction of cleaved Notch1 levels

under hypoxia in the presence of doxycycline **(Figure 3B)**. Similarly, addition of AMPK

inhibitor Compound C to MDA-MB-231 cells exposed to $CoCl_2$ also revealed heightened

reduction in cleaved Notch1 levels compared to vehicle control DMSO **(Figure S3B)**.

Together, these data confirmed a key role for AMPK in regulating cleaved Notch1 stability in

response to hypoxia.

Since cleaved Notch1 is reported to be targeted for degradation through the 26S proteasomal

machinery (Gupta-Rossi et al., 2002), we next investigated if AMPK affects this process. To

address this, we used the proteasomal complex inhibitor MG132, and asked if reduced

cleaved Notch1 levels observed under AMPK inhibited condition can be rescued by

proteasomal inhibition. As expected, treatment with MG132 led to stabilization of cyclinD3,

which served as a positive control (lane 2 vs 1; **Figure 3C**). Further, we found that

Compound C-mediated reduction in cleaved Notch1 levels (lane 3 vs 1) was rescued in the

presence of MG132 (lane 4 vs 3; **Figure 3C**). Thus, AMPK activation plays a critical role in

cleaved Notch1 stabilization by preventing its degradation through the proteasomal complex.

AMPK inhibits Notch1 ubiquitination by modulating its interaction with Itch

Since protein degradation by the proteasomal system involves tagging of target proteins with

ubiquitin, we next investigated if AMPK stabilizes cleaved Notch1 by regulating its

ubiquitination. To address this question, we first asked if AMPK alters Notch1 K48-linked

polyubiquitination which specifically marks proteins for proteasomal degradation. To do so,

we immunoprecipitated Notch1 and probed by immunoblotting using an antibody specific for

K48-linked polyubiquitin. We observed a reduction in K48-linked ubiquitination of cleaved

Notch1 in the presence of AMPK activator A769662 **(Figure 4A)**. In the reverse experiment,

when we inhibited AMPK using Compound C, the K48-linked ubiquitination of cleaved

Notch1 was elevated **(Figure S4A)**. These data suggested that AMPK possibly stabilizes

cleaved Notch1 by regulating its ubiquitination, and thereby degradation by the proteasomal

pathway.

To further explore the mechanism downstream of AMPK in mediating cleaved Notch1

stability, we investigated into post-translational modifications, such as, Ser/Thr

phosphorylation and acetylation that are known to stabilize Notch1 protein levels (Andersson

et al., 2011). We did not detect much change in the overall levels of Ser/Thr phosphorylation

on immuno-precipitated cleaved Notch1 upon AMPK activation **(Figure S4B)**. Further,

bioinformatic analysis revealed Notch1 T1998 as a putative AMPK phosphorylation site. To

test if this site is involved in cleaved Notch1 stability, we carried out site-directed

mutagenesis using an NICD expressing construct at threonine 1998 and changed it to non-

phosphorylatable alanine (T1998A). If cleaved Notch1 stability is mediated by

phosphorylation of this site, we hypothesized that the non-phosphorylatable alanine mutant

would be insensitive to AMPK activation, and thus undergo increased degradation. To

investigate this, wild type NICD and an engineered NICD harbouring T1998A substitution

were transfected into HEK 293T cells, treated with A769662, and cleaved Notch1 levels were

gauged. NICD T1998A responded to AMPK activation in a manner similar to wildtype NICD

(Figure S4C). This indicated that the effect mediated by AMPK is not through

phosphorylation of Notch1 at residue T1998. Further, using an acetyl lysine specific

antibody, we also failed to detect much change in the overall acetylation status of cleaved

Notch1 upon AMPK activation **(Figure S4D)**.

Since we failed to observe changes in the overall Ser/Thr phosphorylation or acetylation

status of cleaved Notch1 in response to AMPK activation, we shifted our focus to ubiquitin

ligases. Notch1 is known to be a substrate for several ubiquitin ligases such as FBXW-7,

Itch/AIP4, NEDD 4, Deltex 1, c-Cbl (Moretti and Brou, 2013). Notably, a study showed that

AMPK activation disrupts the interaction of Itch/AIP4 (henceforth referred to as Itch) with its

substrate p73 (Adamovich et al., 2014). We therefore examined a possible role for Itch in

AMPK-mediated cleaved Notch1 stability. To do so we first examined the levels of Itch upon

AMPK activation. We failed to see changes in the levels of Itch in the presence of AMPK

activator A769662 **(Figure 4B)**. We then queried the activity of Itch by checking its auto-

ubiquitination status. Itch auto-ubiquitinates itself by the non-degradative, K63 linkages

(Scialpi et al., 2008), which also positively regulates its activity. We immunoprecipitated Itch

and checked for its ubiquitination status by immunoblotting. Immunoprecipitated endogenous

Itch revealed highly diminished ubiquitin smear in the presence of AMPK activator A769662

compared to DMSO vehicle control **(Figure 4C)**. We next investigated if AMPK activation

affects the Itch-Notch interaction. Indeed, immunoprecipitation of Notch1 followed by

immunoblotting with Itch revealed a reduced interaction between Notch1 and Itch in the

presence of AMPK activator A769662 **(Figure 4D)**. These data suggested a possible

mechanism downstream of AMPK activation leading to cleaved Notch1 stability by

regulating Itch-Notch1 interaction.

To further confirm a role for AMPK in Itch-mediated regulation of cleaved Notch 1 stability,

we asked if AMPK activation would rescue the effects of Itch overexpression on cleaved

Notch1 levels. As expected, Itch overexpression led to a decrease in the amounts of another

known substrate of Itch, c-Jun, as well as cleaved Notch1, whereas the presence of AMPK

activator A769662 impaired this effect **(Figure 4E)**. Overexpression of Itch in yet another

invasive breast cancer cell line, BT-474, yielded similar down-modulation of cleaved Notch1

which was rescued on AMPK activation **(Figure S4E)**. Taken together, these observations

suggested a protective role for AMPK in Itch-mediated Notch degradation.

We next asked if AMPK plays a similar role in cleaved Notch1 stabilization in response to

hypoxia by modulating Itch-Notch interaction. To address this, we first investigated Itch

levels under hypoxia, and found that Itch levels remained unchanged in hypoxic condition of

3% O_2 compared to normoxia **(Figure 4F)**. We next gauged Notch-Itch interaction in

hypoxia, as well as tested the effect of AMPK depletion in this interaction. To do so, we used

MDA-MB-231 cells harboring the doxycycline inducible AMPK shRNA system. To gauge

Itch-Notch interaction, we immunoprecipitated endogenous Itch and measured the amounts

of associated Notch1 by immunoblotting. Under un-induced (minus doxycycline) condition,

we observed a reduction in cleaved Notch1 levels associated with immunoprecipitated Itch in

hypoxia compared to normoxia (lanes 2 vs 1; **Figure 4G**). Upon induction of AMPK

depletion with doxycycline, we observed higher amount of cleaved Notch1 in the Itch-

immunoprecipitate under hypoxia (lane 4 vs 3), suggesting higher Itch-Notch interaction

upon AMPK depletion. These data supported a role for AMPK in regulating the Itch- Notch1

interaction under hypoxia.

We previously showed a direct role for AMPK in regulating the K-48 linked degradative

ubiquitination of Notch1 (**Figure 4A** and **Figure S4A**). We next inquired if the regulation of

cleaved Notch1-Itch interaction under hypoxia affects the K-48 linked degradative

ubiquitination of cleaved Notch1 in an AMPK-dependent manner. To address this, we

performed immunoprecipitation of Notch1 under hypoxia in the presence and absence of

AMPK inhibitor Compound C, and probed with K-48 linkage specific anti-ubiquitin

antibodies. We found a reduction in the Notch1 K-48 associated ubiquitination in hypoxia

(Figure 4H). In contrast, AMPK inhibition increased the K-48 linked ubiquitination of

cleaved Notch1 in hypoxic conditions **(Figure 4H)**. Thus, these data uncover a novel AMPK-

mediated regulation of cleaved Notch1 stability in hypoxia through modulating its interaction

with the ubiquitin ligase Itch.

AMPK mediates Itch phosphorylation in hypoxia

We next sought to understand how AMPK negatively regulates the Itch-Notch interaction.

Phosphorylation of Itch has been shown to regulate its interaction with substrates (Yang et

al., 2006). In contrast to Ser/Thr phosphorylation, tyrosine phosphorylation of Itch negatively

modulates its activity towards substrates (Yang et al., 2006). To gauge if hypoxia affects

changes in Itch Tyr-phosphorylation, we immunprecipitated Itch and undertook

immunoblotting with phospho-tyrosine (pTyr) specific antibodies. We observed an increase

in tyrosine phosphorylation of immunoprecipitated Itch in hypoxia **(Figure 5A)**. To gauge if

AMPK is involved in this process, we tested the effects of AMPK knockdown on Itch Tyr

phosphorylation. Induction of AMPK knockdown with doxycycline led to a marked decrease

in tyrosine phosphorylation of Itch **(Figure 5A)**, revealing a novel role for AMPK in

regulating Itch Tyr-phosphorylation in hypoxia.

Since AMPK is a Ser/Thr kinase, whereas we noticed a change in Tyr phosphorylation of Itch

in response to AMPK activation, we hypothesized that AMPK might be mediating its effects

indirectly, by affecting a tyrosine kinase. Fyn is a Src family tyrosine kinase which is known

to phosphorylate and inhibit Itch activity (Yang et al., 2006). To check if Fyn-mediated Itch

phosphorylation is involved downstream of AMPK in cleaved Notch1 stabilization in

hypoxia, we used an antibody that specifically recognizes tyrosine phosphorylation of human

Itch (Tyr 420). We observed an increase in the levels of Itch Y420 phosphorylation under

hypoxia **(Figure 5B)**; AMPK knockdown with doxycycline decreased this phosphorylation

(Figure 5B). These results identify a novel role for AMPK in Fyn-mediated phosphorylation

of Itch in hypoxia.

To further confirm a role for Fyn in AMPK-mediated cleaved Notch1 stabilization in hypoxia

we tested the effects of Fyn inhibition on cleaved Notch 1 levels in hypoxia. Treatment of

cells with PP2, a Src family kinase inhibitor, hindered the hypoxia-triggered elevation in

cleaved Notch1 levels **(Figure 5C)**. To more specifically gauge the role of Fyn, we

transfected cells with a dominant negative construct of Fyn and investigated its effect on the

tyrosine phosphorylation of Itch in hypoxia. To do so, we immunoprecipitated endogenous

Itch and probed for Tyr-phosphorylation by immunoblotting. As shown before, hypoxia led

to an increase in Tyr-phosphorylation of Itch, but transfection of DN Fyn impaired this

(Figure 5D). Consistent with this, we noticed reduced cleaved Notch1 levels in DN Fyn

expressing cells in hypoxia (**Figure 5D**, input WB).

To further confirm the role of Fyn in the regulation of active, cleaved Notch1 in hypoxia, we

depleted Fyn using shRNA and tested its effect on NICD levels. Depletion of Fyn led to a

reduction in NICD levels in hypoxia **(Figure S5A)**. Together, these results confirmed the

involvement of Fyn in stabilizing cleaved Notch1 in hypoxia.

The observations so far revealed an AMPK-dependent and Fyn-mediated negative regulation

of Itch in hypoxia. However, a link between AMPK and Fyn activity remains unknown. To

address this, we first investigated the effects of AMPK activation and inhibition on Fyn

activity. The activity of Src-family kinases (SFK), of which Fyn is a member (Parsons and

Parsons, 2004), is regulated by phosphorylation at specific tyrosine residues. Since phospho-

Fyn specific antibodies are not available, we used SFK (Tyr 416) recognizing antibody which

detects Tyrosine 416 specific phosphorylation on Src-family members that leads to their

activation (Roskoski Jr, 2015). AMPK activation with A769662 led to an increase in the

levels of tyrosine 416 phosphorylation on Src family kinases **(Figure 5E)**. To understand the

significance of this AMPK-mediated positive regulation of Fyn activity in hypoxia, we

gauged the effect of AMPK depletion on Src-family tyrosine 416 phosphorylation levels in

hypoxia by immunoblotting. In MDA MB 231 cell harbouring the doxycycline-inducible

AMPK KD system, in the absence of doxycycline, hypoxia increased phosphorylated Src-

family tyrosine 416 levels **(Figure 5F)**. However, addition of doxycycline brought about a

reduction in phosphorylated Src-family tyrosine 416 levels **(Figure 5F)**. These data

suggested a possible role for AMPK in positively regulating the activity of Fyn in hypoxia.

To further understand the mechanism by which AMPK augments Fyn activation, we carried mass spectrometry to detect interacting proteins of Fyn in cells treated with AMPK activator A769662 and vehicle control DMSO. HEK293T cells stably expressing FLAG-tagged Fyn were immunoprecipitated with FLAG specific antibodies and protein complexes were isolated and subjected to LC-MS. We specifically focussed on the association of Fyn with phosphatases that are known to dephosphorylate Fyn and affect its activity. PTPRF (Vacaresse et al., 2008), a phosphatase known to dephosphorylate the inhibitory phosphorylation at Y528 and thus activate Fyn, was associated with Fyn only in AMPK activated condition, but not in control DMSO treated cells **(Supplementary Table 1)**. Further, PTPN5 (Nguyen et al., 2002) and PTPRC (Vacaresse et al., 2008) are phosphatases known to dephosphorylate the activating phosphorylation of Fyn at Y416, thus decreasing its activity. We detected these in DMSO condition, but not in AMPK activated cells **(Supplementary Table 2)**. Thus, our proteomic analysis further substantiated the role of AMPK in positively modulating Fyn activity by facilitating differential association of phosphatases that promote activation or inactivation of Fyn. These results together indicate an important role for Fyn kinase in AMPK-dependent regulation of Itch activity in hypoxia. Together, these data identify a novel AMPK-dependent regulation of tyrosine phosphorylation of Itch by Fyn kinase in the stabilization of cleaved Notch1.

Hypoxia-induced AMPK activation promotes stem-like properties and drug-resistance in breast cancer cells through Notch signaling

Having established an interplay between AMPK and Notch, we next investigated the biological significance of the AMPK-Notch pathway interaction in hypoxia. Hypoxic condition is known to facilitate cancer stem cell (CSC)-like properties through involvement of Notch signaling (Sahlgren et al., 2008; Xing et al., 2011). We investigated if AMPK is

involved in this process. To address this, we first gauged the effect of AMPK inhibition or

depletion on the hypoxia-induced CSC phenotype by measuring the expression of stemness

markers like Bmi1 and Nanog. As expected, MDA-MB-231 cells grown in 3% oxygen

showed elevated levels of Bmi1 and Nanog compared to normoxia **(Figure 6A)**. However,

inhibition of AMPK with compound C in the same experiment prevented the hypoxia-

induced elevated expression of these stemness markers **(Figure 6A)**. Likewise, $CoCl_2$-

induced expression of Bmi1 in these cells was also abrogated by AMPK inhibition **(Figure**

S6A). We also observed similar effect in yet another invasive cancer cell line BT474 **(Figure**

S6B), suggesting a role for AMPK in the hypoxic response leading to elevated stemness.

To further confirm a role for AMPK in hypoxia-triggered stemness properties, we

investigated the effect on sphere formation in defined media condition – an in vitro surrogate

for stemness. As expected, hypoxia led to an increase in the number of spheres generated;

however, AMPK inhibition blocked this **(Figure 6B)**. We obtained similar results with $CoCl_2$

treatment in the presence of AMPK inhibitor **(Figure S6C)**, together indicating the

requirement of AMPK in hypoxia-induced sphere forming potential.

Since AMPK depletion downregulated Notch signaling, we asked if increasing cleaved

Notch1 levels can rescue the effects of AMPK depletion on stemness properties under

hypoxia. To do so, we treated doxycycline induced AMPK depleted MDA-MB-231 cells with

a synthetic short peptide (referred to as DSL peptide) corresponding to the Notch ligand

Jagged1 whose overexpression promotes Notch signaling (Nickoloff et al., 2002). DSL

peptide treatment led to an increase in the levels of cleaved Notch1 in the doxycycline-

induced AMPK depleted cells **(Figure 6C)**. Treatment with DSL also led to an increase in the

stemness marker Bmi1 expression **(Figure 6C)**, as well as increased the number of spheres

formed in these cells **(Figure 6D)**. Integrating our observations with existing literature about

stemness regulators, we devised a mathematical model that predicted that hypoxia triggered

AMPK can drive as well as maintain a stem-like state characterized by high levels of Notch1

and stemness markers such as Oct4 and Bmi1 (**Figure 6E-G**, **Figure S6D**, Supplementary

methods). Put together, these data suggest the involvement of AMPK-Notch signaling axis in

the hypoxia-mediated increase in stemness.

An increase in stemness property is also associated with drug resistance (Prieto-Vila et al.,

2017), and Notch signaling has been implicated in drug resistance under hypoxic condition in

various tumors such as osteosarcoma, ovarian cancer and breast cancer (Morata-Tarifa et al.,

2016; Sansone et al., 2007; Seo et al., 2016). We investigated whether the AMPK-Notch1

signaling axis affected drug sensitivity of breast cancer cells under hypoxia. To do so, we

first gauged the effect of treatment of doxycycline-inducible AMPK knockdown cells with

chemotherapeutic drug doxorubicin under hypoxia. We observed a 2-fold increase in the IC_{50}

value for doxorubicin when the cells were exposed to hypoxia in the absence of doxycycline;

however, addition of doxycycline prevented this response **(Figure 6H)**.

To further confirm a role for AMPK in hypoxia-induced drug resistance phenotype, we

measured the extent of DNA damage accumulated by these cells in hypoxic condition by

undertaking pH2A.X staining. As expected, we observed an increase in pH2A.X staining

intensity upon treatment with doxorubicin (1μM) under normoxic condition **(Figure 6I)**. In

contrast, in hypoxia, treatment with the same concentration of doxorubicin failed to induce

DNA damage response **(Figure 6I)**, whereas AMPK-depletion restored the DNA damage

response in hypoxia **(Figure 6I)**, suggesting that hypoxia-induced AMPK contributes to drug

resistance. To gauge the involvement of Notch in this process, we treated AMPK knockdown

cells with synthetic DSL peptide which increases cleaved Notch1 levels in these cells under

hypoxia **(Figure 6C)** and assessed the DNA damage. We observed a reduction in pH2A.X

staining in DSL peptide-treated cells compared to control untreated cells, indicating

restoration of drug resistance phenotype in AMPK depleted cells by increasing cleaved

Notch1 levels **(Figure 6I)**. Thus, AMPK-Notch axis plays a key role in mediating CSC

phenotype of stemness and drug resistance driven by hypoxia.

Hypoxia-induced AMPK-Notch1 signaling in vivo

Since hypoxia conditions are prevalent in tumors (Gruber et al., 2004), we next sought to

understand the relevance of the AMPK-Notch1 axis in vivo in tumors derived from animal

xenografts. To do so, we measured the levels of cleaved and active forms of Notch1 (NICD)

using Valine 1744 specific antibody in tumors derived from control and AMPK depleted

cells. We used tumors derived from BT 474 cells stably expressing control scrambled

constructs or shRNA constructs against AMPK α2 that were injected sub-cutaneously into

immunocompromised mice (Hindupur et al., 2014). In small tumors generated by AMPK-

depleted cells, we found that NICD levels were highly diminished compared to control

scrambled tumors **(Figure 7A)**.

We next addressed the relevance of AMPK-Notch signaling in human breast cancer. High

grade breast cancers are majorly associated with hypoxia (Vaupel et al., 2002). We sought to

gauge AMPK activity and NICD levels in high grade breast cancer tissue samples. To do so,

we performed immunohistochemical analyses using antibodies against pACC (as a measure

of AMPK activity) and against gamma secretase cleavage-specific Notch1 (Valine 1744)

antibodies (as a measure of NICD) levels. We performed co-staining for pACC and NICD

and immunofluorescence microscopy revealed high co-expression in patient samples **(Figure

S7A)**. Further, we observed a predominant cytoplasmic staining for pACC while NICD

staining was nuclear, as expected **(Figure S7A)**. We then performed DAB-based

immunohistochemistry in serial sections of chemo-naïve, grade III invasive ductal carcinoma

breast cancer patient sample (N=19) for these two antibodies. We observed an association

(p=0.0422, Fischer exact test) between pACC and NICD levels in these samples **(Figure 7B)**.

Together, these data highlighted the importance of the AMPK-Notch1 axis in patient tumors.

Furthermore, we investigated into the response of patient-derived breast cancer cells to

hypoxia. The primary tissue-derived cells exposed to hypoxic condition showed elevated

levels of pACC, cleaved Notch1 and phospho-Itch levels, whereas pharmacological inhibition

of AMPK in these cells under hypoxic condition reduced the cleaved Notch1 and phospho-

Itch levels **(Figure 7C)**, suggesting that the AMPK-Itch-Notch axis might also operate in

patient-derived cells.

In order to further understand the clinical relevance of our study, we analysed the gene

signatures of AMPK (22), Hypoxia (40) and Notch (26) pathways (Supplementary Table 3)

in TCGA primary breast cancer dataset by performing gene-wise correlation across the

signatures by Pearson's product moment correlation **(Figure 8 A-C)** and Spearman

correlation methods **(Figure S8)**. We observed significant correlation (Pearson co-relation)

between AMPK-Hypoxia, Notch-Hypoxia and AMPK-Notch gene signatures **(Figure 8A-C)**.

There was no difference observed between both the correlation methods used (Pearson vs

Spearman's co-relation) **(Figures 8 & S8)**. In 880 (22*40), 1040 (26*40), 572 (22*26)

number of pair-wise gene correlations between AMPK-Hypoxia **(Figure 8A)**, Notch-

Hypoxia **(Figure 8B)**, and AMPK-Notch **(Figure 8C)** respectively, 669, 840 and 515 gene

pairs showed a positive correlation, of which the correlation for 534, 669, 412 gene pairs

were significant (p<0.01), implying a strong interaction between hypoxia-AMPK-Notch

signaling pathways. Further, hierarchical clustering analysis of gene expression profiles from

NCBI-GEO dataset (GSE40206), a microarray dataset of 80 Indian breast cancer patient

cohort, revealed co-expression of hypoxia, AMPK and Notch-responsive genes in majority of

cases **(Figure S9)**, thus underscoring the relevance of the hypoxia-AMPK-Notch axis in

breast cancers.

Discussion

Hypoxia in solid tumors is considered a malice aiding cancer progression and hindering

successful therapy. Notch signaling is implicated as a major regulator of hypoxia-mediated

stemness and drug resistance. Because of its importance in numerous cancers, the Notch

pathway has been a major candidate for developing therapeutic agents. While gamma

secretase inhibitors and antibodies against receptor and ligands have reached clinical trials,

gastrointestinal toxicity and other side effects have led to shifting of focus on druggable

targets that impinge on the regulation of Notch signaling (Andersson and Lendahl, 2014; Guo

et al., 2011). In this study, we identify AMPK as a novel molecular regulator of cleaved

Notch1 stability under hypoxia (Figure 8D), and propose that inhibition of AMPK can be a

potential therapeutic strategy impairing the adaptation of breast cancer cells to hypoxia.

Hypoxia-induced Notch1 signaling has been reported in melanoma (Bedogni et al., 2008),

adenocarcinoma of lungs (Chen et al., 2007), glioma (Qiang et al., 2012), as well as breast

carcinomas (Sahlgren et al., 2008). Mechanistically, it has been shown that HIF1α and NICD

directly interact and augment transcription of Notch responsive genes (Gustafsson et al.,

2005). Another mechanism implicated in this crosstalk involves FIH (factor inhibiting HIF1-

α) which is critical in blocking differentiation in myogenic and neural progenitor cells (Zheng

et al., 2008). HIF-α synergises with Notch co-activator MAML to increase expression of

Snail and *Slug* and reduce *E-cadherin* expression (Chen et al., 2010). Treatment with

mTORC1/2 inhibitors also elevate Notch signaling and maintain a drug-resistant CSC

phenotype (Bhola et al., 2016) in triple-negative breast cancers. Only one study has reported

the stabilization and increased half-life of NICD in hypoxia in a HIF1α-dependent manner

(Gustafsson et al., 2005); however, the molecular mechanisms remained unknown. We show

here that AMPK activation is critical for cleaved Notch1 stabilization under hypoxia. Using

cycloheximide chase assays, we show that AMPK inhibition, depletion or genetic ablation

causes enhanced degradation of cleaved Notch1. Cleaved Notch1 is targeted for proteasomal

degradation after ubiquitination (Öberg et al., 2001; Qiu et al., 2000; Wu et al., 2001). We

show here that the cleaved Notch1-K-48-linked ubiquitination, which marks proteins for

degradation, is reduced under hypoxia as well as upon pharmacological AMPK activation. In

contrast, AMPK inhibition or depletion increases ubiquitination of cleaved Notch1. We

further show that in AMPK DKO MEFs which do not have AMPK activity, cleaved Notch

levels are low, which can be restored by AMPK re-constitution. Our study thus identifies a

novel role for AMPK in cleaved Notch1 stabilization. Our data revealed that $CoCl_2$ treatment

led to both AMPK activation as well as elevated cleaved Notch1 levels. Furthermore, AMPK

inhibition led to heightened decrease of $CoCl_2$-induced elevated cleaved Notch1 levels. Since

$CoCl_2$ mimics hypoxia by stabilizing HIF-1α, these data suggest a possible role for HIF1-α

in this process. However, hypoxia can trigger AMPK both in HIF-1α dependent (Singh et al.,

2013) as well independent fashion (Marin et al., 2016). Thus, AMPK and HIF-1α are likely

to affect cleaved Notch1 stability independently, as well as in a concerted manner. Further

investigations are required to distinguish their individual and combined effects.

Post-translational modifications like phosphorylation and acetylation of the Notch

intracellular domain regulate its ubiquitination (Fryer et al., 2004; Popko-Scibor et al., 2011).

However, we failed to find significant changes in phosphorylation or acetylation of cleaved Notch upon AMPK activation. Notch and its interaction with ubiquitin ligases has been often shown to be influenced by regulators like Numb, which suppresses Notch signaling by recruiting Itch and facilitating degradation of Notch (McGill and McGlade, 2003). We identified AMPK as a negative regulator of Itch-mediated Notch degradation. Several studies have identified phosphorylation of Itch as a major regulator of its activity. Kinases like ATM and JNK activate Itch by increasing its enzymatic activity for a specific set of targets like c-Jun or c-FLIP, whereas other kinases like Fyn reduce interaction of Itch with substrates including Jun-B as well as Notch leading to lesser ubiquitination (Melino et al., 2008). A previous study showed an inhibitory effect of AMPK activator on interaction of Itch with its substrate p73 leading to lesser degradation of the latter (Adamovich et al., 2014). However, the mechanisms that lead to AMPK-dependent alteration of Itch activity was not explored. We show that AMPK activation does not alter Itch levels, but it impairs Itch-Notch interaction by the regulation of Itch tyrosine phosphorylation by Fyn **(Figure 8D)**. We also show that AMPK promotes Fyn activation by altering its interaction with tyrosine phosphatases that regulate its activity. This is in keeping with a report showing LKB1, an upstream kinase of AMPK, promotes protein tyrosine phosphatase activity leading to inhibition of receptor tyrosine kinases in lung and cervical cancer cell lines (Okon et al., 2014). Likewise, we recently reported AMPK-mediated regulation of serine threonine kinase Akt activity by modulating its phosphatase PHLPP2 in breast cancer cells (Saha et al., 2018). Thus, AMPK might alter the activity of a vast number of other kinases by affecting their interaction with or activity of cellular phosphatases. These observations warrant a detailed investigation into AMPK-mediated regulation of signal transduction pathways by regulating cellular phosphatases.

AMPK is traditionally viewed as a tumor suppressor as it negatively regulates mTOR, thus

bringing about slower growth (Gwinn et al., 2008). AMPK is also known to stabilize p53

bringing about cell cycle arrest (Zadra et al., 2015). Both mTOR and p53 pathways have been

shown to cross talk with Notch. mTOR has been reported to regulate Notch1 transcription

(Bhola et al., 2016), but we failed to see changes in Notch1 transcript levels upon AMPK activation.

While p53 was shown to positively regulate Notch1 in epithelial cells (Yugawa et al., 2007),

we have tested the hypoxia-AMPK-Notch signaling axis in p53 mutant cell line (MDA-MB-

231) indicating that p53 is not involved in mediating the observed effects of AMPK on

cleaved Notch1 levels. We show a pro-tumorigenic role of AMPK through positive

regulation of Notch1 stability leading to elevated breast cancer stemness and drug resistance

phenotype **(Figure 8D)**. Interestingly, our study shows a high correlation between AMPK

and Notch activation in grade 3 primary breast cancer samples that are known to contain

hypoxic regions (Zhang et al., 2016). In support of our observations, abundant expression of

activated AMPK has also been reported in human gliomas (Ríos et al., 2013) and breast

cancer (Hart et al., 2015a). Contextual oncogenic properties of AMPK has been reported,

especially in bioenergetic stress conditions such as nutrient deprivation and hypoxia, which

are often faced by cells of a growing tumor (Kishton et al., 2016; Liang and Mills, 2013).

While we have identified a role for AMPK in increasing breast cancer stemness and drug

resistance in hypoxia, previously, a protective role for AMPK has been demonstrated in

androgen-dependent prostate cancer cells in hypoxia (Chhipa et al., 2011) and in other solid

tumors with low oxygen condition (Laderoute et al., 2006). Further, studies from our lab

(Hindupur et al., 2014) and others (Jeon et al., 2012) show in-vivo evidence of tumor

promoting role of AMPK in tumor xenograft models. Another study showed that this role of

AMPK is evident only in a metabolically stressed tumor microenvironment context

(Laderoute et al., 2014). Thus, our report compliments such studies on tumor promoting role

of AMPK in the context of tumor hypoxia.

We have recently reported the role of AMPK in supporting EMT and metastasis in hypoxia

(Saxena et al., 2018). An increased metastatic potential is associated with drug resistance.

Consistent with this, recent studies have implicated AMPK in mediating hypoxia-induced

drug resistance to doxorubicin in osteosarcoma (Zhao et al., 2016). Other studies have

reported AMPK in resistance to cisplatin (Harhaji☐Trajkovic et al., 2009; Shin et al., 2014).

In this study, we show that hypoxia-induced drug resistance phenotype mediated by Notch

signaling in breast cancer cells is supported by AMPK activation.

Our study identifies a novel protective role for AMPK in Itch-mediated Notch degradation.

We further show that AMPK plays an important role in maintaining the stemness and drug

resistance phenotypes induced by hypoxia. Thus, under the stress of hypoxia which is

prevalent in solid tumors, we propose that AMPK inhibition-based strategies can alleviate the

cancer stem cell phenotype induced by hypoxia, thereby rendering cancer cells more

susceptible to existing anti-cancer agents.

Acknowledgement

Funding: This work was majorly supported by grants from the Wellcome Trust-DBT India

Alliance (IA) Senior Research Fellowship (500112/Z/09/Z) to AR. ML acknowledges

Council for Scientific and Industrial Research for CSIR fellowship (19-12/2010(i) EU-IV).

MKJ acknowledges support from Ramanujan Fellowship provided by SERB, DST, Govt. of

India (SB/S2/RJN-049/2019). The authors acknowledge grants from DBT-IISc partnership

programme, support from DST-FIST and UGC, Govt. of India to the Department of MRDG,

IISc.

The authors thank Dr. Benoit Viollet for DKO MEFs and Sai Balaji for help with xenograft

assay. The authors acknowledge the Central Animal Facility and FACS facility at IISc.

Competing interests: The authors declare that they have no competing interests.

Authors' contributions

ML and AR conceived and designed experiments; ML performed majority of the experiments

and analyzed data; SK performed some experiments, and helped with bioinformatics data

analysis. AC performed correlation analyses from TCGA data set. KH and MKJ devised the

mathematical model. ML and AR edited and drafted the manuscript. AR supervised the

study. All authors have read and approved the final version of the manuscript.

Declaration:

Ethics approval and consent to participate: Primary breast tissues (cancer and adjacent

normal) obtained from Kidwai Memorial Institute of Oncology (KMIO), Bangalore, as per

IRB and in compliance with ethical guidelines of KMIO and the Indian Institute of Science

(IISc). All animal experiments were reviewed and approved by the Institutional Animal

Ethics committee of IISc, Bangalore.

Availability of data and material: All data generated or analysed during this study are

included in this published article [and its supplementary information files].

References

Adamovich, Y., Adler, J., Meltser, V., Reuven, N., and Shaul, Y. (2014). AMPK couples p73 with p53 in
cell fate decision. Cell Death & Differentiation *21*, 1451-1459.
Allenspach, E.J., Maillard, I., Aster, J.C., and Pear, W.S. (2002). Notch signaling in cancer. Cancer
biology & therapy *1*, 466-476.
Andersson, E.R., and Lendahl, U. (2014). Therapeutic modulation of Notch signalling—are we there
yet? Nature reviews Drug discovery *13*, 357-378.
Andersson, E.R., Sandberg, R., and Lendahl, U. (2011). Notch signaling: simplicity in design, versatility
in function. Development *138*, 3593-3612.
Artavanis-Tsakonas, S., Rand, M.D., and Lake, R.J. (1999). Notch signaling: cell fate control and signal
integration in development. Science *284*, 770-776.
Aster, J.C., Pear, W.S., and Blacklow, S.C. (2008). Notch signaling in leukemia. Annu Rev Pathol Mech
Dis *3*, 587-613.
Bedogni, B., Warneke, J.A., Nickoloff, B.J., Giaccia, A.J., and Powell, M.B. (2008). Notch1 is an effector
of Akt and hypoxia in melanoma development. The Journal of clinical investigation *118*, 3660-3670.
Bhola, N.E., Jansen, V.M., Koch, J.P., Li, H., Formisano, L., Williams, J.A., Grandis, J.R., and Arteaga,
C.L. (2016). Treatment of triple-negative breast cancer with TORC1/2 inhibitors sustains a drug-
resistant and notch-dependent cancer stem cell population. Cancer research *76*, 440-452.
Blokzijl, A., Dahlqvist, C., Reissmann, E., Falk, A., Moliner, A., Lendahl, U., and Ibáñez, C.F. (2003).
Cross-talk between the Notch and TGF-β signaling pathways mediated by interaction of the Notch
intracellular domain with Smad3. The Journal of cell biology *163*, 723-728.
Bray, S.J. (2006). Notch signalling: a simple pathway becomes complex. Nature reviews Molecular
cell biology *7*, 678-689.
Brou, C., Logeat, F., Gupta, N., Bessia, C., LeBail, O., Doedens, J.R., Cumano, A., Roux, P., Black, R.A.,
and Israël, A. (2000). A novel proteolytic cleavage involved in Notch signaling: the role of the
disintegrin-metalloprotease TACE. Molecular cell *5*, 207-216.
Chen, J., Imanaka, N., and Griffin, J. (2010). Hypoxia potentiates Notch signaling in breast cancer
leading to decreased E-cadherin expression and increased cell migration and invasion. British Journal
of Cancer *102*, 351-360.
Chen, R., Jiang, T., She, Y., Xu, J., Li, C., Zhou, S., Shen, H., Shi, H., and Liu, S. (2017). Effects of cobalt
chloride, a hypoxia-mimetic agent, on autophagy and atrophy in skeletal C2C12 myotubes. BioMed
research international *2017*.
Chen, Y., De Marco, M.A., Graziani, I., Gazdar, A.F., Strack, P.R., Miele, L., and Bocchetta, M. (2007).
Oxygen concentration determines the biological effects of NOTCH-1 signaling in adenocarcinoma of
the lung. Cancer research *67*, 7954-7959.
Chhipa, R.R., Wu, Y., and Ip, C. (2011). AMPK-mediated autophagy is a survival mechanism in
androgen-dependent prostate cancer cells subjected to androgen deprivation and hypoxia. Cellular
signalling *23*, 1466-1472.
Cool, B., Zinker, B., Chiou, W., Kifle, L., Cao, N., Perham, M., Dickinson, R., Adler, A., Gagne, G., and
Iyengar, R. (2006). Identification and characterization of a small molecule AMPK activator that treats
key components of type 2 diabetes and the metabolic syndrome. Cell metabolism *3*, 403-416.
D'Angelo, R.C., Ouzounova, M., Davis, A., Choi, D., Tchuenkam, S.M., Kim, G., Luther, T., Quraishi,
A.A., Senbabaoglu, Y., and Conley, S.J. (2015). Notch reporter activity in breast cancer cell lines
identifies a subset of cells with stem cell activity. Molecular cancer therapeutics *14*, 779-787.
Elenbaas, B., Spirio, L., Koerner, F., Fleming, M.D., Zimonjic, D.B., Donaher, J.L., Popescu, N.C., Hahn,
W.C., and Weinberg, R.A. (2001). Human breast cancer cells generated by oncogenic transformation
of primary mammary epithelial cells. Genes & development *15*, 50-65.
Emerling, B.M., Weinberg, F., Snyder, C., Burgess, Z., Mutlu, G.M., Viollet, B., Budinger, G.S., and
Chandel, N.S. (2009). Hypoxic activation of AMPK is dependent on mitochondrial ROS but
independent of an increase in AMP/ATP ratio. Free Radical Biology and Medicine *46*, 1386-1391.
Fryer, C.J., White, J.B., and Jones, K.A. (2004). Mastermind recruits CycC: CDK8 to phosphorylate the
Notch ICD and coordinate activation with turnover. Molecular cell *16*, 509-520.

Gordon, W.R., Arnett, K.L., and Blacklow, S.C. (2008). The molecular logic of Notch signaling–a
structural and biochemical perspective. Journal of cell science *121*, 3109-3119.
Gruber, G., Greiner, R.H., Hlushchuk, R., Aebersold, D.M., Altermatt, H.J., Berclaz, G., and Djonov, V.
(2004). Hypoxia-inducible factor 1 alpha in high-risk breast cancer: an independent prognostic
parameter? Breast Cancer Research *6*, 1.
Guo, S., Liu, M., and Gonzalez-Perez, R.R. (2011). Role of Notch and its oncogenic signaling crosstalk
in breast cancer. Biochimica et Biophysica Acta (BBA)-Reviews on Cancer *1815*, 197-213.
Gupta-Rossi, N., Le Bail, O., Brou, C., Logeat, F., Six, E., and Israël, A. (2002). Control of Notch Activity
by the Ubiquitin-Proteasome Pathway. In Notch from Neurodevelopment to Neurodegeneration:
Keeping the Fate (Springer), pp. 41-58.
Gusarova, G.A., Trejo, H.E., Dada, L.A., Briva, A., Welch, L.C., Hamanaka, R.B., Mutlu, G.M., Chandel,
N.S., Prakriya, M., and Sznajder, J.I. (2011). Hypoxia leads to Na, K-ATPase downregulation via Ca2+
release-activated Ca2+ channels and AMPK activation. Molecular and cellular biology *31*, 3546-3556.
Gustafsson, M.V., Zheng, X., Pereira, T., Gradin, K., Jin, S., Lundkvist, J., Ruas, J.L., Poellinger, L.,
Lendahl, U., and Bondesson, M. (2005). Hypoxia requires notch signaling to maintain the
undifferentiated cell state. Developmental cell *9*, 617-628.
Gwinn, D.M., Shackelford, D.B., Egan, D.F., Mihaylova, M.M., Mery, A., Vasquez, D.S., Turk, B.E., and
Shaw, R.J. (2008). AMPK phosphorylation of raptor mediates a metabolic checkpoint. Molecular cell
30, 214-226.
Hamilton, S.R., Stapleton, D., O'Donnell, J.B., Kung, J.T., Dalal, S.R., Kemp, B.E., and Witters, L.A.
(2001). An activating mutation in the γ1 subunit of the AMP-activated protein kinase. FEBS letters
500, 163-168.
Hardie, D.G. (2013). AMPK: a target for drugs and natural products with effects on both diabetes and
cancer. Diabetes *62*, 2164-2172.
Hardie, D.G., and Alessi, D.R. (2013). LKB1 and AMPK and the cancer-metabolism link-ten years after.
BMC biology *11*, 36.
Hardie, D.G., Ross, F.A., and Hawley, S.A. (2012). AMPK: a nutrient and energy sensor that maintains
energy homeostasis. Nature reviews Molecular cell biology *13*, 251-262.
Harhaji-Trajkovic, L., Vilimanovich, U., Kravic-Stevovic, T., Bumbasirevic, V., and Trajkovic, V. (2009).
AMPK-mediated autophagy inhibits apoptosis in cisplatin-treated tumour cells. Journal of cellular
and molecular medicine *13*, 3644-3654.
Hart, P., Mao, M., Ansenberger-Fricano, K., Ekoue, D., Ganini, D., Kajdacsy-Balla, A., Diamond, A.,
Minshall, R., Consolaro, M., and Santos, J. (2015a). MnSOD upregulation sustains the Warburg effect
via mitochondrial ROS and AMPK-dependent signalling in cancer. Nature communications *6*, 6053-
6053.
Hart, P.C., Mao, M., De Abreu, A.L.P., Ansenberger-Fricano, K., Ekoue, D.N., Ganini, D., Kajdacsy-
Balla, A., Diamond, A.M., Minshall, R.D., and Consolaro, M.E. (2015b). MnSOD upregulation sustains
the Warburg effect via mitochondrial ROS and AMPK-dependent signalling in cancer. Nature
communications *6*, 6053.
Hindupur, S.K., Balaji, S.A., Saxena, M., Pandey, S., Sravan, G.S., Heda, N., Kumar, M.V., Mukherjee,
G., Dey, D., and Rangarajan, A. (2014). Identification of a novel AMPK-PEA15 axis in the anoikis-
resistant growth of mammary cells. Breast Cancer Research *16*, 1.
Jarriault, S., Brou, C., Logeat, F., Schroeter, E.H., Kopan, R., and Israel, A. (1995). Signalling
downstream of activated mammalian Notch. Nature *377*, 355-358.
Jeon, S.-M., Chandel, N.S., and Hay, N. (2012). AMPK regulates NADPH homeostasis to promote
tumour cell survival during energy stress. Nature *485*, 661-665.
Jeon, S.-M., and Hay, N. (2012). The dark face of AMPK as an essential tumor promoter. Cellular
logistics *2*, 197-202.
Jeon, S.-M., and Hay, N. (2015). The double-edged sword of AMPK signaling in cancer and its
therapeutic implications. Archives of pharmacal research *38*, 346-357.

Jubb, A., Turley, H., Moeller, H., Steers, G., Han, C., Li, J., Leek, R., Tan, E., Singh, B., and Mortensen,
N. (2009). Expression of delta-like ligand 4 (Dll4) and markers of hypoxia in colon cancer. British
journal of cancer *101*, 1749-1757.
Kim, J.-Y., and Lee, J.-Y. (2017). Targeting tumor adaption to chronic hypoxia: implications for drug
resistance, and how it can be overcome. International journal of molecular sciences *18*, 1854.
Kishton, R.J., Barnes, C.E., Nichols, A.G., Cohen, S., Gerriets, V.A., Siska, P.J., Macintyre, A.N.,
Goraksha-Hicks, P., De Cubas, A.A., and Liu, T. (2016). AMPK is essential to balance glycolysis and
mitochondrial metabolism to control T-ALL cell stress and survival. Cell metabolism *23*, 649-662.
Kunz, M., and Ibrahim, S.M. (2003). Molecular responses to hypoxia in tumor cells. Molecular cancer
2, 23.
Laderoute, K.R., Amin, K., Calaoagan, J.M., Knapp, M., Le, T., Orduna, J., Foretz, M., and Viollet, B.
(2006). 5′-AMP-activated protein kinase (AMPK) is induced by low-oxygen and glucose deprivation
conditions found in solid-tumor microenvironments. Molecular and cellular biology *26*, 5336-5347.
Laderoute, K.R., Calaoagan, J.M., Chao, W.-r., Dinh, D., Denko, N., Duellman, S., Kalra, J., Liu, X.,
Papandreou, I., and Sambucetti, L. (2014). 5′-AMP-activated protein kinase (AMPK) supports the
growth of aggressive experimental human breast cancer tumors. Journal of Biological Chemistry *289*,
22850-22864.
Liang, J., and Mills, G.B. (2013). AMPK: a contextual oncogene or tumor suppressor? Cancer research
73, 2929-2935.
Marin, J.J., Lozano, E., and Perez, M.J. (2016). Lack of mitochondrial DNA impairs chemical hypoxia-
induced autophagy in liver tumor cells through ROS-AMPK-ULK1 signaling dysregulation
independently of HIF-1α. Free Radical Biology and Medicine *101*, 71-84.
McGill, M.A., and McGlade, C.J. (2003). Mammalian numb proteins promote Notch1 receptor
ubiquitination and degradation of the Notch1 intracellular domain. Journal of Biological Chemistry
278, 23196-23203.
Melino, G., Gallagher, E., Aqeilan, R., Knight, R., Peschiaroli, A., Rossi, M., Scialpi, F., Malatesta, M.,
Zocchi, L., and Browne, G. (2008). Itch: a HECT-type E3 ligase regulating immunity, skin and cancer.
Cell Death & Differentiation *15*, 1103-1112.
Mihaylova, M.M., and Shaw, R.J. (2011). The AMPK signalling pathway coordinates cell growth,
autophagy and metabolism. Nature cell biology *13*, 1016-1023.
Mittal, S., Subramanyam, D., Dey, D., Kumar, R.V., and Rangarajan, A. (2009). Cooperation of Notch
and Ras/MAPK signaling pathways in human breast carcinogenesis. Molecular cancer *8*, 1.
Morata-Tarifa, C., Jiménez, G., García, M.A., Entrena, J.M., Griñán-Lisón, C., Aguilera, M., Picon-Ruiz,
M., and Marchal, J.A. (2016). Low adherent cancer cell subpopulations are enriched in tumorigenic
and metastatic epithelial-to-mesenchymal transition-induced cancer stem-like cells. Scientific
reports *6*, 18772.
Moretti, J., and Brou, C. (2013). Ubiquitinations in the notch signaling pathway. International journal
of molecular sciences *14*, 6359-6381.
Mungai, P.T., Waypa, G.B., Jairaman, A., Prakriya, M., Dokic, D., Ball, M.K., and Schumacker, P.T.
(2011). Hypoxia triggers AMPK activation through reactive oxygen species-mediated activation of
calcium release-activated calcium channels. Molecular and cellular biology *31*, 3531-3545.
Nguyen, T.-H., Liu, J., and Lombroso, P.J. (2002). Striatal enriched phosphatase 61 dephosphorylates
Fyn at phosphotyrosine 420. Journal of Biological Chemistry *277*, 24274-24279.
Nickoloff, B., Qin, J., Chaturvedi, V., Denning, M., Bonish, B., and Miele, L. (2002). Jagged-1 mediated
activation of notch signaling induces complete maturation of human keratinocytes through NF-κB
and PPARγ. Cell Death & Differentiation *9*, 842-855.
Öberg, C., Li, J., Pauley, A., Wolf, E., Gurney, M., and Lendahl, U. (2001). The Notch intracellular
domain is ubiquitinated and negatively regulated by the mammalian Sel-10 homolog. Journal of
Biological Chemistry *276*, 35847-35853.

Okon, I.S., Coughlan, K.A., and Zou, M.-H. (2014). Liver kinase B1 expression promotes phosphatase
activity and abrogation of receptor tyrosine kinase phosphorylation in human cancer cells. Journal of
Biological Chemistry *289*, 1639-1648.
Parsons, S.J., and Parsons, J.T. (2004). Src family kinases, key regulators of signal transduction.
Oncogene *23*, 7906.
Pece, S., Serresi, M., Santolini, E., Capra, M., Hulleman, E., Galimberti, V., Zurrida, S., Maisonneuve,
P., Viale, G., and Di Fiore, P.P. (2004). Loss of negative regulation by Numb over Notch is relevant to
human breast carcinogenesis. The Journal of cell biology *167*, 215-221.
Popko-Scibor, A.E., Lindberg, M.J., Hansson, M.L., Holmlund, T., and Wallberg, A.E. (2011).
Ubiquitination of Notch1 is regulated by MAML1-mediated p300 acetylation of Notch1. Biochemical
and biophysical research communications *416*, 300-306.
Prieto-Vila, M., Takahashi, R.-u., Usuba, W., Kohama, I., and Ochiya, T. (2017). Drug resistance driven
by cancer stem cells and their niche. International journal of molecular sciences *18*, 2574.
Qiang, L., Wu, T., Zhang, H., Lu, N., Hu, R., Wang, Y., Zhao, L., Chen, F., Wang, X., and You, Q. (2012).
HIF-1α is critical for hypoxia-mediated maintenance of glioblastoma stem cells by activating Notch
signaling pathway. Cell Death & Differentiation *19*, 284-294.
Qiu, L., Joazeiro, C., Fang, N., Wang, H.-Y., Elly, C., Altman, Y., Fang, D., Hunter, T., and Liu, Y.-C.
(2000). Recognition and ubiquitination of Notch by Itch, a hect-type E3 ubiquitin ligase. Journal of
Biological Chemistry *275*, 35734-35737.
Radtke, F., and Raj, K. (2003). The role of Notch in tumorigenesis: oncogene or tumour suppressor?
Nat Rev Cancer *3*, 756-767.
Reedijk, M., Odorcic, S., Chang, L., Zhang, H., Miller, N., McCready, D.R., Lockwood, G., and Egan, S.E.
(2005). High-level coexpression of JAG1 and NOTCH1 is observed in human breast cancer and is
associated with poor overall survival. Cancer research *65*, 8530-8537.
Ríos, M., Foretz, M., Viollet, B., Prieto, A., Fraga, M., Costoya, J.A., and Señarís, R. (2013). AMPK
activation by oncogenesis is required to maintain cancer cell proliferation in astrocytic tumors.
Cancer research *73*, 2628-2638.
Roskoski Jr, R. (2015). Src protein-tyrosine kinase structure, mechanism, and small molecule
inhibitors. Pharmacological research *94*, 9-25.
Saha, M., Kumar, S., Bukhari, S., Balaji, S.A., Kumar, P., Hindupur, S.K., and Rangarajan, A. (2018).
AMPK–Akt double-negative feedback loop in breast cancer cells regulates their adaptation to matrix
deprivation. Cancer research *78*, 1497-1510.
Sahlgren, C., Gustafsson, M.V., Jin, S., Poellinger, L., and Lendahl, U. (2008). Notch signaling mediates
hypoxia-induced tumor cell migration and invasion. Proceedings of the National Academy of
Sciences *105*, 6392-6397.
Samanta, D., Park, Y., Andrabi, S.A., Shelton, L.M., Gilkes, D.M., and Semenza, G.L. (2016). PHGDH
expression is required for mitochondrial redox homeostasis, breast cancer stem cell maintenance,
and lung metastasis. Cancer research *76*, 4430-4442.
Sansone, P., Storci, G., Giovannini, C., Pandolfi, S., Pianetti, S., Taffurelli, M., Santini, D., Ceccarelli, C.,
Chieco, P., and Bonafe, M. (2007). p66Shc/Notch-3 Interplay Controls Self-Renewal and Hypoxia
Survival in Human Stem/Progenitor Cells of the Mammary Gland Expanded In Vitro as
Mammospheres. Stem Cells *25*, 807-815.
Saxena, K., and Jolly, M.K. (2019). Acute vs. chronic vs. cyclic hypoxia: Their differential dynamics,
molecular mechanisms, and effects on tumor progression. Biomolecules *9*, 339.
Saxena, M., Balaji, S.A., Deshpande, N., Ranganathan, S., Pillai, D.M., Hindupur, S.K., and Rangarajan,
A. (2018). AMP-activated protein kinase promotes epithelial-mesenchymal transition in cancer cells
through Twist1 upregulation. Journal of cell science *131*, jcs208314.
Schaffer, B.E., Levin, R.S., Hertz, N.T., Maures, T.J., Schoof, M.L., Hollstein, P.E., Benayoun, B.A.,
Banko, M.R., Shaw, R.J., and Shokat, K.M. (2015). Identification of AMPK phosphorylation sites
reveals a network of proteins involved in cell invasion and facilitates large-scale substrate prediction.
Cell metabolism *22*, 907-921.

Schroeter, E.H., Kisslinger, J.A., and Kopan, R. (1998). Notch-1 signalling requires ligand-induced
proteolytic release of intracellular domain. Nature *393*, 382-386.
Scialpi, F., Malatesta, M., Peschiaroli, A., Rossi, M., Melino, G., and Bernassola, F. (2008). Itch self-
polyubiquitylation occurs through lysine-63 linkages. Biochemical pharmacology *76*, 1515-1521.
Seo, E.J., Kim, D.K., Jang, I.H., Choi, E.J., Shin, S.H., Lee, S.I., Kwon, S.-M., Kim, K.-H., Suh, D.-S., and
Kim, J.H. (2016). Hypoxia-NOTCH1-SOX2 signaling is important for maintaining cancer stem cells in
ovarian cancer. Oncotarget *7*, 55624.
Shin, D.H., Choi, Y.-J., and Park, J.-W. (2014). SIRT1 and AMPK mediate hypoxia-induced resistance of
non–small cell lung cancers to cisplatin and doxorubicin. Cancer research *74*, 298-308.
Singh, P., Satriano, J., Li, H., and Hallows, K. (2013). HIF-1alpha and AMPK interactions regulate
cellular hypoxia adaptation in the subtotal nephrectomy model of CKD (Federation of American
Societies for Experimental Biology).
Stylianou, S., Clarke, R.B., and Brennan, K. (2006). Aberrant activation of notch signaling in human
breast cancer. Cancer research *66*, 1517-1525.
Sundararaman, A., Amirtham, U., and Rangarajan, A. (2016). Calcium-oxidant signaling network
regulates AMP-activated protein kinase (AMPK) activation upon matrix deprivation. Journal of
Biological Chemistry *291*, 14410-14429.
Vacaresse, N., Møller, B., Danielsen, E.M., Okada, M., and Sap, J. (2008). Activation of c-Src and Fyn
kinases by protein-tyrosine phosphatase RPTPα is substrate-specific and compatible with lipid raft
localization. Journal of Biological Chemistry *283*, 35815-35824.
Vaupel, P., Briest, S., and Höckel, M. (2002). Hypoxia in breast cancer: pathogenesis, characterization
and biological/therapeutic implications. Wiener Medizinische Wochenschrift *152*, 334-342.
Wang, Z., Li, Y., Ahmad, A., Azmi, A.S., Banerjee, S., Kong, D., and Sarkar, F.H. (2010). Targeting Notch
signaling pathway to overcome drug resistance for cancer therapy. Biochimica et Biophysica Acta
(BBA)-Reviews on Cancer *1806*, 258-267.
Wenger, R.H., and Gassmann, M. (1997). Oxygen (es) and the hypoxia-inducible factor-1. Biological
chemistry *378*, 609-616.
Winder, W., and Hardie, D. (1996). Inactivation of acetyl-CoA carboxylase and activation of AMP-
activated protein kinase in muscle during exercise. American Journal of Physiology-Endocrinology
And Metabolism *270*, E299-E304.
Wu, G., Lyapina, S., Das, I., Li, J., Gurney, M., Pauley, A., Chui, I., Deshaies, R.J., and Kitajewski, J.
(2001). SEL-10 is an inhibitor of notch signaling that targets notch for ubiquitin-mediated protein
degradation. Molecular and cellular biology *21*, 7403-7415.
Xi, L., Taher, M., Yin, C., Salloum, F., and Kukreja, R.C. (2004). Cobalt chloride induces delayed cardiac
preconditioning in mice through selective activation of HIF-1α and AP-1 and iNOS signaling.
American Journal of Physiology-Heart and Circulatory Physiology *287*, H2369-H2375.
Xiang, L., Gilkes, D.M., Hu, H., Takano, N., Luo, W., Lu, H., Bullen, J.W., Samanta, D., Liang, H., and
Semenza, G.L. (2014). Hypoxia-inducible factor 1 mediates TAZ expression and nuclear localization to
induce the breast cancer stem cell phenotype. Oncotarget *5*, 12509.
Xing, F., Okuda, H., Watabe, M., Kobayashi, A., Pai, S.K., Liu, W., Pandey, P.R., Fukuda, K., Hirota, S.,
and Sugai, T. (2011). Hypoxia-induced Jagged2 promotes breast cancer metastasis and self-renewal
of cancer stem-like cells. Oncogene *30*, 4075-4086.
Yang, C., Zhou, W., Jeon, M.-s., Demydenko, D., Harada, Y., Zhou, H., and Liu, Y.-C. (2006). Negative
regulation of the E3 ubiquitin ligase itch via Fyn-mediated tyrosine phosphorylation. Molecular cell
21, 135-141.
Yugawa, T., Handa, K., Narisawa-Saito, M., Ohno, S.-i., Fujita, M., and Kiyono, T. (2007). Regulation of
Notch1 gene expression by p53 in epithelial cells. Molecular and cellular biology *27*, 3732-3742.
Zadra, G., Batista, J.L., and Loda, M. (2015). Dissecting the dual role of AMPK in cancer: from
experimental to human studies. Molecular cancer research *13*, 1059-1072.
Zhang, C., Samanta, D., Lu, H., Bullen, J.W., Zhang, H., Chen, I., He, X., and Semenza, G.L. (2016).
Hypoxia induces the breast cancer stem cell phenotype by HIF-dependent and ALKBH5-mediated

m6A-demethylation of NANOG mRNA. Proceedings of the National Academy of Sciences *113*, E2047-
E2056.

Zhang, H., Lu, H., Xiang, L., Bullen, J.W., Zhang, C., Samanta, D., Gilkes, D.M., He, J., and Semenza,
G.L. (2015). HIF-1 regulates CD47 expression in breast cancer cells to promote evasion of
phagocytosis and maintenance of cancer stem cells. Proceedings of the National Academy of
Sciences *112*, E6215-E6223.

Zhao, C., Zhang, Q., Yu, T., Sun, S., Wang, W., and Liu, G. (2016). Hypoxia promotes drug resistance in
osteosarcoma cells via activating AMP-activated protein kinase (AMPK) signaling. Journal of bone
oncology *5*, 22-29.

Zheng, X., Linke, S., Dias, J.M., Zheng, X., Gradin, K., Wallis, T.P., Hamilton, B.R., Gustafsson, M., Ruas,
J.L., and Wilkins, S. (2008). Interaction with factor inhibiting HIF-1 defines an additional mode of
cross-coupling between the Notch and hypoxia signaling pathways. Proceedings of the National
Academy of Sciences *105*, 3368-3373.

Zhou, G., Myers, R., Li, Y., Chen, Y., Shen, X., Fenyk-Melody, J., Wu, M., Ventre, J., Doebber, T., and
Fujii, N. (2001). Role of AMP-activated protein kinase in mechanism of metformin action. The Journal
of clinical investigation *108*, 1167-1174.

Figure legends

Figure 1: AMPK regulates cleaved Notch1 levels in breast cancer cells:

(A) MDA-MB-231 cells were cultured in the presence of A769662 (A76; 100 µM) or DMSO

as control and immunoblotted for indicated proteins. α Tubulin was used as loading control

(n=3). Graphs in all immunoblotting experiments represent the densitometric analysis for

quantification of relative amount of indicated protein and error bars represent ±SEM. (B)

MDA-MB-231 cells transfected with constitutively active gamma subunit of AMPK (γ

R70Q) or vector control and immunoblotted for indicated proteins (n=3). (C) MDA-MB-231

cells were treated with Compound C (Comp C; 10 µM) or DMSO as control for 24 h and

immunoblotted for indicated proteins (n=6). (D) MDA-MB-231 cells stably expressing

shAMPK α2 (#4) were induced with doxycycline (5µg/µl) for 48 h and immunoblotted for

indicated proteins (n=4). (E) Immortalized wild type MEF and AMPKα1/2 DKO MEFs

(DKO) were immunoblotted for indicated proteins (n=4). (F) MDA-MB-231 cells were

cultured in the presence of A76 (100 µM) or DMSO as control for 24 h and subjected to

immunoblotting for indicated protein (n=3). (G) MDA-MB-231 cells were transfected with

12xCSL-Luciferase (FFL) and Renilla TK (RL) and treated with A76 (100 µM) or DMSO as

control for 48 h. Luciferase activity is represented as a ratio of Firefly (FFL) to Renila (RL)

luciferase (n=4). (H) MDA-MB-231 cells were treated with A76 (100 µM) or DMSO as

control for 48 h and RT-PCR analysis was carried out for *HES*; *GAPDH* served as

housekeeping gene. Graph represents the fold change in *HES1* transcript levels normalized to

GAPDH (n=4).

Figure 2: AMPK inhibition or knockdown impairs hypoxia-induced cleaved Notch1

generation

(A) MDA-MB-231 cells were subjected to $CoCl_2$ (150 µM) treatment for 24 h and

immunoblotted for indicated protein; lanes from the same run were assembled together.

Graphs in all immunoblots represent the densitometric analysis for quantification of relative

amount of indicated protein (n=4). (B) MDA-MB-231 cells were subjected to $CoCl_2$ (150

µM) with DMSO or Comp C (10 µM) treatment as indicated for 24 h and immunoblotted for

indicated proteins (n=4). (C) MDA-MB-231 cells stably expressing shAMPK α2 (#1) and

vector control pTRIPZ were induced with doxycycline (5µg/µl) for 24 h followed by

treatment with $CoCl_2$ (150 µM) and immunoblotted for indicated proteins (n=3). (D) MDA-

MB-231 cells were grown in 21% or 3% O_2 for 24 h and harvested for immunoblotting for

indicated proteins. (E) MDA-MB-231 cells were grown in 21% or in 3% O_2 in the presence

or absence of Comp C (10 µM) and immunoblotted for indicated proteins (n=3). (F) MDA-

MB-231 cells stably expressing shAMPK α2 (#4) were induced with doxycycline (5µg/µl)

for 24 h and subsequently grown in 3% O_2 and immunoblotted for indicated proteins (n=3).

(G) MDA-MB-231 cells stably expressing shRNA against AMPK α2 (#4) were induced with

doxycycline (5µg/µl) for 24 h and subsequently grown in 3% O_2 for 24 h and were

immunostained for valine 1744 cleavage-specific NICD protein (Scale bar: 50 µm) (n=3).

Graph represents the mean intensity of valine 1744 specific NICD protein expression in 3

independent experiments. (H) HEK-293T cells were transfected with 12xCSL-Luciferase

(FFL) and Renilla TK (RL) and NICD expression construct. The cells were treated with

CoCl$_2$ for 48 h with DMSO or Compound C. Luciferase activity is represented as a ratio of

Firefly (FFL) to Renila (RL) luciferase (n=3). Error bars represent ±SEM in all graphs.

Figure 3: AMPK inhibition affects stability of cleaved Notch1 under hypoxia:

(A-B) MDA-MB-231 cells stably expressing shAMPK α2 (#4) were induced by doxycycline

(5µg/µl) for 48 h and (A) subsequently treated with cycloheximide for 4 h and 8 h and

immunoblotted for indicated proteins; and graphs represent the densitometric analysis for

quantification of relative amount of indicated protein and (B) grown in 3% O$_2$ for another 24

h followed by cycloheximide treatment for 8 h and immunoblotted for indicated; (n=3 each).

(C) MDA-MB-231 cells were treated with Comp C (10 µM) or DMSO control for 24 h and

subsequently treated with or without MG-132 (5 µM) for 8 h. Cells were immunoblotted for

indicated proteins (n=3); Error bars represent ±SEM in all graphs.

Figure 4: AMPK inhibits Notch1 ubiquitination by modulating its interaction with Itch:

(A) MDA-MB-231 cells were cultured in the presence of A76 (100 µM) or DMSO control

for 24 h and subsequently treated with MG132 (5 µM) for 2 h prior to Notch1 immuno-

precipitation using the C-terminus specific Notch1 antibody followed by immunoblotting for

anti-K-48 ubiquitin linkage specific antibody (n=3). (B) MDA-MB-231 cells were cultured in

the presence of A76 (100 µM) or DMSO control for 24 h and subsequently immunoblotted

for Itch protein (n=3). (C) MDA-MB-231 cells were cultured in the presence of A76 (100

µM) or DMSO control for 24 h and subsequently treated with MG132 (5 µM) for 2 h prior to

Itch immuno-precipitation and immunoblotted for pan-ubiquitin. (D) MDA-MB-231 cells

were cultured in the presence of A76 (100 µM) or DMSO control for 24 h and subsequently

treated with MG132 (5 µM) for 2 h prior to Notch1 immuno-precipitation and immunoblotted

for Itch protein (n=3). (E) MDA-MB-231 cells were transfected with vector control or Myc

tagged Itch plasmid. Cells were cultured in the presence of A76 (100 µM) or DMSO control

and immunoblotted for indicated proteins. Graph represents the densitometric analysis for

quantification of relative amount of indicated protein (n=3). (F) MDA-MB-231 cells grown

in 21% O_2 or 3% O_2 were immunoblotted for indicated proteins (n=3). (G) MDA-MB-231

cells stably expressing shAMPK α2 (#1) were grown for 24 h in 21% O_2 or 3% O_2 with and

without induction by doxycycline (5µg/µl) (as indicated) and Itch was immuno-precipitated

and immunoblotted for Notch1 protein. Same lysates were used for both the panels.

Immunoprecipitation was carried out separately for the 2 panels and immunoblotted

separately. (n=3) (H) MDA-MB-231 cells grown in 21% O_2 or 3% O_2 were treated with

DMSO or Comp C (10 µM) for 24 h and with MG132 (5 µM) for 2 h prior to harvesting.

Cleaved Notch1 was immuno-precipitated followed by immunoblotting using K-48 linked

ubiquitin-specific antibodies (n=3). Lysates were prepared separately and

immunoprecipitation and immunoblotting were carried out separately for the 2 panels.

Figure 5: AMPK mediates Itch phosphorylation in hypoxia:

(A) MDA-MB-231 cells stably expressing shAMPK α2 (#1) were grown for 24 h in 21% O_2,

or in 3% O_2 with and without induction by doxycycline (5µg/µl) and Itch was immuno-

precipitated and immunoblotted for phospho-tyrosine levels (n=3) (B) MDA-MB-231 cells

stably expressing shAMPK α2 (#1) were grown in 21% O_2 or 3% O2 for 24 h with and

without induction by doxycycline (5µg/µl) and were immunoblotted for phospho-Itch Y420

levels (this blot was performed by reprobing the blot in Figure 2F, and hence the tubulin

panel has been recalled from Figure 2F. (C) MDA-MB-231 cells grown in 21% O_2 or 3% O_2

and were treated with PP2 (10 µM) for 24 h and harvested for immunoblotting of specified

proteins. (D) MDA-MB-231 cells transfected with control vector or dominant negative Fyn

(DN Fyn) were grown in 21% O_2 or 3% O_2 for 24 h and harvested for Itch immuno-

precipitation followed by immunoblotting for phospho-tyrosine levels (n=3). (E) MDA-MB-

231 cells were treated with A76 (100 µM) or DMSO control for 24 h and harvested for

immunoblotting and probed for phospho Src family kinase levels (pSFK) (n=3). (F) MDA-

MB-231 cells stably expressing shRNA against AMPK α2 (#1) with and without induction by

doxycycline (5µg/µl) for 24 h were grown in 3% O_2 for 24 h harvested for immunoblotting

for pSFK levels (n=3).

Figure 6: Hypoxia-induced AMPK activation promotes stem-like properties and drug-

resistance in breast cancer cells through Notch signaling:

(A-B) MDA-MB-231 cells grown in 21% O_2 or 3% O_2 were treated with DMSO or Comp C

(10 µM) for 48 h and (A) immunoblotted for indicated proteins and (B) subjected to sphere

formation assay. The plot represents the average number of day-7 spheres in 5 fields in two

replicates of 3 independent experiments. (C-D) MDA-MB-231 cells stably expressing

shAMPK α2 (#1) were grown in 3% O_2 for 24 h with doxycycline induction and subsequently

treated with and without DSL ligand (100 nM) and (C) immunoblotted for indicated proteins

and (D) subjected to sphere formation assay. The plot represents the average number of day-7

spheres of 3 independent experiments. (E) Bifurcation diagram showing the switch (arrows)

from a differentiated state (lower blue curve) to a more stem-like state (upper blue curve)

upon increase in hypoxia. (F-G) Bifurcation diagrams for BMI1 and OCT4 showing

transitions from a more differentiated state (lower blue curve) to a more stem-like one (upper

blue curve) (H) MDA-MB-231 cells stably expressing inducible shAMPK α2 were treated

with indicated concentrations of doxorubicin for 48 h under normoxic and hypoxic conditions

with and without with doxycycline induced AMPK α2 knockdown (n=3). Cell viability

evaluated by MTT assay and fold change in IC_{50} values have been indicated. (I) MDA-MB-

231 cells stably expressing inducible shAMPK α2 were treated with 1μM of doxorubicin for

24 h under normoxic and hypoxic conditions with and without with doxycycline induced

AMPK α2 knockdown and additional Notch activation by DSL ligand. DNA damage was

assessed by pH2A.X staining. Graphs represent the mean intensity of pH2A.X staining from

3 experiments.

Figure7: Hypoxia-induced AMPK-Notch1 signaling in vivo:

(A) Representative images from immunohistochemical analysis performed on xenograft

tumors generated using BT-474 cells stably expressing scrambled or shAMPKα2 to assess

cleaved Notch1 (Valine 1744) NICD levels (Scale bar: 100 μm). Graphs represent the

staining intensity of cleaved Notch1 (Valine 1744) NICD. (B) Representative images from

immunohistochemical analysis of pACC and cleaved Notch1 (Valine 1744) NICD in breast

cancer patient samples (n=19). (C) Patient derived breast cancer epithelial cells grown in 21%

or 3% O_2 were treated with DMSO or Comp C (10 μM) for 24 h and immunoblotted for

indicated proteins (n=3).

Figure 8: Hypoxia-AMPK-Notch gene signature correlations in TCGA Breast cancer

dataset:

(A-C) Pearson's correlation plots between AMPK-Hypoxia (A), Notch-Hypoxia (B) and

AMPK-Notch (C) analyzed using AMPK, Notch and Hypoxia gene signature in TCGA-

BRCA (all subtypes) dataset; Pearson's correlation value (cor) of each gene pair is

represented as the size of the circle and filled with corresponding color from the color palette

represented below the ranging from -1(red) to +1(blue). Boxes highlighted by the black

squares represent insignificant (p>0.01) Pearson cor values. (D) Proposed model depicting

AMPK and Notch crosstalk under normoxia and hypoxia: In conditions of normoxia, basal

activity of AMK does not interfere with interaction of E3-ubiquitin ligase Itch with its

substrate cleaved Notch1, which is ubiquitinated and targeted for proteasomal degradation. In

conditions of hypoxia, activated AMPK levels enhance Fyn activation, which phosphorylates

and hinders binding of ligase Itch with its substrate cleaved Notch1, thereby resulting in

cleaved Notch1 stabilization. This results in elevated stemness and drug resistance of the

tumors.

Figure 1: AMPK regulates cleaved Notch1 levels in breast cancer cells

Figure 2: AMPK inhibition or knockdown impairs hypoxia-induced cleaved Notch1 generation

Figure 3: AMPK inhibition affects stability of cleaved Notch1 under hypoxia

Figure 4 AMPK inhibits Notch1 ubiquitination by modulating its interaction with Itch

Figure 5 AMPK mediates Itch phosphorylation in hypoxia

Figure 6
Hypoxia induced AMPK activation promotes stem-like properties in breast cancer cells

Figure 7 Hypoxia-induced AMPK-Notch1 signalling in vivo

Figure 8 Hypoxia-AMPK-Notch1 gene signature correlations in TCGA breast cancer dataset