Combined linkage and association mapping of putative QTLs controlling black tea
quality and drought tolerance traits

Robert. K. Koech[1, 2], Richard Mose[3], Samson M. Kamunya[2] and Zeno Apostolides[1]*
[1]Department of Biochemistry, Genetics and Microbiology, University of Pretoria, Pretoria
0002, South Africa
[2]Kenya Agriculture and Livestock Research Organization, Tea Research Institute, P.O. Box
820, Kericho 20200, Kenya
[3]James Finlay (Kenya) Limited, P.O. Box 223, Kericho 20200, Kenya.
*Corresponding author: za@up.ac.za

Abstract

The advancements in genotyping have opened new approaches for identification and precise
mapping of Quantitative Trait Loci (QTLs) in plants, particularly by combining linkage and
association mapping (AM) analysis. In this study, a combination of linkage and the AM
approach was used to identify and authenticate putative QTLs associated with black tea
quality traits and percent relative water content (%RWC). The population structure analysis
clustered two parents and their respective 261 F1 progenies from the two reciprocal crosses
into two clusters with 141 tea accessions in cluster one and 122 tea accessions in cluster two.
The two clusters were of mixed origin with tea accessions in population TRFK St. 504
clustering together with tea accessions in population TRFK St. 524. A total of 71 putative
QTLs linked to black tea quality traits and %RWC were detected in interval mapping (IM)
method and were used as cofactors in multiple QTL model (MQM) mapping where 46
putative QTLs were detected. The phenotypic variance for each QTL ranged from 2.8-23.3%
in IM and 4.1-23% in MQM mapping. Using Q-model and Q+K-model in AM, a total of 49
DArTseq markers were associated with 16 phenotypic traits. Significant marker-trait
association in AM were similar to those obtained in IM, and MQM mapping except for six
more putative QTLs detected in AM which are involved in biosynthesis of secondary
metabolites, carbon fixation and abiotic stress. The combined linkage and AM approach
appears to have great potential to improve the selection of desirable traits in tea breeding.

Keywords: Quantitative Trait Loci; Tea quality; Drought tolerance; Linkage mapping;
Association mapping

Introduction
Tea plant (*Camellia sinensis* (L.) O. Kuntze) is an important economic crop grown in 36
tropical and semi-tropical countries around the world for production of tea beverages. In
Africa, tea is mainly grown in Kenya, Malawi, Uganda, Tanzania, Zimbabwe, Rwanda, South
Africa, Burundi, and Mauritius, although some tea is also grown in Ethiopia, Nigeria and
Cameroon (FAO 2015a). Globally, Kenya is ranked third in terms of annual volume
production of black tea after China and India and therefore, tea is Kenya's top foreign
exchange earner and export commodity among agricultural produces (Kenya National Bureau
of Statistics 2012). Tea being universally the most popular beverage, the need to develop tea
cultivars with optimum potential for black or green tea quality has recently become the single
most important breeding objective. Currently, Kenya produces cut, tear and curl (CTC) black
tea as the only product, for which the prices have declined owing to a global glut. This calls
for concerted action, the chief being the diversification and added value of tea products, not
only to help Kenya maintain its position as a leading exporter but also to enhance foreign
exchange earnings.

Tea breeding involves the creation of genetic variability through controlled hybridisation
between selected progenitors and evaluation of the newly generated progeny. This is to

identify promising individuals with desirable traits with the aim of improving tea
productivity. However, conventional tea breeding requires at least 20-25 years (Chen et al.
2013), from individual selection to local adaptability testing and to the final release of new
tea cultivars to the farmers, hence requires considerable investment in terms of land, labour,
time and money. Therefore, precise identification and screening of genetic resources, the
development of molecular breeding technologies and the deep understanding of the genetics
of important agronomic traits should be given more attention. Currently, the tea breeding
program is primarily focused on high-yielding tea, better quality processed tea, genetic
resistance to biotic and abiotic stresses prevailing in different tea growing regions and the
regional adaptability. However, these agronomic traits in tea plant are quantitative in nature
and not amenable to easy manipulations in breeding programs without elaborate and long-
term field testing (Wachira et al. 2001; Kamunya et al. 2004). Thus, tea breeding
improvement process would be more enhanced with the application of reliable and stable
selection tools that are not prone to the continually changing environment. For these reasons,
the selection of breeding materials by the evaluation of DNA markers for important
agronomic traits that are controlled by a few loci could be preferable method for marker-
assisted selection (MAS) for many traits in many different populations (Chen et al. 2013).

Linkage mapping and association mapping (AM) and are two methods widely used for
locating genes or QTL in plant breeding (Xu et al. 2017). Linkage mapping has been used
extensively for identifying the genetic basis of quantitative traits in plants (Kulwal 2018).
However, linkage mapping is usually limited by low polymorphism or small population size
and few recombination events, which are considered to estimate the genetic distances
between marker loci causative genomic regions for QTLs (Collard et al. 2005). However,
AM also known as Linkage Disequilibrium mapping has extensively been used to circumvent
the limitations posed by linkage mapping in plants in the last few years (Kraakman et al.
2004). Association mapping identifies QTL by examining the trait-marker associations has
been used to overcome the limitations of linkage mapping (Kraakman et al. 2004). In linkage
mapping, only two alleles at any given locus can be studied in bi-parental crosses and gives
low mapping resolution (Flint-Garcia et al. 2003). Thus, AM generally offers higher
resolution mapping due to a greater number of recombination events. However, AM has a
lower power to detect the effect of rare alleles compared to linkage analysis which has a
higher statistical power due to greater allele replication (Korte and Farlow 2013). Moreover,
population structure may cause false positives in AM, but this has been overcome by using a
mixed-model, which take both population structure (Q) and kinship (K) into account to
reduce false positives (Yu et al. 2006). Although linkage mapping and AM each offer more
advantages over the other, they are often applied in conjunction to validate the QTL
identified, thus reducing spurious associations.

The integrated approach of linkage-AM has been used in other crop plants such as
Arabidopsis (Sterken et al. 2012), wheat (Shi et al. 2017) and maize (Li et al. 2016) to
dissect quantitatively inherited traits. In tree species, an integrated method of linkage mapping and
AM to decipher the nature of genetic architecture of potential QTLs for growth traits has
been reported in poplar hybrids (Du et al. 2016) and maritime pine (Bartholomé et al. 2016).
In grapes, (Fournier-Level et al. 2010) a combined linkage mapping and AM has been used to
study the genetic patterns of anthocyanin content, which is a determinant of berry colour, in
grapes. Recently, AM has been used to study the genetic relationship between tea caffeine
synthase gene (TCS1) and the caffeine content in tea plant and its related species (Jin et al.
2016). In this current study, we integrated the high QTL detection power of the linkage
mapping with the high-resolution power of AM. The linkage-AM approach will not only
accelerate the pace of QTL mapping in tea breeding improvement but will also precisely

identify reliable QTLs linked to black tea quality and drought tolerance in tea. The objective
of this study was to integrate linkage mapping and AM to precisely identify and authenticate
putative QTLs linked to caffeine, catechins, theaflavins, tea tasters' scores, and %RWC.

Material and methods
Population type
Two pseudo-testcross populations used in this current study consisted of 109 F_1 progeny from
TRFK St. 504 (TRFK 303/577 x GW Ejulu) and 152 F_1 progeny from TRFK St. 524 (GW
Ejulu x TRFK 303/577) as described previously (Koech et al. 2018). In brief, the two parental
clones, TRFK 303/577 and GW Ejulu, are of Assamica and China varieties, respectively. The
two clones were chosen on the basis of their contrasting attributes. GW Ejulu is a low-
yielding, high black tea quality and moderate levels of caffeine, but high in total catechins
and individual catechins contents. Clone TRFK 303/577, which is an open-pollinated
progeny of clone TRFK 6/8, is high-yielding, drought-tolerant, medium in black tea quality,
caffeine and individual catechins.

Phenotypic data
A total of 16 phenotypic traits as described in the previous work by Koech et al. (2018) were
used in this study. In brief, all the 16 phenotypic traits were assessed for a normal
distribution. The mean and the standard deviation for each phenotypic trait in each parent and
the progenies were calculated. The significance difference between parental values and
progenies was analysed using Student's t test.

Population structure
The population structure was inferred from the 1,421 DArT markers data previously used to
construct genetic linkage map using the STRUCTURE software version 2.3.4 (Pritchard et al.
2010). Twenty independent runs were carried out using the following parameters; number of
populations (K) from 1 to 10, burn-in time and Markov Chain Monte Carlo (MCMC)
replication number were both set to 100,000 for model of admixture with correlated allele
frequencies. The natural logarithms of the probability data (LnP(K)) and the *ad hoc* delta K
statistical were calculated using STRUCTURE Harvester (Earl 2012), and the optimal K
according to the delta K value was then selected (Mezmouk et al. 2011). Finally, the
population structure matrix (Q) was obtained by integrating 20 independent replicate runs and
applying CLUMPP software (Jakobsson and Rosenberg 2007). STRUCTURE bar plots were
plotted using STRUCTURE Plot v 2.0 (Ramasamy et al. 2014).

Linkage mapping, QTL analyses and allelic effects
One thousand four hundred and twenty-one DArTseq markers derived from *C. sinensis*
genomic DNA of the two parents and 261 F_1 progeny were subjected to linkage mapping
analyses using JoinMap 4.0 (Van Ooijen 2006) and QTL mapping analyses using MapQTL
6.0 software (Van Ooijen *et al.*, 2000) as described previously (Koech et al. 2018; Koech et
al., 2019).
The allelic effects were estimated as $A_f = [(\mu ac + \mu ad) - (\mu bc + \mu bd)]/4$ for female additivity,
$A_m = [(\mu ac + \mu bc) - (\mu ad + \mu bd)]/4$ for male additivity and $D = [(\mu ac + \mu bd) - (\mu ad + \mu bc)]/4$
for dominance where μac, μad, μbc and μbd are estimated phenotypic means associated to
each of the four possible genotypic classes ac, bc, ad and bd, derived from ab×cd cross
(Sadok et al, 2013).

Association mapping
Using the Q and K matrices data files from STRUCTURE software as a covariates, 1,421
DArT markers of the 261 F_1 progeny were tested for association with each phenotype using a

general linear model (GLM), mixed linear model (MLM) that included kinship, phylogenetic
tree in TASSEL (trait analysis by association, evolution and linkage) Version 5.2.43 software
(Bradbury et al. 2007). Principal component analysis (PCA) was performed using JMP Pro
14 to visualise the dispersion of the association panel in a graph. The Q and kinship (K)
matrices were used to correct the effects of population substructure in the association panel
which can cause false-positive associations. The p-value and R^2 were used to determine
whether a QTL is associated with the marker and the magnitude of QTL effects, respectively.

Results
Frequency distribution
A wide range of variation in individual catechins, theaflavins, and caffeine contents was
observed in the two parents and F_1 progeny for all 16 phenotypic traits as described
previously by Koech et al. (2018).

Population structure
A population structure analysis showed that the value of Evanno's ΔK presented a sharp spike
at K = 2, which suggested that this population panel was clustered into two groups (Figure 1).
The two groups or clusters are represented by two clades with parental clone TRFK 303/577
clustering with 141 progenies in cluster one (blue) and parental clone GW Ejulu (not shown)
clustering with 122 progenies in cluster two (black), respectively (Figure 2). The two clusters
were accession with mixed origin with tea accessions in TRFK St. 504 population clustering
together with tea accessions in TRFK St. 524 population. Furthermore, the results from PCA
(Figure 3) and a phylogenetic tree-based on Nei's genetic distance was in agreement with the
structure analysis results (Figure 4). The population structure of the two tea populations was
examined with the neighbour-joining algorithm using Euclidian distance on the DArTseq
marker band intensities.

Linkage mapping and association mapping
Interval mapping and allelic effects
The genetic linkage map was constructed using a total of 1,421 DArT markers as described
previously (Koech et al. 2018). A total of 71 putative QTLs were identified for the 16
phenotypic traits measured using the interval mapping (Table 1). Of all the putative QTLs
identified, most QTLs were found in almost all linkage groups, except LG03, LG05 and
LG11. For black tea quality traits, 67 putative QTLs were detected and were located on
almost all linkage groups, except on LG03, LG05 and LG11. The remaining three putative
QTLs which are associated with drought tolerance were identified in LG02, LG06 and LG09.
Eight putative QTLs were identified on LG01, seven in LG02 and LG12, 10 in LG04, five in
LG06 and LG15, three in LG07, LG08, LG09, two in LG10, 15 in LG13, four in LG15,
respectively. The highest number of putative QTLs were associated with catechin (13), EGC
(12), tea tasters' score (12), caffeine (10) and EGC (9), respectively. The phenotypic variance
for all the identified putative QTLs ranged from 2.8% for EGC to 23.3% for qECG,
respectively. The phenotypic variance for %RWC ranged from 5.7 to 7.3%, while for the tea
tasters' score, it ranged from 5.8 to 9.1%. Half of the allelic effects identified to be associated
with black tea quality and drought tolerance traits had a positive additive effect. There were
eight allelic effects associated with female additivity, and 18 were associated with male
additive effects, respectively. The highest positive female and male additive effects were
qEGC and qECG, respectively. Although, some putative QTLs also exhibited high negative

allelic effects, which were from either female or male additive effects or both, such as
qEGCG, qCAT, qEC and qCaffeine (Table 1). All the putative QTLs identified in IM were
used as cofactors in MQM mapping.

Multiple QTL model mapping and allelic effects
A total of 46 putative QTLs (LOD > 3.0) were identified in 16 phenotypic traits using MQM
mapping method from the selected cofactors in IM. Similarly, no putative QTLs in LG03,
LG05 and LG11, respectively were associated with any of the phenotypic traits. Most of the
putative QTLs previously identified in IM were also identified MQM mapping. Of these 46
putative QTLs identified, 43 putative QTLs that were associated with black tea quality traits
while three were associated with qRWC (Table 2). The three putative QTLs associated with
qRWC were similar to those identified in IM. Four putative QTLs were identified in LG01
and LG12, seven in LG02, five in LG04, three in LG06, LG09, LG13, LG14 and LG15, two
in LG07 and LG08 and one in LG01, respectively. A similar trend was observed in the
number of putative QTLs for each trait as in IM with a high putative QTL number for
catechin, ECG, EGC and caffeine, respectively. The phenotypic variance for all the identified
putative QTL ranged from 4.1% for qCAT to 23% for qECG, respectively. The phenotypic
variance for qRWC was similar to IM results, and it ranged from 5.7 to 7.3%, while for the
tea tasters' score, it ranged from 6.0 to 9.1%. For allelic effects, similar results to IM with
qEGC and qECG showing high positive female additive and male additive effects,
respectively, this trend was also observed in MQM mapping. The putative QTLs identified
using both IM, and MQM mapping were almost similar although the efficiency and the
accuracy of detecting QTL is achieved by employing MQM instead of the single QTL model
used in IM.

Association mapping
Four models GLM, GLM (Q), mixed linear model with kinship matrix (MLM) (K) and mixed
linear model with kinship matix and population structure matrix (MLM) (Q+K) in TASSEL
software v5.2.43 were used to determine AM and the effects of population structure on the
AM to reduce the inflation of false-positive associations. The p-values were plotted
cumulatively for each model, and the distribution examined. However, no further analysis
was done using GLM model since marker-trait association results were characterised by
excess small p-values, which indicated an abundance of spurious associations
(Supplementary Figure 1). The association analysis using the Q-model and Q+K-model
detected a total of 49 DArT markers associated with 15 different black tea quality traits and
one drought-tolerant trait. The GLM (Q) model showed associations between DArT markers
and traits (p < 0.01) and was confirmed using the MLM (Q+K) model (Table 3,
Supplementary Figures 2-4). The MLM Q+K-model (MLM with Q-matrix and K-matrix as a
correction for population structure) showed a good fit for the p-values (p < 0.01), as
compared to GLM (Q) models, which were characterised by a few associations with excess
of small p-values, which indicates abundance of spurious associations (Table 3,
Supplementary Figures 2 and 4). Also, the MLM Q+K-model also showed better small p-
values than the K-model (MLM with K-matrix as a correction for population structure)
(Supplementary Figures 3 and 4). The GLM (Q) model may not have accounted for the
heterogeneity of the genetic background in some tea cultivars under study, which may have
resulted in false-positive associations. Ideally, the distribution of p-values should follow a
uniform distribution with less deviation from the expected p-values. However, the two
models were used to compare the effect of population matrix and kinship matrix on GLM and
MLM, respectively, in associating mapping. Therefore, taking into account the performance
of the two different models, the results from the MLM (Q+K) model appeared to have

controlled better population structure and kinship relationships than GLM (Q) model. While it might be tempting to consider p-values that remain extreme after MLM (Q+K) model correction as true associations, we need to keep in mind that minor allele frequency also influences p-values.

A total of 49 DArT markers were associated with 16 different phenotypic traits in six out of 15 linkage groups (Table 3). Since, all 1,421 DArT markers in the genotypic dataset had a known map location, the loci associated with particular traits could be allocated to a linkage group. The numbers of marker loci associated with traits together with their location and positions in different linkage groups at a test level of p< 0.01 in the two test models are shown in Table 3. Choosing a significance level of p< 0.01 which involves multiple testing corrections in both GLM (Q) and MLM (Q+K) models, 32 DArT marker loci were detected for all individual catechins (C, EC, ECG, EGC, EGCG), scattered over five linkage groups namely LG01, LG02, LG04, LG12 and LG15 (Table 3). For caffeine trait, three DArT markers were only located in LG02 while 11 DArT markers for all individual theaflavins traits (TF1, TF2, TF3, TF4) were detected in LG04 and LG15 (Table 3). The trait for tea liquor colour which is an indicator for black tea quality in terms of tea tasters' score appeared to be associated with only one DArT marker locus at the significance level of p< 0.01 in LG07 (Table 3). Drought tolerance trait (%RWC) was also associated with only one DArT marker locus detected in LG02 (Table 3). The majority of the DArT markers loci associated with the traits were found in LG04, LG12, LG15, LG02, and LG01. The three DArT markers associated with catechin trait in LG04, namely, 5136945, 5064764 and 5134490 had the highest proportion of the percent explained phenotypic variation of 41%, 38%, and 25%, respectively. The DArT markers that also showed high proportion of percent explained phenotypic variation were also observed in LG04 for qEGCG, qEGC, and qTF1 traits, respectively. In this study, a pleiotropic locus which is associated with a single locus affecting two or more distinct phenotypic traits was observed. Several pleiotropic loci were identified including DArT markers 5136945, 5112438, 5136058, 5123257, 5106022 and 5064764 which were associated with qC, qEC, qEGC, qTF1, qTF2 and qTF4 phenotypic traits (Table 3). All the pleiotropic loci detected and associated with the respective phenotypic traits were located in LG04.

A comparison of linkage mapping and association mapping
The putative QTLs detected using IM, and MQM mapping in MapQTL 6.0 were compared with marker-trait association results obtained using GLM (Q) and MLM (Q+K) models in AM. The significant marker-trait association results obtained using AM were similar to with the results obtained using MapQTL 6.0 (Tables 4 and 5). The IM analysis produced 14 significant QTLs (LOD > 3.0) that were also found to be associated with the phenotypic traits in AM analysis (Table 4). From the 14 QTLs, a total of 11 markers were associated with four individual catechins (qCAT, qEC, qECG, and qEGC) and one marker each was associated with qTF2, qCL, and qRWC, respectively. On the other hand, the MQM mapping analysis produced eight significant QTLs (LOD > 3.0) which were also found to be significantly associated with the phenotypic traits in AM analysis (Table 5). Of these, five markers were associated with four individual catechins (qCAT, qEC, qECG, and qEGC) and one marker each was associated with qTF2, qCL, and qRWC, respectively. Furthermore, the four mapping models were compared with each other to find markers that could associate with phenotypic traits in all four mapping models (Table 6). A total of 13 markers were found to be associated with four individual catechins (qCAT, qEC, qECG, and qEGC) and one marker each was associated with qTF2, qCL, and qRWC, respectively. Most of the markers that were associated with different phenotypic traits were only located in LG01, LG02, LG04, LG12

and LG15 with a majority of markers showing both positive and negative male additive
effects.

Functional annotation of putative QTLs in linkage and association mapping
The putative QTLs detected in both IM, and MQM mapping were identified by the BLAST
algorithm, searched on the reference tea genome (NCBI) and assigned functions on Blast2GO
database. A total of 53 QTLs were detected in both interval and MQM mapping methods, of
which 45 proteins were assigned functions. An additional six markers in LG02, LG04, and
LG12 were found to be associated with caffeine, EC, ECG and EGC traits in AM (Table 7).
The six additional QTLs detected in AM were also functionally annotated using the above
databases. The putative candidate genes identified were involved in purine or thiamine
biosynthesis, phenylalanine biosynthesis, tyrosine and tryptophan biosynthesis, carbon
fixation (photosynthesis) and abiotic stress. The putative QTL, qCaffeine in LG02 was
putatively annotated N-(5'phosphoribosyl) anthranilate (PRA) isomerase enzyme. The two
other putative QTLs identified for qEC and qEGC located in LG04 were annotated as
phosphoribulokinase or uridine kinase family proteins. Two putative QTLs, for qEC in LG04,
were annotated as autophagy-related protein 11 and protein tyrosine kinase, respectively.

Discussion
In this current study, the integration of linkage mapping and AM was used to provide a
mutual and precise authentication of QTL, which will enable more reliable results to be
obtained. Linkage mapping method has a relatively low genome resolution while AM method
is affected by population structure and individual relationships. The integration of linkage
mapping and AM has successfully been applied to many plant studies (Fan and Xiong 2003;
Jung et al. 2005; Lu et al. 2010; Lou et al. 2015; Cao et al. 2017).

The population structure analysis in discovery mapping panel used in the current study
identified two groups or clusters. The dendrogram demonstrated two subspecies of the two
parental clones, TRFK 303/577 and GW Ejulu, which are of *C. sinensis* var. *assamica* and *C.*
sinensis var. *sinensis* variety, respectively. The neighbour-joining tree for the 261 tea
accessions grouped them into two major clusters on the base of Nei's genetic distance. The
phylogenetic relationships for the parental cultivars and their progenies clustering into their
respective groups in separate clades based on their parental genotypes were consistent with
their genetic backgrounds (Kamunya et al. 2010). The parental clone TRFK 303/577, which
is an open-pollinated progeny of clone TRFK 6/8, is high-yielding, drought-tolerant, medium
in black tea quality, caffeine, and individual catechins. On the other hand, parental clone GW
Ejulu is a low-yielding, high black tea quality and moderate levels of caffeine, but high in
total catechins and individual catechin contents (TRFK 2012). The bulk of tea cultivars used
for green and black tea production have been derived through individual selection,
hybridisation and molecular breeding of *C. sinensis* var. *assamica* and *C. sinensis* var.
*sinensi*s varieties to produce tea with desirable characteristics (Yao et al. 2008). Therefore,
developing a few markers that have the capability of discriminating the two subspecies and
their cultivars is of paramount interest. The results on the linkage mapping for all the
phenotypic traits and the QTL positions in different linkage groups in this study were similar
as reported in (Koech et al. 2018).

In this study, AM was conducted with three different models, Q, K, and Q+K. The observed
–log10 (P) values for QTL deviated from the expected –log10 (P) values in the Q method
(GLM), indicated that there might be a few false-positive. However, the addition of genetic
relatedness K (kinship) used in a mixed linear model has proven to be more powerful and
reduces the number of false-positive associations (Finno et al. 2014). This is because a lot
more QTLs were identified in this current study using Q-model as compared with K and Q +

K-model. Therefore, there could be higher chances of false-positive or negative errors in AM due to complex population structure (Müller et al. 2011; George 2013). In this study, a mixed linear model approach using the K-matrix or a combination (Q+K) performed better than a GLM (Q). However, the K or the Q + K methods did show to be too strict and resulted in the missing of some possibly useful QTLs. Nearly all the significant QTLs identified in AM using the Q, and Q+K models were in line with those identified in MapQTL 6.0 (Koech et al. 2018). The percentages of phenotypic variation explained (R^2) by various markers analysed using AM were significant as those analysed using MapQTL 6.0. This is in agreement with previous studies reported in plants on efficiency and robustness of combining linkage mapping and AM for precise identification of QTL (Mammadov et al. 2015; Li et al. 2016).

The six additional QTLs identified in AM were found to be associated with purine or thiamine biosynthesis, phenylalanine, tyrosine and tryptophan biosynthesis, carbon fixation (photosynthesis) and abiotic stress. The putative QTL, qCaffeine located in LG02 was putatively annotated N-(5'phosphoribosyl) anthranilate (PRA) isomerase enzyme. The enzyme is involved in the first and intermediate pathway in the purine or thiamine biosynthesis and tryptophan biosynthesis pathway (Li et al. 1995). Caffeine (1,3,7-trimethylxanthine) and other methylxanthines such as theobromine (3,7-dimethylxanthine) and methyluric acids present in tea are classified as purine alkaloids (Ashihara et al. 2008). Therefore, the putative QTL, qCaffeine could be associated with the biosynthesis of caffeine, theobromine and methyluric acids in tea. Also, phenylalanine, tyrosine, and tryptophan are products of shikimate pathway which is involved in the biosynthesis of plant flavonoids, including catechins (Ghasemzadeh and Ghasemzadeh 2011).

The putative QTLs, qEC, and qEGC in LG04 were annotated as phosphoribulokinase/uridine kinase family proteins which are involved in carbon fixation or photosynthesis in plants (Miziorko 2000). Photosynthesis is an important process in plants for provision of carbon skeletons to the shikimate pathway (Janacek et al. 2009). Catechins are synthesised in the leaves of the tea plant through the acetic-malonic acid and shikimic-cinnamic acid metabolic pathways (Vuong et al. 2011). The chalcone and gallic acid are produced from the shikimic acid pathway, which then produce the different catechins (Chu and Juneja 1997). The two putative QTLs, qEC annotated putatively as autophagy-related protein 11 and protein tyrosine kinase are involved in abiotic stress response through a process of stomatal closure and ABA signalling (Sah et al. 2016; Batoko et al. 2017). Stress conditions can interfere with photosynthetic energy production in plants, which leads to stomatal closure, which inhibits CO_2 intake and thereby reduces photosynthetic activity in leaves (Chaves et al. 2009). Abscisic acid has been reported to confer abiotic stress-tolerance in crop plants (Sah et al. 2016). In stress conditions like drought, extreme temperature, and high salinity, content in plants increases considerably, inspiring stress-tolerance effects that help plants, adapt, and survive under these stressful situations (Ng et al. 2014). Also, ABA is also required for plant growth and development under non-stress conditions (Nakashima and Yamaguchi-Shinozaki 2013).

Conclusion

The tea plant is a woody perennial crop with long generation intervals, which usually hinders its genetic improvement by conventional breeding methods. The approach of combining the high QTL detection power of genetic linkage mapping with the high-resolution power of AM allowed identification and precise authentication of putative QTL controlling black tea quality and drought tolerance traits. Based on the two mapping approaches, the putative QTLs associated with caffeine, catechins, theaflavins, tea tasters' scores, and %RWC detected in linkage mapping and AM were not significantly different except that an additional six more QTLs were detected using AM method. Although, conventional QTL mapping

through bi- or multi- parental populations is a powerful method, it suffers from a limited
amount of recombination. However, AM can partly overcome the limitation by using a
diverse germplasm but may lead to a number of false positive or negative associations.
Therefore, the two different methods can be complementary to each other and benefit can be
achieved by mitigating the other's limitations. In this study, the combination of AM and QTL
mapping has led to successful identification of QTL with potential application in breeding
programs. However, the QTL identified by AM requires further confirmation in bi- or multi-
parental populations. This will accelerate tea breeding progress in terms of reduced breeding
time, reduced cost, increased efficiency, and precision of selection.

Author contributions
ZA, SK and RM were involved with the design of the experiment and plant material used.
RK was involved in collection of plant material. RK performed the experiments, analyzed
samples and interpreted the data. RK wrote the manuscript and revised by RM, SK and ZA.
The final manuscript was reviewed and approved by all the authors. The TRI of Kenya and
the University of Pretoria had role in study design, data collection and analysis, decision to
publish, or preparation of the manuscript.

Conflict of interest statement
All the authors declare that there is no commercial or financial relationships that can
precedence to conflict of interest in research conducted.

Acknowledgement
The authors acknowledge the financial support to conduct this research, and study grants for
RK and PM from James Finlay (Kenya) Ltd., George Williamson (Kenya) Ltd., Sotik Tea
Company (Kenya) Ltd., Mcleod Russell (Uganda) Ltd., the TRI of Kenya and Southern
African Biochemistry and Informatics for Natural Products (SABINA). The *C. sinensis*
cultivars used in this study were provided by the TRI of Kenya. Supplementary funding was
provided by, the Technology and Human Resources for Industry Programme (THRIP), an
initiative of the Department of Trade and Industries of South Africa (dti), the National
Research Foundation (NRF) of South Africa, and the University of Pretoria (South Africa).

References
Ashihara H, Sano H, Crozier A (2008) Caffeine and related purine alkaloids: biosynthesis,
catabolism function and genetic engineering. Phytochemistry 69(4): 841-856.
Bartholomé J, Bink MC, van Heerwaarden J, Chancerel E, Boury C, Lesur I, Isik F, Bouffier
L, Plomion C (2016) Linkage and association mapping for two major traits used in the
maritime pine breeding program: height growth and stem straightness. PloS one
11(11): e0165323.
Batoko H, Dagdas Y, Baluska F, Sirko A (2017) Understanding and exploiting autophagy
signaling in plants. Essays Biochem 61(6): 675-685.
Bradbury PJ, Zhang Z, Kroon DE, Casstevens TM, Ramdoss Y, Buckler ES (2007) TASSEL:
software for association mapping of complex traits in diverse samples. Bioinformatics
23(19): 2633-2635.
Cao Y, Li S, Wang Z, Chang F, Kong J, Gai J, Zhao T (2017) Identification of major
quantitative trait loci for seed oil content in soybeans by combining linkage and
genome-wide association mapping. Front plant sci 8: 1222.
Chaves MM, Flexas J, Pinheiro C (2009) Photosynthesis under drought and salt stress:
regulation mechanisms from whole plant to cell. Ann Bot 103(4): 551-560.
Chen L, Apostolides Z, Chen ZM (2013) Global tea breeding: achievements challenges and
perspectives. Springer Science, Business Media.

Chu DC, Juneja LR (1997) General Chemical Composition of Green Tea and Its Infusion. In: Yamamoto T, Juneja LR, Chu DC, Kim M, Eds., Chemistry and Applications of Green Tea, CRC Press, Boca Raton, 13-22.

Collard B, Jahufer M, Brouwer J, Pang E (2005) An introduction to markers quantitative trait loci (QTL) mapping and marker-assisted selection for crop improvement: the basic concepts. Euphytica 142(1-2): 169-196.

Du Q, Gong C, Wang Q, Zhou D, Yang H, Pan W, Li B, Zhang D (2016) Genetic architecture of growth traits in Populus revealed by integrated quantitative trait locus (QTL) analysis and association studies. New Phytol 209 (3): 1067-1082.

Earl DA (2012) STRUCTURE HARVESTER: a website and program for visualizing STRUCTURE output and implementing the Evanno method. Conserv Genet Resour 4(2): 359-361.

Fan R, Xiong M (2003). Combined high resolution linkage and association mapping of quantitative trait loci. Eur J Hum Genet 11(2): 125.

FAO (2015a) World tea production and trade current and future development In *FAO Intergovernmental Group on Tea A Subsidiary Body of the FAO Committee on Commodity Problems (CCP)* 1-13 (Ed. K Chang) Rome Italy.

Finno CJ, Aleman M, Higgins RJ, Madigan JE, Bannasch DL (2014) Risk of false positive genetic associations in complex traits with underlying population structure: a case study. Vet J 202(3): 543-549,

Flint-Garcia SA, Thornsberry JM, Buckler IV ES (2003) Structure of linkage disequilibrium in plants. Annu Rev Plant Biol 54(1): 357-374.

Fournier-Level A, Lacombe T, Le Cunff L, Boursiquot J, This P (2010) Evolution of the VvMybA gene family the major determinant of berry colour in cultivated grapevine (*Vitis vinifera* L.). Heredity 104(4): 351.

George A (2013) Controlling type 1 error rates in genome-wide association studies in plants. Heredity 111(1): 86.

Ghasemzadeh A, Ghasemzadeh N (2011) Flavonoids and phenolic acids: Role and biochemical activity in plants and human. J Med Plants Res 5(31): 6697-6703.

Jakobsson M, Rosenberg NA (2007) CLUMPP: a cluster matching and permutation program for dealing with label switching and multimodality in analysis of population structure. Bioinformatics 23(14): 1801-1806.

Janacek SH, Trenkamp S, Palmer B, Brown NJ, Parsley K, Stanley S, Astley HM, Rolfe SA, Paul Quick W, Fernie AR (2009) Photosynthesis in cells around veins of the C3 plant *Arabidopsis thaliana* is important for both the shikimate pathway and leaf senescence as well as contributing to plant fitness. Plant J 59(2): 329-343.

Jin JQ, Yao MZ, Ma CL, Ma JQ, Chen L (2016) Association mapping of caffeine content with TCS1 in tea plant and its related species. Plant Physiol Biochem 105: 251-259.

Jung J, Fan R, Jin L (2005) Combined linkage and association mapping of quantitative trait loci by multiple markers. Genetics 170(2): 881-898.

Kamunya S, Wachira F, Muoki R (2004) Evaluation of newly developed clones of tea (*Camellia sinensis* (L.) O. Kuntze) for yields drought tolerance and quality: preliminary indicators. Tea 25(1): 12-19.

Kamunya S, Wachira F, Pathak R, Korir R, Sharma V, Kumar R, Bhardwaj P, Chalo R, Ahuja P, Sharma R (2010) Genomic mapping and testing for quantitative trait loci in tea (*Camellia sinensis* (L.) O. Kuntze). Tree Genet Genomes 6(6): 915-929.

Kenya National Bureau of Statistics (2012) Kenya facts and Figureures 2012. Kenya.

Koech RK, Malebe PM, Nyarukowa C, Mose R, Kamunya SM, Apostolides Z (2018) Identification of novel QTL for black tea quality traits and drought tolerance in tea plants (*Camellia sinensis*). Tree Genet Genomes 14(1): 9.

Koech, R.K., Malebe, P.M., Nyarukowa, C., Mose, R., Kamunya, S.M., Joubert, F. and Apostolides, Z. (2019) Functional annotation of putative QTL associated with black tea quality and drought tolerance traits. Sci Rep 9(1): 1465.

Korte A, Farlow A (2013) The advantages and limitations of trait analysis with GWAS: a review. Plant Methods 9(1): 29.

Kraakman AT, Niks RE, Van den Berg PM, Stam P, Van Eeuwijk FA (2004) Linkage disequilibrium mapping of yield and yield stability in modern spring barley cultivars. Genetics 168(1): 435-446.

Kulwal PL (2018) Trait Mapping Approaches Through Linkage Mapping in Plants.

Li J, Zhao J, Rose AB, Schmidt R, Last RL (1995) *Arabidopsis* phosphoribosylanthranilate isomerase: molecular genetic analysis of triplicate tryptophan pathway genes. Plant Cell 7(4): 447-461.

Li X, Zhou Z, Ding J, Wu Y, Zhou B, Wang R, Ma J, Wang S, Zhang X, Xia Z (2016) Combined linkage and association mapping reveals QTL and candidate genes for plant and ear height in maize. Front Plant Sci 7: 833.

Lou Q, Chen L, Mei H, Wei H, Feng F, Wang P, Xia H, Li T, Luo L (2015) Quantitative trait locus mapping of deep rooting by linkage and association analysis in rice. J Exp Bot 66(15): 4749-4757.

Lu Y, Zhang S, Shah T, Xie C, Hao Z, Li X, Farkhari M, Ribaut JM, Cao M, Rong T (2010) Joint linkage–linkage disequilibrium mapping is a powerful approach to detecting quantitative trait loci underlying drought tolerance in maize. Proc Natl Acad Sci 107(45): 19585-19590.

Mammadov J, Sun X, Gao Y, Ochsenfeld C, Bakker E, Ren R, Flora J, Wang X, Kumpatla S, Meyer D (2015) Combining powers of linkage and association mapping for precise dissection of QTL controlling resistance to gray leaf spot disease in maize (Zea mays L.). BMC genomics 16(1): 916.

Mezmouk S, Dubreuil P, Bosio M, Décousset L, Charcosset A, Praud S, Mangin B (2011) Effect of population structure corrections on the results of association mapping tests in complex maize diversity panels. Theor Appl Genet 122(6): 1149-1160.

Miziorko HM (2000) Phosphoribulokinase: current perspectives on the structure/function basis for regulation and catalysis. Adv Enzymol Relat Areas Mol Biol 74: 95-127.

Müller B, Stich B, Piepho H (2011) A general method for controlling the genome-wide type I error rate in linkage and association mapping experiments in plants. Heredity 106(5): 825.

Nakashima K, Yamaguchi-Shinozaki K (2013). ABA signaling in stress-response and seed development. Plant Cell Rep 32(7): 959-970.

Ng LM, Melcher K, Teh BT, Xu HE (2014) Abscisic acid perception and signaling: structural mechanisms and applications. Acta Pharmacol Sin 35(5): 567.

Pritchard J, Wen X, Falush D (2010) Documentation for STRUCTURE software. [Documentation file] *Available with the program at http://pritchbsduchicagoedu/structurehtml*.

Ramasamy RK, Ramasamy S, Bindroo BB, Naik VG (2014) STRUCTURE PLOT: a program for drawing elegant STRUCTURE bar plots in user friendly interface. SpringerPlus 3(1): 431.

Sadok IB, Celton JM, Essalouh L, El Aabidine AZ, Garcia G, Martinez S, Grati-Kamoun N, Rebai A, Costes E, Khadari B (2013) QTL mapping of flowering and fruiting traits in olive. PloS one 8(5): e62831.

Sah SK, Reddy KR, Li J (2016) Abscisic acid and abiotic stress tolerance in crop plants. Front Plant Sci 7: 571.

Shi W, Hao C, Zhang Y, Cheng J, Zhang Z, Liu J, Yi X, Cheng X, Sun D, Xu Y (2017) A Combined Association Mapping and Linkage Analysis of Kernel Number Per Spike in Common Wheat (*Triticum aestivum* L.). Front Plant Sci 8: 1412.

Sterken R, Kiekens R, Boruc J, Zhang F, Vercauteren A, Vercauteren I, De Smet L, Dhondt S, Inzé D, De Veylder L (2012) Combined linkage and association mapping reveals CYCD5; 1 as a quantitative trait gene for endoreduplication in *Arabidopsi*s. Proc Natl Acad Sci 109(12): 4678-4683,

TRFK (2012) TRFK clonal catalogue (Ed. TRFK).

Van Ooijen J (2006) JoinMap® 4: Software for the calculation of genetic linkage maps in experimental populations. Kyazma BV Wageningen 33: 10,1371.

Van Ooijen JW, Boer M, Jansen R, Maliepaard C (2000) MapQTL 4.0: software for the calculation of QTL positions on genetic maps (user manual). Plant Research International.

Vuong QV, Stathopoulos CE, Nguyen MH, Golding JB, Roach PD(2011). Isolation of green tea catechins and their utilization in the food industry. Food Rev Int 27(3): 227-247.

Wachira F, Tanaka J, Takeda Y (2001) Genetic variation and differentiation in tea (*Camellia sinensis*) germplasm revealed by RAPD and AFLP variation. J Hortic Sci Biotechnol 76(5): 557-563.

Xu Y, Li P, Yang Z, Xu C (2017) Genetic mapping of quantitative trait loci in crops. Crop J l 5(2): 175-184.

Yao M, Chen L, Liang Y (2008) Genetic diversity among tea cultivars from China Japan and Kenya revealed by ISSR markers and its implication for parental selection in tea breeding programmes. Plant Breeding 127(2): 166-172.

Yu J, Pressoir G, Briggs WH, Bi IV, Yamasaki M, Doebley JF, McMullen MD, Gaut BS, Nielsen DM, Holland JB (2006) A unified mixed-model method for association mapping that accounts for multiple levels of relatedness. Nature Genet 38(2): 203.

Figure 1. Delta K values plotted from 1 to 10 for 261 tea accessions panel.

Figure 2. Population structure of 261 tea accessions panel (K = 2)

Figure 3. Principal component analysis of 261 tea accessions. ⬤ represents cluster 1, ▲ represents cluster 2

Figure 4. Phylogenetic tree-based on Nei's genetic distance. Blue clade represents cluster 1, black clade represents cluster 2.

Table 1. Linkage mapping results for putative QTLs in black tea quality and drought tolerance traits in St. 504 and St. 524 using interval mapping in MapQTL 6.0

Nr	[a]Trait	Locus	LG	Position (cM)	LOD	PEV	Af	Am	D	Allelic effect
1	AST	5115441_E-26	1	96.6	4.5	9.1	0.224	-0.061	-0.015	Af
2	AST	5135087	1	97.3	4.5	9.1	0.234	-0.057	-0.014	Af
3	BRK	5115441_E-26	1	96.6	3.6	7.3	0.196	-0.082	-0.020	Af
4	BRK	5135087	1	97.3	3.5	7.2	0.204	-0.078	-0.019	Af
5	ECG	5128890	1	96.4	6.8	11.7	0.326	0.046	0.011	Af
6	ECG	5135087	1	97.3	6.6	11.3	0.347	0.053	0.013	Af
7	ECG	5115441_E-26	1	96.6	6.5	11.1	0.331	0.051	0.013	Af
8	EGC	5133866	1	87.1	3.3	5.8	0.109	0.218	0.054	Am
9	CAFF	5064585	2	50.9	3.3	5.8	5.849	-9.702	-2.425	Af; Am
10	CAT	5135436	2	0.0	3.6	6.1	-0.838	-11.195	-2.799	Am
11	EC	5072338	2	2.2	4.3	7.2	-14.648	-12.555	-3.139	Af; Am
12	EGC	5124128	2	7.7	3.3	5.6	-36.447	1.544	0.386	Af
13	RWC	5136794	2	60.9	4.3	7.3	0.482	1.130	0.283	Am
14	TF1	5084595	2	4.5	3.2	5.7	-0.952	8.158	2.039	Am
15	TF2	5084595	2	4.5	4.1	7.3	2.290	9.140	2.285	Am
16	AR	100158044|F|0	4	70.1	2.9	6	-0.268	0.090	0.023	Af
17	CAFF	5112599	4	68.6	3.8	6.7	2.413	11.852	2.963	Am
18	CAT	5063001	4	30.3	5.3	4.1	-17.604	-53.467	-13.367	Am
19	EC	5123475	4	27.2	14.0	21.9	3.485	44.620	11.155	Am
20	EC	5123257	4	26.7	13.4	21.1	1.703	43.813	10.953	Am
21	EC	5134490	4	26.4	12.4	19.6	0.260	41.569	7.005	Am
22	ECG	5087113	4	17.7	3.8	5.8	-2891.857	3715.230	928.808	Af; Am
23	EGC	5123475	4	27.2	3.7	2.8	7071.043	-6846.394	-1711.598	Af; Am
24	EGCG	5087017	4	37.2	4.9	6	-320.975	-200.380	-50.095	Af; Am
25	ECG	5098382	6	57.0	5.2	8.7	35.650	-32.760	-8.190	Af; Am
26	EGC	5073424	6	66.4	3.9	6.7	-13.905	14.749	3.687	Af; Am
27	EGC	5124993	6	66.6	3.9	6.6	-12.321	15.635	3.909	Af; Am
28	RWC	5082606	6	66.2	3.3	5.7	-2.094	-0.814	-0.203	Af
29	TF1	5136045	6	69.6	4.3	7.7	2.795	3.782	0.946	Af; Am
30	CAFF	5064391	7	48.1	3.4	6	-12.264	-3.397	-0.849	Af
31	CAFF	5056614	7	48.3	3.3	5.9	-12.029	-3.985	-0.996	Af
32	CL	5132432	7	72.7	3.2	6.5	-0.138	-0.118	-0.029	Af; Am
33	CAFF	5134558	8	18.8	4.3	7.5	17.375	-12.332	-3.083	Af; Am
34	CAFF	5111497	8	18.8	4.1	7.1	11.542	-12.927	-3.232	Af; Am
35	CAT	5130194	8	12.5	3.9	6.6	-29.556	-9.720	-2.430	Af
36	AST	5123950	9	87.6	3.2	6.6	-0.036	0.178	0.045	Am
37	BRK	5123950	9	87.6	3.6	7.4	-0.012	0.190	0.047	Am

38	RWC	5130531	9	6.7	4.1	7	1.933	2.876	0.719	Am
39	AR	5075627	10	26.9	3.4	6.9	-0.064	0.110	0.027	Am
40	ECG	5136108	10	20.6	3.3	5.7	4.297	34.819	8.705	Am
41	CAT	5123751	12	43.0	4.2	6.5	-36.478	-3.162	-0.791	Af
42	CAT	5127224	12	42.9	4.2	6.5	-36.467	-3.045	-0.761	Af
43	ECG	5136790	12	50.4	15.0	23.3	-22.365	60.821	15.205	Am
44	ECG	5135536	12	50.4	15.0	23.2	-22.316	60.808	15.202	Am
45	EGC	5123751	12	43.0	5.0	8.5	50.465	-30.542	-7.635	Af; Am
46	EGC	5127224	12	42.9	5.0	8.4	50.241	-30.563	-7.641	Af; Am
47	EGCG	5104630	12	48.2	4.2	7.2	18.115	-52.080	-13.020	Am
48	BRT	5088162	13	29.8	3.0	6.1	0.006	-0.142	-0.035	Am
49	BRT	5135810	13	26.0	2.8	5.8	0.007	-0.137	-0.034	Am
50	CAFF	5088162	13	29.8	3.4	5.4	7.183	-29.136	-7.284	Am
51	CAFF	5080631	13	35.6	2.9	3.1	-14.651	23.420	5.855	Af; Am
52	CAT	5122819	13	10.2	4.3	7.3	-27.341	-14.521	-3.630	Af; Am
53	CAT	5111268	13	58.1	4.6	7.2	14.200	17.804	4.451	Af; Am
54	CAT	5129729	13	9.8	4.2	7.2	-26.752	-14.467	-3.617	Af; Am
55	CAT	5115793	13	58.7	4.3	6.8	12.635	17.248	4.312	Af; Am
56	CAT	5136782	13	58.5	4.2	6.6	12.029	17.237	4.309	Af; Am
57	CAT	5103784	13	50.6	3.8	6.1	4.188	21.966	5.491	Am
58	CAT	5122899	13	50.5	3.8	6	3.656	21.921	5.480	Am
59	ECG	5123761	13	60.7	3.3	5.8	-0.469	-0.112	-0.028	Af
60	ECG	5072523	13	60.3	3.0	5.2	-0.261	-0.231	-0.058	Af; Am
61	ECG	5114985	13	61.6	2.8	5	-0.439	-0.088	-0.022	Af
62	EGC	5136623	13	50.6	3.7	6.4	8.946	-29.764	-7.441	Am
63	BRT	5125626	14	71.1	3.3	6.8	0.000	0.143	0.036	Am
64	CAFF	5123053	14	5.5	4.0	7.1	-7.622	-11.572	-2.893	Af; Am
65	CAFF	5054639	14	6.4	3.9	6.9	-7.260	-11.533	-2.883	Af; Am
66	CAT	5132370	14	60.7	6.5	10.8	-19.945	35.937	8.984	Af; Am
67	EGC	5132370	14	60.7	3.7	6.4	46.492	-7.514	-1.878	Af
68	EC	5085963	15	25.4	4.4	7.5	11.674	-24.472	-6.118	Am
69	EC	5099958_E-25	15	25.1	4.4	7.4	11.686	-24.101	-6.025	Am
70	ECG	5111164	15	75.1	4.2	7.2	22.426	34.202	8.550	Af; Am
71	EGCG	5114089	15	32.1	4.0	6.8	-1.178	56.567	14.142	Am

[a] Putative QTL identified in Interval Mapping using MapQTL 6.0. LOD- Logarithm of odds for putative QTL in IM (LOD>2); PEV- percentage phenotypic variation explained by markers in IM; Af- Female additivity; Am- Male additivity; D- Dominance

Table 2. Linkage mapping for putative QTLs in black tea quality and drought tolerance traits in St. 504 and St. 524 using multiple QTL model mapping in MapQTL 6.0

Nr	[b]Trait	Locus	LG	Position (cM)	LOD	PEV	Af	Am	D	Allelic effect
1	AST	5135087	1	97.3	4.5	9.1	0.234	-0.057	-0.014	Af
2	BRK	5115441_E-26	1	96.6	3.6	7.3	0.196	-0.082	-0.020	Af
3	ECG	5128890	1	96.4	6.8	11.7	0.326	0.046	0.011	Af
4	EGC	5133866	1	87.1	3.3	5.8	0.109	0.218	0.054	Am
5	CAFF	5064585	2	50.9	3.3	5.8	5.849	-9.702	-2.425	Af; Am
6	CAT	5135436	2	0.0	3.6	6.1	-0.838	-11.195	-2.799	Am
7	EC	5072338	2	2.2	4.3	7.2	-14.648	-12.555	-3.139	Af; Am
8	EGC	5124128	2	7.7	3.3	5.6	-36.447	1.544	0.386	Af
9	RWC	5136794	2	60.9	4.3	7.3	0.482	1.130	0.283	Am
10	TF1	5084595	2	4.5	3.2	5.7	-0.952	8.158	2.039	Am
11	TF2	5084595	2	4.5	4.1	7.3	2.290	9.140	2.285	Am
12	AR	100158044|F|0	4	70.1	2.9	6	-0.268	0.090	0.023	Af
13	CAFF	5112599	4	68.6	3.8	6.7	2.413	11.852	2.963	Am
14	CAT	5063001	4	30.3	5.3	4.1	-17.604	-53.467	-13.367	Am
15	EC	5123475	4	27.2	14.0	21.9	3.485	44.620	11.155	Am
16	ECG	5087113	4	17.6	4.7	8.2	-0.133	0.141	-0.090	Af; Am
17	EGC	5123475	4	27.1	3.7	2.8	7071.043	-6846.394	7330.160	Af; Am; D
18	EGCG	5087017	4	37.2	4.9	6	-320.975	-200.380	-50.095	Af; Am
19	ECG	5098382	6	57.0	5.2	8.7	35.650	-32.760	-8.190	Af; Am
20	EGC	5073424	6	66.4	3.9	6.7	-13.905	14.749	3.687	Af; Am
21	TF1	5136045	6	69.6	4.3	7.7	2.795	3.782	6.425	D
22	RWC	5082606	6	66.2	3.3	5.7	-2.094	-0.814	-0.203	Af
23	CAFF	5064391	7	48.1	3.4	6	-12.264	-3.397	-0.849	Af
24	CL	5132432	7	72.7	3.2	6.5	-0.138	-0.118	-0.029	Af; Am
25	CAFF	5134558	8	18.8	4.3	7.5	17.375	-12.332	-3.083	Af; Am
26	CAT	5130194	8	12.5	3.9	6.6	-29.556	-9.720	-2.430	Af
27	AST	5123950	9	87.6	3.2	6.6	-0.036	0.178	0.045	Am
28	BRK	5123950	9	87.6	3.6	7.4	-0.012	0.190	0.047	Am
29	RWC	5130531	9	6.7	4.1	7	1.933	2.876	0.719	Am
30	ECG	5136108	10	20.6	3.3	5.7	4.297	34.819	8.705	Am
31	AR	5075627	10	26.9	3.4	6.9	-0.064	-0.064	-0.064	Am
32	CAT	5123751	12	43.0	4.2	6.5	-36.478	-3.162	-0.791	Af
33	ECG	5136790	12	50.4	15.0	23.3	-22.365	60.821	15.205	Am

34	EGC	5123751	12	43.0	5.0	8.5	50.465	-30.542	-7.635	Af; Am
35	EGCG	5104630	12	48.2	4.2	7.2	18.115	-52.080	-13.020	Am
36	BRK	5114985	13	61.6	3.1	6.3	-0.366	-0.071	0.374	Af; D
37	CAFF	5088162	13	29.8	3.4	5.4	7.183	-29.136	-7.284	Am
38	CAT	5111268	13	58.1	4.6	7.2	14.200	17.804	4.451	Af; Am
39	EGC	5136623	13	50.6	3.7	6.4	8.946	-29.764	-7.441	Am
40	BRT	5125626	14	71.1	3.3	6.8	0.000	0.143	0.036	Am
41	CAFF	5123053	14	5.5	4.0	7.1	-7.622	-11.572	-2.893	Af; m
42	CAT	5132370	14	60.7	6.5	10.8	-19.945	35.937	8.984	Af; Am
43	EGC	5132791	14	51.4	3.0	3.0	-14.572	66.511	-43.428	Am
44	EC	5085963	15	25.4	4.4	7.5	11.674	-24.472	-6.118	Am
45	ECG	5111164	15	75.1	4.2	7.2	22.426	34.202	8.550	Af; Am
46	EGCG	5114089	15	32.1	4.0	6.8	-1.178	56.567	14.142	Am

[b] Putative QTL identified in Multiple QTL Model Mapping using MapQTL 6.0

LOD- Logarithm of odds for putative QTL in MQM (LOD>3); PEV- percentage phenotypic variation explained by markers in MQM; Af- Female additivity; Am- Male additivity; D- Dominance

CAFF- Caffeine; CAT- Catechin; EC- Epicatechin; ECG- Epicatechin gallate; EGC- Epigallocatechin gallate; EGCG- Epigallocatechin gallate; TF1-Theaflavin; TF2- Theaflavin-3-gallate; TF3- Theaflavin-3'-gallate; TF4- Theaflavin-3,3'-digallate; CL- Colour; BRT- Brightness; AST- Astringency; BRK- Briskness; AR- Aroma; RWC- percent relative water content

Table 3. Association mapping results for black tea quality and drought tolerance traits in St. 504 and St.524 using the GLM (Q) and MLM (Q+K) method, respectively (p <0.01)

Nr	[c]Trait	Marker	LG	Position (cM)	p-Value	Marker R^2 GLM (Q)	p-Value	Marker R^2 MLM (Q+K)
1	ECG	5128890	1	96.4	1.18E-03	0.042	5.31E-03	0.031
2	EGC	5086476	1	87.5	8.20E-03	0.031	8.54E-04	0.043
3	EGC	5122985	1	87.5	1.47E-04	0.056	2.41E-03	0.038
4	EGC	5133866	1	87.1	8.94E-03	0.030	6.62E-04	0.046
5	CAFF	5134416	2	44.4	2.46E-04	0.057	5.58E-04	0.050
6	CAFF	5136551	2	43.3	9.60E-04	0.048	7.51E-03	0.030
7	CAFF	5137282	2	50.7	2.27E-03	0.042	9.72E-03	0.026
8	EC	5132500	2	7.9	2.91E-03	0.036	1.37E-04	0.052
9	RWC	5136794	2	60.9	8.58E-03	0.032	3.02E-03	0.033
10	CAT	5063001	4	30.3	2.34E-09	0.098	6.67E-08	0.109
11	CAT	5064764	4	28.8	6.42E-37	0.326	7.80E-22	0.379
12	CAT	5136945	4	28.1	6.39E-40	0.345	1.81E-21	0.406
13	EC	5112438	4	27.4	4.90E-07	0.087	1.50E-05	0.067
14	EC	5123257	4	26.7	3.89E-08	0.101	7.16E-07	0.089
15	EC	5134490	4	26.4	5.15E-19	0.228	6.21E-15	0.246
16	EC	5136058	4	27.6	1.00E-06	0.083	5.31E-06	0.073
17	EC	5136410	4	29.9	2.39E-05	0.065	1.77E-05	0.065
18	ECG	5087113	4	17.7	3.74E-03	0.035	2.92E-04	0.048
19	EGC	5112438	4	27.4	1.50E-11	0.148	1.95E-08	0.134
20	EGC	5123257	4	26.7	9.59E-13	0.162	6.95E-10	0.167
21	EGC	5136058	4	27.6	3.43E-11	0.143	3.60E-09	0.146
22	EGCG	5074553	4	39.5	3.26E-14	0.210	1.86E-12	0.221
23	EGCG	5123463	4	28.9	3.38E-17	0.250	1.09E-13	0.266
24	EGCG	5134853	4	37.6	7.03E-14	0.205	2.00E-09	0.170
25	EGCG	5136554	4	38.7	1.46E-16	0.242	2.08E-11	0.222
26	EGCG	100011901|F|0	4	38.7	1.24E-04	0.066	3.13E-03	0.039
27	TF1	5064764	4	28.8	2.35E-10	0.149	5.78E-06	0.088
28	TF1	5106352	4	26.0	1.95E-04	0.061	8.45E-03	0.032
29	TF1	5123463	4	28.9	1.33E-10	0.153	2.66E-06	0.097
30	TF1	5136945	4	28.1	1.69E-11	0.166	8.20E-07	0.107
31	TF2	5063001	4	30.3	9.76E-04	0.050	4.22E-03	0.035
32	TF2	5064764	4	28.8	1.85E-08	0.122	7.36E-05	0.063
33	TF2	5106022	4	29.1	2.99E-09	0.134	3.06E-04	0.052

34	TF4	5064764	4	28.8	8.64E-06	0.082	1.33E-03	0.042
35	TF4	5106022	4	29.1	1.38E-06	0.094	1.57E-03	0.040
36	TF4	5136945	4	28.1	6.26E-07	0.099	1.59E-04	0.060
37	CL	5132432	7	72.7	1.31E-03	0.047	7.30E-04	0.049
38	ECG	5059563	12	49.0	8.91E-09	0.111	2.29E-05	0.068
39	ECG	5136077	12	42.8	1.91E-07	0.094	6.34E-03	0.027
40	ECG	5136790	12	50.4	7.87E-11	0.137	1.58E-04	0.051
41	EGC	5135536	12	50.4	2.25E-04	0.053	2.82E-06	0.086
42	EGCG	5087581	12	48.3	1.51E-04	0.065	1.75E-03	0.040
43	EGCG	5088456	12	47.9	7.70E-04	0.053	2.46E-04	0.060
44	EGCG	5133629	12	47.7	7.17E-04	0.054	1.95E-03	0.041
45	EC	5085963	15	25.4	3.52E-04	0.049	9.09E-03	0.029
46	EC	5132852	15	24.8	1.09E-04	0.056	2.50E-03	0.036
47	EC	5099958_E-25	15	25.1	1.00E-04	0.056	6.37E-03	0.031
48	TF3	5083854	15	15.2	3.52E-03	0.043	2.77E-03	0.035
49	TF3	5136247	15	16.3	1.57E-03	0.049	5.40E-04	0.046

[c] Putative QTL identified in General Linear Model and Mixed Linear Model methods using TASSEL software v5.2.43

LG- Linkage group

Marker R^2- percentage phenotypic variation explained by markers in both GLM (Q) and MLM (Q+K) methods

Q- Population structure matrix

K- Kinship matrix

CAFF- Caffeine; CAT- Catechin; EC- Epicatechin; ECG- Epicatechin gallate; EGC- Epigallocatechin gallate; EGCG- Epigallocatechin gallate; TF1- Theaflavin; TF2- Theaflavin-3-gallate; TF3- Theaflavin-3'-gallate; TF4- Theaflavin-3,3'-digallate; CL- Colour; BRT- Brightness; AST- Astringency; BRK- Briskness; AR- Aroma; RWC- percent relative water content

Table 4. Association mapping and linkage mapping results for putative QTLs in black tea quality and drought tolerance traits in St.504 and St.524 using GLM (Q), MLM (Q + K) methods (p < 0.01) in TASSEL software v5.2.43 and interval mapping in MapQTL 6.0

Nr	[ac]Trait	Marker	LG	Position (cM)	p-Value	Marker R^2 GLM (Q)	p-Value	Marker R^2 MLM (Q+K)
1	ECG	5128890	1	96.4	1.18E-03	0.042	5.31E-03	0.031
2	EGC	5133866	1	87.1	8.94E-03	0.030	6.62E-04	0.046
3	RWC	5136794	2	60.9	8.58E-03	0.032	3.02E-03	0.033
4	CAT	5063001	4	30.3	2.34E-09	0.098	6.67E-08	0.109
5	EC	5123257	4	26.7	3.89E-08	0.101	7.16E-07	0.089
6	EC	5134490	4	26.4	5.15E-19	0.228	6.21E-15	0.246
7	ECG	5087113	4	17.7	3.74E-03	0.035	2.92E-04	0.048
8	EGC	5123257	4	26.7	9.59E-13	0.162	6.95E-10	0.167
9	TF2	5063001	4	30.3	9.76E-04	0.050	4.22E-03	0.035
10	CL	5132432	7	72.7	1.31E-03	0.047	7.30E-04	0.049
11	ECG	5136790	12	50.4	7.87E-11	0.137	1.58E-04	0.051
12	EGC	5135536	12	50.4	2.25E-04	0.053	2.82E-06	0.086
13	EC	5085963	15	25.4	3.52E-04	0.049	9.09E-03	0.029
14	EC	5099958_E-25	15	25.1	1.00E-04	0.056	6.37E-03	0.031

[ac] Putative QTL identified in both Interval Mapping (MapQTL 6.0), General Linear Model and Mixed Linear Model methods (TASSEL software v5.2.43)

Marker R^2- percentage phenotypic variation explained by markers in GLM and MLM

CAT- Catechin; EC- Epicatechin; ECG- Epicatechin gallate; EGC- Epigallocatechin gallate; TF2- Theaflavin-3-gallate; CL- Colour; RWC- Percent relative water content

Table 5. Association mapping and linkage mapping results for putative QTLs in black tea quality and drought tolerance traits in St.504 and St.524 using GLM (Q), MLM (Q + K) methods (p < 0.01) in TASSEL software v5.2.43 and multiple QTL model mapping in MapQTL 6.0

Nr	[bc]Trait	Marker	LG	Position (cM)	p-Value	Marker R2 GLM (Q)	p-Value	Marker R2 MLM (Q+K)
1	ECG	5128890	1	96.4	1.18E-03	0.042	5.31E-03	0.031
2	EGC	5133866	1	87.1	8.94E-03	0.030	6.62E-04	0.046
3	RWC	5136794	2	60.9	8.58E-03	0.032	3.02E-03	0.033
4	CAT	5063001	4	30.3	2.34E-09	0.098	6.67E-08	0.109
5	TF2	5063001	4	30.3	9.76E-04	0.050	4.22E-03	0.035
6	CL	5132432	7	72.7	1.31E-03	0.047	7.30E-04	0.049
7	ECG	5136790	12	50.4	7.87E-11	0.137	1.58E-04	0.051
8	EC	5085963	15	25.4	3.52E-04	0.049	9.09E-03	0.029

[bc] Putative QTL identified in both multiple QTL model mapping (MapQTL 6.0), General Linear Model and Mixed Linear Model methods (TASSEL software v5.2.43)

Marker R^2- percentage phenotypic variation explained by markers in GLM and MLM

CAT- Catechin; EC- Epicatechin; ECG- Epicatechin gallate; EGC- Epigallocatechin gallate; TF2- Theaflavin-3-gallate; CL- Colour; RWC- Percent relative water content

Table 6. Association mapping and linkage mapping results for putative QTLs in black tea quality and drought tolerance traits in St.504 and St.524 using GLM (Q), MLM (Q + K) methods (p < 0.01) in TASSEL software v5.2.43 and interval mapping and multiple QTL model mapping in MapQTL 6.0

Nr	[abc] Trait	Marker	LG	Position (cM)	p-Value	Marker R2 GLM (Q)	p-Value	Marker R2 MLM (Q+K)	LOD	PEV	Af	Am	D	Allelic effects
1	ECG	5128890	1	96.4	1.18E-03	0.042	5.31E-03	0.031	6.8	11.7	0.326	0.046	0.011	Af
2	EGC	5133866	1	87.1	8.94E-03	0.030	6.62E-04	0.046	3.3	5.8	0.109	0.218	0.054	Am
3	RWC	5136794	2	60.9	8.58E-03	0.032	3.02E-03	0.033	4.3	7.3	0.482	1.13	0.283	Am
4	CAT	5063001	4	30.3	2.34E-09	0.098	6.67E-08	0.109	5.3	4.1	-17.6	-53.45	-13.37	Am
5	EC	5123257	4	26.7	3.89E-08	0.101	7.16E-07	0.089	13.4	21.1	1.703	43.81	10.95	Am
6	ECG	5087113	4	17.7	3.74E-03	0.035	2.92E-04	0.048	3.8	5.8	-2891.86	3715.23	928.81	Af; Am
7	EGC	5123257	4	26.7	9.59E-13	0.162	6.95E-10	0.167	13.4	21.1	1.703	43.81	10.95	Am
8	TF2	5063001	4	30.3	9.76E-04	0.050	4.22E-03	0.035	5.3	4.1	-17.604	-53.47	-13.37	Am
9	CL	5132432	7	72.7	1.31E-03	0.047	7.30E-04	0.049	3.2	6.5	-0.138	-0.118	-0.029	Af; Am
10	ECG	5136790	12	50.4	7.87E-11	0.137	1.58E-04	0.051	15	23.3	-22.37	60.82	15.21	Am
11	EGC	5135536	12	50.4	2.25E-04	0.053	2.82E-06	0.086	15	23.2	-22.32	60.81	15.20	Am
12	EC	5085963	15	25.4	3.52E-04	0.049	9.09E-03	0.029	4.4	7.5	11.67	-24.47	-6.12	Am
13	EC	5099958_E-25	15	25.1	1.00E-04	0.056	6.37E-03	0.031	4.4	7.4	11.69	-24.1	-6.03	Am

[abc] Putative QTL identified using both Interval Mapping and Multiple QTL Model Mapping (MapQTL 6.0) and General Linear Model and Mixed Linear Model method (TASSEL software v5.2.43)

Marker R^2- percentage phenotypic variation explained by markers in GLM and MLM

LG- Linkage group

LOD- Logarithm of odds ratio for putative QTL

PEV- percentage phenotypic variation explained by markers in IM and MQM

Af- Female additivity

Am- Male additivity

D- Dominance

CAT- Catechin; EC- Epicatechin; ECG- Epicatechin gallate; EGC- Epigallocatechin gallate; TF2- Theaflavin-3-gallate; CL- Colour; RWC- Percent relative water content

Table 7. Association mapping, linkage mapping and functional annotation protein of putative QTLs in black tea quality traits in St.504 and St.524 based on IM and MLM (Q + K) method (p < 0.01) in MapQTL 6.0 and TASSEL software v5.2.43, respectively.

Nr	Trait	Marker	LG	Pos (cM)	p-Value	A	p-Value	B	p-Value	Annotated protein	Function
1	[abc]ECG	5128890	1	96.4	1.2E-03	0.042	5.3E-03	0.031	2.0E-25	['Actin']	Cell signalling (response to cold stress, dehydration)
2	[c]CAFF	5137282	2	50.7	2.3E-03	0.042	9.7E-03	0.026	8.0E-09	["N-(5'phosphoribosyl) anthranilate (PRA) isomerase"]	Involved in the first and intermediate pathway in the purine/thiAmine biosynthesis; It also participates in the phenylalanine, tyrosine and tryptophan biosynthesis pathway
3	[abc]ECG	5087113	4	17.7	3.7E-03	0.035	2.9E-04	0.048	6.0E-22	['impB/mucB/sAmB fAmily']	Involved in UV protection through DNA repair
4	[ac]TF1	5106352	4	26	2.0E-04	0.061	8.5E-03	0.032	2.0E-06	['Thiolase, C-terminal domain']	Benzoic acid biosynthesis (volatile compounds)
5	[c]EC	5112438	4	27.4	4.9E-07	0.087	1.5E-05	0.067	9.0E-21	['Phosphoribulokinase / Uridine kinase fAmily']	Carbon fixation (photosynthesis) which are products for shikimate pathway (phenolics)
6	[c]EGC	5112438	4	27.4	1.5E-11	0.148	2.0E-08	0.134	9.0E-21	['Phosphoribulokinase / Uridine kinase fAmily']	Carbon fixation (photosynthesis) which are productsfor shikimate pathway (phenolics)
7	[ac]EC	5134490	4	26.4	5.2E-19	0.228	6.2E-15	0.246	3.0E-08	['Aminotransferase class I and II']	Phenylalanine, tyrosine and tryptophan biosynthesis
8	[ac]EGCG	5134853	4	37.6	7.0E-14	0.205	2.0E-09	0.17	2.0E-06	['Diacylglycerol kinase catalytic domain']	Water, salt, ROS, cold and freezing stress response
9	[c]EC	5136058	4	27.6	1.0E-06	0.083	5.3E-06	0.073	8.0E-09	['Autophagy-related protein 11']	Abiotic stress response (stomatal closure)
10	[ac]EGC	5136058	4	27.6	3.4E-11	0.143	3.6E-09	0.146	8.0E-09	['Autophagy-related protein 11']	Abiotic stress response (stomatal closure)
11	[c]EC	5136410	4	29.9	2.4E-05	0.065	1.8E-05	0.065	7.0E-06	['Protein tyrosine kinase']	ABA signalling
12	[ac]EGCG	5088456	12	47.9	7.7E-04	0.053	2.5E-04	0.06	4.0E-23	['Protein kinase domain']	Abiotic stress response
13	[c]ECG	5136077	12	42.8	1.9E-07	0.094	6.3E-03	0.027	5.0E-13	CSA026168	-

[abc] Putative QTL identified using both Interval Mapping and Multiple QTL Model Mapping (MapQTL 6.0), General Linear Model and Mixed Linear Model (TASSEL software v5.2.43), respectively.

[c] Putative QTL identified in General Linear Model and Mixed Linear Model (MLM) using TASSEL software v5.2.43; A- Marker R^2 GLM (Q); B- Marker R^2 MLM (Q+K)