Combined linkage and association mapping of putative QTLs controlling black tea quality and drought tolerance traits

Robert. K. Koech[1, 2], Richard Mose[3], Samson M. Kamunya[2] and Zeno Apostolides[1]*

[1]Department of Biochemistry, Genetics and Microbiology, University of Pretoria, Pretoria 0002, South Africa

[2]Kenya Agriculture and Livestock Research Organization, Tea Research Institute, P.O. Box 820, Kericho 20200, Kenya

[3]James Finlay (Kenya) Limited, P.O. Box 223, Kericho 20200, Kenya.

*Corresponding author: za@up.ac.za

Abstract

The advancements in genotyping have opened new approaches for identification and precise mapping of quantitative trait loci (QTL) in plants, particularly by combining linkage and association mapping (AM) analysis. In this study, a combination of linkage and AM approach was used to precisely identify and authenticate putative QTLs associated with black tea quality traits and %RWC. The population structure analysis clustered two parents and their respective 261 F_1 progenies from the two reciprocal crosses into two clusters with 141 tea accessions in cluster one and 122 tea accessions in cluster two. The two clusters were of mixed origin with tea accessions in population TRFK St. 504 clustering together with tea accessions in population TRFK St. 524. A total of 70 putative QTLs linked to black tea quality traits and percent relative water contents (%RWC) were detected in interval mapping (IM) method. Cofactors obtained in IM were used in multiple QTL model (MQM) mapping where 40 putative QTLs were detected. The phenotypic variance for each QTL ranged from 2.8-23.3% in IM and 4.1-23% in MQM mapping. Using Q-model and Q+K-model in AM, a total of 49 DArTseq markers were

associated with 16 phenotypic traits. Significant marker-trait association results obtained using

AM were similar to those obtained in IM, and MQM mapping with six more QTLs detected in

AM. The functional annotations of six additional putative QTLs detected in AM were involved

in purine/thiamine biosynthesis, phenylalanine, tyrosine and tryptophan biosynthesis, carbon

fixation and abiotic stress. The application of a combination of linkage and AM approach

appears to have great potential and could reduce the cost and improve the efficacy and accuracy

of selection of desirable traits in tea breeding.

Introduction

Linkage mapping has been a key tool for identifying the genetic basis of quantitative traits in

plants. However, linkage mapping is usually limited by low polymorphism or small population

size, which is required for suitable crosses. Also, the mapping resolution is limited by few

recombination events, which are considered to estimate the genetic distances between marker

loci causative genomic regions for QTLs. AM also known as LD mapping has extensively been

used to circumvent the limitations posed by linkage mapping in plants in the last few years (1).

AM which exploits broader genetic diversity offers more advantages over linkage mapping in

terms of mapping resolution, allele number, saves time in establishing a marker-trait

association and its application in a breeding program (2). Although linkage mapping and AM

each offer more advantages over the other, they are often applied in conjunction to validate the

QTL identified, thus reducing spurious associations.

The integrated approach of linkage-AM has been used in other plants such as *Arabidopsis* (3),

wheat (4) and maize (5) to dissect quantitatively inherited traits. In tree species, an integrated

method of linkage mapping and AM to decipher the nature of genetic architecture of potential

QTLs for growth traits has been reported in poplar hybrids (6) and maritime pine (7). In grapes,

(8) combined linkage mapping and AM to study the genetic patterns of anthocyanin content, a

determinant of berry colour, in grapes. In a study to investigate genetic relationship between

tea caffeine synthase gene (TCS1) and the caffeine content in tea plant and its related species,

AM was used (9). In this current study, we combined the high QTL detection power of the

linkage mapping with the high-resolution power of AM. The linkage-AM approach will not

only accelerate the pace of QTL mapping in tea breeding improvement but will also precisely

identify reliable QTLs linked to black tea quality and drought tolerance in tea. The objective

of this study was to combine linkage mapping and AM to precisely identify and authenticate

putative QTLs linked to caffeine content, catechins fractions, theaflavins fractions, tea tasters'

scores, and %RWC.

Material and methods

Population type

60 Two pseudo-testcross populations used in this current study consisted of 109 F_1 progeny from

61 TRFK St. 504 (TRFK 303/577 x GW Ejulu) and 152 F_1 progeny from TRFK St. 524 (GW

62 Ejulu x TRFK 303/577) as described previously (10).

Phenotypic data

64 A total of 16 phenotypic traits were used as described previously (10).

Population structure

The population structure was inferred from the 1,421 DArT markers data previously used to

construct genetic linkage map using the STRUCTURE software version 2.3.4 (11). Twenty

independent runs were carried out using the following parameters; number of populations (K)

from 1 to 10, burn-in time and Markov Chain Monte Carlo (MCMC) replication number were

both set to 100,000 for model of admixture with correlated allele frequencies. The natural

logarithms of the probability data (LnP(K)) and the *ad hoc* delta K statistical were calculated

using STRUCTURE Harvester (12), and the optimal K according to the delta K value was then selected (13). Finally, the population structure matrix (Q) was obtained by integrating 20 independent replicate runs and applying CLUMPP software (14). STRUCTURE bar plots were plotted using STRUCTURE Plot v 2.0 (15).

Linkage mapping, QTL analyses and allelic effects

Six thousand five hundred and eighty-eight DArTseq markers derived from *C. sinensis* genomic DNA of the two parents and 261 F_1 progeny were subjected to linkage mapping analyses using JoinMap 4.0 (16) and QTL mapping analyses using MapQTL 6.0 software (17) as described previously (Koech *et al.*, 2018).

The allelic effects were estimated as $A_f=[(\mu ac+\mu ad)-(\mu bc+\mu bd)]/4$ for female additivity, $A_m=[(\mu ac+\mu bc)-(\mu ad+\mu bd)]/4$ for male additivity and $D=[(\mu ac+\mu bd)-(\mu ad+\mu bc)]/4$ for dominance where μac, μad, μbc and μbd are estimated phenotypic means associated to each of the four possible genotypic classes ac, bc, ad and bd, derived from ab×cd cross (18).

Association mapping

Using the Q and K matrices data files from STRUCTURE software as a covariates, 1,421 DArT markers of the 261 F_1 progeny were tested for association with each phenotype using a general linear model (GLM), mixed linear model (MLM) that included kinship, phylogenetic tree in TASSEL (trait analysis by association, evolution and linkage) Version 5.2.43 software (19). Principal component analysis (PCA) was performed using JMP Pro 14 to visualise the dispersion of the association panel in a graph. The Q and kinship (K) matrices were used to correct the effects of population substructure in the association panel which can cause false-positive associations. The p-value and R^2 were used to determine whether a QTL is associated with the marker and the magnitude of QTL effects, respectively.

Results

Frequency distribution

The frequency distribution results for 16 phenotypic traits were described previously (10).

Population structure

A total of 261 tea accessions, 109 for population TRFK St. 504 and 152 for population TRFK St.524 were combined to construct a panel to study their genetic structure. The two tea populations, TRFK St. 504 and TRFK St. 524 are a reciprocal cross of two parents TRFK 303/577 and GW Ejulu. A population structure analysis showed that the value of Evanno's ΔK presented a sharp spike at $K = 2$, which suggested that this population panel was clustered into two groups (Fig 1). The two groups or clusters are represented by two clades with parental clone TRFK 303/577 clustering with 141 progenies in cluster one (blue) and parental clone GW Ejulu (not shown) clustering with 122 progenies in cluster two (black), respectively (Fig 2). The two clusters were accession with mixed origin with tea accessions in TRFK St. 504 population clustering together with tea accessions in TRFK St. 524 population. Furthermore, the results from PCA (Fig 3) and a phylogenetic tree-based on Nei's genetic distance was in agreement with the structure analysis results (Fig 4). The population structure of the two tea populations was examined with the neighbour-joining algorithm using Euclidian distance on the DArTseq marker band intensities.

Fig 1. Delta K values plotted from 1 to 10 for 261 tea accessions panel.

Fig 2. Population structure of 261 tea accessions panel (K = 2)

Fig 3. Principal component analysis of 261 tea accessions. represents cluster 1, represents cluster 2

Fig 4. Phylogenetic tree-based on Nei's genetic distance. Blue clade represents cluster 1,

black clade represents cluster 2.

Linkage mapping and association mapping

Interval mapping and allelic effects

The genetic linkage map was constructed using a total of 1,421 DArT markers as described

previously (10). A total of 70 putative QTLs were identified for the 16 phenotypic traits

measured using the interval mapping (Table 1). Of all the putative QTLs identified, most QTLs

were found in almost all linkage groups, except LG03, LG05 and LG11. For black tea quality

traits, 67 putative QTLs were detected and were located on almost all linkage groups, except

on LG03, LG05 and LG11. The remaining three putative QTLs which are associated with

drought tolerance were identified in LG02, LG06 and LG09. Eight putative QTLs were

identified on LG01, seven in LG02 and LG12, 10 in LG04, five in LG06 and LG15, three in

LG07, LG08, LG09, two in LG10, 15 in LG13, four in LG15, respectively. The highest number

of putative QTLs were associated with catechin (13), EGC (12), tea tasters' score (12), caffeine

(10) and EGC (9), respectively. The phenotypic variance for all the identified putative QTLs

ranged from 2.8% for EGC to 23.3% for qECG, respectively. The phenotypic variance for

%RWC ranged from 5.7 to 7.3%, while for the tea tasters' score, it ranged from 5.8 to 9.1%.

Half of the allelic effects identified to be associated with black tea quality and drought tolerance

traits had a positive additive effect. There were eight allelic effects associated with female

additivity, and 18 were associated with male additive effects, respectively. The highest positive

female and male additive effects were qEGC and qECG, respectively. Although, some putative

QTLs also exhibited high negative allelic effects, which were from either female or male

additive effects or both, such as qEGCG, qCAT, qEC and qCaffeine (Table 1). All the putative

QTLs identified in IM were used as cofactors in MQM mapping.

141 Table 1. Linkage mapping results for putative QTLs in black tea quality and drought tolerance traits in St. 504 and St. 524 using interval
142 mapping in MapQTL 6.0

Nr	[a]Trait	Locus	LG	Position (cM)	LOD	PEV	Af	Am	D	Allelic effect
1	AST	5115441_E-26	1	96.6	4.5	9.1	0.224	-0.061	-0.015	Af
2	AST	5135087	1	97.3	4.5	9.1	0.234	-0.057	-0.014	Af
3	BRK	5115441_E-26	1	96.6	3.6	7.3	0.196	-0.082	-0.020	Af
4	BRK	5135087	1	97.3	3.5	7.2	0.204	-0.078	-0.019	Af
5	ECG	5128890	1	96.4	6.8	11.7	0.326	0.046	0.011	Af
6	ECG	5135087	1	97.3	6.6	11.3	0.347	0.053	0.013	Af
7	ECG	5115441_E-26	1	96.6	6.5	11.1	0.331	0.051	0.013	Af
8	EGC	5133866	1	87.1	3.3	5.8	0.109	0.218	0.054	Am
9	CAFF	5064585	2	50.9	3.3	5.8	5.849	-9.702	-2.425	Af; Am
10	CAT	5135436	2	0.0	3.6	6.1	-0.838	-11.195	-2.799	Am
11	EC	5072338	2	2.2	4.3	7.2	-14.648	-12.555	-3.139	Af; Am
12	EGC	5124128	2	7.7	3.3	5.6	-36.447	1.544	0.386	Af
13	RWC	5136794	2	60.9	4.3	7.3	0.482	1.130	0.283	Am
14	TF1	5084595	2	4.5	3.2	5.7	-0.952	8.158	2.039	Am
15	TF2	5084595	2	4.5	4.1	7.3	2.290	9.140	2.285	Am
16	AR	100158044|F|0	4	70.1	2.9	6	-0.268	0.090	0.023	Af
17	CAFF	5112599	4	68.6	3.8	6.7	2.413	11.852	2.963	Am
18	CAT	5063001	4	30.3	5.3	4.1	-17.604	-53.467	-13.367	Am
19	EC	5123475	4	27.2	14.0	21.9	3.485	44.620	11.155	Am
20	EC	5123257	4	26.7	13.4	21.1	1.703	43.813	10.953	Am
21	EC	5134490	4	26.4	12.4	19.6	0.260	41.569	7.005	Am
22	ECG	5087113	4	17.7	3.8	5.8	-2891.857	3715.230	928.808	Af; Am
23	EGC	5123475	4	27.2	3.7	2.8	7071.043	-6846.394	-1711.598	Af; Am
24	EGCG	5087017	4	37.2	4.9	6	-320.975	-200.380	-50.095	Af; Am
25	ECG	5098382	6	57.0	5.2	8.7	35.650	-32.760	-8.190	Af; Am
26	EGC	5073424	6	66.4	3.9	6.7	-13.905	14.749	3.687	Af; Am
27	EGC	5124993	6	66.6	3.9	6.6	-12.321	15.635	3.909	Af; Am
28	RWC	5082606	6	66.2	3.3	5.7	-2.094	-0.814	-0.203	Af
29	TF1	5136045	6	69.6	4.3	7.7	2.795	3.782	0.946	Af; Am
30	CAFF	5064391	7	48.1	3.4	6	-12.264	-3.397	-0.849	Af
31	CAFF	5056614	7	48.3	3.3	5.9	-12.029	-3.985	-0.996	Af
32	CL	5132432	7	72.7	3.2	6.5	-0.138	-0.118	-0.029	Af; Am
33	CAFF	5134558	8	18.8	4.3	7.5	17.375	-12.332	-3.083	Af; Am
34	CAFF	5111497	8	18.8	4.1	7.1	11.542	-12.927	-3.232	Af; Am
35	CAT	5130194	8	12.5	3.9	6.6	-29.556	-9.720	-2.430	Af

36	AST	5123950	9	87.6	3.2	6.6	-0.036	0.178	0.045	Am
37	BRK	5123950	9	87.6	3.6	7.4	-0.012	0.190	0.047	Am
38	RWC	5130531	9	6.7	4.1	7	1.933	2.876	0.719	Am
39	AR	5075627	10	26.9	3.4	6.9	-0.064	0.110	0.027	Am
40	ECG	5136108	10	20.6	3.3	5.7	4.297	34.819	8.705	Am
41	CAT	5123751	12	43.0	4.2	6.5	-36.478	-3.162	-0.791	Af
42	CAT	5127224	12	42.9	4.2	6.5	-36.467	-3.045	-0.761	Af
43	ECG	5136790	12	50.4	15.0	23.3	-22.365	60.821	15.205	Am
44	ECG	5135536	12	50.4	15.0	23.2	-22.316	60.808	15.202	Am
45	EGC	5123751	12	43.0	5.0	8.5	50.465	-30.542	-7.635	Af; Am
46	EGC	5127224	12	42.9	5.0	8.4	50.241	-30.563	-7.641	Af; Am
47	EGCG	5104630	12	48.2	4.2	7.2	18.115	-52.080	-13.020	Am
48	BRT	5088162	13	29.8	3.0	6.1	0.006	-0.142	-0.035	Am
49	BRT	5135810	13	26.0	2.8	5.8	0.007	-0.137	-0.034	Am
50	CAFF	5088162	13	29.8	3.4	5.4	7.183	-29.136	-7.284	Am
51	CAFF	5080631	13	35.6	2.9	3.1	-14.651	23.420	5.855	Af; Am
52	CAT	5122819	13	10.2	4.3	7.3	-27.341	-14.521	-3.630	Af; Am
53	CAT	5111268	13	58.1	4.6	7.2	14.200	17.804	4.451	Af; Am
54	CAT	5129729	13	9.8	4.2	7.2	-26.752	-14.467	-3.617	Af; Am
55	CAT	5115793	13	58.7	4.3	6.8	12.635	17.248	4.312	Af; Am
56	CAT	5136782	13	58.5	4.2	6.6	12.029	17.237	4.309	Af; Am
57	CAT	5103784	13	50.6	3.8	6.1	4.188	21.966	5.491	Am
58	CAT	5122899	13	50.5	3.8	6	3.656	21.921	5.480	Am
59	ECG	5123761	13	60.7	3.3	5.8	-0.469	-0.112	-0.028	Af
60	ECG	5072523	13	60.3	3.0	5.2	-0.261	-0.231	-0.058	Af; Am
61	ECG	5114985	13	61.6	2.8	5	-0.439	-0.088	-0.022	Af
62	EGC	5136623	13	50.6	3.7	6.4	8.946	-29.764	-7.441	Am
63	BRT	5125626	14	71.1	3.3	6.8	0.000	0.143	0.036	Am
64	CAFF	5123053	14	5.5	4.0	7.1	-7.622	-11.572	-2.893	Af; Am
65	CAFF	5054639	14	6.4	3.9	6.9	-7.260	-11.533	-2.883	Af; Am
66	CAT	5132370	14	60.7	6.5	10.8	-19.945	35.937	8.984	Af; Am
67	EGC	5132370	14	60.7	3.7	6.4	46.492	-7.514	-1.878	Af
68	EC	5085963	15	25.4	4.4	7.5	11.674	-24.472	-6.118	Am
69	EC	5099958_E-25	15	25.1	4.4	7.4	11.686	-24.101	-6.025	Am
70	ECG	5111164	15	75.1	4.2	7.2	22.426	34.202	8.550	Af; Am
71	EGCG	5114089	15	32.1	4.0	6.8	-1.178	56.567	14.142	Am

143   [a] Putative QTL identified in Interval Mapping using MapQTL 6.0. LOD- Logarithm of odds for putative QTL in IM (LOD>2); PEV- percentage phenotypic

144   variation explained by markers in IM; Af- Female additivity; Am- Male additivity; D- Dominance

Multiple QTL model mapping and allelic effects

A total of 40 putative QTLs (LOD > 3.0) were identified in 16 phenotypic traits using MQM mapping method from the selected cofactors in IM. Similarly, no putative QTLs in LG03, LG05 and LG11, respectively were associated with any of the phenotypic traits. Most of the putative QTLs previously identified in IM were also identified MQM mapping. Of these 40 putative QTLs identified, 37 putative QTLs that were associated with black tea quality traits while three were associated with qRWC (Table 2). The three putative QTLs associated with qRWC were similar to those identified in IM. Four putative QTLs were identified in LG01 and LG12, seven in LG02, five in LG04, three in LG06, LG09, LG13, LG14 and LG15, two in LG07 and LG08 and one in LG01, respectively. A similar trend was observed in the number of putative QTLs for each trait as in IM with a high putative QTL number for catechin, ECG, EGC and caffeine, respectively. The phenotypic variance for all the identified putative QTL ranged from 4.1% for qCAT to 23% for qECG, respectively. The phenotypic variance for qRWC was similar to IM results, and it ranged from 5.7 to 7.3%, while for the tea tasters' score, it ranged from 6.0 to 9.1%. For allelic effects, similar results to IM with qEGC and qECG showing high positive female additive and male additive effects, respectively, this trend was also observed in MQM mapping. The putative QTLs identified using both IM, and MQM mapping were almost similar although the efficiency and the accuracy of detecting QTL is achieved by employing MQM instead of the single QTL model used in IM.

Table 2. Linkage mapping for putative QTLs in black tea quality and drought tolerance traits in St. 504 and St. 524 using multiple QTL model mapping in MapQTL 6.0

Nr	bTrait	Locus	LG	Position (cM)	LOD	PEV	Af	Am	D	Allelic effect
1	AST	5135087	1	97.3	4.5	9.1	0.234	-0.057	-0.014	Af
2	BRK	5115441_E-26	1	96.6	3.6	7.3	0.196	-0.082	-0.020	Af
3	ECG	5128890	1	96.4	6.8	11.7	0.326	0.046	0.011	Af
4	EGC	5133866	1	87.1	3.3	5.8	0.109	0.218	0.054	Am
5	CAFF	5064585	2	50.9	3.3	5.8	5.849	-9.702	-2.425	Af; Am
6	CAT	5135436	2	0.0	3.6	6.1	-0.838	-11.195	-2.799	Am
7	EC	5072338	2	2.2	4.3	7.2	-14.648	-12.555	-3.139	Af; Am
8	EGC	5124128	2	7.7	3.3	5.6	-36.447	1.544	0.386	Af
9	RWC	5136794	2	60.9	4.3	7.3	0.482	1.130	0.283	Am
10	TF1	5084595	2	4.5	3.2	5.7	-0.952	8.158	2.039	Am
11	TF2	5084595	2	4.5	4.1	7.3	2.290	9.140	2.285	Am
12	AR	100158044|F|0	4	70.1	2.9	6	-0.268	0.090	0.023	Af
13	CAFF	5112599	4	68.6	3.8	6.7	2.413	11.852	2.963	Am
14	CAT	5063001	4	30.3	5.3	4.1	-17.604	-53.467	-13.367	Am
15	EC	5123475	4	27.2	14.0	21.9	3.485	44.620	11.155	Am
16	EGCG	5087017	4	37.2	4.9	6	-320.975	-200.380	-50.095	Af; Am
17	ECG	5098382	6	57.0	5.2	8.7	35.650	-32.760	-8.190	Af; Am
18	EGC	5073424	6	66.4	3.9	6.7	-13.905	14.749	3.687	Af; Am
19	RWC	5082606	6	66.2	3.3	5.7	-2.094	-0.814	-0.203	Af
20	CAFF	5064391	7	48.1	3.4	6	-12.264	-3.397	-0.849	Af
21	CL	5132432	7	72.7	3.2	6.5	-0.138	-0.118	-0.029	Af; Am
22	CAFF	5134558	8	18.8	4.3	7.5	17.375	-12.332	-3.083	Af; Am
23	CAT	5130194	8	12.5	3.9	6.6	-29.556	-9.720	-2.430	Af
24	AST	5123950	9	87.6	3.2	6.6	-0.036	0.178	0.045	Am
25	BRK	5123950	9	87.6	3.6	7.4	-0.012	0.190	0.047	Am
26	RWC	5130531	9	6.7	4.1	7	1.933	2.876	0.719	Am
27	ECG	5136108	10	20.6	3.3	5.7	4.297	34.819	8.705	Am
28	CAT	5123751	12	43.0	4.2	6.5	-36.478	-3.162	-0.791	Af
29	ECG	5136790	12	50.4	15.0	23.3	-22.365	60.821	15.205	Am
30	EGC	5123751	12	43.0	5.0	8.5	50.465	-30.542	-7.635	Af; Am
31	EGCG	5104630	12	48.2	4.2	7.2	18.115	-52.080	-13.020	Am
32	CAFF	5088162	13	29.8	3.4	5.4	7.183	-29.136	-7.284	Am

33	CAT	5111268	13	58.1	4.6	7.2	14.200	17.804	4.451	Af; Am
34	EGC	5136623	13	50.6	3.7	6.4	8.946	-29.764	-7.441	Am
35	BRT	5125626	14	71.1	3.3	6.8	0.000	0.143	0.036	Am
36	CAFF	5123053	14	5.5	4.0	7.1	-7.622	-11.572	-2.893	Af; m
37	CAT	5132370	14	60.7	6.5	10.8	-19.945	35.937	8.984	Af; Am
38	EC	5085963	15	25.4	4.4	7.5	11.674	-24.472	-6.118	Am
39	ECG	5111164	15	75.1	4.2	7.2	22.426	34.202	8.550	Af; Am
40	EGCG	5114089	15	32.1	4.0	6.8	-1.178	56.567	14.142	Am

[b] Putative QTL identified in Multiple QTL Model Mapping using MapQTL 6.0

LOD- Logarithm of odds for putative QTL in MQM (LOD>3); PEV- percentage phenotypic variation explained by markers in MQM; Af- Female additivity;

Am- Male additivity; D- Dominance

CAFF- Caffeine; CAT- Catechin; EC- Epicatechin; ECG- Epicatechin gallate; EGC- Epigallocatechin gallate; EGCG- Epigallocatechin gallate; TF1-

Theaflavin; TF2- Theaflavin-3-gallate; TF3- Theaflavin-3'-gallate; TF4- Theaflavin-3,3'-digallate; CL- Colour; BRT- Brightness; AST- Astringency; BRK-

Briskness; AR- Aroma; RWC- percent relative water content

Association mapping

Four models GLM, GLM (Q), mixed linear model with kinship matrix (MLM) (K) and mixed linear model with kinship matix and population structure matrix (MLM) (Q+K) in TASSEL software v5.2.43 were used to determine AM and the effects of population structure on the AM to reduce the inflation of false-positive associations. The p-values were plotted cumulatively for each model, and the distribution examined. However, no further analysis was done using GLM model since marker-trait association results were characterised by excess small p-values, which indicated an abundance of spurious associations (S1 Fig). The association analysis using the Q-model and Q+K-model detected a total of 49 DArT markers associated with 15 different black tea quality traits and one drought-tolerant trait. The GLM (Q) model showed associations between DArT markers and traits ($p < 0.01$) and was confirmed using the MLM (Q+K) model (Table 3, S2-S4 Figs). The MLM Q+K-model (MLM with Q-matrix and K-matrix as a correction for population structure) showed a good fit for the p-values ($p < 0.01$), as compared to GLM (Q) models, which were characterised by a few associations with excess of small p-values, which indicates abundance of spurious associations (Table 3, S2 and S4 Figs). Also, the MLM Q+K-model also showed better small p-values than the K-model (MLM with K-matrix as a correction for population structure) (S3 and S4 Figs). The GLM (Q) model may not have accounted for the heterogeneity of the genetic background in some tea cultivars under study, which may have resulted in false-positive associations. Ideally, the distribution of p-values should follow a uniform distribution with less deviation from the expected p-values. However, the two models were used to compare the effect of population matrix and kinship matrix on GLM and MLM, respectively, in associating mapping. Therefore, taking into account the performance of the two different models, the results from the MLM (Q+K) model appeared to have controlled better population structure and kinship relationships than GLM (Q) model. While it might be tempting to consider p-values that remain extreme after MLM (Q+K) model

correction as true associations, we need to keep in mind that minor allele frequency also

influences p-values.

A total of 49 DArT markers were associated with 16 different phenotypic traits in six out of 15

linkage groups (Table 3). Since, all 1,421 DArT markers in the genotypic dataset had a known

map location, the loci associated with particular traits could be allocated to a linkage group.

The numbers of marker loci associated with traits together with their location and positions in

different linkage groups at a test level of $p< 0.01$ in the two test models are shown in Table 3.

Choosing a significance level of $p< 0.01$ which involves multiple testing corrections in both

GLM (Q) and MLM (Q+K) models, 32 DArT marker loci were detected for all individual

catechins (C, EC, ECG, EGC, EGCG), scattered over five linkage groups namely LG01, LG02,

LG04, LG12 and LG15 (Table 3). For caffeine trait, three DArT markers were only located in

LG02 while 11 DArT markers for all individual theaflavins traits (TF1, TF2, TF3, TF4) were

detected in LG04 and LG15 (Table 3). The trait for tea liquor colour which is an indicator for

black tea quality in terms of tea tasters' score appeared to be associated with only one DArT

marker locus at the significance level of $p< 0.01$ in LG07 (Table 3). Drought tolerance trait

(%RWC) was also associated with only one DArT marker locus detected in LG02 (Table 3).

The majority of the DArT markers loci associated with the traits were found in LG04, LG12,

LG15, LG02, and LG01. The three DArT markers associated with catechin trait in LG04,

namely, 5136945, 5064764 and 5134490 had the highest proportion of the percent explained

phenotypic variation of 41%, 38%, and 25%, respectively. The DArT markers that also showed

high proportion of percent explained phenotypic variation were also observed in LG04 for

qEGCG, qEGC, and qTF1 traits, respectively. In this study, a pleiotropic locus which is

associated with a single locus affecting two or more distinct phenotypic traits was observed.

Several pleiotropic loci were identified including DArT markers 5136945, 5112438, 5136058,

5123257, 5106022 and 5064764 which were associated with qC, qEC, qEGC, qTF1, qTF2 and

qTF4 phenotypic traits (Table 3). All the pleiotropic loci detected and associated with the

respective phenotypic traits were located in LG04.

Table 3. Association mapping results for black tea quality and drought tolerance traits in St. 504 and St.524 using the GLM (Q) and MLM (Q+K) method, respectively (p <0.01)

Nr	cTrait	Marker	LG	Position (cM)	p-Value	Marker R^2 GLM (Q)	p-Value	Marker R^2 MLM (Q+K)
1	ECG	5128890	1	96.4	1.18E-03	0.042	5.31E-03	0.031
2	EGC	5086476	1	87.5	8.20E-03	0.031	8.54E-04	0.043
3	EGC	5122985	1	87.5	1.47E-04	0.056	2.41E-03	0.038
4	EGC	5133866	1	87.1	8.94E-03	0.030	6.62E-04	0.046
5	CAFF	5134416	2	44.4	2.46E-04	0.057	5.58E-04	0.050
6	CAFF	5136551	2	43.3	9.60E-04	0.048	7.51E-03	0.030
7	CAFF	5137282	2	50.7	2.27E-03	0.042	9.72E-03	0.026
8	EC	5132500	2	7.9	2.91E-03	0.036	1.37E-04	0.052
9	RWC	5136794	2	60.9	8.58E-03	0.032	3.02E-03	0.033
10	CAT	5063001	4	30.3	2.34E-09	0.098	6.67E-08	0.109
11	CAT	5064764	4	28.8	6.42E-37	0.326	7.80E-22	0.379
12	CAT	5136945	4	28.1	6.39E-40	0.345	1.81E-21	0.406
13	EC	5112438	4	27.4	4.90E-07	0.087	1.50E-05	0.067
14	EC	5123257	4	26.7	3.89E-08	0.101	7.16E-07	0.089
15	EC	5134490	4	26.4	5.15E-19	0.228	6.21E-15	0.246
16	EC	5136058	4	27.6	1.00E-06	0.083	5.31E-06	0.073
17	EC	5136410	4	29.9	2.39E-05	0.065	1.77E-05	0.065
18	ECG	5087113	4	17.7	3.74E-03	0.035	2.92E-04	0.048
19	EGC	5112438	4	27.4	1.50E-11	0.148	1.95E-08	0.134
20	EGC	5123257	4	26.7	9.59E-13	0.162	6.95E-10	0.167
21	EGC	5136058	4	27.6	3.43E-11	0.143	3.60E-09	0.146
22	EGCG	5074553	4	39.5	3.26E-14	0.210	1.86E-12	0.221
23	EGCG	5123463	4	28.9	3.38E-17	0.250	1.09E-13	0.266
24	EGCG	5134853	4	37.6	7.03E-14	0.205	2.00E-09	0.170
25	EGCG	5136554	4	38.7	1.46E-16	0.242	2.08E-11	0.222
26	EGCG	100011901|F|0	4	38.7	1.24E-04	0.066	3.13E-03	0.039
27	TF1	5064764	4	28.8	2.35E-10	0.149	5.78E-06	0.088
28	TF1	5106352	4	26.0	1.95E-04	0.061	8.45E-03	0.032
29	TF1	5123463	4	28.9	1.33E-10	0.153	2.66E-06	0.097
30	TF1	5136945	4	28.1	1.69E-11	0.166	8.20E-07	0.107
31	TF2	5063001	4	30.3	9.76E-04	0.050	4.22E-03	0.035
32	TF2	5064764	4	28.8	1.85E-08	0.122	7.36E-05	0.063

33	TF2	5106022	4	29.1	2.99E-09	0.134	3.06E-04	0.052
34	TF4	5064764	4	28.8	8.64E-06	0.082	1.33E-03	0.042
35	TF4	5106022	4	29.1	1.38E-06	0.094	1.57E-03	0.040
36	TF4	5136945	4	28.1	6.26E-07	0.099	1.59E-04	0.060
37	CL	5132432	7	72.7	1.31E-03	0.047	7.30E-04	0.049
38	ECG	5059563	12	49.0	8.91E-09	0.111	2.29E-05	0.068
39	ECG	5136077	12	42.8	1.91E-07	0.094	6.34E-03	0.027
40	ECG	5136790	12	50.4	7.87E-11	0.137	1.58E-04	0.051
41	EGC	5135536	12	50.4	2.25E-04	0.053	2.82E-06	0.086
42	EGCG	5087581	12	48.3	1.51E-04	0.065	1.75E-03	0.040
43	EGCG	5088456	12	47.9	7.70E-04	0.053	2.46E-04	0.060
44	EGCG	5133629	12	47.7	7.17E-04	0.054	1.95E-03	0.041
45	EC	5085963	15	25.4	3.52E-04	0.049	9.09E-03	0.029
46	EC	5132852	15	24.8	1.09E-04	0.056	2.50E-03	0.036
47	EC	5099958_E-25	15	25.1	1.00E-04	0.056	6.37E-03	0.031
48	TF3	5083854	15	15.2	3.52E-03	0.043	2.77E-03	0.035
49	TF3	5136247	15	16.3	1.57E-03	0.049	5.40E-04	0.046

[c] Putative QTL identified in General Linear Model and Mixed Linear Model methods using TASSEL software v5.2.43

LG- Linkage group

Marker R^2- percentage phenotypic variation explained by markers in both GLM (Q) and MLM (Q+K) methods

Q- Population structure matrix

K- Kinship matrix

CAFF- Caffeine; CAT- Catechin; EC- Epicatechin; ECG- Epicatechin gallate; EGC- Epigallocatechin gallate; EGCG- Epigallocatechin gallate; TF1-

Theaflavin; TF2- Theaflavin-3-gallate; TF3- Theaflavin-3'-gallate; TF4- Theaflavin-3,3'-digallate; CL- Colour; BRT- Brightness; AST- Astringency;

BRK- Briskness; AR- Aroma; RWC- percent relative water content

A comparison of linkage mapping and association mapping

The putative QTLs detected using IM, and MQM mapping in MapQTL 6.0 were compared with marker-trait association results obtained using GLM (Q) and MLM (Q+K) models in AM. The significant marker-trait association results obtained using AM were similar to with the results obtained using MapQTL 6.0 (Tables 4 and 5). The IM analysis produced 14 significant QTLs (LOD > 3.0) that were also found to be associated with the phenotypic traits in AM analysis (Table 4). From the 14 QTLs, a total of 11 markers were associated with four individual catechins (qCAT, qEC, qECG, and qEGC) and one marker each was associated with qTF2, qCL, and qRWC, respectively. On the other hand, the MQM mapping analysis produced eight significant QTLs (LOD > 3.0) which were also found to be significantly associated with the phenotypic traits in AM analysis (Table 5). Of these, five markers were associated with four individual catechins (qCAT, qEC, qECG, and qEGC) and one marker each was associated with qTF2, qCL, and qRWC, respectively. Furthermore, the four mapping models were compared with each other to find markers that could associate with phenotypic traits in all four mapping models (Table 6). A total of 13 markers were found to be associated with four individual catechins (qCAT, qEC, qECG, and qEGC) and one marker each was associated with qTF2, qCL, and qRWC, respectively. Most of the markers that were associated with different phenotypic traits were only located in LG01, LG02, LG04, LG12 and LG15 with a majority of markers showing both positive and negative male additive effects.

261 Table 4. Association mapping and linkage mapping results for putative QTLs in black tea quality and drought tolerance traits in St.504 and St.524
262 using GLM (Q), MLM (Q + K) methods (p < 0.01) in TASSEL software v5.2.43 and interval mapping in MapQTL 6.0

Nr	[ac]Trait	Marker	LG	Position (cM)	p-Value	Marker R^2 GLM (Q)	p-Value	Marker R^2 MLM (Q+K)
1	ECG	5128890	1	96.4	1.18E-03	0.042	5.31E-03	0.031
2	EGC	5133866	1	87.1	8.94E-03	0.030	6.62E-04	0.046
3	RWC	5136794	2	60.9	8.58E-03	0.032	3.02E-03	0.033
4	CAT	5063001	4	30.3	2.34E-09	0.098	6.67E-08	0.109
5	EC	5123257	4	26.7	3.89E-08	0.101	7.16E-07	0.089
6	EC	5134490	4	26.4	5.15E-19	0.228	6.21E-15	0.246
7	ECG	5087113	4	17.7	3.74E-03	0.035	2.92E-04	0.048
8	EGC	5123257	4	26.7	9.59E-13	0.162	6.95E-10	0.167
9	TF2	5063001	4	30.3	9.76E-04	0.050	4.22E-03	0.035
10	CL	5132432	7	72.7	1.31E-03	0.047	7.30E-04	0.049
11	ECG	5136790	12	50.4	7.87E-11	0.137	1.58E-04	0.051
12	EGC	5135536	12	50.4	2.25E-04	0.053	2.82E-06	0.086
13	EC	5085963	15	25.4	3.52E-04	0.049	9.09E-03	0.029
14	EC	5099958_E-25	15	25.1	1.00E-04	0.056	6.37E-03	0.031

[ac] Putative QTL identified in both Interval Mapping (MapQTL 6.0), General Linear Model and Mixed Linear Model methods (TASSEL software v5.2.43)

Marker R^2- percentage phenotypic variation explained by markers in GLM and MLM

CAT- Catechin; EC- Epicatechin; ECG- Epicatechin gallate; EGC- Epigallocatechin gallate; TF2- Theaflavin-3-gallate; CL- Colour; RWC- Percent
relative water content

Table 5. Association mapping and linkage mapping results for putative QTLs in black tea quality and drought tolerance traits in St.504 and St.524
using GLM (Q), MLM (Q + K) methods (p < 0.01) in TASSEL software v5.2.43 and multiple QTL model mapping in MapQTL 6.0

Nr	[bc]Trait	Marker	LG	Position (cM)	p-Value	Marker R2 GLM (Q)	p-Value	Marker R2 MLM (Q+K)
1	ECG	5128890	1	96.4	1.18E-03	0.042	5.31E-03	0.031
2	EGC	5133866	1	87.1	8.94E-03	0.030	6.62E-04	0.046
3	RWC	5136794	2	60.9	8.58E-03	0.032	3.02E-03	0.033
4	CAT	5063001	4	30.3	2.34E-09	0.098	6.67E-08	0.109
5	TF2	5063001	4	30.3	9.76E-04	0.050	4.22E-03	0.035
6	CL	5132432	7	72.7	1.31E-03	0.047	7.30E-04	0.049
7	ECG	5136790	12	50.4	7.87E-11	0.137	1.58E-04	0.051
8	EC	5085963	15	25.4	3.52E-04	0.049	9.09E-03	0.029

[bc] Putative QTL identified in both multiple QTL model mapping (MapQTL 6.0), General Linear Model and Mixed Linear Model methods (TASSEL
software v5.2.43)

Marker R^2- percentage phenotypic variation explained by markers in GLM and MLM

CAT- Catechin; EC- Epicatechin; ECG- Epicatechin gallate; EGC- Epigallocatechin gallate; TF2- Theaflavin-3-gallate; CL- Colour; RWC- Percent
relative water content

Table 6. Association mapping and linkage mapping results for putative QTLs in black tea quality and drought tolerance traits in St.504 and St.524
using GLM (Q), MLM (Q + K) methods (p < 0.01) in TASSEL software v5.2.43 and interval mapping and multiple QTL model mapping in
MapQTL 6.0

Nr	[abc] Trait	Marker	LG	Position (cM)	p-Value	Marker R2 GLM (Q)	p-Value	Marker R2 MLM (Q+K)	LOD	PEV	Af	Am	D	Allelic effects
1	ECG	5128890	1	96.4	1.18E-03	0.042	5.31E-03	0.031	6.8	11.7	0.326	0.046	0.011	Af
2	EGC	5133866	1	87.1	8.94E-03	0.030	6.62E-04	0.046	3.3	5.8	0.109	0.218	0.054	Am
3	RWC	5136794	2	60.9	8.58E-03	0.032	3.02E-03	0.033	4.3	7.3	0.482	1.13	0.283	Am
4	CAT	5063001	4	30.3	2.34E-09	0.098	6.67E-08	0.109	5.3	4.1	-17.6	-53.45	-13.37	Am
5	EC	5123257	4	26.7	3.89E-08	0.101	7.16E-07	0.089	13.4	21.1	1.703	43.81	10.95	Am
6	ECG	5087113	4	17.7	3.74E-03	0.035	2.92E-04	0.048	3.8	5.8	-2891.86	3715.23	928.81	Af; Am
7	EGC	5123257	4	26.7	9.59E-13	0.162	6.95E-10	0.167	13.4	21.1	1.703	43.81	10.95	Am
8	TF2	5063001	4	30.3	9.76E-04	0.050	4.22E-03	0.035	5.3	4.1	-17.604	-53.47	-13.37	Am
9	CL	5132432	7	72.7	1.31E-03	0.047	7.30E-04	0.049	3.2	6.5	-0.138	-0.118	-0.029	Af; Am
10	ECG	5136790	12	50.4	7.87E-11	0.137	1.58E-04	0.051	15	23.3	-22.37	60.82	15.21	Am
11	EGC	5135536	12	50.4	2.25E-04	0.053	2.82E-06	0.086	15	23.2	-22.32	60.81	15.20	Am
12	EC	5085963	15	25.4	3.52E-04	0.049	9.09E-03	0.029	4.4	7.5	11.67	-24.47	-6.12	Am
13	EC	5099958_E-25	15	25.1	1.00E-04	0.056	6.37E-03	0.031	4.4	7.4	11.69	-24.1	-6.03	Am

[abc] Putative QTL identified using both Interval Mapping and Multiple QTL Model Mapping (MapQTL 6.0) and General Linear Model and Mixed Linear
Model method (TASSEL software v5.2.43)
Marker R^2- percentage phenotypic variation explained by markers in GLM and MLM
LG- Linkage group
LOD- Logarithm of odds ratio for putative QTL
PEV- percentage phenotypic variation explained by markers in IM and MQM
Af- Female additivity
Am- Male additivity
D- Dominance
CAT- Catechin; EC- Epicatechin; ECG- Epicatechin gallate; EGC- Epigallocatechin gallate; TF2- Theaflavin-3-gallate; CL- Colour; RWC- Percent
relative water content
296

Functional annotation of putative QTLs in linkage and association mapping

The putative QTLs detected in both IM, and MQM mapping were identified by the BLAST algorithm, searched on the reference tea genome (NCBI) and assigned functions on Blast2GO database. A total of 53 QTLs were detected in both interval and MQM mapping methods, of which 45 proteins were assigned functions. An additional six markers in LG02, LG04, and LG12 were found to be associated with caffeine, EC, ECG and EGC traits in AM (Table 7). The six additional QTLs detected in AM were also functionally annotated using the above databases. The putative candidate genes identified were involved in purine or thiamine biosynthesis, phenylalanine biosynthesis, tyrosine and tryptophan biosynthesis, carbon fixation (photosynthesis) and abiotic stress. The putative QTL, qCaffeine in LG02 was putatively annotated N-(5'phosphoribosyl) anthranilate (PRA) isomerase enzyme. The two other putative QTLs identified for qEC and qEGC located in LG04 were annotated as phosphoribulokinase or uridine kinase family proteins. Two putative QTLs, for qEC in LG04, were annotated as autophagy-related protein 11 and protein tyrosine kinase, respectively.

312 Table 7. Association mapping, linkage mapping and functional annotation protein of putative QTLs in black tea quality traits in St.504 and St.524
313 based on IM and MLM (Q + K) method (p < 0.01) in MapQTL 6.0 and TASSEL software v5.2.43, respectively.

Nr	Trait	Marker	LG	Pos (cM)	p-Value	A	p-Value	B	p-Value	Annotated protein	Function
1	[abc]ECG	5128890	1	96.4	1.2E-03	0.042	5.3E-03	0.031	2.0E-25	['Actin']	Cell signalling (response to cold stress, dehydration)
2	[c]CAFF	5137282	2	50.7	2.3E-03	0.042	9.7E-03	0.026	8.0E-09	["N-(5'phosphoribosyl) anthranilate (PRA) isomerase"]	Involved in the first and intermediate pathway in the purine/thiAmine biosynthesis; It also participates in the phenylalanine, tyrosine and tryptophan biosynthesis pathway
3	[abc]ECG	5087113	4	17.7	3.7E-03	0.035	2.9E-04	0.048	6.0E-22	['impB/mucB/sAmB fAmily']	Involved in UV protection through DNA repair
4	[ac]TF1	5106352	4	26	2.0E-04	0.061	8.5E-03	0.032	2.0E-06	['Thiolase, C-terminal domain']	Benzoic acid biosynthesis (volatile compounds)
5	[c]EC	5112438	4	27.4	4.9E-07	0.087	1.5E-05	0.067	9.0E-21	['Phosphoribulokinase / Uridine kinase fAmily']	Carbon fixation (photosynthesis) which are products for shikimate pathway (phenolics)
6	[c]EGC	5112438	4	27.4	1.5E-11	0.148	2.0E-08	0.134	9.0E-21	['Phosphoribulokinase / Uridine kinase fAmily']	Carbon fixation (photosynthesis) which are productsfor shikimate pathway (phenolics)
7	[ac]EC	5134490	4	26.4	5.2E-19	0.228	6.2E-15	0.246	3.0E-08	['Aminotransferase class I and II']	Phenylalanine, tyrosine and tryptophan biosynthesis
8	[ac]EGCG	5134853	4	37.6	7.0E-14	0.205	2.0E-09	0.17	2.0E-06	['Diacylglycerol kinase catalytic domain']	Water, salt, ROS, cold and freezing stress response
9	[c]EC	5136058	4	27.6	1.0E-06	0.083	5.3E-06	0.073	8.0E-09	['Autophagy-related protein 11']	Abiotic stress response (stomatal closure)
10	[ac]EGC	5136058	4	27.6	3.4E-11	0.143	3.6E-09	0.146	8.0E-09	['Autophagy-related protein 11']	Abiotic stress response (stomatal closure)
11	[c]EC	5136410	4	29.9	2.4E-05	0.065	1.8E-05	0.065	7.0E-06	['Protein tyrosine kinase']	ABA signalling
12	[ac]EGCG	5088456	12	47.9	7.7E-04	0.053	2.5E-04	0.06	4.0E-23	['Protein kinase domain']	Abiotic stress response
13	[c]ECG	5136077	12	42.8	1.9E-07	0.094	6.3E-03	0.027	5.0E-13	CSA026168	-

314 [abc] Putative QTL identified using both Interval Mapping and Multiple QTL Model Mapping (MapQTL 6.0), General Linear Model and Mixed Linear Model (TASSEL software
315 v5.2.43), respectively.
316 [c] Putative QTL identified in General Linear Model and Mixed Linear Model (MLM) using TASSEL software v5.2.43; A- Marker R^2 GLM (Q); B- Marker R^2 MLM (Q+K)

Discussion

In this current study, a combination of linkage mapping and AM was used to provide a mutual and precise authentication of QTL, which will enable more reliable results to be obtained. Linkage mapping method has a relatively low genome resolution while AM method is affected by population structure and individual relationships. The combination of linkage mapping and AM has successfully been applied to many plant studies (20-24).

The population structure analysis in discovery mapping panel used in the current study identified two groups or clusters. The dendrogram demonstrated two subspecies of the two parental clones, TRFK 303/577 and GW Ejulu, which are of *C. sinensis* var. *assamica* and *C. sinensis* var. *sinensis* variety, respectively. The neighbour-joining tree for the 261 tea accessions grouped them into two major clusters on the base of Nei's genetic distance. The phylogenetic relationships for the parental cultivars and their progenies clustering into their respective groups in separate clades based on their parental genotypes were consistent with their genetic backgrounds (25). The parental clone TRFK 303/577, which is an open-pollinated progeny of clone TRFK 6/8, is high-yielding, drought-tolerant, medium in black tea quality, caffeine, and individual catechins. On the other hand, parental clone GW Ejulu is a low-yielding, high black tea quality and moderate levels of caffeine, but high in total catechins and individual catechin contents (26). The bulk of tea cultivars used for green and black tea production have been derived through individual selection, hybridisation and molecular breeding of *C. sinensis* var. *assamica* and *C. sinensis* var. *sinensi*s varieties to produce tea with desirable characteristics (27). Therefore, developing a few markers that have the capability of discriminating the two subspecies and their cultivars is of paramount interest. The results on the linkage mapping for all the phenotypic traits and the QTL positions in different linkage groups in this study were similar as reported in (10).

In this study, AM was conducted with three different models, Q, K, and Q+K. The observed –

log10 (P) values for QTL deviated from the expected –log10 (P) values in the Q method

(GLM), indicated that there might be a few false-positive. However, the addition of genetic

relatedness K (kinship) used in a mixed linear model has proven to be more powerful and

reduces the number of false-positive associations (28). This is because a lot more QTLs were

identified in this current study using Q-model as compared with K and Q + K-model. Therefore,

there could be higher chances of false-positive or negative errors in AM due to complex

population structure (29, 30). In this study, a mixed linear model approach using the K-matrix

or a combination (Q+K) performed better than a GLM (Q). However, the K or the Q + K

methods did show to be too strict and resulted in the missing of some possibly useful QTLs.

Nearly all the significant QTLs identified in AM using the Q, and Q+K models were in line

with those identified in MapQTL 6.0 (10). The percentages of phenotypic variation explained

(R^2) by various markers analysed using AM were significant as those analysed using MapQTL

6.0. This is in agreement with previous studies reported in plants on efficiency and robustness

of combining linkage mapping and AM for precise identification of QTL (5, 31).

The six additional QTLs identified in AM were found to be associated with purine or thiamine

biosynthesis, phenylalanine, tyrosine and tryptophan biosynthesis, carbon fixation

(photosynthesis) and abiotic stress. The putative QTL, qCaffeine located in LG02 was

putatively annotated N-(5'phosphoribosyl) anthranilate (PRA) isomerase enzyme. The enzyme

is involved in the first and intermediate pathway in the purine or thiamine biosynthesis and

tryptophan biosynthesis pathway (32). Caffeine (1,3,7-trimethylxanthine) and other

methylxanthines such as theobromine (3,7-dimethylxanthine) and methyluric acids present in

tea are classified as purine alkaloids (33). Therefore, the putative QTL, qCaffeine could be

associated with the biosynthesis of caffeine, theobromine and methyluric acids in tea. Also,

phenylalanine, tyrosine, and tryptophan are products of shikimate pathway which is involved

in the biosynthesis of plant flavonoids, including catechins (34).

The putative QTLs, qEC, and qEGC in LG04 were annotated as phosphoribulokinase/uridine

kinase family proteins which are involved in carbon fixation or photosynthesis in plants (35).

Photosynthesis is an important process in plants for provision of carbon skeletons to the

shikimate pathway (36). Catechins are synthesised in the leaves of the tea plant through the

acetic-malonic acid and shikimic-cinnamic acid metabolic pathways (37). The chalcone and

gallic acid are produced from the shikimic acid pathway, which then produce the different

catechins (38). The two putative QTLs, qEC annotated putatively as autophagy-related protein

11 and protein tyrosine kinase are involved in abiotic stress response through a process of

stomatal closure and ABA signalling (39, 40). Stress conditions can interfere with

photosynthetic energy production in plants, which leads to stomatal closure, which inhibits

CO_2 intake and thereby reduces photosynthetic activity in leaves (41). Abscisic acid has been

reported to confer abiotic stress-tolerance in crop plants (39). In stress conditions like drought,

extreme temperature, and high salinity, content in plants increases considerably, inspiring

stress-tolerance effects that help plants, adapt, and survive under these stressful situations (42).

Also, ABA is also required for plant growth and development under non-stress conditions (43).

Conclusion

The tea plant is a woody perennial crop with long generation intervals, which usually hinders

its genetic improvement by conventional breeding methods. The approach of combining the

high QTL detection power of genetic linkage mapping with the high-resolution power of AM

allowed identification and precise authentication of putative QTL controlling black tea quality

and drought tolerance traits. Based on the two mapping approaches, the putative QTLs

associated with caffeine content, catechins fractions, theaflavins fractions, tea tasters' scores,

and %RWC detected in linkage mapping and AM were not significantly different except that

an additional six more QTLs were detected using AM method. Therefore, the combination of

linkage and association strategies can be used to identify closely linked molecular markers for

robust MAS and trait introgression. This will accelerate tea breeding progress in terms of

reduced breeding time, reduced cost, increased efficiency, and precision of selection.

Acknowledgments

395 The authors are grateful to KALRO-Tea Research Institute (Kenya) and the University of

396 Pretoria (South Africa) for providing space for field experiments, facilitating data collection

397 and analysis and laboratory work.

References

1.   Kraakman AT, Niks RE, Van den Berg PM, Stam P, Van Eeuwijk FA. Linkage

disequilibrium mapping of yield and yield stability in modern spring barley cultivars. Genetics.

2004;168(1):435-46.

2.   Soto-Cerda BJ, Cloutier S. Association mapping in plant genomes. Genetic diversity

in plants: InTech; 2012.

3.   Sterken R, Kiekens R, Boruc J, Zhang F, Vercauteren A, Vercauteren I, et al. Combined

linkage and association mapping reveals CYCD5; 1 as a quantitative trait gene for

endoreduplication in Arabidopsis. Proceedings of the National Academy of Sciences.

2012;109(12):4678-83.

4.   Shi W, Hao C, Zhang Y, Cheng J, Zhang Z, Liu J, et al. A Combined Association

Mapping and Linkage Analysis of Kernel Number Per Spike in Common Wheat (Triticum

aestivum L.). Frontiers in Plant Science. 2017;8:1412.

5.   Li X, Zhou Z, Ding J, Wu Y, Zhou B, Wang R, et al. Combined linkage and association

mapping reveals QTL and candidate genes for plant and ear height in maize. Frontiers in plant

science. 2016;7:833.

6.   Du Q, Gong C, Wang Q, Zhou D, Yang H, Pan W, et al. Genetic architecture of growth

traits in Populus revealed by integrated quantitative trait locus (QTL) analysis and association

studies. New Phytologist. 2016;209(3):1067-82.

7.   Bartholomé J, Bink MC, van Heerwaarden J, Chancerel E, Boury C, Lesur I, et al.

Linkage and association mapping for two major traits used in the maritime pine breeding

program: height growth and stem straightness. PloS one. 2016;11(11):e0165323.

8.   Fournier-Level A, Lacombe T, Le Cunff L, Boursiquot J, This P. Evolution of the

VvMybA gene family, the major determinant of berry colour in cultivated grapevine (Vitis

vinifera L.). Heredity. 2010;104(4):351.

9.   Jin J-Q, Yao M-Z, Ma C-L, Ma J-Q, Chen L. Association mapping of caffeine content

with TCS1 in tea plant and its related species. Plant Physiology and Biochemistry.

2016;105:251-9.

10.   Koech RK, Malebe PM, Nyarukowa C, Mose R, Kamunya SM, Apostolides Z.

Identification of novel QTL for black tea quality traits and drought tolerance in tea plants

(Camellia sinensis). Tree Genetics & Genomes. 2018;14(1):9.

11.   Pritchard J, Wen X, Falush D. Documentation for STRUCTURE

software.[Documentation file]. Available with the program at http://pritch bsd uchicago

edu/structure html. 2010.

12.   Earl DA. STRUCTURE HARVESTER: a website and program for visualizing

STRUCTURE output and implementing the Evanno method. Conservation genetics resources.

2012;4(2):359-61.

13.   Mezmouk S, Dubreuil P, Bosio M, Décousset L, Charcosset A, Praud S, et al. Effect of

population structure corrections on the results of association mapping tests in complex maize

diversity panels. Theoretical and applied genetics. 2011;122(6):1149-60.

14.   Jakobsson M, Rosenberg NA. CLUMPP: a cluster matching and permutation program

for dealing with label switching and multimodality in analysis of population structure.

Bioinformatics. 2007;23(14):1801-6.

15.   Ramasamy RK, Ramasamy S, Bindroo BB, Naik VG. STRUCTURE PLOT: a program

for drawing elegant STRUCTURE bar plots in user friendly interface. SpringerPlus.

2014;3(1):431.

16.   Van Ooijen J. JoinMap® 4, Software for the calculation of genetic linkage maps in

experimental populations. Kyazma BV, Wageningen. 2006;33:10.1371.

17.   Van Ooijen JW, Boer M, Jansen R, Maliepaard C. MapQTL 4.0: software for the

calculation of QTL positions on genetic maps (user manual). Plant Research International;

2000.

18.   Sadok IB, Celton J-M, Essalouh L, El Aabidine AZ, Garcia G, Martinez S, et al. QTL

mapping of flowering and fruiting traits in olive. PLoS One. 2013;8(5):e62831.

19.   Bradbury PJ, Zhang Z, Kroon DE, Casstevens TM, Ramdoss Y, Buckler ES. TASSEL:

software for association mapping of complex traits in diverse samples. Bioinformatics.

2007;23(19):2633-5.

20.   Fan R, Xiong M. Combined high resolution linkage and association mapping of

quantitative trait loci. European Journal of Human Genetics. 2003;11(2):125.

21.   Jung J, Fan R, Jin L. Combined linkage and association mapping of quantitative trait

loci by multiple markers. Genetics. 2005;170(2):881-98.

22.   Lu Y, Zhang S, Shah T, Xie C, Hao Z, Li X, et al. Joint linkage–linkage disequilibrium

mapping is a powerful approach to detecting quantitative trait loci underlying drought tolerance

in maize. Proceedings of the National Academy of Sciences. 2010;107(45):19585-90.

23.   Lou Q, Chen L, Mei H, Wei H, Feng F, Wang P, et al. Quantitative trait locus mapping

of deep rooting by linkage and association analysis in rice. Journal of experimental botany.

2015;66(15):4749-57.

24.   Cao Y, Li S, Wang Z, Chang F, Kong J, Gai J, et al. Identification of major quantitative

trait loci for seed oil content in soybeans by combining linkage and genome-wide association

mapping. Frontiers in plant science. 2017;8:1222.

25.   Kamunya S, Wachira F, Pathak R, Korir R, Sharma V, Kumar R, et al. Genomic

mapping and testing for quantitative trait loci in tea (Camellia sinensis (L.) O. Kuntze). Tree

genetics & genomes. 2010;6(6):915-29.

26.   TRFK. TRFK clonal catalogue. In: TRFK, editor. 2012.

27.   Yao M, Chen L, Liang Y. Genetic diversity among tea cultivars from China, Japan and

Kenya revealed by ISSR markers and its implication for parental selection in tea breeding

programmes. Plant Breeding. 2008;127(2):166-72.

28.   Finno CJ, Aleman M, Higgins RJ, Madigan JE, Bannasch DL. Risk of false positive

genetic associations in complex traits with underlying population structure: a case study. The

Veterinary Journal. 2014;202(3):543-9.

29.   Müller B, Stich B, Piepho H. A general method for controlling the genome-wide type I

error rate in linkage and association mapping experiments in plants. Heredity.

2011;106(5):825.

30.   George A. Controlling type 1 error rates in genome-wide association studies in plants.

Heredity. 2013;111(1):86.

31.   Mammadov J, Sun X, Gao Y, Ochsenfeld C, Bakker E, Ren R, et al. Combining powers

of linkage and association mapping for precise dissection of QTL controlling resistance to gray

leaf spot disease in maize (Zea mays L.). BMC genomics. 2015;16(1):916.

32.   Li J, Zhao J, Rose AB, Schmidt R, Last RL. Arabidopsis phosphoribosylanthranilate

isomerase: molecular genetic analysis of triplicate tryptophan pathway genes. The Plant Cell.

1995;7(4):447-61.

33.   Ashihara H, Sano H, Crozier A. Caffeine and related purine alkaloids: biosynthesis,

catabolism, function and genetic engineering. Phytochemistry. 2008;69(4):841-56.

34.   Ghasemzadeh A, Ghasemzadeh N. Flavonoids and phenolic acids: Role and

biochemical activity in plants and human. Journal of medicinal plants research.

2011;5(31):6697-703.

35.   Miziorko HM. Phosphoribulokinase: current perspectives on the structure/function

basis for regulation and catalysis. Advances in Enzymology and Related Areas of Molecular

Biology, Volume 74. 2000:95-127.

36.   Janacek SH, Trenkamp S, Palmer B, Brown NJ, Parsley K, Stanley S, et al.

Photosynthesis in cells around veins of the C3 plant Arabidopsis thaliana is important for both

the shikimate pathway and leaf senescence as well as contributing to plant fitness. The Plant

Journal. 2009;59(2):329-43.

37.   Vuong QV, Stathopoulos CE, Nguyen MH, Golding JB, Roach PD. Isolation of green

tea catechins and their utilization in the food industry. Food Reviews International.

2011;27(3):227-47.

38.   Chu D, Juneja L. General chemical composition of green tea and its infusion. Chemistry

and applications of green tea. 1997:13-22.

39.   Sah SK, Reddy KR, Li J. Abscisic acid and abiotic stress tolerance in crop plants.

Frontiers in plant science. 2016;7:571.

40.   Batoko H, Dagdas Y, Baluska F, Sirko A. Understanding and exploiting autophagy

signaling in plants. Essays in biochemistry. 2017;61(6):675-85.

41.   Chaves MM, Flexas J, Pinheiro C. Photosynthesis under drought and salt stress:

regulation mechanisms from whole plant to cell. Annals of botany. 2009;103(4):551-60.

42.   Ng LM, Melcher K, Teh BT, Xu HE. Abscisic acid perception and signaling: structural

mechanisms and applications. Acta Pharmacologica Sinica. 2014;35(5):567.

43.   Nakashima K, Yamaguchi-Shinozaki K. ABA signaling in stress-response and seed

development. Plant cell reports. 2013;32(7):959-70.

Supporting information

S1 Fig. Trait-wise association Q-Q plots for GLM. AR- Aroma; AST- Astringency; BRK-

Briskness; BRT- Brightness; C- Catechin; CAFF- Caffeine; CL- Colour; EC- Epicatechin;

ECG- Epicatechin gallate; EGC- Epicatechin gallate; EGCG- Epigallocatechin gallate; RWC-

Percent relative water content; TF1- Theaflavin; TF2- Theaflavin-3-gallate; TF3- Theaflavin-

3'-gallate; TF4- Theaflavin-3,3'-digallate

S2 Fig. Trait-wise association Q-Q plots for GLM (Q). AR- Aroma; AST- Astringency;

BRK- Briskness; BRT- Brightness; C- Catechin; CAFF- Caffeine; CL- Colour; EC-

Epicatechin; ECG- Epicatechin gallate; EGC- Epicatechin gallate; EGCG- Epigallocatechin

gallate; RWC- Percent relative water content; TF1- Theaflavin; TF2- Theaflavin-3-gallate;

TF3- Theaflavin-3'-gallate; TF4- Theaflavin-3,3'-digallate

S3 Fig. Trait-wise Q-Q association plots for mixed linear model MLM (K). AR- Aroma;

AST- Astringency; BRK- Briskness; BRT- Brightness; C- Catechin; CAFF- Caffeine; CL-

Colour; EC- Epicatechin; ECG- Epicatechin gallate; EGC- Epicatechin gallate; EGCG-

Epigallocatechin gallate; RWC- Percent relative water content; TF1- Theaflavin; TF2-

Theaflavin-3-gallate; TF3- Theaflavin-3'-gallate; TF4- Theaflavin-3,3'-digallate

S4 Fig. Trait-wise Q-Q association plots for mixed linear model MLM (Q+K). AR- Aroma;

AST- Astringency; BRK- Briskness; BRT- Brightness; C- Catechin; CAFF- Caffeine; CL-

Colour; EC- Epicatechin; ECG- Epicatechin gallate; EGC- Epicatechin gallate; EGCG-

Epigallocatechin gallate; RWC- Percent relative water content; TF1- Theaflavin; TF2-

Theaflavin-3-gallate; TF3- Theaflavin-3'-gallate; TF4- Theaflavin-3,3'-digallate

Figure

Figure

Figure

Figure