A triterpenoid saponin, Spergulin-A from *Glinus oppositifolius* is a potent immunostimulator and antileishmanial agent

Saswati Banerjee [a¥], Niladri Mukherjee [a¥], Rahul L. Gajbhiye [b], Parasuraman Jaisankar [b], Sriparna Datta [c], Krishna Das Saha [a]*

[a]Cancer Biology and Inflammatory Disorder Division, CSIR-Indian Institute of Chemical Biology, 4 Raja S. C. Mullick Road, Kolkata-700032, India

[b]Organic and Medicinal Chemistry Division, CSIR-Indian Institute of Chemical Biology, 4, Raja S.C. Mullick Road, Kolkata 700032, India

[c]Department of Chemical Technology, University of Calcutta, Kolkata 700009, India

Email address: krishna@iicb.res.in (Krishna Das Saha); ¥ Equal contribution, *Corresponding Author

ABSTRACT

Present chemotherapeutics are inadequate against visceral leishmaniasis (VL) an immunosuppressive ailment caused by *Leishmania donovani*. Despite the interest in plant-based drug development, an antileishmanial drug from plant source is unavailable. *Glinus oppositifolius* had been reported in favor of being immunomodulators along with other traditional uses. Novel anti-VL therapies can rely on host immune-modulation with associated leishmanicidal action. With this rationale, an n-BuOH fraction of the methanolic extract of the plant and isolated triterpenoid saponin Spergulin-A were evaluated against acellular and intracellular *L. donovani*. Immunostimulatory activity of them was confirmed by elevated TNF-α and extracellular NO production from treated MΦs and was found nontoxic to the host cells. Identification and structure confirmation for isolated Spergulin-A was performed by ESI-MS,[13]C, and [1]H NMR. Spergulin-A was found ineffective against the acellular forms while, against the intracellular parasites at 30μg/ml, the reduction was 92.6% after 72h. Spergulin-A enhanced ROS and nitric oxide (NO) release and changes in Gp91-phox, i-NOS, and pro and anti-inflammatory cytokines elaborated its intracellular anti-leishmanial activity. The results supported that *G. oppositifolius* and Spergulin-A can potentiate new lead molecules for the development of alternative drugs against VL.

Author summary

Visceral Leishmaniasis (VL) is prevalent in tropical and temperate regions and is endemic in 88 countries with an estimated yearly incidence of 1–1.5 million cases of cutaneous leishmaniasis and 500,000 cases of VL, of which 70,000 people die every year. More than 90% of cases occur in India, Bangladesh, Nepal, Sudan, Ethiopia, and Brazil with an estimated incidence of at least 500,000 new cases and 50,000 deaths each year.

This work highlights the anti-leishmanial effect of a triterpenoid saponin that was extracted from a tropical plant, *Glinus oppositifolius*, commonly found in Asian and parts of African countries. The drug showed potent Leishmanicidal activity against *Leishmania donovani* at very low concentration. As the application of the conventional drugs (pentavalent antimoniates, amphotericin B, and pentamidine) is limited due to low efficacy, life-threatening side effects, high toxicity, induction of parasite resistance, length of treatment and high cost and anti-leishmanial vaccines may not become available in the near future, so plant extracts or plant compounds, represent a valuable source of new medicinal agents and alternative, effective and less toxic treatments, leading to a search for natural products with a potential leishmanicidal action. Being an immunosuppressive disease efficacy against VL can be demonstrated against intracellular parasites in the lights of emphasizing host MΦ immunostimulation as elaborated precisely in the present study.

Introduction

Visceral Leishmaniasis (VL) considered as the most severe form of leishmaniasis caused by *Leishmania donovani* and untreated patients nearly always die [1]. Leishmaniasis is endemic in 88 countries with approximately 350 million people are at risk with an estimated yearly incidence of 500,000 cases and almost 70,000 deaths [2]. Leishmaniasis is liable for the ninth most substantial infectious diseases burden, however, is mostly disregarded among tropical disease priorities [3]. Sodium antimony gluconate, Miltefosine, Pentamidine, and Amphotericin B are the primary therapeutics though associated with toxicity or resistance [2].

Pathogen recognition by neutrophils, macrophages, dendritic cells, and natural killer cells activates intracellular signaling pathways leading into inflammatory responses like inflammasome activation and IL-1β production, which is not the case for leishmanial infection [4]. *L. donovani* infection is characterized by the parasite-induced active subversion of the host immune system and also immune deviation which additively favors infection establishment [5]. Additionally, *Leishmania* prevents inflammatory response by the impaired release of different pro-inflammatory cytokines (IL-1, TNF-α, IL-12) and enhanced releases of immunosuppressive signaling molecules, such as arachidonic acid metabolites and the cytokines TGF-β and IL-10 [6]. Involvement of co-stimulatory molecule B7-CTLA-4 is associated with increased TGF-ß production in VL with increased apoptosis of $CD4^+$ T cells and decreased macrophage apoptosis [7]. Leishmanial infection also undermines a generation of microbicidal macrophage nitric oxide (NO) and reactive oxygen species (ROS) with the hindrance of antigenic peptide display to T cells, and permeation of IL-10 producing T regulatory cells [8]. In recent years increased instances of VL has been reported in connection with immune-suppressed AIDS patients [8].

92   Discovery of novel compounds which intercedes host immune-modulation associated

93 with leishmanicidal function having permissible side effects is a precise research objective

94 [9,10]. It is also of vital importance that the drug required for parasite elimination in immune-

95 stimulated cells was significantly less than the immune-suppressed ones [11].

96   The anti-parasitic effectiveness of plant extracts majorly relies upon secondary

97 metabolites of diverse chemical groups including alkaloids, polyphenol, flavonoids,

98 terpenoids, phenylpropanoids, etc. [12]. For isolation and characterization of a herbal extract

99 or an active compound, different research strategies can be employed mainly in the extraction

100 steps. However, bioactivity-guided fractionation considered as simple, rapid, cost-effective

101 and reproducible [1]. For obtaining a potent anti-leishmanial agent with immune-modulation,

102 different studies were conducted with compounds like aslicarin A, niranthrin, skimmianine,

103 quassin, tannins, linalool, etc. nevertheless with varying degree of effectiveness and

104 satisfaction [8].

105   Aerial parts of *Glinus oppositifolius* (Family Molluginaceae) are used for treating

106 abdominal pain and jaundice, while decoction is used against malaria [13]. Plants of this

107 genus were previously documented for the presence of triterpenoid saponins [14], and

108 isolated pectin polysaccharides are antiprotozoal [15] and immunomodulators [14]. *G.*

109 *oppositifolius* is indicated for wound healing and used in traditional medicine for

110 treating diarrhea, joint pains, inflammations, intestinal parasites, fever, boils and skin

111 disorders [16]. Some triterpenoid saponins, 3-O-(β-D-xylopyranosyl)-spergulagenin-A,

112 Spergulacin, Spergulacin-A Spergulin-A, and Spergulin-B had been isolated from *G.*

113 *oppositifolius* [17]. Application of plant-based immunomodulators is smart as they mediate

114 their effectiveness by enhancing the inherent host-derived protective machinery without the

115 involvement of specific microbicidal agents namely antibiotics [13].

116   Most of the available antimalarial drugs are plant-derived; regrettably, there is no

117 anti-leishmanial drug present which is of plant origin. The recent efforts to achieve this goal

118 also were restricted against the promastigotes [18]. In the present work, an attempt has been

119 taken to evaluate the intracellular anti-leishmanial activity of this plant and its bioactive

120 component, Spergulin-A emphasizing on the immunostimulatory activity.

Materials and methods

Chemicals and Reagents

Cell culture media, serum, antibiotics, HEPES (Gibco, USA), CFSE, DAPI

(Invitrogen, USA), MTT, DMSO (Sigma Chemical Co. USA), cytokine Assay kit (Thermo

Scientific, USA), DAF-2 DA, nitric oxide assay kit, DCFDA (Calbiochem, USA) and all

other chemicals were of the highest grade commercially available. Primary and secondary

antibodies were obtained from Santa Cruz Biotechnology (USA) or Cell signaling

technologies (USA).

Isolation of methanolic fractions from G. oppositifolius

The aerial parts of *G. oppositifolius* were shade dried (1kg) and were first defatted

using petroleum ether (60-80ºC, 3.5L X 3) at room temperature for 48h. The marc was

than subjected to extraction using MeOH (3.5L X 3) at room temperature for 48h. The

extract was then filtered and MeOH was evaporated under reduced pressure and finally was

lyophilized to obtain the crude MeOH extract (13g). A part (10g) of ME was then

suspended in milli-Q water and partitioned sequentially with EtOAc and n-BuOH. Each

fraction was evaporated under vacuum and lyophilized to yield the EtOAc fraction (EAF;

3.6g), n-BuOH fraction (NBF; 4.1g) and aqueous fraction (AF; 2.3g). All the fractions were

stored at 4ºC till further use.

Among these four fractions, n-BuOH showed promising antileishmanial activity.

Around 4g of NBF was subjected to column chromatography of Diaion HP 20 (100g) and

the column washed with water followed by 30, 40, 60, 80 and 100% of MeOH to obtain a total of six fractions. Fractions eluted with 50% MeOH showed similar spots on TLC and were mixed and then re-chromatographed on Dianion HP 20 column to furnish 7mg of Spergulin-A.

Isolation and characterization of Spergulin-A

General experimental procedures: ESI Mass spectra were recorded on an Agilent 6545 Q-TOF mass spectrometer System. ^{1}H and ^{13}C NMR were recorded on a Bruker Ultrashield NMR (600 MHz) in pyridine-d5 with TMS as an internal standard. Diaion HP 20 was used for column chromatography; silica gel (60 F254) was used for TLC and spots were visualized by spraying with Lieberman–Burchard reagent followed by heating.

Parasite culture and maintenance

Passage in BALB/c mice maintained *Leishmania donovani* [MHOM/IN/1983/AG83] infection. Complete M199 media (10% FBS, pH 7.4, 100U/ml penicillin G-sodium, 100μg/ml streptomycin sulfate, 25mM HEPES) was used to culture the promastigotes.

Axenic amastigote from the log phase culture of promastigotes was prepared as described by Saar et al. [19].

Macrophage culture, parasite infection, and treatment

RAW 264.7 MΦ cell line was obtained from the ATCC and maintained in complete RPMI 1640 (10% FBS, 100μg/ml streptomycin sulfate, 100U/ml penicillin G sodium, 0.2% sodium bicarbonate, 25 mM HEPES) in a humidified atmosphere and 5% CO_2 at 37°C.

Infection of MΦs with *L. donovani* promastigotes was performed in a ratio of 1:10 (MΦ: parasite) for 4h then washed twice with media. Normal and parasitized macrophages were treated with *G. oppositifolius* fractions or Spergulin-A at required concentrations up to 72h. Infection was measured by counting the intracellular parasites and expressed as parasite count/20 MΦs.

ELISA assay

Cell-free supernatants from MΦs (1×10^5 cells/well) were collected, and the concentration of TNF-α was estimated by sandwich ELISA, using a commercially available assay kit [20].

MTT assay

20μl of MTT (5mg/ml in PBS) was added to each well of the control and treated MΦs (4×10^3) and 5×10^3 parasites (promastigote and axenic amastigote) in a 96 well plate and incubated for 4h at 37°C and the formazan crystals were dissolved in 150μl of DMSO. Absorption was measured at 595nm by an ELISA reader [21].

Measurement of extracellular NO

The collected supernatants from control, infected and treated MΦs were incubated with equal volumes of Griess reagent, NED (0.1% in distilled water) and Sulphanilamide (1% in 5% H_3PO_4) at room temperature for 10min. The absorbance was measured at 540nm on a microplate reader. NO concentration was determined using a dilution of sodium nitrite as the standard [20].

FACS and Confocal microscopy with CFSE tagged *L. donovani*

MΦs were parasitized with CFSE-tagged ($25μM/1\times10^6$ promastigote in 1ml media for 30min) promastigotes followed by Spergulin-A (10,20,30μg/ml) treatment for 24h.

For confocal microscopy, MΦs (1×10^5/glass coverslips) after infection and treatment, washed twice with PBS, fixed in chilled 70% ethanol and nuclei were stained with DAPI and observed under Olympus Fluoview FV10i confocal microscope with 60X objective lens. For FACS, MΦs (1×10^5/well in a six-well plate) after infection and treatment, washed twice with PBS then harvested by scraping and resuspended in 400μl PBS and analyzed by BD FACS LSR Fortessa with excitation at 494nm and emission at 518nm.

FACS analyses of intracellular ROS and NO

The level of intracellular ROS was determined based on the change in fluorescence of H2DCFDA. DAF-2 DA was used for detecting intracellular NO [2]. Briefly, after infection and treatment, MΦs were scrapped and incubated in PBS containing DAF-2DA (7.0μM) at 37°C for 30min or 20μM DCF-DA at 37°C for 15min and then analyzed by BD FACS LSR Fortessa with excitation at 480nm and emission at 515nm for both.

Western blot analysis

40μg of proteins harvested from MΦs were electrophoretically separated in SDS-polyacrylamide gel and transferred to PVDF membrane, blocked with BSA and incubated with respective primary antibodies overnight. The membranes were then incubated with HRP conjugated secondary antibodies, and immunoreactive bands were visualized by adding proper substrates. β-Actin was used as loading controls [22].

Statistical analysis

All values were expressed as mean ± SEM obtained from at least three replicate experiments. Statistical significance and differences among groups were assessed with One-Way analysis of variance (ANOVA) followed by Dunnett's test. P values $\leq$ 0.05 (*) or $\leq$ 0.01 (**) were considered as indicative of significance.

Results

Immunostimulatory effect of methanolic extract of *G. oppositifolius*

Of the aqueous, ethyl acetate and n-BuOH (50:50) fractions of *G. oppositifolius* methanolic extract, only the n-BuOH fraction (50μg/ml) showed a considerable increase in TNF-α and extracellular NO production (Supple. Figure 1) from the treated (24h) MΦs. The six subfractions (50μg/ml) of the n-BuOH fraction were also checked for the same parameters (Figure 1A and B), and highest augmentation was noticed in fraction 4 for being a worthy immunostimulatory agent and the lead fraction.

Dose-dependent immunostimulatory effect of the lead fraction

A dose-dependent increase in TNF-α production was monitored in treated (24h) MΦs in two sets (Figure 2B and C). In first set the lead fraction was applied at the doses of 10, 20, 40 and 80μg/ml (Figure 2B) and in the second set at the doses of 20, 30, 40 and 50μg/ml and most increments was noticed at 30μg/ml (Figure 2C) and selected as the significant dose for this study.

Assessment of macrophage survival and extracellular NO release

Survival of the MΦs after exposing them to increasing concentrations (10,20,30,40 and 50μg/ml) of the lead fraction for 24h was monitored by MTT assay (Figure 2D). The lead fraction was found safe for this dose range, and survival was about 81.45% even at 50μg/ml. Most increment (5fold) in extracellular NO was noticed for 30μg/ml of the lead fraction (Figure 2E). The increase was also significant for most of the doses.

Identification and characterization of Spergulin-A

Among those six n-BuOH bioactive sub-fractions, fraction 4 have shown most significant upliftment in TNF-α and extracellular NO production. After purification of this fraction by using column chromatography, a white amorphous compound was obtained. Molecular formula $C_{35}H_{58}O_{11}S$ (Figure 3A) was assigned from the ESI-MS (Supple. Figure 2)**.** Structural elucidation of this compound was achieved by critical analysis of the ^{1}H NMR, ^{13}C NMR and ^{13}C NMR DEPT results (Supple. Results and Supple. Figure 3-6). All the spectral data of this compound were found to be in complete agreement with reported ones and structure was then confirmed as Spergulin-A [17].

Assessment of anti-leishmanial activity of Spergulin-A

Effect on different forms of *L. donovani*. Anti-leishmanial effect of Spergulin-A (10,20,30,50,100μg/ml) was first evaluated by MTT assay against promastigote and axenic amastigote, the acellular forms (Figure 3B) after 24h. Viability was 87.6% and 77.8%

respectively for promastigote and axenic forms even at 100μg/ml. At the lower doses (10, 20 and 30μg/ml) the viability reduction was found to be non-significant (Figure 3B).

After that, the effect of Spergulin-A was evaluated against the MΦ internalized parasites (Figure 3C). Here, a significant reduction of the parasite was noticed as low as 10μg/ml and further increased with the increment of doses (Figure 3C). However, above 30μg/ml MΦ internalized antileishmanial effect of Spergulin-A had reached a plateau, and thus further experiments were mostly conducted with 30μg/ml.

Time-dependent intracellular leishmanicidal effect. Efficacy of Spergulin-A (30μg/ml) against the intracellular parasites was evaluated at every 24h interval up to 72h (Figure 3D). Inside the infected MΦs parasite count increased significantly. The treated MΦs exhibited significantly reduced numbers of parasites from the initial infection as well as the corresponding observation point of infected macrophages (Figure 3D). The parasite reduction at 72h was 86.2% less than the initial point of infection and 92.6% less than the 72h count of infected MΦs.

Dose-dependent quantitative and qualitative anti-leishmanial effect. After infecting the MΦs with CFSE-tagged promastigotes and treatment with Spergulin-A (10, 20 and 30μg/ml, 24h), the cells were evaluated by FACS in FITC filter (Figure 4A). Highest intensity was noticed in the infected panels (Figure 4A.ii) and with the increment of Spergulin-A doses the intensity gradually reduces which is reciprocal to the reduced parasite count (Figure 4A.iii-v). Interestingly, no significant changes in number of parasitized MΦs observed within the treatment panel (Figure 4A.vi) and what changes were the intensity denoted the parasite count (Figure 4A.vii). Thus, possibly Spergulin-A reduced the parasite by immune-stimulating the host MΦs.

Confocal micrographs depicted the reduction in parasite emitting green fluorescence

with the increment of Spergulin-A doses. Moreover, most decreases were noticed when the

parasitized MΦs was treated with 30μg/ml of it (Figure 4B).

Intracellular ROS and NO status due to infection and treatment. The

fluorescence intensity for H2DCFDA reciprocal to ROS production was found to be minimal

for the infected panel (Figure 4C.ii) and significantly higher in the treated panel (Figure

4C.iii). The fluorescence intensity of DAF-2DA for NO was also found to be highest for the

treated panel (Figure 4D.iii) compared to the control and infected MΦs (Figure 4D.i and ii).

Estimation of alteration of different cytokines and immunostimulators. In

western blot analysis, it was observed that Gp91-phox liable for anti-microbial ROS

production in MΦs got upregulated in treated panel (Figure 4E). Level of i-NOS which

regulates anti-leishmanial NO production also increased with Spergulin-A treatment

compared to control and infected MΦs (Figure 4F).

Expression of pro-inflammatory TNF-α and IL-12β got down-regulated with infection

and considerably up-regulated in treatment panel (Figure 4G). Anti-inflammatory IL-10 and

TGF-β expression were up-regulated in infected MΦs compared to the control, and with the

treatment of Spergulin-A (30μg/ml) their appearance got normalized like that of control

(TGF-β) or even less (IL-10) (Figure 4G).

Discussion

Precisely an immune-suppressive ailment, VL inhabit and modulate the microbicidal

function of the macrophages and create a microenvironment favoring parasite growth inside

the visceral organs by modulating pro and anti-inflammatory cytokines and impairing ROS

and NO release [6]. The present *in vitro* study aimed to assess the immunostimulatory

property of *G. oppositifolius* and isolation of a compound Spergulin-A which can reduce the

intracellular parasites by immune-stimulation within a safe dose range for the host cells.

Primary appraisal of an extract that can have an immunostimulatory effect was based upon

the increased release of TNF-α and extracellular NO from treated MΦs. It was observed that

at an applied same dose (50μg/ml, 24h) only the n-BuOH fraction showed a considerable

increase in TNF-α and extracellular NO production. The selection of TNF-α and extracellular

NO as the potent immunostimulatory index in the present context is vital because endogenous

TNF-α produced by infected macrophages can elicit the release of L-arginine-derived

nitrogen intermediates detrimental for the intracellular parasites [9]. The six n-BuOH sub-

fractions were again verified for immune-stimulation, and fraction 4 emerged as the most

promising one. The fraction was also considered to be safe for the MΦs even at 50μg/ml

(81.5% viability) having CC_{50} well above 100μg/ml.

Interested by the initial observations on elevated immunostimulation by the n-BuOH

sub-fraction 4 and based upon the previous inspection of the presence of triterpenoid

saponins in this fraction [17] a triterpenoid saponin Spergulin-A was isolated and purified

following the well-established method (Figure 3A and Supple. information) [17,23].

Evaluation of the anti-leishmanial property of Spergulin-A was the principal research

interest after identification and isolation of it. The possible anti-leishmanial effect was first

checked against the acellular forms of the parasite but without major success even at

100μg/ml (Figure 3B). Therefore, if Spergulin-A has any leishmanicidal effect on the

intracellular parasites that will possibly mediate by altering the microenvironment of the host

MΦs. After 24h of incubation Spergulin-A (30μg/ml) was found to be pretty useful in

reducing the intracellular parasite count and when applied for a prolonged period of 72h

reduction of intracellular *Leishmania* was significant in respect to both the infected MΦs of

corresponding time point and initial parasitized MΦs (Figure 3C,D). It is also of great

significance that no considerable alteration in number of parasitized MΦs was found in

treatment panel while reduction of CFSE-stained internalized parasites was notable (Figure

4A). This specific observation strongly advocated in favor of immunostimulation by Spergulin-A which mediated parasite killing. For being leishmanicidal, Spergulin-A should be useful in enhancing the release of NO and ROI, well recognized for their worth against *Leishmania* [6,24]. NO is especially critical for intracellular parasite clearance as it was reported that mice with impaired inducible nitric oxide synthase (i-NOS) and thus restricted production and release of NO are incapable of *Leishmania* control even for isolated macrophages *in vitro* [25]. At the dose of 30μg/ml, Spergulin-A showed enhanced production of intracellular NO and ROI (Figure 4C,D) which directly reciprocal to the reduction of intracellular parasites and signified their connection. In the western blot analysis also increment in i-NOS production (Figure 4F) was noticed in Spergulin-A treated parasitized MΦs in amendable enhanced NO production and parasite control. The same interpretation is also pertinent for an increase of Gp91-phox, a subunit of the NADPH oxidase and enhancement of ROI (Figure 4 E,C) in Spergulin-A mediated intracellular parasite killing. It was previously established that intracellular leishmanial killing proceeds by NO production from arginine by i-NOS, and superoxide (O_2^-) generated by the NADPH oxidase [26]. Several Leishmania species induce immunosuppressive TGF-β production, and diminution of TGF-β secretion is in direct correlation with enhanced i-NOS production [6] and can lead to internalized parasite removal as found with Spergulin-A treatment. TGF-β also found to increase the VL progression and averts disease cure in murine models [26]. IL-10 is another anti-inflammatory cytokine which makes macrophages indifferent to various activation signals and thus causes impairment of intracellular parasite killing also by down-regulating the production of TNF-α and NO [27]. IL-10 production increased in infected macrophages *in vitro*, apparently via interaction with the Fc receptor [28] and which down-regulated with Spergulin-A treatment (Figure 4G) in connection with parasite reduction. Parasite infection is responsible for the suppression of macrophage microbicidal activity that relies upon NO, ROI

339 and cytokines like IL-1, IL-12β, and TNF-α [29] as with infection the levels of TNF-α IL-12β

340 got down-regulated, but with the application of Spergulin-A, these cytokines significantly

341 elevated which signifies their role in intracellular *Leishmania* parasite control.

Conclusion

The inconsistency of direct leishmanicidal effect of *G. oppositifolius* and Spergulin-A

isolated from it against promastigotes and axenic amastigote on the one hand and the proved

efficacy of them against intracellular parasites was explained by emphasizing host MΦ

immunostimulation as elaborated precisely in the present study. Bio-guided fractionation was

employed to isolate and identify Spergulin-A as the immunostimulant, and its anti-

leishmanial property was evaluated at a dose best suited for the research and found to be safe

against the host cells assessed *in vitro* in RAW 264.7 MΦs infected with a virulent strain of

L. donovani.

Supporting information

S1 NMR data of Spergulin A: Detailed NMR data for structure confirmation of Spergulin-

A. (DOC)

S1 Fig. 1. Evaluation of immunostimulation by fractions of *G. oppositifolius* MeOH

extract. The fractions, aqueous, ethanolic and n-BuOH (50μg/ml) were checked for altered

TNF-α production and extracellular NO production in the culture supernatant of treated RAW

264.7 MΦs. (TIF)

S1 Fig. 2. ESI-Mass Spectra of Compound 1 (Spergulin A). (TIF)

S1 Fig. 3. [1]H NMR Spectra of Compound 1 (Spergulin A) in pyridine-d5. (TIF)

S1 Fig. 4. [13]C NMR Spectra of Compound 1 (Spergulin A) in pyridine-d5**.** (TIF)

S1 Fig. 5. DEPT-135 of Compound 1 (Spergulin A) in pyridine-d5. (TIF)

S1 Fig. 6. DEPT-90 of Compound 1 (Spergulin A) in pyridine-d5. (TIF)

References

1) Tiuman TS, Santos AO, Ueda-Nakamura T, Filho BP, Nakamura CV. Recent advances in

leishmaniasis treatment. Int J Infect Dis 2011;15:e525–32.

2) Banerjee S, Bose D, Chatterjee N, Das S, Chakraborty S, Das T, et al. Attenuated

Leishmania induce pro- inflammatory mediators and influence leishmanicidal activity by

p38 MAPK dependent phagosome maturation in *Leishmania donovani* co-infected

macrophages. Sci Rep 2016;6:22335.

3) Alvar J, Ve´lez ID, Bern C, Herrero M, Desjeux P, Cano J, et al. Leishmaniasis

Worldwide and Global Estimates of Its Incidence. PLoS One 2012;7:e35671.

4) Freitas EO, Leoratti FMS, Freire-de-Lima CG, Morrot A, Feijó DF. The Contribution of

Immune Evasive Mechanisms to Parasite Persistence in Visceral Leishmaniasis. Front

Immunol 2016;7:153.

5) Bhardwaj S, Srivastava N, Sudan R, Saha B. *Leishmania* interferes with host cell signaling

to devise a survival strategy. J Biomed Biotechnol 2010;2010:109189.

6) Olivier M, Gregory DJ, Forget G. Subversion mechanisms by which *Leishmania* parasites

can escape the host immune response: a signaling point of view. Clin Microbiol Rev

2005;18:293–305.

7) Goto H, Lindoso JAL. Immunity and immunosuppression in experimental visceral

leishmaniasis. Braz J Med Biol Res 2004;37:615–23.

8) Islamuddin M, Chouhan G, Want MY, Ozbak HA, Hemeg HA, Afrin F.

Immunotherapeutic potential of eugenol emulsion in experimental visceral leishmaniasis.

PLoS Negl Trop Dis 2016;10:e0005011.

9) Maran N, Gomes PS, Freire-de-Lima L, Freitas EO, Freire-de-Lima CG, Morrot A. Host

resistance to visceral leishmaniasis: prevalence and prevention. Expert Rev Anti Infect

Ther 2016;14:435–42.

10) Flora R, Aghazadeh-Dibavar S, Bandyopadhyay M, Dasgupta S. Immunosuppression

during *Leishmania donovani* infection: a potential target for the development of therapy.

Ann Parasitol 2014;60:239–45.

11) Haidaris CG, Bonventre PF. Efficacy of combined immunostimulation and chemotherapy

in experimental visceral Leishmaniasis. Am J Trop Med Hyg 1983;32:286–95.

12) Mukherjee N, Mukherjee S, Saini P, Roy P, Sinha Babu SP. Phenolics and Terpenoids; the

Promising New Search for Anthelmintics: A Critical Review. Mini Rev Med Chem

2016;16:1415–41.

13) Inngjerdingen KT, Kiyohara H, Matsumoto T, Petersen D, Michaelsen TE, Diallo D, et al.

An immunomodulating pectic polymer from *Glinus oppositifolius*. Phytochemistry

2007;68:1046–58.

14) Sahakitpichan P, Disadee W, Ruchirawat S, Kanchanapoom T. L-()-(N-trans-Cinnamoyl)-

arginine, an acylamino acid from *Glinus oppositifolius* (L.) Aug. DC. Molecules

2010;15:6186–92.

15) Traore F, Faure R, Ollivier E, Gasquet M, Azas N, Debrauwer L, et al. Structure and

antiprotozoal activity of triterpenoid saponins from *Glinus oppositifolius*, Planta Med

2000;66:368–71.

16) Sahu SK, Das D, Tripathy NK, Dinda SC, Sundeep Kumar HK. Evaluation of

hypoglycemic activity of *Mollugo pentaphylla* and *Glinus oppositifolius* (L). Rasayan J

Chem 2012;5:57–62.

17) Kumar D, Shah V, Ghosh R, Pal BC. A new triterpenoid saponins from *Glinus*
oppositifolius with a-glucosidase inhibitory activity. Nat Prod Res 2013;27:624–29.

18) Kaur H, Thakur A, Kaur S. Immunoprophylactic Potential of a Cocktail of Three Low
Molecular Weight Antigens of *Leishmania donovani* along with Various Adjuvants
Against Experimental Visceral leishmaniasis. Iran J Parasitol 2018;13:11–23.

19) Saar Y, Ransford A, Waldman E, Mazareb S, Amin-Spector S, Plumblee J, et al.
Characterization of developmentally-regulated activities in axenic amastigotes of
Leishmania donovani. Mol Biochem Parasitol 1998;95:9–20.

20) Chatterjee N, Das S, Bose D, Banerjee S, Das S, Chattopadhyay D, et al., Exploring the
anti-inflammatory activity of a novel 2-phenylquinazoline analog with protection against
inflammatory injury. Toxicol Appl Pharmacol 2012;264:182–91.

21) Mosmann T. Rapid colorimetric assay for cellular growth and survival: application to
proliferation and cytotoxicity assays. J Immunol Methods 1983;65:55–63.

22) Sikdar Y, Modak R, Bose D, Banerjee S, Bieńko D, Zierkiewicz W, et al. Doubly chloro
bridged dimeric copper(II) complex: magneto-structural correlation and anticancer
activity. Dalton Trans 2015;44:8876.

23) Kumar D, Mallick S, Vedasiromoni JR, Pal BC. Anti-leukemic activity of *Dillenia indica*
L. fruit extract and quantification of betulinic acid by HPLC. Phytomedicine 2010;17:431–
35.

24) Matte C, Maion G, Mourad W, Olivier M. *Leishmania donovani*-induced macrophages
cyclooxygenase-2 and prostaglandin E2 synthesis. Parasite Immunol 2001;23:177–84.

25) Wei XQ, Charles IG, Smith A, Ure J, Feng GJ, Huang FP, et al. Altered immune
responses in mice lacking inducible nitric oxide synthase, Nature 1995;375:408–11.

26) Wilson ME, Jeronimo SMB, Pearson RD. Immunopathogenesis of infection with the
visceralizing Leishmania species. Microb Pathogenesis 2005;38:147–60.

27) Nylen S, Sacks D. Interleukin-10 and the pathogenesis of human visceral leishmaniasis.

Trends Immunol 2007;28:378–84.

28) Sutterwala FS, Noel GJ, Salgame P, Mosser DM. Reversal of proinflammatory responses

by ligating the macrophage Fcgamma receptor type I. J Exp Med 1998;188:217–22.

29) Kane MM, Mosser DM. Leishmania parasites and their ploys to disrupt macrophage

activation. Curr Opin Hematol 2000;7:26–31.

Figure Captions

Fig 1. Evaluation of immunostimulation by subfractions of n-BuOH fraction of *G.*

oppositifolius MeOH extract.

(A) The subfractions (50μg/ml) were checked for altered TNF-α production after 24h

treatment of RAW 264.7 MΦs from the culture supernatant. (B) Extracellular NO production

was monitored in the culture supernatant of treated (six subfractions) and control after 24h by

Griess reagent. All values are expressed as mean ± SEM from triplicate assays from three

independent experiments (P values ≤ 0.05 (*) or ≤ 0.01 (**) vs. control).

Fig 2. Dose-dependent evaluation of the immunostimulatory effect of the n-BuOH lead

fraction and impact on RAW 264.7 MΦ survival.

(A) TLC profile of n-BuOH lead fraction. (B, C) The dose-dependent release of TNF-α was

monitored in treated (24h) RAW 264.7 MΦs, and highest increment was found in 30μg/ml.

(D) Survival of RAW 264.7 MΦs at different doses of an n-BuOH lead fraction after 24h

exposure measured by MTT assay. (E) Dose-dependent evaluation of extracellular NO

production was monitored in the culture supernatant of 24h treated RAW 264.7 MΦs by

Griess reagent. All values are denoted as mean ± SEM from triplicate assays from three

independent experiments (P values ≤ 0.05 (*) or ≤ 0.01 (**) vs. control).

Fig 3. Anti-leishmanial activity of Spergulin-A.

(A) Structure of the triterpenoid saponins Spergulin-A. (B) Dose-dependent effects of Spergulin-A against acellular forms namely the promastigotes and axenic amastigote of *L. donovani* after 24h. (C) Dose-dependent effect of Spergulin-A against the intracellular amastigote of *L. donovani* after 24h of treatment and expressed as number of parasites/20 MΦs. (D) 24h time lapse evaluation of anti amastigote effect of Spergulin-A at 30μg/ml for 72h. All values are expressed as mean ± SEM from triplicate assays from three independent experiments (P values ≤ 0.05 (*) or ≤ 0.01 (**) vs. control).

Fig 4. Elaboration of the Anti-leishmanial activity of Spergulin-A.

(A) Quantitative dose-dependent assessment of intracellular CFSE stained antileishmanial activity of Spergulin-A by flow cytometry after 24h treatment. (B) Confocal laser-scanning micrographs to evaluate the reduction in intracellular CFSE stained *L. donovani* after 24h of Spergulin-A treatment (magnification 120X). (C) FACS monitored the changes in ROS production in control, infected, and treated (30μg/ml Spergulin-A) MΦs with H2DCFDA. (D) The changes in intracellular NO production were monitored by FACS with fluorescent probe DAF-2 DA. (E, F, G) Changes in the expression of Gp91-Phox, i-NOS and cytokines namely pro-inflammatory TNF-α, IL-12β and anti-inflammatory IL-10 and TGF-β with β-actin as loading control and densitometry analyses was provided after band normalization. Images are representative of three separate experiments, and all values are expressed as mean ± SEM from triplicate assays from three independent experiments (P values ≤ 0.05 (*) or ≤ 0.01 (**) vs. control).

Figure 1

Figure 2

Figure 3

Figure 4

Graphical Abstract