# Intracellular anti-leishmanial effect of Spergulin-A, a triterpenoid saponin of *Glinus oppositifolius*

Saswati Banerjee [a¥], Niladri Mukherjee [a¥], Rahul L. Gajbhiye [b], Parasuraman Jaisankar [b], Sriparna Datta[c], Krishna Das Saha[a]*

[a]Cancer Biology and Inflammatory Disorder Division, CSIR-Indian Institute of Chemical Biology, 4 Raja S. C. Mullick Road, Kolkata-700032, India

[b]Organic and Medicinal Chemistry Division, CSIR-Indian Institute of Chemical Biology, 4, Raja S.C. Mullick Road, Kolkata 700032, India

[c]Department of Chemical Technology, University of Calcutta, Kolkata 700009, India

Email address: krishna@iicb.res.in (Krishna Das Saha); [¥] Equal contribution, *Corresponding Author

## ABSTRACT

Many of present chemotherapeutics are inadequate and also resistant against visceral

leishmaniasis (VL) an immunosuppressive ailment caused by *Leishmania donovani*.

Despite the interest in plant-based drug development, very few number of

antileishmanial drugs from plant source is available. *Glinus oppositifolius* had been

reported in favor of being immunomodulators along with other traditional uses. Novel

anti-VL therapies can rely on host immune-modulation with associated leishmanicidal

action. With this rationale, an n-BuOH fraction of the methanolic extract of the plant

and obtained triterpenoid saponin Spergulin-A were evaluated against acellular and

intracellular *L. donovani*. Immunostimulatory activity of them was confirmed by

elevated TNF-α and extracellular NO production from treated MΦs and was found

nontoxic to the host cells. Identification and structure confirmation for isolated

Spergulin-A was performed by ESI-MS, $^{13}$C, and $^{1}$H NMR. Spergulin-A was found

ineffective against the acellular forms while, against the intracellular parasites at

30μg/ml, the reduction was 92.6% after 72h. Spergulin-A enhanced ROS and nitric

oxide (NO) release and changes in Gp91-phox, i-NOS, and pro and anti-inflammatory

cytokines elaborated its intracellular anti-leishmanial activity. The results supported

that *G. oppositifolius* and Spergulin-A can potentiate new lead molecules for the

development of alternative drugs against VL.

## Key Words

## Introduction

Visceral Leishmaniasis (VL) considered as the most severe form of
leishmaniasis caused by *Leishmania donovani* and untreated patients nearly always die
[1]. Leishmaniasis is endemic in nearly 100 countries with approximately 350 million
people are at risk with an estimated yearly incidence of 500,000 cases and almost
70,000 deaths.  Leishmaniasis is liable for the ninth most substantial infectious diseases
burden, however, is mostly disregarded among tropical disease priorities [2]. Sodium
antimony gluconate, Miltefosine, Pentamidine, and Amphotericin B are the primary
therapeutics though associated with toxicity or resistance [3].

Pathogen recognition by neutrophils, macrophages, dendritic cells, and natural
killer cells activates intracellular signaling pathways leading into inflammatory
responses like inflammasome activation and IL-1β production, which is not the case for
leishmanial infection [4]. *L. donovani* infection is characterized by the parasite-induced
active subversion of the host immune system and also immune deviation which
additively favors infection establishment [5]. Additionally, *Leishmania* prevents
inflammatory response by the impaired release of different pro-inflammatory cytokines
(IL-1, TNF-α, IL-12) and enhanced releases of immunosuppressive signaling molecules,
such as arachidonic acid metabolites and the cytokines TGF-β and IL-10 [6].
Involvement of co-stimulatory molecule B7-CTLA-4 is associated with increased TGF-
ß production in VL with increased apoptosis of $CD4^+$ T cells and decreased macrophage
apoptosis [7]. Leishmanial infection also undermines a generation of microbicidal
macrophage nitric oxide (NO) and reactive oxygen species (ROS) with the hindrance of
antigenic peptide display to T cells, and permeation of IL-10 producing T regulatory
cells [8]. In recent years increased instances of VL has been reported in connection with
immune-suppressed AIDS patients [8].

Discovery of novel compounds which intercedes host immune-modulation

associated with leishmanicidal function having permissible side effects is a precise

research objective [9,10]. It is also of vital importance that the drug required for parasite

elimination in immune-stimulated cells was significantly less than the immune-

suppressed ones [11].

The anti-parasitic effectiveness of plant extracts majorly relies upon secondary

metabolites of diverse chemical groups including alkaloids, polyphenol, flavonoids,

terpenoids, phenylpropanoids, etc. [12]. For isolation and characterization of a herbal

extract or an active compound, different research strategies can be employed mainly in

the extraction steps. However, bioactivity-guided fractionation considered as simple,

rapid, cost-effective and reproducible [1]. For obtaining a potent anti-leishmanial agent

with immune-modulation, different studies were conducted with compounds like

aslicarin A, niranthrin, skimmianine, quassin, tannins, linalool, etc. nevertheless with

varying degree of effectiveness and satisfaction [8].

Aerial parts of *Glinus oppositifolius* (Family Molluginaceae) are used for

treating abdominal pain and jaundice, while decoction is used against malaria [13].

Plants of this genus were previously documented for the presence of triterpenoid

saponins [14], and isolated pectin polysaccharides are antiprotozoal [15] and

immunomodulators [14]. *G. oppositifolius* is indicated for wound healing and used in

traditional medicine for treating diarrhea, joint pains, inflammations, intestinal

parasites, fever, boils and skin disorders [16]. Some triterpenoid saponins, 3-O-(β-

D-xylopyranosyl)-spergulagenin-A, Spergulacin, Spergulacin-A Spergulin-A, and

Spergulin-B had been isolated from *G. oppositifolius* [17]. Application of plant-based

immunomodulators is smart as they mediate their effectiveness by enhancing the

inherent host-derived protective machinery without the involvement of specific microbicidal agents namely antibiotics [13].

Most of the available antimalarial drugs are plant-derived; regrettably, there is no anti-leishmanial drug present which is of plant origin. The recent efforts to achieve this goal also were restricted against the promastigotes [18]. In the present work, an attempt has been taken to evaluate the intracellular anti-leishmanial activity of this plant and its bioactive component, Spergulin-A emphasizing on the immunostimulatory activity.

## Materials and methods

### *Chemicals and Reagents*

Cell culture media, serum, antibiotics, HEPES (Gibco, USA), CFSE, DAPI (Invitrogen, USA), MTT, Miltefosine, DMSO (Sigma Chemical Co. USA), cytokine Assay kit (Thermo Scientific, USA), DAF-2 DA, nitric oxide assay kit, DCFDA (Calbiochem, USA) and all other chemicals were of the highest grade commercially available. Primary and secondary antibodies were obtained from Santa Cruz Biotechnology (USA) or Cell signaling technologies (USA).

### *Isolation of methanolic fractions from G. oppositifolius*

The aerial parts of *G. oppositifolius* were shade dried (1kg) and were first defatted using petroleum ether (60-80$^{o}$C, 3.5L X 3) at room temperature for 48h. The marc was than subjected to extraction using MeOH (3.5L X 3) at room temperature for 48h. The extract was then filtered and MeOH was evaporated under reduced pressure and finally was lyophilized to obtain the crude MeOH extract (13g). A part (10g) of ME was then suspended in milli-Q water and partitioned sequentially with EtOAc and n-BuOH. Each fraction was evaporated under vacuum and lyophilized to

yield the EtOAc fraction (EAF; 3.6g), n-BuOH fraction (NBF; 4.1g) and aqueous

fraction (AF; 2.3g). All the fractions were stored at 4ºC till further use.

Among these four fractions, n-BuOH showed promising antileishmanial activity.

Around 4g of NBF was subjected to column chromatography of Diaion HP 20 (100g)

and the column washed with water followed by 30, 40, 60, 80 and 100% of MeOH to

obtain a total of six fractions.  Fractions eluted with 50% MeOH showed similar spots

on TLC and were mixed and then re-chromatographed on Dianion HP 20 column to

furnish 7mg of Spergulin-A. The compound isolation procedures were performed as

followed by Kumar et al. [19].

*Isolation and characterization of Spergulin-A*

ESI Mass spectra were recorded on an Agilent 6545 Q-TOF mass spectrometer

System. [1]H and [13]C NMR were recorded on a Bruker Ultrashield NMR (600 MHz) in

pyridine-d5 with TMS as an internal standard. Diaion HP 20 was used for column

chromatography; silica gel (60 F254) was used for TLC and spots were visualized by

spraying with Lieberman–Burchard reagent followed by heating.

*Parasite culture and maintenance*

Passage    in    BALB/c    mice    maintained    *Leishmania    donovani*

[MHOM/IN/1983/AG83] infection. Complete M199 media (10% FBS, pH 7.4,

100U/ml penicillin G-sodium, 100μg/ml streptomycin sulfate, 25mM HEPES) was used

to culture the promastigotes. Axenic  amastigote  from  the  log  phase  culture  of

promastigotes was prepared as described by Saar et al. [20].

*Macrophage culture, parasite infection, and treatment*

RAW 264.7 MΦ cell line was obtained from the ATCC and maintained in

complete RPMI 1640 (10% FBS, 100μg/ml streptomycin sulfate, 100U/ml penicillin G

sodium, 0.2% sodium bicarbonate, 25 mM HEPES) in a humidified atmosphere and 5%

$CO_2$ at 37°C.

Infection of MΦs with *L. donovani* promastigotes was performed in a ratio of

1:10 (MΦ: parasite) for 4h then washed twice with media. Normal and parasitized

macrophages were treated with *G. oppositifolius* fractions or Spergulin-A at required

concentrations up to 72h. Infection was measured by counting the intracellular parasites

and expressed as parasite count/20 MΦs. Miltefosine served as positive anti-leishmanial

reference.

***ELISA assay***

Cell-free supernatants from MΦs ($1\times10^5$ cells/well) were collected, and the

concentration of TNF-α was estimated by sandwich ELISA, using a commercially

available assay kit [21].

***MTT assay***

20μl of MTT (5mg/ml in PBS) was added to each well of the control and treated

MΦs ($4\times10^3$) and $5\times10^3$ parasites (promastigote and axenic amastigote) in a 96 well

plate and incubated for 4h at 37°C and the formazan crystals were dissolved in 150μl of

DMSO. Absorption was measured at 595nm by an ELISA reader [22].

***Measurement of extracellular NO***

The collected supernatants from control, infected and treated MΦs were

incubated with equal volumes of Griess reagent, NED (0.1% in distilled water) and

Sulphanilamide (1% in 5% $H_3PO_4$) at room temperature for 10min. The absorbance was

measured at 540nm on a microplate reader. NO concentration was determined using a

dilution of sodium nitrite as the standard [21].

*FACS and Confocal microscopy with CFSE tagged L. donovani*

MΦs were parasitized with CFSE-tagged ($25\mu M/1\times10^6$ promastigote in 1ml

media for 30min) promastigotes followed by Spergulin-A (10,20,30μg/ml) treatment

for 24h.

For confocal microscopy, MΦs ($1\times10^5$/glass coverslips) after infection and

treatment, washed twice with PBS, fixed in chilled 70% ethanol and nuclei were stained

with DAPI and observed under Olympus Fluoview FV10i confocal microscope with

60X objective lens. For FACS, MΦs ($1\times10^5$/well in a six-well plate) after infection

and treatment, washed twice with PBS then harvested by scraping and re-suspended in

400μl PBS and analyzed by BD FACS LSR Fortessa with excitation at 494nm and

emission at 518nm.

*FACS analyses of intracellular ROS and NO*

The level of intracellular ROS was determined based on the change in

fluorescence of H2DCFDA. DAF-2 DA was used for detecting intracellular NO [3].

Briefly, after infection and treatment, MΦs were scrapped and incubated in PBS

containing DAF-2DA (7.0μM) at 37°C for 30min or 20μM DCF-DA at 37°C for 15min

and then analyzed by BD FACS LSR Fortessa with excitation at 480nm and emission at

515nm for both.

*Western blot analysis*

40μg of proteins harvested from MΦs were electrophoretically separated in

SDS-polyacrylamide gel and transferred to PVDF membrane, blocked with BSA and

incubated with respective primary antibodies overnight. The membranes were then

incubated with HRP conjugated secondary antibodies, and immunoreactive bands were

visualized by adding proper substrates. β-actin was used as loading controls [23].

*Statistical analysis*

All values were expressed as mean ± SEM obtained from at least three replicate

experiments. Statistical significance and differences among groups were assessed with

One-Way analysis of variance (ANOVA) followed by Dunnett's test. P values ≤ 0.05 (*)

or ≤ 0.01 (**) were considered as indicative of significance.

**Results**

*Immunostimulatory effect of methanolic extract of G. oppositifolius*

Of the aqueous, ethyl acetate and n-BuOH (50:50) fractions of *G. oppositifolius*

methanolic extract, only the n-BuOH fraction (50µg/ml) showed a considerable increase

in TNF-α and extracellular NO production (Supple. Figure 1) from the treated (24h)

MΦs. The six sub-fractions (50µg/ml) of the n-BuOH fraction were also checked for the

same parameters (Figure 1A and B), and highest augmentation was noticed in fraction 4

for being a worthy immunostimulatory agent and the lead fraction.

*Dose-dependent immunostimulatory effect of the lead fraction*

A dose-dependent increase in TNF-α production was monitored in treated (24h)

MΦs in two sets (Figure 2B and C). In first set the lead fraction was applied at the

doses of 10, 20, 40 and 80µg/ml (Figure 2B) and in the second set at the doses of 20,

30, 40 and 50µg/ml and most increments was noticed at 30µg/ml (Figure 2C) and

selected as the significant dose for this study.

*Assessment of macrophage survival and extracellular NO release*

Survival of the MΦs after exposing them to increasing concentrations

(10,20,30,40 and 50µg/ml) of the lead fraction for 24h was monitored by MTT assay

(Figure 2D). The lead fraction was found safe for this dose range, and survival was

about 81.45% even at 50µg/ml. Most increment (5fold) in extracellular NO was noticed

for 30μg/ml of the lead fraction (Figure 2E). The increase was also significant for most of the doses.

### *Identification and characterization of Spergulin-A*

Among those six n-BuOH bioactive sub-fractions, fraction 4 have shown most significant upliftment in TNF-α and extracellular NO production. After purification of this fraction by using column chromatography, a white amorphous compound was obtained. Molecular formula $C_{35}H_{58}O_{11}S$ (Figure 3A) was assigned from the ESI-MS (Supple. Figure 2)**.** Structural elucidation of this compound was achieved by critical analysis of the $^1$H NMR, $^{13}$C NMR and $^{13}$C NMR DEPT results (Supple. results and Supple. Figure 3-6). All the spectral data of this compound were found to be in complete agreement with reported ones and structure was then confirmed as Spergulin-A [24].

### *Assessment of anti-leishmanial activity of Spergulin-A*

#### *Effect on different forms of L. donovani*

Anti-leishmanial effect of Spergulin-A (10,20,30,50,100μg/ml) was first evaluated by MTT assay against promastigote and axenic amastigote, the acellular forms (Figure 3B) after 24h. Viability was 87.6% and 77.8% respectively for promastigote and axenic forms even at 100μg/ml. At the lower doses (10, 20 and 30μg/ml) the viability reduction was found to be non-significant (Figure 3B).

After that, the effect of Spergulin-A was evaluated against the MΦ internalized parasites (Figure 3C). Here, a significant reduction of the parasite was noticed as low as 10μg/ml and further increased with the increment of doses (Figure 3C). However, above 30μg/ml MΦ internalized antileishmanial effect of Spergulin-A had reached a plateau, and thus further experiments were mostly conducted with 30μg/ml.

*Time-dependent intracellular leishmanicidal effect*

Efficacy of Spergulin-A (30μg/ml) against the intracellular parasites was

evaluated at every 24h interval up to 72h (Figure 3D). Inside the infected MΦs parasite

count increased significantly. The treated MΦs exhibited significantly reduced numbers

of parasites from the initial infection as well as the corresponding observation point of

infected macrophages (Figure 3D). The parasite reduction at 72h was 86.2% less than

the initial point of infection and 92.6% less than the 72h count of infected MΦs.

Miltefosine was used in parallel as a positive reference anti-leishmanial compound and

$LC_{50}$/24h and $LC_{90}$/24h values against these three forms were expressed as histogram

(Figure 3E).

*Dose-dependent quantitative and qualitative anti-leishmanial effect*

After infecting the MΦs with CFSE-tagged promastigotes and treatment with

Spergulin-A (10, 20 and 30μg/ml, 24h), the cells were evaluated by FACS in FITC filter

(Figure 4A). Highest intensity was noticed in the infected panels (Figure 4A.ii) and

with the increment of Spergulin-A doses the intensity gradually reduces which is

reciprocal to the reduced parasite count (Figure 4A.iii-v). Interestingly, no significant

changes in number of parasitized MΦs observed within the treatment panel (Figure

4A.vi) and what changes were the intensity denoted the parasite count (Figure 4A.vii).

Thus, possibly Spergulin-A reduced the parasite by immune-stimulating the host MΦs.

Confocal micrographs depicted the reduction in parasite emitting green

fluorescence with the increment of Spergulin-A doses. Moreover, most decreases were

noticed when the parasitized MΦs was treated with 30μg/ml of it (Figure 4B).

 *Intracellular ROS and NO status due to infection and treatment*

The fluorescence intensity for H2DCFDA reciprocal to ROS production was

found to be minimal for the infected panel (Figure 4C.ii) and significantly higher in the

treated panel (Figure 4C.iii). The fluorescence intensity of DAF-2DA for NO was also

found to be highest for the treated panel (Figure 4D.iii) compared to the control and

infected MΦs (Figure 4D.i and ii).

*Estimation of alteration of different cytokines and immunostimulators*

In western blot analysis, it was observed that Gp91-phox liable for anti-

microbial ROS production in MΦs got up regulated in treated panel (Figure 4E). Level

of i-NOS which regulates anti-leishmanial NO production also increased with

Spergulin-A treatment compared to control and infected MΦs (Figure 4F).

Expression of pro-inflammatory TNF-α and IL-12β got down-regulated with

infection and considerably up-regulated in treatment panel (Figure 4G). Anti-

inflammatory IL-10 and TGF-β expression were up-regulated in infected MΦs

compared to the control, and with the treatment of Spergulin-A (30μg/ml) their

appearance got normalized like that of control (TGF-β) or even less (IL-10) (Figure

4G).

## Discussion

Precisely an immune-suppressive ailment, VL inhabit and modulate the

microbicidal function of the macrophages and create a microenvironment favoring

parasite growth inside the visceral organs by modulating pro and anti-inflammatory

cytokines and impairing ROS and NO release [6]. The present *in vitro* study aimed to

assess the immunostimulatory property of *G. oppositifolius* and isolation of a compound

Spergulin-A which can reduce the intracellular parasites by immune-stimulation within

a safe dose range for the host cells. Primary appraisal of an extract that can have an

immunostimulatory effect was based upon the increased release of TNF-α and

extracellular NO from treated MΦs. It was observed that at an applied same dose

(50μg/ml, 24h) only the n-BuOH fraction showed a considerable increase in TNF-α and

extracellular NO production. The selection of TNF-α and extracellular NO as the potent

immunostimulatory index in the present context is vital because endogenous TNF-α

produced by infected macrophages can elicit the release of L-arginine-derived nitrogen

intermediates detrimental for the intracellular parasites [9]. The six n-BuOH sub-

fractions were again verified for immune-stimulation, and fraction 4 emerged as the

most promising one. The fraction was also considered to be safe for the MΦs even at

50μg/ml (81.5% viability) having $CC_{50}$ well above 100μg/ml.

Interested by the initial observations on elevated immunostimulation by the n-

BuOH sub-fraction 4 and based upon the previous inspection of the presence of

triterpenoid saponins in this fraction [24] a triterpenoid saponin Spergulin-A was

isolated and purified following the well-established method (Fig. 3A and Supple.

information) [24,19].

Evaluation of the anti-leishmanial property of Spergulin-A was the principal

research interest after identification and isolation of it. The possible anti-leishmanial

effect was first checked against the acellular forms of the parasite but without major

success even at 100μg/ml (Figure 3B). Therefore, if Spergulin-A has any leishmanicidal

effect on the intracellular parasites that will possibly mediate by altering the

microenvironment of the host MΦs. After 24h of incubation Spergulin-A (30μg/ml) was

found to be pretty useful in reducing the intracellular parasite count and when applied

for a prolonged period of 72h reduction of intracellular *Leishmania* was significant in

respect to both the infected MΦs of corresponding time point and initial parasitized

MΦs (Figure 3C,D). It is also of great significance that no considerable alteration in

number of parasitized MΦs was found in treatment panel while reduction of CFSE-

stained internalized parasites was notable (Figure 4A). This specific observation

strongly advocated in favor of immunostimulation by Spergulin-A which mediated

parasite killing. For being leishmanicidal, Spergulin-A should be useful in enhancing

the release of NO and ROI, well recognized for their worth against *Leishmania* [6,25].

NO is especially critical for intracellular parasite clearance as it was reported that mice

with impaired inducible nitric oxide synthase (i-NOS) and thus restricted production

and release of NO are incapable of *Leishmania* control even for isolated macrophages *in*

*vitro* [26]. At the dose of 30μg/ml, Spergulin-A, showed enhanced production of

intracellular NO and ROI (Fig. 4C,D) which was found to be directly reciprocal to the

reduction of intracellular parasites and signified their connection. In the western blot

analysis also increment in i-NOS production (Figure 4F) was noticed in Spergulin-A

treated parasitized MΦs in amendable enhanced NO production and parasite control.

The same interpretation is also pertinent for an increase of Gp91-phox, a subunit of the

NADPH oxidase and enhancement of ROI (Figure 4 E,C) in Spergulin-A mediated

intracellular parasite killing. It was previously established that intracellular leishmanial

killing proceeds by NO production from arginine by i-NOS, and superoxide ($O_2^-$)

generated by the NADPH oxidase [27]. Several Leishmania species induce

immunosuppressive TGF-β production, and diminution of TGF-β secretion is in direct

correlation with enhanced i-NOS production [6] and can lead to internalized parasite

removal as found with Spergulin-A treatment. TGF-β also found to increase the VL

progression and averts disease cure in murine models [27]. IL-10 is another anti-

inflammatory cytokine which makes macrophages indifferent to various activation signals and thus causes impairment of intracellular parasite killing also by down-regulating the production of TNF-α and NO [28]. IL-10 production increased in infected macrophages *in vitro*, apparently via interaction with the Fc receptor [29] and which down-regulated with Spergulin-A treatment (Figure 4G) in connection with parasite reduction. Parasite infection is responsible for the suppression of macrophage microbicidal activity that relies upon NO, ROI and cytokines like IL-1, IL-12β, and TNF-α [30] as with infection the levels of TNF-α IL-12β got down-regulated, but with the application of Spergulin-A, these cytokines significantly elevated which signifies their role in intracellular *Leishmania* parasite control.

## Conclusion

The inconsistency of direct leishmanicidal effect of *G. oppositifolius* and Spergulin-A isolated from it against promastigotes and axenic amastigote on the one hand and the proved efficacy of them against intracellular parasites was explained by emphasizing host MΦ immunostimulation as elaborated precisely in the present study. Bio-guided fractionation was employed to isolate and identify Spergulin-A as the immunostimulant, and its anti-leishmanial property was evaluated at a dose best suited for the research and found to be safe against the host cells assessed *in vitro* in RAW 264.7 MΦs infected with a virulent strain of *L. donovani*.

## Acknowledgment

Sincere thanks are given to Science and Engineering Research Board, Govt. of India (Grant No. PDF/2016/001437 and EMR/2015/001674) for financial assistance. The manuscript has been checked critically by the full version of the Grammarly software

353 and corrected by Dr. Basudeb Achari, ex-scientist of CSIR-Indian Institute of Chemical

354 Biology.

## References

1) Tiuman TS, Santos AO, Ueda-Nakamura T, Filho BP, Nakamura CV. Recent

advances in leishmaniasis treatment. Int J Infect Dis 2011;15:e525–32.

2) Alvar J, Ve´lez ID, Bern C, Herrero M, Desjeux P, Cano J, et al. Leishmaniasis

Worldwide and Global Estimates of Its Incidence. PLoS One 2012;7:e35671.

3) Banerjee S, Bose D, Chatterjee N, Das S, Chakraborty S, Das T, et al.

Attenuated *Leishmania* induce pro- inflammatory mediators and influence

leishmanicidal activity by p38 MAPK dependent phagosome maturation in

*Leishmania donovani* co-infected macrophages. Sci Rep 2016;6:22335.

4) Freitas EO, Leoratti FMS, Freire-de-Lima CG, Morrot A, Feijó DF. The

Contribution of Immune Evasive Mechanisms to Parasite Persistence in Visceral

Leishmaniasis. Front Immunol 2016;7:153.

5) Bhardwaj S, Srivastava N, Sudan R, Saha B. *Leishmania* interferes with host

cell signaling to devise a survival strategy. J Biomed Biotechnol 2010;2010:109189.

6) Olivier M, Gregory DJ, Forget G. Subversion mechanisms by which *Leishmania*

parasites can escape the host immune response: a signaling point of view. Clin

Microbiol Rev 2005;18:293–305.

7) Goto H, Lindoso JAL. Immunity and immunosuppression in experimental

visceral leishmaniasis. Braz J Med Biol Res 2004;37:615–23.

8) Islamuddin M, Chouhan G, Want MY, Ozbak HA, Hemeg HA, Afrin F.

Immunotherapeutic potential of eugenol emulsion in experimental visceral

leishmaniasis. PLoS Negl Trop Dis 2016;10:e0005011.

9)  Maran N, Gomes PS, Freire-de-Lima L, Freitas EO, Freire-de-Lima CG, Morrot

A. Host resistance to visceral leishmaniasis: prevalence and prevention. Expert Rev

Anti Infect Ther 2016;14:435–42.

10) Flora  R,  Aghazadeh-Dibavar  S,  Bandyopadhyay  M,  Dasgupta  S.

Immunosuppression during *Leishmania donovani* infection: a potential target for the

development of therapy. Ann Parasitol 2014;60:239–45.

11) Haidaris CG, Bonventre PF. Efficacy of combined immunostimulation and

chemotherapy  in  experimental  visceral  Leishmaniasis.  Am  J  Trop  Med  Hyg

1983;32:286–95.

12) Mukherjee N, Mukherjee S, Saini P, Roy P, Sinha Babu SP. Phenolics and

Terpenoids; the Promising New Search for Anthelmintics: A Critical Review.

Mini Rev Med Chem 2016;16:1415–41.

13) Inngjerdingen KT, Kiyohara H, Matsumoto T, Petersen D, Michaelsen TE,

Diallo D, et al. An immunomodulating pectic polymer from *Glinus oppositifolius.*

Phytochemistry 2007;68:1046–58.

14) Sahakitpichan P, Disadee W, Ruchirawat S, Kanchanapoom T. L-()-(N-trans-

Cinnamoyl)-arginine, an acylamino acid from *Glinus oppositifolius* (L.) Aug. DC.

Molecules 2010;15:6186–92.

15) Traore F, Faure R, Ollivier E, Gasquet M,  Azas N,  Debrauwer L, et al.

Structure  and  antiprotozoal  activity  of  triterpenoid  saponins  from  *Glinus*

*oppositifolius*, Planta Med 2000;66:368–71.

16) Sahu SK, Das D, Tripathy NK, Dinda SC, Sundeep Kumar HK. Evaluation of

hypoglycemic  activity  of  *Mollugo pentaphylla*  and  *Glinus oppositifolius*  (L).

Rasayan J Chem 2012;5:57–62.

17) Kumar D, Shah V, Ghosh R, Pal BC. A new triterpenoid saponins from *Glinus*

*oppositifolius* with a-glucosidase inhibitory activity. Nat Prod Res 2013;27:624–29.

18) Kaur H, Thakur A, Kaur S. Immunoprophylactic Potential of a Cocktail of

Three Low Molecular Weight Antigens of *Leishmania donovani* along with Various

Adjuvants Against Experimental Visceral leishmaniasis. Iran J Parasitol 2018;13:11–

23.

19) Kumar D, Mallick S, Vedasiromoni JR, Pal BC. Anti-leukemic activity of

*Dillenia indica* L. fruit extract and quantification of betulinic acid by HPLC.

Phytomedicine 2010;17:431–35.

20) Saar Y, Ransford A, Waldman E, Mazareb S, Amin-Spector S, Plumblee J, et al.

Characterization of developmentally-regulated activities in axenic amastigotes of

*Leishmania donovani*. Mol Biochem Parasitol 1998;95:9–20.

21) Chatterjee N, Das S, Bose D, Banerjee S, Das S, Chattopadhyay D, et al.,

Exploring the anti-inflammatory activity of a novel 2-phenylquinazoline analog with

protection against inflammatory injury. Toxicol Appl Pharmacol 2012;264:182–91.

22) Mosmann T. Rapid colorimetric assay for cellular growth and survival:

application to proliferation and cytotoxicity assays. J Immunol Methods 1983;65:55–

63.

23) Sikdar Y, Modak R, Bose D, Banerjee S, Bieńko D, Zierkiewicz W, et al. Doubly

chloro bridged dimeric copper(II) complex: magneto-structural correlation and

anticancer activity. Dalton Trans 2015;44:8876.

24) Sahu NP, Koike K, Banerjee S, Achari B, Nikaido T. Triterpenoid saponins

from *Mollugo spergula*. Phytochemistry 2001;58:1177-82.

25) Matte C, Maion G, Mourad W, Olivier M. *Leishmania donovani*-induced

macrophages cyclooxygenase-2 and prostaglandin E2 synthesis. Parasite Immunol

2001;23:177–84.

26) Wei XQ, Charles IG, Smith A, Ure J, Feng GJ, Huang FP, et al. Altered

immune responses in mice lacking inducible nitric oxide synthase, Nature

1995;375:408–11.

27) Wilson ME, Jeronimo SMB, Pearson RD. Immunopathogenesis of infection

with the visceralizing Leishmania species. Microb Pathogenesis 2005;38:147–60.

28) Nylen S, Sacks D. Interleukin-10 and the pathogenesis of human visceral

leishmaniasis. Trends Immunol 2007;28:378–84.

29) Sutterwala FS, Noel GJ, Salgame P, Mosser DM. Reversal of proinflammatory

responses by ligating the macrophage Fcgamma receptor type I. J Exp Med

1998;188:217–22.

30) Kane MM, Mosser DM. Leishmania parasites and their ploys to disrupt

macrophage activation. Curr Opin Hematol 2000;7:26–31.

**Figure Captions**

**Fig. 1.** Evaluation of immunostimulation by subfractions of n-BuOH fraction of *G.*

*oppositifolius* MeOH extract.

(A) The subfractions (50μg/ml) were checked for altered TNF-α production after 24h

treatment of RAW 264.7 MΦs from the culture supernatant. (B) Extracellular NO

production was monitored in the culture supernatant of treated (six subfractions) and

control after 24h by Griess reagent. All values are expressed as mean ± SEM from

triplicate assays from three independent experiments (P values ≤ 0.05 (*) or ≤ 0.01 (**)

vs. control).

**Fig. 2.** Dose-dependent evaluation of the immunostimulatory effect of the n-BuOH lead

fraction and impact on RAW 264.7 MΦ survival.

(A) TLC profile of n-BuOH lead fraction. (B, C) The dose-dependent release of TNF-α

was monitored in treated (24h) RAW 264.7 MΦs, and highest increment was found in

30μg/ml. (D) Survival of RAW 264.7 MΦs at different doses of an n-BuOH lead

fraction after 24h exposure measured by MTT assay. (E) Dose-dependent evaluation of

extracellular NO production was monitored in the culture supernatant of 24h treated

RAW 264.7 MΦs by Griess reagent. All values are denoted as mean ± SEM from

triplicate assays from three independent experiments (P values $\leq 0.05$ (*) or $\leq 0.01$ (**)

vs. control).

**Fig. 3.** Anti-leishmanial activity of Spergulin-A.

(A) Structure of the triterpenoid saponins Spergulin-A. (B) Dose-dependent effects of

Spergulin-A against acellular forms namely the promastigotes and axenic amastigote of

*L. donovani* after 24h. (C) Dose-dependent effect of Spergulin-A against the

intracellular amastigote of *L. donovani* after 24h of treatment and expressed as number

of parasites/20 MΦs. (D) 24h time lapse evaluation of anti amastigote effect of

Spergulin-A at 30μg/ml for 72h. (E) Miltefosine $LC_{50}$/24h and $LC_{90}$/24h values against

promastigote, axenic amastigote and intracellular amastigote forms of *L. donovani*. All

values are expressed as mean ± SEM from triplicate assays from three independent

experiments (P values $\leq 0.05$ (*) or $\leq 0.01$ (**) vs. control).

**Fig. 4.** Elaboration of the Anti-leishmanial activity of Spergulin-A.

(A) Quantitative dose-dependent assessment of intracellular CFSE stained

antileishmanial activity of Spergulin-A by flow cytometry after 24h treatment. (B)

Confocal laser-scanning micrographs to evaluate the reduction in intracellular CFSE

stained *L. donovani* after 24h of Spergulin-A treatment (magnification 120X). (C)

FACS monitored the changes in ROS production in control, infected, and treated

(30μg/ml Spergulin-A) MΦs with H2DCFDA. (D) The changes in intracellular NO

production were monitored by FACS with fluorescent probe DAF-2 DA. (E, F, G)

Changes in the expression of Gp91-Phox, i-NOS and cytokines namely pro-

inflammatory TNF-α, IL-12β and anti-inflammatory IL-10 and TGF-β with β-actin as

loading control and densitometry analyses was provided after band normalization.

Images are representative of three separate experiments, and all values are expressed as

mean ± SEM from triplicate assays from three independent experiments (P values ≤

0.05 (*) or ≤ 0.01 (**) vs. control).