Parallel patterns of development between independent cases of hybrid seed

inviability in *Mimulus*

Jenn M. Coughlan[1,2], John H. Willis[1]

1. *Biological Sciences, Duke University 25 Science Drive, Durham NC*

2. *Biology Department, University of North Carolina, Chapel Hill*

Author Contributions: *JMC designed this project design, performed data collection and*

analysis, and wrote the manuscript. JHW contributed substantially to the ideas presented

here and the project design.

Key Words: *Cryptic Species, Endosperm, Parental Conflict, Reproductive Isolation*

Running headline: *Parental conflict and hybrid seed inviability in Mimulus*

Summary

Rationale: The evolutionary drivers of many reproductive barriers are unknown. Hybrid seed inviability (HSI) is a common barrier in angiosperms. We test the idea that conflict between maternal and paternal interests in resource allocation to developing offspring (i.e. parental conflict) drives HSI in two independent incidences of HSI.

Methods: We quantified HSI between *M. guttatus* and two clades of *M. decorus* with oppositely asymmetric HSI. In the northern, weaker-conflict clade, seeds are inviabile when *M. decorus* is the mother, while in the southern, stronger-conflict clade, inviability occurs when *M. decorus* is the father. We quantified whole-seed development for hybrids and parents to assess whether reciprocal F1 seeds show parent-of-origin dependent growth phenotypes, as predicted by conflict.

Key Results: Crosses between *M. guttatus* and both clades of *M. decorus* show parent-of-origin effects on reciprocal F1 seed development. Inviable hybrid seeds exhibit paternal-excess phenotypes, wherein endosperm is large and chaotic. In contrast, viable hybrid seeds show maternal-excess phenotypes, wherein endosperm development is precocious.

Main Conclusions: We find strong parent-of-origin effects on development in reciprocal F1s that are repeated across independent incidences of HSI in *Mimulus.* These patterns suggest that parental conflict may play a strong role in shaping the evolution of HSI.

Introduction

The repeatability of speciation is a contentious topic in evolutionary biology. Much pre-zygotic reproductive isolation can be attributed to adaptations to the abiotic or biotic environment (Coyne & Orr 2004), and the repeatability of extrinsic reproductive isolation may be directly related to the repeatability of adaptation (i.e. Rundle *et al.* 2000; Rolán-Alvarez *et al.* 2004; Stankowski 2013; Soria-Carrasco *et al.* 2014; Stuart *et al.* 2017). Yet, intrinsic post-zygotic reproductive isolation also often manifests as canonical hybrid phenotypes (i.e. abdominal ablation in hybrid *Drosophila* species pairs- Orr *et al.* 1997; Gavin-Smyth & Matute 2013; CMS hybrid sterility phenotypes in plants- reviewed in Hanson & Bentolila 2004; Chase 2007; Rieseberg & Blackman 2010; hybrid necrosis in multiple plant and fungi species pairs- reviewed in Bomblies & Weigel 2007), and much less is known about the evolutionary mechanisms that cause this repeatability at the phenotypic level. Moreover, in many systems it is unclear whether these common and repeated incompatibility phenotypes are developmentally similar, and whether this can inform our understanding of the drivers of hybrid incompatibilities. A classic hybrid incompatibility phenotype in plants is that of hybrid seed inviability (HSI).

Proper seed development is achieved through the effective coordination of three tissues with differing genetic compositions: the maternally derived seed coat, the developing embryo, and a nutritive tissue that sustains the embryo until germination- the endosperm. In diploid species, seed development begins with a double fertilization event; one in which the ovule is fertilized and the developing embryo is formed, while the other involves the fertilization of a diploid central cell that gives rise to a triploid endosperm tissue. The endosperm is thus comprised of 2:1 maternal:paternal genetic composition. This ratio is essential for appropriate seed development, because many genes exhibit parent-of-origin specific expression in the endosperm (i.e. imprinting; Köhler & Weinhofer-Molisch 2010). In *Arabidopsis*, this is caused primarily by DNA methylation and histone methylation (Köhler *et al.* 2012), the later of which are controlled by the Fertilization Independent Seed- Polycomb Group Complex (FIS-PcG complex; Köhler *et al.* 2003; Köhler *et al.* 2005; Gehring *et al.* 2006; Makarevich *et al.* 2006; Köhler &

Makarevic 2006; Gehring *et al.* 2009; Hsieh *et al.* 2009; Wolff *et al.* 2011; Waters *et al.*

2011; Derkacheva *et al.* 2016). Indeed, imprinting in plants is thought to have evolved as

a mechanism to decrease conflict over paternal and maternal interests in nutritive tissue

growth (i.e. parental conflict; Trivers 1974; Haig & Westoby 1989).

Parental conflict is the hypothesis that maternal and paternal interests in resources

allocation to offspring differ in non-monogamous systems (Trivers 1974; Haig &

Westoby 1989). As maternity is guaranteed, selection should favor equal distribution of

resources from mothers to offspring, but when paternity is variable, selection should

favor the evolution of paternally derived alleles which increase resource expenditure on

developing offspring (Trivers 1974; Haig & Westoby 1989). If populations have evolved

differentially in this co-evolutionary arms race, then mismatches between resource-

restrictive maternal alleles and resource-acquiring paternal alleles in hybrids can lead to

developmental defects, and ultimately, death. One prediction of parental conflict is that

reciprocal F1s should show significantly different developmental trajectories, wherein

strong-conflict populations should produce growth-restrictive phenotypes when the

maternal contributor, and growth-excessive phenotypes when the paternal contributor in

crosses with weaker conflict populations.

In *Arabidopsis* hybrid seed inviability in interploidy crosses, maternal-restrictive

and paternal-excess phenotypes are observed (Scott *et al.* 1998). Inviability in this case

arises because of de-repression of normally imprinted genes, particularly that of *ADM,*

SUVH7, PEG2, and *PEG9*, which are all associated with carbohydrate sequestration and

cellularization in normally developing seeds (Kradofler *et al.* 2013; Wolff *et al.* 2015).

Overexpression of these alleles causes excessive growth and improper cellularization of

the endosperm associated with paternal excess, while maternal excess is often associated

with precocious endosperm development (Scott *et al.* 1998; reviewed in Lafon-Placette &

Köhler 2016). These defects in the endosperm are causative to hybrid inviability; both

embryo rescues and dosage rescue experiments strongly suggest that in these interploidy

crosses, hybrid seed inviability is due specifically to inappropriate development and

growth of the endosperm, rather than incompatibilities arising in the embryo itself

(Bushell *et al.* 2003; Josefsson *et al.* 2006; Dilkes *et al.* 2008; Erilova *et al.* 2009;

Kradolfer *et al.* 2013).

While hybrid seed inviability is quite common in diploid species pairs (Tiffin *et*

al. 2001), far fewer studies have determined the mechanistic underpinnings of hybrid

seed inviability among diploids. In plants with nuclear endosperm development, such as

Arabidopsis and *Capsella,* hybrid seed inviability mimic interploidy crosses with a

delayed or premature endosperm cellularization, depending on the maternal and paternal

donor in the cross, and endosperm defects are intrinsically linked with hybrid seed

inviability (Joseffson *et al.* 2006; Burkart-Waco *et al.* 2012; Rebernig *et al.* 2015; Lafon-

Placette *et al.* 2017l Lafon-Placette *et al.* 2018). In plants with cellular endosperm

development (i.e. *Solanum* and *Mimulus*), hybrid seed inviability is much less explored,

although there is some evidence to suggest that inappropriate replication of endosperm is

at least partially causative of hybrid seed inviability in both *M. nudatus* and *M. guttatus*

inviable hybrid seeds (Oneal *et al.* 2016), and in wild tomato varieties (Lafon-Placette *et*

al. 2016; Roth *et al.* 2017).

We previously reported a wide diversity of hybrid seed barriers between *M.*

guttatus and its morphological variant, *M. decorus* which confer relatively strong

reproductive isolation (Coughlan *et al.* 2018). We found that *M. decorus* is in fact

multiple biological species; a northern clade exhibits strong asymmetric hybrid seed

inviability when it is the maternal donor in crosses to *M. guttatus*, a southern clade that

exhibits strong asymmetric hybrid seed inviability when it is the paternal donor in crosses

to *M. guttatus,* and at least one tetraploid taxon that exhibits variable levels of

reproductive isolation via hybrid seed inviability. Here, we test whether hybrid seed

inviability between different diploid accessions of *M. decorus* and *M. guttatus* are

associated with similar developmental defects, whether these defects are endosperm

related, and whether they are consistent with paternal excess/maternal restriction,

hypothesized by parental conflict. In addition, we assess the level of developmental

parallelism of inviable and viable hybrid seeds between diploid species pairs that have

strongly asymmetric hybrid seed inviability, but in the opposite direction.

Methods

Growth conditions and plant crossing

Based on previous results, *Mimulus decorus* exhibits two distinct crossing patterns: the northern clade of *M. decorus* produces inviable seeds when they are the maternal plant in crosses with *M. guttatus*, while for the southern clade produces inviable seeds when they are the paternal plant in crosses with *M. guttatus*. To determine the extent of developmental parallelism, we chose a representative diploid population from each crossing type of *M. decorus* (northern clade: *IMP*; southern clade: *Odell Creek*) to cross with an annual population of *M. guttatus*- Iron Mountain (*IM*). For each population, we planted seed from 3-4 maternal families or inbred lines. For all plants, seeds were sprinkled onto moist Fafard 4-P soil, left to stratify at 4°C for 1 week, then moved to the Duke greenhouses. Plants were transferred within the first 3 days after germination to one plant per 2" pot, then were grown in long days (18h days, 21C day/ 18C nights).

Crossing Assays

Plants were emasculated and crossed reciprocally, as well as self pollinated to yield an average of 6 crosses for each pairwise combination of maternal families between IM and Odell Creek, and IM and IMP (IMxOdell range: 2-13 replicate crosses, IMxIMP: 2-10 replicate crosses for a total of 298 crosses). Fruits were allowed to develop until ripe but before dehiscence, then seeds for each cross were counted and categorized as inviable or viable based on morphology for each fruit.

Microscopy and measuring growth

We sought to determine the extent of developmental parallelism across hybrid seeds which show opposite patterns of inviability between these two populations of *M. decorus*. We next crossed plants, as in above, but collected fruits at 4,6,8,10 and 14 Days

After Pollination (DAP). Fruits were collected and immediately placed in FAA fixative.

Plants were processed in a similar manner to White & Turner (2012). In brief, after at

least 24 hours in fixative, fruits were gradually dehydrated in a Tetrabutyl-Alcohol

dehydration series, then mounted in paraffin wax with ~5% gum elemi resin to increase

hardness. Paraffin mounted specimens were then sliced to 8 micron ribbons and mounted

onto slides. We performed a staining series using Safranin-O and Fast-Green, which stain

for nucleic acids and carbohydrates, respectively. Cover slips were then adhered for

permanent storage.

167   We visualized and photographed at least three developing seeds from a single

168   fruit for each cross type and time point combination (40 unique combinations, 120 seeds

169   measured). At least 25 endosperm cell widths were measured per seed using ImageJ

170   (Scheider *et al.* 2012).

172   *Embryo Rescues*

173   To assess whether hybrid seed inviability was due largely to inappropriate

development of the endosperm tissue, we also performed embryo rescue assays on both

IMxOdell seeds and IMPxIM seeds, which would normally be inviable if left to develop

fully in the seed pod. For this, we crossed IM reciprocally to both IMP and Odell creek,

as well as performed self fertilizations for IMP, Odell Creek, and IM. Fruits were left to

mature for 7 or 9 days. Based on our developmental time series, embryos appear to die

between 10-14 DAP for both sets of crosses. We plated whole developing seeds for each

cross type on a sucrose rich growth media (1/2 MS salts, 0.7% agar, 4% sucrose, adjusted

to pH 5.9) at 7 and 9 DAP. Plates were cold stratified for 2 days, then placed in the Duke

Greenhouses (18h days, 21C day/ 18C nights).

Statistical Analyses

185   Cross incompatibilities were determined with a linear mixed model using the

package *lme4* and a type III ANOVA using *car* (Bates *et al.* 2015; Fox & Weisberg

2011) in the statistical interface R. All analyses were completed separately for IMxOdell

and IMxIMP crosses. For both total seed number, and proportion viable seed, we treated

cross type as a fixed factor and replicate maternal and paternal lines as random effects

(i.e. *Proportion viable seed ~ Cross Type + (1|maternal replicate) + (1|paternal*

replicate) or total seed count ~ Cross Type + (1|maternal replicate) + (1|paternal

replicate)). Pairwise T-tests with Holm correction for multiple testing were performed to

determine which cross types varied significantly from one another. Differences in

developmental trajectories were also assessed using a linear mixed model, as above. In

this case, cross type, DAP, and their interaction were the fixed effects and the seed

replicate was treated as a random effect.

Results

Post-Mating Reproductive Isolation

200   IMP and Odell Creek show consistent patterns of strong, and asymmetric hybrid

seed inviability when crossed with individuals from Iron Mountain, but in opposing

directions (Figure 1; Odell Creek: X^2= 4598, *df*=3, *p*<0.0001; IMP: X^2= 2969, *df*=3,

p<0.0001). IMP produces entirely inviable seeds when the maternal donor in these

crosses, while Odell Creek produces entirely inviable seeds when the paternal donor (and

some inviable seeds when the maternal donor; Figure 1). Inviable hybrid seeds appear

larger, disc-like, and shriveled, while viable hybrid seeds appear smaller overall (similar

to Coughlan *et al.* 2018).

208    There is some evidence of pollen-pistil incompatibilities between annual *M.*

209 *guttatus* and Odell Creek (Odell Creek: X^2= 49.33, *df*=3, *p*<0.0001; Figure S1), wherein

210 IM lines set less seed when pollinated by Odell Creek than any other cross type. We find

211 no evidence of pollen-pistil interactions between IM and IMP, although we note that IMP

212 has substantially larger ovaries and produces more seeds, regardless of the pollen donor

213 (X^2= 23.7, *df*=3, *p*<0.0001; Figure S1).

215   **Figure 1: Proportion viable seed between three lines of *M. guttatus* (IM) and**

216 **(A) four maternal families of northern *M. decorus* (IMP) and (B) three maternal**

217 **families of southern *M. decorus* (Odell Creek). Maternal parent is listed first.**

219 *Developmental Trajectories*

220    The three parental species surveyed here display similar seed development. In all

221 three parents, endosperm is tightly packed around the embryo and continues to grow until

222 around 8-10 DAP. At this point, the endosperm is depleted, presumably to feed the

223 growing embryo. Embryo development among the three parents is also quite similar, with

224 IM displaying a slightly accelerated growth, relative to either Odell or IMP (i.e. they

225 reach a heart-shaped embryo around 10 DAP, while this is not achieved in Odell or IMP

226 until around 14 DAP; Figure 2; Figure 3). By 14 DAP the embryo comprises most of the

227 embryo sac, with a thin layer of endosperm remaining.

Hybrid seeds from both the IM/IMP and IM/Odell crosses show distinct

developmental trajectories based on the maternal versus paternal donor of the cross, but

both hybrid crosses display similar seed trajectories, depending on the fate of the seed

(i.e. whether it will remain viable or not). In both cross types, seeds which remain viable

(e.g. IMxIMP and OdellxIM) display a precocious development, with several small

endosperm cells that are quickly degraded by the embryo (Figure 2; Figure 3) . By 14

DAP, the only remaining endosperm is a thin layer which surrounds the embryo (Figure

2; Figure 3). The overall seed size is also much smaller than either parent or the

reciprocal hybrid (Coughlan *et al.* 2018). In contrast, the hybrid seeds which will

eventually become inviable (i.e. IMPxIM and IMxOdell) exhibit a chaotic endosperm

growth, producing fewer, but larger and more diffuse endosperm cells (Figure 2; Figure

3). Embryos in this direction of the cross remain small, relative to both parents and the

reciprocal hybrid (Figure 2; Figure 3). By 14 DAP, most embryos were entirely degraded.

Figure 2: Seed development of IM62 x IMP crosses at 4, 6, 8, 10, and 14 Days After Pollination (DAP). Maternal parent is listed first.

These patterns are recapitulated when looking simply at endosperm cell width. In both

crosses, the type of cross significantly effected the endosperm cell width (Table 1; Figure

4, Figure S2). Reciprocal F1 hybrids display the most extreme endosperm cell sizes, with

inviable seeds exhibiting the largest endosperm cell size, and viable seeds displaying the

smallest endosperm cells. For both IMxIMP and IMxOdell Creek crosses, the

relationship between cell size and cross type was dependent on DAP (although we note

this relationship is not linear; significant interaction term: Table 1; Figure S2).

Figure 3: Seed development of IM62 x Odell Creek crosses at 4, 6, 8, 10, and 14 Days After Pollination (DAP). Maternal parent is listed first.

While the two different crosses display quite similar overall patterns of development, there are some distinctions between hybrid seeds formed between IMxIMP, and IMxOdell Creek. For example, in IMPxIM crosses, the embryo remains comparatively small, and does not appear to grow past the 6 or 8 DAP. In contrast, seeds formed from a IMxOdell cross often exhibit a developing embryo up until 10 DAP, suggesting that the seed is aborted slightly later in this cross.

Figure 4: Cell width at 8 Days After Pollination (DAP) for each cross type (maternal parent listed first), for both *M. guttatus* crossed to IMP (yellow) and Odell Creek (blue).

Overall, we see striking parallelism between two distinct seed incompatibilities with reciprocal F1 seeds displaying either excessive and chaotic growth patterns or precocious and stunted patterns, depending on whether the seeds are eventually aborted or not. We suggest that hybrid seed inviability in both of these cases is likely a result, at least in part, of defective endosperm development which manifests as chaotic and excessive growth.

Table 1: Mixed model for type III ANOVAs for seed cell size across developmental time for IMP and Odell Creek crossed reciprocally to IM62. Cross Type= both reciprocal F1 crosses and self-fertilizations from each species treated as a separate level; treated as a fixed effect), DAP= Days After Pollination (fixed effect). Seed replicate was treated as a random effect.

	IMP			Odell Creek		
	X^2	DF	p	X^2	DF	p
Intercept	295.04	1	<0.0001	123.74	1	<0.0001
Cross Type	119.848	3	<0.0001	14.92	3	0.0018
DAP	36.033	1	<0.0001	0.0029	1	0.96
Cross Type*DAP	42.583	3	<0.0001	32.95	3	<0.0001

Embryo Rescues

Embryo rescues were generally unsuccessful in these crosses. No seeds germinated when collected 7 DAP. Only a small fraction of IM62 and hybrids which were viable (i.e. OdellxIM and IMxIMP) germinated in the 9 DAP collections (i.e. 5-10%). Overall, this suggests that 7 DAP may be too premature for proper endosperm development of viable and inviable seeds to occur outside of the maternal environment, while only rapidly developing seeds are able survive transplants at 9 DAP. Unfortunately, determining whether hybrid seed inviability is due solely to endosperm malformation is yet to be determined in this group.

Discussion

As previously reported, the northern and southern clades of *M. decorus* exhibit strong, and asymmetric reproductive isolation with *M. guttatus*, but in opposite directions of the cross. This observation is true for all maternal family/line combinations tested here (as well as multiple populations tested in Coughlan *et al.* 2018), suggesting that HSI is not geographically restricted, as other intrinsic post-zygotic barriers have been found to be (Sweigart *et al.* 2007; Case &

Willis 2008; Martin & Willis 2010; Sicard *et al.* 2015; Barnard-Kubow & Galloway 2017). Total

reproductive isolation in these crosses was ~52% for both both species pairs, and thus may

represent a significant barrier to reproduction. Hybrid seed inviability has been shown to be a

formidable barrier to reproduction in other systems (i.e. Tiffin *et al.* 2001; Lowry *et al.* 2008),

but the developmental causes of this incompatibility have really only been quantified for a small

number of diploid species pairs.

Here we show that hybrid seed inviability is strongly associated with developmental

defects in endosperm deposition. In inviable directions of the cross, endosperm is chaotically

distributed, and the cells are few and large, suggesting a failure to proliferate. In the viable

directions of the cross, seeds develop precociously, and endosperm deposition is limited, as

witnessed by the general lack of endosperm remaining when seeds are fully developed, as well as

the generally smaller size of both the cells and final seeds (Coughlan *et al.* 2018). Overall, these

developmental patterns fit with a paternal excess/maternal repression model of hybrid inviability,

wherein paternal excess is more likely to lead to seed abortion. Unfortunately, given the

generally low success rate of the embryo rescues, it remains unclear whether hybrid seed

inviability is due strictly to malformation of the endosperm, or whether genetic incompatibilities

in the embryo also contribute.

These patterns have been shown in wild species pairs of both *Arabidopsis* and *Capsella*,

which have a nuclear-type endosperm development (Rebernig *et al.* 2015; Lafon-Placette *et al.*

2017). In these crosses, as well as interploidy crosses, endosperm over-proliferation is

accompanied by a failure of endosperm cells to cellularize (Scott *et al.* 1998; Bushell *et al.* 2003;

Rebernig *et al.* 2015; Lafon-Placette *et al.* 2017), and at least in *Capsella,* paternal excess is

associated with inviability, but maternal excess still produces viable seed (Rebernig *et al.* 2015).

Initial findings from species with cellular-type endosperm development (e.g. wild tomatoes and

other *Mimulus* species pairs) also show endosperm defects as a major cause of hybrid inviability

(Roth *et al.* 2017; Oneal *et al.* 2016). However, these defects to not represent defects in rates of

cellularization, but rather rates of proliferation, cell size and distribution, and optical density

(Oneal *et al.* 2016; Roth *et al.* 2017).

The developmental trajectories of viable vs non-viable hybrid seeds are strikingly similar

between OdellxIM, and IMPxIM crosses, despite the fact that inviability occurs in the opposite

cross direction. This suggests that similar evolutionary pressures may indeed be causing hybrid

seed inviability within this complex, namely that of parental conflict. Despite considerable

developmental parallelism, it remains to be seen how constrained the genetic targets of selection

are between these species pairs. In other post-zygotic reproductive barriers, canonical hybrid

incompatibilities often share similar genetic bases. For example, hybrid necrosis is often caused

by interactions involving R genes (reviewed in Bomblies & Weigel 2007), and CMS almost

always involves PPR genes (polcentric peptide repeat genes; reviewed in Rieseberg & Blackman

2010). Yet, whether this remains true for hybrid seed inviability remains to be seen. Indeed, one

may predict that the genomic target size for influencing resource provisioning to seeds is

presumably larger than the number of genes responsible for auoto-immune function or managing

organelle expression, as is the case with hybrid necrosis and CMS, respectively. However, this

prediction remains to be tested.

Acknowledgements

We thank members of the Willis lab- particularly Elean Oneal, as well as Miguel Florez and Bob

Franks for useful discussions of this project. We are deeply grateful to Melvin Turner for

teaching JMC histological techniques, and to Dick White for allowing us access to his histology

lab. This project was funded from NSF grants EF-0328636 and EF-0723814 to JHW, a DDIG

(DEB- 1501758), ASN Student research award and SSE student research award to JMC. The

Duke Graduate School also provided funding to JMC with the Myra and William Waldo Boone

fellowship, and financial support was provided from the Duke Biology Department.

Data Availability

Data for the manuscript will be made publically available.

357   **Supplemental Figures**

359   **Figure S1: Total number of seeds in crosses for (A) three lines from IM and four maternal**

360   **families of IMP selfed and crossed reciprocally and (B) three lines of IM and three**

361   **maternal families of Odell Creek (Od) selfed and crossed reciprocally.**

Figure S2: Cell lengths across all cross types for 5 developmental stages for both IMxIMP

and IMxOdell crosses.