HIV-1 subtype C with PYxE insertion has enhanced binding of gag-p6 to host cell protein ALIX and increased replication fitness

Robert van Domselaar[1], Duncan Njenda[2], Rohit Rao[3], Anders Sönnerborg[1,2,3,4,5], Kamalendra Singh[2,5], Ujjwal Neogi[2,3*]

[1]Division of Infectious Diseases and Dermatology, Department of Medicine Huddinge, Karolinska Institutet, Stockholm, Sweden

[2]Division of Clinical Microbiology, Department of Laboratory Medicine, Karolinska Institutet, Stockholm, Sweden

[3]Department of Clinical Microbiology, Karolinska University Laboratory, Karolinska University Hospital, Stockholm, Sweden

[4]Department of Infectious Diseases, Karolinska University Hospital, Stockholm, Sweden

[5]Molecular Microbiology and Immunology, University of Missouri, Columbia, MO 65211, USA

*Correspondence: Ujjwal Neogi, Division of Clinical Microbiology, Department of Laboratory Medicine, Karolinska Institutet, Stockholm, Sweden, e-mail- ujjwal.neogi@ki.se, Phone: +46 852483680

Abstract

Human immunodeficiency virus type 1 subtype C (HIV-1C) has a natural deletion of an YPxL

motif in its gag-p6 late domain. This domain mediates the binding of gag to host cell protein

ALIX and subsequently facilitates viral budding. In a subset of HIV-1C infected individuals, the

tetrapeptide insertion PYxE has been identified at the deleted YPxL motif site. Here, we report

the consequences of PYxE insertion on the interaction with ALIX and the relevance in terms of

replication fitness and drug sensitivity. In our three HIV-1C cohorts, PYKE and PYQE were

most prevalent among PYxE variants. Through *in silico* predictions and *in vitro* experiments, we

showed that HIV-1C gag has an increased binding to ALIX when PYxE motif is present. To go

more into the clinical relevance of the PYxE insertion, we obtained patient-derived gag-pol

sequences from HIV-1C$_{PYxEi}$ viruses and inserted them in a reference HIV-1. Viral growth was

increased and the sensitivity to protease inhibitor (PI) lopinavir (LPV) and nucleoside reverse

transcriptase inhibitor tenofovir alafenamide (TAF) was decreased for some of the HIV-1C PYxE

variants compared to wild-type variants. Our data suggest that PYxE insertion in gag restores the

ability of gag to bind ALIX and correlates with enhanced viral fitness in the absence or presence

of LPV and TAF. The high prevalence and increased replication fitness of the HIV-1C virus with

PYxE insertion could indicate the clinical importance of these viral variants.

Importance

Genomic differences within HIV-1 subtypes is associated with varying degree of viral spread,

disease progression, and clinical outcome. Viral budding is essential in the HIV-1 life cycle and

mainly mediated through the interaction of gag with host proteins. Two motifs within gag-p6

mediate binding of host cell proteins and facilitate budding. HIV-1 subtype C (HIV-1C) has a

natural deletion of one of these two motifs resulting in an inability to bind to host cell protein

ALIX. Previously, we have identified a tetrapeptide (PYxE) insertion at this deleted motif site in

a subset of HIV-1C patients. Here, we report the incidence of PYxE insertions in three different

HIV-1C cohorts, and the insertion restores the binding of gag to ALIX. It also increases viral

growth even in the presence of antiretroviral drugs lopinavir and tenofovir alafenamide. Hence,

PYxE insertion in HIV-1C might be biologically relevant for viruses and clinically significant

among patients.

Introduction

Human immunodeficiency virus type 1 (HIV-1) is a global threat with an estimated 36.7 million HIV-1 infected individuals and 1.0 million deaths in 2016. Virulence of HIV-1 is determined by its capacity to replicate within infected cells and its ability to infect new cells. Among different HIV-1 subtypes, more than 50% of all infections are caused by HIV-1 subtype C (HIV-1C) that is prevalent in South Africa, Zimbabwe, Mozambique, Botswana, Ethiopia, Eritrea, India, and in some parts of Brazil. However, HIV-1C has also become highly prevalent in several European countries, including Sweden (1, 2). The *gag* gene encodes for a polyprotein that is proteolytically processed by viral and cellular proteases into six final products: p17 matrix protein (MA), p24 capsid protein (CA), spacer peptide 1 (SP1), p7 nucleocapsid protein (NC), spacer peptide 2 (SP2), p6 protein (P6). These proteins are required for the assembly and release of new virions (3-6). The p6 late domain of gag contains two conserved short peptide motifs that bind host factors to facilitate viral budding (6-8). This PTAP motif is present in both HIV-1 and HIV-2, and sometimes it is duplicated in HIV-1 genomes (9, 10). It mediates binding to ESCRT-I subunit tumour susceptibility gene 101 (TSG101) (11). The YPx_nL (where x_n represents a random number of any possible amino acids) motif binds to ESCRT-III adaptor protein ALG-2-interacting protein X (ALIX) (12). However, this latter motif is deleted exclusively in HIV-1C(13). This deletion is associated with the loss of binding of p6 late domain to ALIX and a decrease in virus release from infected cells. Of note, HIV-2 lacks the YPx_nL motif but contains an alternative ALIX-binding motif, namely PYKEVTEDL, that originates from simian immunodeficiency virus (SIV) in rhesus macaques (14).

Recently, we identified a tetrapeptide insertion PYxE [where x represents lysine (K), glutamine (Q) or arginine (R)] within the gag protein in a subgroup of HIV-1C infected individuals (C_{PYxEi}-strains) (15). This C_{PYxEi}-strain was preferentially identified in East African HIV-1C infected patients, but was less common among HIV-1C infected patients from South Africa, India, and Germany (15). Furthermore, the insertion appears more frequent among patients on antiretroviral therapy, e.g. ritonavir-boosted protease inhibitors (PI), compared to therapy naïve patients in India, and more frequent in therapy-failure patients in South Africa (15, 16). Moreover, lower pre-therapy $CD4^+$ T cell counts, higher plasma viral loads, and reduced increase in $CD4^+$ T cell

counts were noted to be associated with PYxE insertion in HIV-1C patients compared to patients
with wild type HIV-1C from East Africa (17). Furthermore, we observed that increased
replication fitness in PYQE inserted HIV-1C viruses was polymerase independent (17). A more
recent study also claimed that gag-protease is the major determinant of subtype differences in
disease progression among HIV-1 subtypes (18).

As the tetrapeptide PYxE insertion was found at the site of the lost YPx_nL motif of HIV-1C gag-
p6, we hypothesized that this PYxE insertion may restore the interaction of the gag-p6 late
domain with ALIX and thereby increases replication fitness and reduces sensitivity to PIs. A
recent report showed that the PYRE insertion could rescue viral growth of a PTAP-deleted HIV-1
variant similar to the YPx_nL motif in the absence of a PTAP sequence (19). However, since the
PTAP motif is always present in clinical HIV-1 isolates, it is not known if and how PYxE
insertion affects HIV-1C pathogenicity in a clinical relevant context. To address this, we assessed
the distribution and frequency of this PYxE insertion in different cohorts of HIV-1C infected
individuals. We also evaluated the ability of PYxE motif to reconstitute the interaction of gag
with ALIX, its association with an increased viral growth of clinical isolates, and its association
with drug responses against all three drug classes; reverse transcriptase inhibitors (RTIs), PIs, and
integrase strand transfer inhibitors (INSTIs).

Material and Methods

Cell culture and plasmids

Cells were cultured in 5% CO_2 at 37°C. HEK293T cells were maintained in Dulbecco's modified
Eagle medium (DMEM, Sigma, USA) supplemented with 10% fetal calf serum (Sigma,USA), 2
mM L-glutamine (Sigma, USA), 0.1 mM MEM Non-Essential Amino Acids (Gibco/Thermo
Fisher Scientific, USA), and 10 units/mL penicillin combined with 10 μg/mL streptomycin
(Sigma). TZM-bl reporter cells were also maintained in above-mentioned medium. This reporter
cell line is a derivative of HeLa cells, but stably expresses high levels of CD4 and CCR5 on the
cell surface, and has integrated copies of luciferase gene under the control of the HIV-1 promoter
that allows simple and quantitative analysis of HIV-1 infection. MT-4 cells were maintained in
Roswell Park Memorial Institute 1640 (RPMI, Sigma, USA) medium supplemented with 10%
fetal calf serum, 20 units/mL penicillin and 20 μg/mL streptomycin. Cells were transfected using

FuGene HD according to the manufacturer's instructions (Promega, USA) in a 3:1 ratio with

DNA.

The pEGFPN1 and mCherry plasmids were obtained from Addgene (Addgene, USA). The

plasmid encoding for HA-tagged ubiquitin was a kind gift from Soham Gupta (Karolinska

Institutet, Sweden). pCR3.1/HIV-gag-mCherry (HIV-1B gag) and pCR3.1-GFP-ALIX plasmids

were kind gifts from Dr. Paul Bieniasz (The Rockefeller University, USA). A codon-optimized

HIV-1 gag with PYQE motif gene of East African HIV-1C viruses was cloned into the

pCR3.1/HIV-gag-mCherry plasmid that contained a mCherry protein C-terminal tag sequence.

The gag-PYQE-mCherry plasmid was modified by site directed mutagenesis to make a base pair

substitution mutation (C to A) that changes the PYQE motif to PYKE using the Q5 SDM kit

(New England Biolabs, USA). Gag-PYQE-mCherry plasmid was linearized by PCR using 5'-

<u>A</u>AGGAGCCTCTGACGAGCC-3' as forward primer (with the C to A base pair substitution

underlined) and 5'-ATAGGGACCCTGGTCTTTCAGC-3' as reverse primer. Linear products

were re-circularized using DpnI, a polynucleotide kinase and a ligase from the SDM kit following

the manufacturer's protocol. DNA sequences were verified by Sanger sequencing.

Gag-p6 sequences and clinical specimens

To identify the distribution of PYQE, PYRE and PYKE motifs, we re-analysed the HIV-1C gag-

p6 sequences that were collected from three cohorts, the Swedish InfCare Cohort (n=140), the

German Cohort (n=127), and the Ethiopian Cohort (n=73), as reported previously with respect to

naturally occurring polymorphisms in PYxE motif (15). Additional HIV-1B (n=2754) and HIV-

1C (n=1432) gag-p6 sequences were collected from HIV-1 Los Alamos Database. Stored plasma

samples from therapy naïve patients infected with HIV-1C (n=10) were randomly selected based

on gag-p6 sequences from the HIV-cohort at Karolinska University Hospital, Stockholm,

Sweden.

Recombinant virus production with patient-derived gag-pol

The recombinant viruses were produced as described by us recently (20). Briefly, the *gag-pol*

fragment (HXB2:0702-5798) was cloned into pNL4-3 plasmid following digestion with *BssHII*

and *SalI* (New England Biolab, USA) and ligation using T4 DNA ligase (New England Biolabs,

USA). The chimeric viruses were produced by transient transfection of the plasmids into the

293T cell line using FuGene HD and harvested 72 hours later by a collection of the cell-free

supernatant cleared by centrifugation and stored in aliquots at -80°C.

In silico analysis: Molecular modelling, docking and Ubiquitin binding motif prediction

The structure of HIV-1C gag late domain with the PYKE insertion was generated by homology

modeling techniques. The crystal structures of the gag late domain in complex with ALIX (PDB

entries 2XS1 and 2R02) were used as template molecules. For this purpose, the 'Prime' utility of

Schrödinger Suite (Schrödinger Inc., USA) was used. The structure was subjected to restricted

minimization using OPLS_2005 force field. This structure was docked into the crystal structure

of ALIX (PDB file 2XS1) using protein-protein docking program of 'BioLuminate' (Schrödinger

Inc., USA), after deleting SIV_{mac239} PYKEVTEDL late domain structure. The resulting structure

was further minimized for 1000 iterations to remove steric clashes. A similar protocol was used

to form the complex among ALIX, gag late domain and ubiquitin. The crystal structure of

ubiquitin in complex with the TSG101-binding PTAP domain (PDB file 1S1Q) was used after

exclusion of TSG101 coordinates. The entire complex was then subjected to molecular dynamics

simulations for 1,000,000 steps with step size of 50 pico seconds using OPLS3 force field. The

most energetically stable model of the complex was used for further analysis. In order to obtain

energetically favoured polar interactions, the conformational search of polar sidechains at the

interface of proteins was also conducted.

The prediction of ubiquitin binding with the HIV-1C gag late domain with the PYKE motif P6

was performed by adding the amino acid sequence in the prediction tools presented at the website

http://www.ubpred.org/.

Immunofluorescence

HEK293T cells were cultured on poly-L-lysine-coated glass coverslips and co-transfected with

GFP-tagged ALIX and mCherry-tagged codon-optimized gag variants. At 24 hours post

transfection, cells were washed twice with PBS, fixed in 10% formalin (Sigma, USA) for 20 min

at room temperature, washed three times with PBS, and stored in PBS at 4°C. Later, nuclear

counterstaining with DAPI was performed followed by mounting of the glass coverslips on glass

slides. When HA-ubiquitin was also co-transfected, cells were fixed in ice-cold methanol for 15

min at -20°C, washed twice with PBS, and stored in PBS at 4°C. For ubiquitin detection, fixed

cells were incubated with anti-HA antibody (mouse monoclonal, clone 12CA5, Sigma, USA) for

1 hour at room temperature followed by goat-anti-mouse-Alexa Fluor 647-Plus-conjugated

secondary antibody (Invitrogen, USA) for 1 hour at room temperature before nuclear

counterstaining with DAPI. Fluorescence was analyzed by confocal laser scanning microscopy

using a Nikon Single point scanning confocal microscope with ×60/1.4 oil objective (Nikon,

Japan).

Microscale Thermophoresis (MST)

The microscale thermophoresis experiments were conducted by Monolith NT0.115 instrument

(NanoTemper Technologies) at 40% MST and LED power at 25°C. Hexahistidine-tagged ALIX

protein was labelled by NTA-dye (NanoTemper Technologies) using manufacturer's protocol.

The peptides were synthesized at the Molecular Interaction Core (University of Missouri).

Thermophoresis was induced by mixing peptides (0.001-1µM) and fixed NTA-labelled ALIX (50

nM) in a buffer 50 mM Tris-Cl pH 7.8, 100 mM NaCl and 0.1% pluronic-F127. The binding

isotherms were obtained by plotting the difference in normalized fluorescence against increasing

peptide concentration. The binding affinities were determined fitting the data points to a

quadratic equation (equation 1) using non-linear regression using MO Affinity software

(NanoTemper Technologies), Prism (GraphPad Inc. version 6.0) or OriginLab (version 18,

OriginLab Corp. Northampton, MA, USA).

$$F_b = \frac{(K_{d1}+[PEP_0]+[ALIX_0]) - \sqrt{(K_{d1}+[PEP_0]+[ALIX_0])^2 - 4[ALIX_0][PEP_0]}}{2[ALIX_0]} \quad \text{(Eq. 1)}$$

Where F_b is fraction of ALIX/peptide complex, where K_d = [ALIX][PEP]/[ALIX-PEP], [ALIX]

is the concentration of free ALIX, $[ALIX_0]$ is the concentration of total ALIX, [PEP] is the

concentration of free peptide and $[PEP_0]$ is the total concentration of peptide.

Co-immunoprecipitation assay

HEK293T cells were co-transfected with GFP-tagged ALIX and codon-optimized gag variants

from HIV-1B or HIV-1C. At 24 hours post transfection, cells were collected, washed twice with

ice-cold PBS and immunoprecipitation of GFP-tagged proteins was performed using GFP-

Trap®_A kit (Chromotek, Germany) according to manufacturer's protocol. Briefly, cells were

lysed in ice-cold lysis buffer (10 mM Tris pH 7.5, 150 mM NaCl, 0.5 mM EDTA, 0.5% NP-40,

supplemented with a protease cocktail inhibitor from Roche, Switzerland) for 30 minutes on ice.

Then lysates were diluted in ice-cold wash buffer (10 mM Tris pH 7.5, 150 mM NaCl, 0.5 mM

EDTA) supplemented with a protease cocktail inhibitor from Roche, input samples (10% of total)

were saved for immunoblot analysis, and the remaining (90%) part of the diluted lysates were

incubated with equilibrated GFP-Trap®_A beads end-over-end for 1 hour at 4°C. Beads were

washed three times with ice-cold wash buffer before resuspending in 2x Leammli buffer

(Invitrogen, 4x buffer diluted 1:1 with PBS) and heated for 10 minutes at 95°C. Protein

concentration of input samples was determined by DC protein assay (Biorad, USA) according to

the manufacturer's microplate assay protocol. Equal amounts of protein for input samples and

equal volumes of pull-down samples were subjected to immunoblotting using primary antibodies

against GFP (rabbit monoclonal, clone EPR14104, Abcam, UK), gag (rabbit polyclonal to HIV1

p55 + p24 + p17, Abcam, ab63917), and β-actin (rabbit polyclonal, Abcam, ab8227) and

secondary horseradish peroxidase-conjugated antibodies (polyclonal goat anti-rabbit,

DAKO/Agilent, USA). Immunoblotted proteins were detected using the enhanced

chemiluminescence detection system (Pierce/Thermo Fisher Scientific, USA) and Hyperfilms

(Amersham/GE Healthcare Life Sciences, UK).

Viral growth kinetics assay

First, $TCID_{50}$ for each propagated recombinant viral clone was determined in TZM-bl reporter

cells. For each recombinant viral clone, ten-fold serial dilutions were prepared ranging from 10^2

to 10^7 in dilution factor from a single aliquot in medium containing DEAE (20 μg/μL). TZM-bl

cells were plated in 96-well plates and six replicates were incubated with the various virus

dilutions for 48 hours at 37°C in a 5% CO_2 humidified incubator. Virus infectivity was quantified

by measuring Renilla luciferase activity (relative light units [RLU]) using Bright-Glo™

Luciferase Assay System (Promega, USA) on the Tecan microplate reader (Tecan Infinite® 200

Pro, Tecan Group Ltd, Switzerland). The Spearman-Karber method was used to calculate the

$TCID_{50}$ for each recombinant viral clone. Then, MT-4 cells (2×10^5) were seeded in a 12-wells

plate and infected with each recombinant virus at a multiplicity of infection (MOI) of 0.05

plaque-forming units (pfu) per cell in triplicates for each condition. Supernatants were collected

on day 0, 3, 5 and 7 days post infection and HIV-1 p24 levels in supernatants were measured on

the HIV COBAS 8000 platform (Roche diagnostics, Switzerland). Viral growth kinetics (VGK)

were analysed as a relationship between signal to cut-off ratio (i.e., electrochemiluminescence

signal of sample relative to calibrated negative samples for the HIV combi PT kit on the COBAS

8000 system) for each time point and replicate. Graphical representation of the result was made

using Graphpad prism v6 (Graphpad Inc., USA).

Drug sensitivity assay (DSA)

The following drugs were purchased from Selleckchem, USA: Atazanavir sulfate (ATV),

darunavir ethanoate (DRV), lopinavir (LPV), efavirenz (EFV), nevirapine (NVP), rilpivirine

(RPV), etravirine (ETR), tenofovir alafenamide (TAF), elvitegravir (EVG), dolutegravir (DTG),

and cabotegravir (CAB). The DSA was performed by determining the extent to which the

antiretroviral drug inhibited the replication of the reference virus (pNL4-3), wild-type, PTAPP

duplicated and PYxE derived recombinant viruses, respectively. One round was required for all

drugs as previously reported (20), except for the protease inhibitor (PI) drugs that required two

rounds of infection. Briefly for the for two-round infection using PI DA, TZM-bl cells ($1x10^4$)

were seeded in 96-wells plates and cultured for 24 hours. Drugs were serially diluted in culture

media (ranging from 0.1 μM to 0.01 pM) and added in triplicate to the cells. The following day,

viruses were added to each well at an MOI of 0.05 pfu/cell in the presence of DEAE (10μg/ml).

After 48 hours, the viral supernatant from respective 96-wells plates with each drug was

transferred to newly TZM-bl pre-seeded 96-wells plates without adding new drugs. Thereafter,

virus replication was quantified by measuring Renilla luciferase activity (relative light units

[RLU]) using Bright-Glo™ Luciferase Assay System (Promega, USA) 48 hours post-reinfection.

Drug concentrations required for inhibiting virus replication by 50% (EC_{50}) were calculated by a

dose-response curve using non-linear regression analysis (GraphPad Prism, version 6.07;

GraphPad Software, USA). The DSA experiments were performed with three technical replicates

for each virus with the specified dynamic concentration range of the drug, and at least two

independent analyses were performed. The reproducibility of the DSA was assessed by the 95%

confidence interval obtained for the drug EC_{50} and the degree of correlation between technical

replicates. The output for the drug EC_{50} results was used to compute the fold change value (FCV)

for each virus relatively to pNL4-3.

Results

Evolution of gag-p6 late domain sequences in HIV-1B and HIV-1C

The sequence analysis of HIV-1B and South African and Indian HIV-1C gag-p6 sequences

(obtained from the Los Alamos Database) identified conserved a $LYPx_nL$ motif in HIV-1B which

was missing in HIV-1C (Fig 1a). We noted a unique evolution of the lysine (K) residue in the

HIV-1B and HIV-1C gag-p6 late domain, which is the target of ubiquitination. While K475

residue was conserved in both HIV-1B and C, there was an evolution of a lysine residue at amino

acid position 479 along with K481R mutation in the majority of HIV-1C sequences. Interestingly,

the multiple sequence alignment of HIV-1C$_{PYxEi}$ strains identified conservation of HIV-1C

specific 479K residue, but residue R481 was reverted to K481 in HIV-1B (Fig 1a). When we

analyzed gag-p6 sequences from HIV-1C patients from the Swedish InfCareHIV cohort (n=140),

we found that 68% (95/140) of HIV-1C patients had a wildtype strain whereas 32% (45/140) had

a PYxE insertion; 22% (31/140) had a PYKE insertion, and 9% (13/140) had a PYQE insertion

(Fig.1b). All of the PYxE sequences in HIV-1C of patients from the Swedish cohort belonged to

HIV-1C strains of Ethiopian and Eritrean origin. Similar trend was observed in the German HIV-

1C cohort with 12% PYKE insertion (15/127) (Fig.1c) and the Ethiopian HIV-1C cohort with

37% (27/73) PYKE insertion (Fig. 1d). Thus, our analysis indicated that PYKE and PYQE are

predominating in HIV-1C$_{PYxEi}$ viruses of all cohorts analysed. Subsequently, we restricted our

analysis to these two variants.

In silico analysis predicts binding of ALIX to the PYKE motif in gag

As reported before, the lysine residues in the gag-p6 are subjected to ubiquination (21), we first

predicted the likelihood of K_{ins} ubiquitination compared to lysine residues adjacent to the PYKE

motif. Indeed, K_{ins} was predicted to be more likely ubiquitinated than lysine residues in the

vicinity of the PYKE motif (Fig.2a). Since the PYKE motif is present at the missing ALIX-

binding site and contains three charged amino acids that can facilitate interaction with other

proteins, we used the PYKE insertion (amino acids 483 to 486) in the molecular modelling to

predict the interaction of the gag-p6 late domain with ALIX. Amino acids E482, Y484 and E486

of gag could directly facilitate binding to ALIX (Fig.2b and Table 1). Ubiquitin was predicted to

bind K501 in ALIX and thereby affect the interaction between E486 in gag-PYKE with K501 in

ALIX. In addition, ubiquitin could also interact with E387 of ALIX and K_{ins} of gag-PYKE

through its residues K63 and E64, respectively. Thus, our *in silico* analysis predicted that the

PYxE motif can restore the interaction of gag with host cell ESCRT-III adaptor protein ALIX and

that, in case of a PYKE insertion, ubiquitination of the K_{ins} residue could further facilitate this

interaction.

PYxE insertion enhances type C gag: ALIX interaction

Prior to testing the interaction between gag and ALIX, we evaluated the cellular localization of

both proteins. 293T cells were co-transfected with GFP-tagged ALIX with or without three

codon-optimized variants of mCherry-tagged HIV-1C gag, namely wild type gag or gag that

includes the tetrapeptide insertion PYKE or PYQE. Confocal analysis showed cytoplasmic

localization of ALIX and predominant plasma membrane localization of all three gag variants

(HIV-1C$_{wildtype}$, HIV-1C$_{PYQEi}$ and HIV-1C$_{PYKEi}$) (Fig.3a). We also tested the localization of

ubiquitin in this context. Since both gag and ALIX are known to be ubiquitinated, it is not

surprising that ubiquitin co-localized with ALIX and gag (Fig.3b). Thus, the localization of gag

was not due to the tetrapeptide insertion, but most likely due to PTAP:TSG101-facilitated

transport of gag to and viral budding from the plasma membrane.

Next, we evaluated the binding affinity of the four HIV-1C gag variants, HIV-1B$_{WT}$ (Fig 4a),

HIV-1C$_{WT}$ (Fig 4b), HIV-1C$_{PYKEi}$ (Fig 4c) and HIV-1C$_{PYQEi}$ (Fig 4d), to ALIX. The late domain

motifs' peptides were selected based on crystal structure PDB:2XS1 and PDB:2R02. The binding

affinity of the gag peptide with ALIX were statistically significantly higher in HIV-1B$_{WT}$ (32 ± 3

nM) compared to all HIV-1C variants, HIV-1C$_{WT}$ (290 ± 6 nM), HIV-1C$_{PYKEi}$ (110 ± 4 nM),

HIV-1C$_{PYQEi}$ (96 ± 4 nM) (Fig 4e). However statistically significant higher binding affinity to

ALIX was observed with HIV-1C$_{PYKEi}$ and HIV-1C$_{PYQEi}$ compared to HIV-1C$_{WT}$ (Fig 4e).

As the peptides may not retain their conformal structures as in the complete protein, we

confirmed the binding by co-immunoprecipitation pull-down assays. GFP-tagged ALIX and

HIV-1C gag variants were transfected either alone or together in 293T cells. A plasmid encoding

for HIV-1B gag was used together with GFP-ALIX as a positive control for gag:ALIX

interaction. Lysates were prepared 24h post-transfection, GFP-tagged proteins were pulled down

and co-immunoprecipitated proteins were analysed by immunoblotting (Fig.4b). GFP-tagged

ALIX was efficiently and the amount of HIV-1B gag was increased when it was co-transfected

with GFP-ALIX confirming that HIV-1B gag binds to ALIX. HIV-1C gag wild type was also

detected when co-transfected with GFP-ALIX, but HIV-1C gag PYKE was more prominently

detected. The intensity of HIV-1C gag PYQE was consistently lower than that of the PYKE

variant, but slightly increased compared to the wild type HIV-1C gag, indicating that potential

ubiquitination of the lysine residue is not a necessity, although it can facilitate a stronger

interaction of gag with ALIX. Thus, the tetrapeptide insertion PYxE in HIV-1C gag-p6 late

domain promotes binding of gag towards ALIX.

Differences in viral growth and drug responses of clinical strains of HIV-1C with PYxE motif

Finally, we assessed the consequence of the PYxE insertion on HIV-1C replication fitness using

clinical strains obtained from HIV-1C infected patients. Gag-pol sequences were cloned from one

wildtype HIV-1C virus with a single PTAP motif and no PYxE insertion (PT01), three HIV-1C

strains with a PYKE insertion (PT04-06), one with a PYQE insertion (PT07). We also included

two HIV-1C strains with a PTAP duplication (PT02 and PT03) as it has been shown to increase

the viral fitness and affect the drug sensitivity (Fig.5a). Then, these sequences were cloned into

the genome of a reference virus (NL4-3, HIV-1B wild type) replacing its gag-pol sequence. The

various recombinant viruses were tested for viral replication in an *ex vivo* viral growth kinetic

assay and compared to the reference virus (Fig.5b). In concordance with a previous report (22),

viral replication was reduced for the HIV-1C wild type clone compared to the HIV-1B reference

virus. However, when a PTAP-duplication or a PYxE insertion was present, viral growth was

increased compared to the HIV-1C wild type variant and was comparable to HIV-1B wild type.

This correlation suggests that HIV-1C strains with a PYxE insertion have a growth advantage

compared to wild type HIV-1C.

Several reports have shown that a PYxE insertion is more frequent among HIV-1C therapy

failure patients (16, 23). To examine whether our various recombinant viral clones without any

known drug resistance mutations to RTIs, PIs and INSTIs responded differently to various ART

drugs, the target cells were infected with individual viral clones in the presence of a specific drug.

The viral replication was assessed and the EC_{50} was calculated (Fig. 5c). No differences in EC_{50}

values were observed among the various recombinant viral clones in the presence of non-

nucleoside RTIs (NNRTIs) efavirenz (EFV), rilpivirine (RPV) or etravirine (ETR), or INSTIs

raltegravir (RAL), elvitegravir (EVG), dolutegravir (DTG) or cabotegravir (CAB). In contrast,

with the PI lopinavir (LPV) one out of two viral clones with a PTAP-duplication and two out of

four with PYxE insertion showed a >4-fold decreased sensitivity towards the drug. However,

there was no change in susceptibility against DRV and ATV. One of the other viral clones with a

PYxE insertion showed a 3.5-fold decreased sensitivity towards nucleoside reverse transcriptase

inhibitor (NRTI) tenofovir alafenamide (TAF). Thus, our data indicates that PYxE insertion may

potentially alter the sensitivity towards the PI LPV and NRTI TAF and that may be due to the

increased replication fitness. However, this is not universal across all the HIV-1$_{PYxEi}$-strains and

could be due to co-evolution of other amino acids in the gag region.

Discussion

Genetic diversity among HIV-1 strains and their adaptation can provide advantages for individual
viral strains in different biological contexts. Here, we describe the distribution of HIV-1C
containing tetra-peptide PYxE insertion within the gag and assessed the binding efficiency of gag
with host cell protein ALIX. Our results showed that the PYxE insertion enhances the binding
capacity of HIV-1C gag to ALIX in ubiquitin dependent or independent manner. In addition, the
insertion increased the replication fitness of the virus and seemed to alter the sensitivity against
the PI LPV and NRTI TAF in some of the strains.

Although the PYxE insertion is quite prevalent among HIV-1C from Ethiopia and Eritrea
individuals, it is not as prevalent in South African or Indian HIV-1C strains (16, 23). The twelve
base pair sequence encoding the PYxE motif is not present in non-subtype C M group HIV-1
sequences, but is present within the gag-p6 late domains of SIV_{mac239}, SIV_{smE543}, and their close
relative HIV-2 (14, 24). As HIV-1C has a natural deletion of the YPx_nL motif and an insertion of
PYxE has replication advantages, we hypothesize that these strains have evolved through a
recombination event of HIV-1C with either SIV or HIV-2. It is well described that the HIV-1
epidemic in Ethiopia geographically clusters very strongly (25) and the Ethiopian HIV-1C has
been proposed to originate from either a single lineage or multiple descendants (26). However to
explain why the PYxE insertion is significantly more prevalent in HIV-1C from Ethiopia and
Eritrea as compared to South African and Indian strains need further evolutionary study to prove
this hypothesis.

Viral budding is mediated by the gag-p6 late domain of gag through the PTAP motif, the YPx_nL
motif, and the PYxE motif. The PTAP motif mediates binding of gag to ESCRT protein TSG101
and is present in all HIV and SIV variants. Therefore, this domain appears to be necessary for
viral budding. Although one report shows that PTAP duplication only increases HIV-1
replication in the presence of a PI (27), others and we observed increased viral growth when
HIV-1 contained a PTAP duplication (28). Although deletion of the PTAP motif has not been
identified in clinical strains so far, artificial deletion or mutation of this PTAP motif severely
abrogates release of HIV-1 from infected cells (11, 29, 30). However, the presence of either one
of the two other motifs that both facilitate binding to host cell protein ALIX still allows for virion
release (12, 14, 19, 31, 32). This indicates that both TSG101- and ALIX-mediated pathways for

viral budding are not dependent on each other and each pathway is sufficient to mediate viral

budding. The first ALIX-binding motif is YPx_nL and is present in most HIV-1 subtypes but not in

HIV-1C or HIV-2 (13, 14). HIV-2 gag can mediate binding to ALIX through a distinct

PYKEVTEDL motif, and mutations in this motif can abrogate ALIX binding and decrease virion

release (14, 24). Thus, the PYxE motif in a subgroup of HIV-1C strains is similar but shorter than

the ALIX-binding motif found in HIV-2. We showed that this insertion of PYxE in HIV-1C gag

at the natural deletion site of the YPx_nL domain could reconstitute binding of HIV-1C gag to

ALIX. Hence, together with our observed PYxE insertion in HIV-1C, all HIV-1 subtypes have an

ALIX-binding motif next to their PTAP-motif within the gag late domain.

Our data suggests that ubiquitin may play a role in facilitating the binding of ALIX to the PYKE

motif. This is not surprising, as ubiquitin has been shown to be involved in viral budding of HIV.

Although dispensable for virus budding, ubiquitination of gag and the ESCRT-proteins are

essential for efficient release of HIV-1 from infected cells (21, 33, 34). Ubiquitination of the

lysine of the PYKE motif could increase the binding between ALIX and gag and thereby mediate

more efficient viral budding. Indeed, immunoprecipitation of HIV-1C gag-PYKE was more

efficient compared to the gag-PYQE variant (Fig.4b). As the PYKE motif insertion seems to be

the original insertion within HIV-1C gag, ubiquitination of its lysine might not be essential since

other variants of the PYKE motif, i.e. PYQE and PYRE, are identified among HIV-1C strains

along with evolution of other lysine (K) residues in the late domain. Because we only have few

clinical strains with a PYxE insertion, we could not identify differences in viral growth between

HIV-1C gag-PYKE and other PYxE variants (Fig.5). Nonetheless, whereas the other three

residues seem to be required for the binding of ALIX to gag, the lysine residue might provide

increased stability for this interaction.

Although the ALIX-mediated viral budding mechanism appeared to be lost in HIV-1C, it is

surprising to observe that HIV-1C PYxE strains were prevalent among HIV-1C infected

individuals. This suggests that the interaction of gag with ALIX is of biological importance for

the virus. In our limited cohort of HIV-1C infected patients, PYxE insertion was always co-

present with the PTAP motif. This suggests that the PYxE motif is not redundant from the PTAP

motif-mediated viral budding process through TSG101 nor a compensatory mechanism for loss

of the PTAP-motif and HIV-1 requires both motifs for efficient viral replication. Importantly,

viruses with HIV-1C gag PYKE or PYQE insertion showed enhanced viral growth compared to

viruses with HIV-1C wild type gag (Fig.5). This finding is in line with a recent report that

showed that a PYRE insertion within gag enhances HIV-1C viral growth (19).

Decreased sensitivity towards PI LPV or NRTI TAF was found in several HIV-1C strains with a

PTAP duplication or PYxE insertion. For the PTAP duplication it has previously been reported

that it could enhance HIV-1 growth in the presence of PIs lopinavir and ritonavir (27). For the

PYxE insertion, our earlier studies have shown that the PYxE insertion is more frequent in HIV-

1C therapy-failure patients (15, 16). Nonetheless, since our clinical HIV-1C PYxE strains did not

have any known drug resistance mutations, it seems likely that the correlation of these viruses

with advanced viral growth and reduced sensitivity towards the PI LPV and TAF could increase

the risk for therapy failure in HIV-1C PYxE-infected patients. Though mechanistic studies shows

that at the time of reverse transcription of HIV-1C from the RNA template, the nucleotide in

HIV-1C K65 position (AAG) in RT favoured pausing, which increases the probability for the

development of TDF/TAF mutation K65R (35, 36). However no studies have shown any co-

relation with gag-mutation on susceptibility of TAF. We are presently performing research to

understand the role of gag mutations on sensitivity for TAF as both tenofovir (TDF) and TAF are

used globally.

In conclusion, our study showed that PYxE insertion in HIV-1C of patients originating from

Ethiopia and Eritrea restored the interaction of gag with ALIX, which is mediated in an ubiquitin-

dependent or -independent way. Importantly, the insertion was positively correlated with the

replication fitness that could affect the sensitivity against LPV in the absence of any PI drug

resistance mutation. Most importantly, reduced sensitivity against TAF in the absence of any

TAF-mutation needs further studies to analyse any role of gag-mutations in reducing

susceptibility towards TAF. Altogether, it will be important to follow up on whether HIV-1C$_{PYxEi}$

strains are emerging following the failure of antiretroviral therapy with LPV and TAF based

regimens due to increased replication fitness. This information could provide important insights

in the clinical significance of the PYxE insertion within HIV-1C gag.

Acknowledgements

The study is funded by grants from the Swedish Research Council (2017-01330, UN; 2016-

01675, AS), Karolinska Institutet Doctoral Student Funding (KID2015-154, UN) and the

Stockholm County Council (ALF 20160074, AS). We thank S.G. Aralaguppe, W. Zhang and S.

Svensson Akusjärvi for technical assistance. The microscopic part of the study was performed at the Live Cell Imaging facility, Karolinska Institutet, Sweden, supported by grants from the Knut and Alice Wallenberg Foundation, the Swedish Research Council, the Centre for Innovative Medicine and the Jonasson center at the Royal Institute of Technology, Sweden. The following reagents were obtained through the AIDS Reagent Program, Division of AIDS, NIAID, NIH: TZM-bl cells (Cat#8129) from Dr. John C. Kappes, and Dr. Xiaoyun Wu, MT-4 cells from Dr. Douglas Richman, and HIV-1 NL4-3 Infectious Molecular Clone (pNL4-3) from Dr. Malcolm Martin (Cat#114).

References

1. **Siddik AB, Haas A, Rahman MS, Aralaguppe SG, Amogne W, Bader J, Klimkait T, Neogi U.** 2018. Phenotypic co-receptor tropism and Maraviroc sensitivity in HIV-1 subtype C from East Africa. Sci Rep **8:**2363.

2. **Neogi U, Haggblom A, Santacatterina M, Bratt G, Gisslen M, Albert J, Sonnerborg A.** 2014. Temporal trends in the Swedish HIV-1 epidemic: increase in non-B subtypes and recombinant forms over three decades. PLoS One **9:**e99390.

3. **Sundquist WI, Krausslich HG.** 2012. HIV-1 assembly, budding, and maturation. Cold Spring Harb Perspect Med **2:**a006924.

4. **Bieniasz PD.** 2006. Late budding domains and host proteins in enveloped virus release. Virology **344:**55-63.

5. **Demirov DG, Freed EO.** 2004. Retrovirus budding. Virus Res **106:**87-102.

6. **Votteler J, Sundquist WI.** 2013. Virus budding and the ESCRT pathway. Cell Host Microbe **14:**232-241.

7. **Martin-Serrano J, Neil SJ.** 2011. Host factors involved in retroviral budding and release. Nat Rev Microbiol **9:**519-531.

8. **Scourfield EJ, Martin-Serrano J.** 2017. Growing functions of the ESCRT machinery in cell biology and viral replication. Biochem Soc Trans **45:**613-634.

9. **Flys T, Marlowe N, Hackett J, Parkin N, Schumaker M, Holzmayer V, Hay P, Eshleman SH.** 2005. Analysis of PTAP duplications in the gag-p6 region of subtype C HIV type 1. AIDS Res Hum Retroviruses **21:**739-741.

10. **Sharma S, Aralaguppe SG, Abrahams MR, Williamson C, Gray C, Balakrishnan P, Saravanan S, Murugavel KG, Solomon S, Ranga U.** 2017. The PTAP sequence duplication in HIV-1 subtype C Gag-p6 in drug-naive subjects of India and South Africa. BMC Infect Dis **17:**95.

11. **Garrus JE, von Schwedler UK, Pornillos OW, Morham SG, Zavitz KH, Wang HE, Wettstein DA, Stray KM, Cote M, Rich RL, Myszka DG, Sundquist WI.** 2001. Tsg101 and the vacuolar protein sorting pathway are essential for HIV-1 budding. Cell **107:**55-65.

12. **Fisher RD, Chung HY, Zhai Q, Robinson H, Sundquist WI, Hill CP.** 2007. Structural and biochemical studies of ALIX/AIP1 and its role in retrovirus budding. Cell **128:**841-852.

13. **Patil A, Bhattacharya J.** 2012. Natural deletion of L35Y36 in p6 gag eliminate LYPXnL/ALIX auxiliary virus release pathway in HIV-1 subtype C. Virus Res **170:**154-158.

14.   **Zhai Q, Landesman MB, Robinson H, Sundquist WI, Hill CP.** 2011. Identification and structural
characterization of the ALIX-binding late domains of simian immunodeficiency virus SIVmac239
and SIVagmTan-1. J Virol **85:**632-637.
15.   **Neogi U, Rao SD, Bontell I, Verheyen J, Rao VR, Gore SC, Soni N, Shet A, Schulter E, Ekstrand**
ML, Wondwossen A, Kaiser R, Madhusudhan MS, Prasad VR, Sonnerborg A. 2014. Novel tetra-
peptide insertion in Gag-p6 ALIX-binding motif in HIV-1 subtype C associated with protease
inhibitor failure in Indian patients. Aids **28:**2319-2322.
16.   **Neogi U, Engelbrecht S, Claassen M, Jacobs GB, van Zyl G, Preiser W, Sonnerborg A.** 2016.
Mutational Heterogeneity in p6 Gag Late Assembly (L) Domains in HIV-1 Subtype C Viruses from
South Africa. AIDS Res Hum Retroviruses **32:**80-84.
17.   **Aralaguppe SG, Winner D, Singh K, Sarafianos SG, Quinones-Mateu ME, Sonnerborg A, Neogi**
U. 2017. Increased replication capacity following evolution of PYxE insertion in Gag-p6 is
associated with enhanced virulence in HIV-1 subtype C from East Africa. J Med Virol **89:**106-111.
18.   **Kiguoya MW, Mann JK, Chopera D, Gounder K, Lee GQ, Hunt PW, Martin JN, Ball TB, Kimani J,**
Brumme ZL, Brockman MA, Ndung'u T. 2017. Subtype-Specific Differences in Gag-Protease-
Driven Replication Capacity Are Consistent with Intersubtype Differences in HIV-1 Disease
Progression. J Virol **91**.
19.   **Chaturbhuj D, Patil A, Gangakhedkar R.** 2018. PYRE insertion within HIV-1 subtype C p6-Gag
functions as an ALIX-dependent late domain. Sci Rep **8:**8917.
20.   **Neogi U, Singh K, Aralaguppe SG, Rogers LC, Njenda DT, Sarafianos SG, Hejdeman B,**
Sonnerborg A. 2018. Ex-vivo antiretroviral potency of newer integrase strand transfer inhibitors
cabotegravir and bictegravir in HIV type 1 non-B subtypes. Aids **32:**469-476.
21.   **Sette P, Nagashima K, Piper RC, Bouamr F.** 2013. Ubiquitin conjugation to Gag is essential for
ESCRT-mediated HIV-1 budding. Retrovirology **10:**79.
22.   **Wright JK, Brumme ZL, Carlson JM, Heckerman D, Kadie CM, Brumme CJ, Wang B, Losina E,**
Miura T, Chonco F, van der Stok M, Mncube Z, Bishop K, Goulder PJ, Walker BD, Brockman MA,
Ndung'u T. 2010. Gag-protease-mediated replication capacity in HIV-1 subtype C chronic
infection: associations with HLA type and clinical parameters. J Virol **84:**10820-10831.
23.   **Singh K, Flores JA, Kirby KA, Neogi U, Sonnerborg A, Hachiya A, Das K, Arnold E, McArthur C,**
Parniak M, Sarafianos SG. 2014. Drug resistance in non-B subtype HIV-1: impact of HIV-1 reverse
transcriptase inhibitors. Viruses **6:**3535-3562.
24.   **Bello NF, Wu F, Sette P, Dussupt V, Hirsch VM, Bouamr F.** 2011. Distal leucines are key
functional determinants of Alix-binding simian immunodeficiency virus SIV(smE543) and
SIV(mac239) type 3 L domains. J Virol **85:**11532-11537.
25.   **Amogne W, Bontell I, Grossmann S, Aderaye G, Lindquist L, Sonnerborg A, Neogi U.** 2016.
Phylogenetic Analysis of Ethiopian HIV-1 Subtype C Near Full-Length Genomes Reveals High
Intrasubtype Diversity and a Strong Geographical Cluster. AIDS Res Hum Retroviruses **32:**471-
474.
26.   **Tully DC, Wood C.** 2010. Chronology and evolution of the HIV-1 subtype C epidemic in Ethiopia.
Aids **24:**1577-1582.
27.   **Martins AN, Waheed AA, Ablan SD, Huang W, Newton A, Petropoulos CJ, Brindeiro RD, Freed**
EO. 2016. Elucidation of the Molecular Mechanism Driving Duplication of the HIV-1 PTAP Late
Domain. J Virol **90:**768-779.
28.   **Sharma S, Arunachalam PS, Menon M, Ragupathy V, Satya RV, Jebaraj J, Ganeshappa**
Aralaguppe S, Rao C, Pal S, Saravanan S, Murugavel KG, Balakrishnan P, Solomon S, Hewlett I,
Ranga U. 2018. PTAP motif duplication in the p6 Gag protein confers a replication advantage on
HIV-1 subtype C. J Biol Chem doi:10.1074/jbc.M117.815829.

29.   **Demirov DG, Orenstein JM, Freed EO.** 2002. The late domain of human immunodeficiency virus
539   type 1 p6 promotes virus release in a cell type-dependent manner. J Virol **76:**105-117.

540   30.   **Huang M, Orenstein JM, Martin MA, Freed EO.** 1995. p6Gag is required for particle production
541   from full-length human immunodeficiency virus type 1 molecular clones expressing protease. J
542   Virol **69:**6810-6818.

543   31.   **Strack B, Calistri A, Craig S, Popova E, Gottlinger HG.** 2003. AIP1/ALIX is a binding partner for
544   HIV-1 p6 and EIAV p9 functioning in virus budding. Cell **114:**689-699.

545   32.   **Dussupt V, Javid MP, Abou-Jaoude G, Jadwin JA, de La Cruz J, Nagashima K, Bouamr F.** 2009.
546   The nucleocapsid region of HIV-1 Gag cooperates with the PTAP and LYPXnL late domains to
547   recruit the cellular machinery necessary for viral budding. PLoS Pathog **5:**e1000339.

548   33.   **Gottwein E, Jager S, Habermann A, Krausslich HG.** 2006. Cumulative mutations of ubiquitin
549   acceptor sites in human immunodeficiency virus type 1 gag cause a late budding defect. J Virol
550   **80:**6267-6275.

551   34.   **Zhadina M, McClure MO, Johnson MC, Bieniasz PD.** 2007. Ubiquitin-dependent virus particle
552   budding without viral protein ubiquitination. Proc Natl Acad Sci U S A **104:**20031-20036.

553   35.   **Coutsinos D, Invernizzi CF, Moisi D, Oliveira M, Martinez-Cajas JL, Brenner BG, Wainberg MA.**
554   2011. A template-dependent dislocation mechanism potentiates K65R reverse transcriptase
555   mutation development in subtype C variants of HIV-1. PLoS One **6:**e20208.

556   36.   **Coutsinos D, Invernizzi CF, Xu H, Moisi D, Oliveira M, Brenner BG, Wainberg MA.** 2009.
557   Template usage is responsible for the preferential acquisition of the K65R reverse transcriptase
558   mutation in subtype C variants of human immunodeficiency virus type 1. J Virol **83:**2029-2033.

561 **Tables**

562 Table 1. List of amino acids that are involved in the interaction between gag-P6, ALIX and
563 ubiquitin (Ub).

Interacting atoms	Residue 1	Residue 2	Distance (Å)	Interaction type
OE1 – NH1	E482 (LD)	R386 (ALIX)	2.96	Charge-charge
OE2 – NH2	E482 (LD)	R386 (ALIX)	2.83	Charge-charge
OE1 – NZ	E387 (ALIX)	K63 (Ub)	3.05	Charge-charge
OE2 – NZ	E387 (ALIX)	K63 (Ub)	2.97	Charge-charge
OE2 – NZ	E64 (Ub)	K485 (LD)	2.93	Charge-charge
OE2 – NZ	E486 (LD)	K501 (ALIX)	2.95	Charge-charge
OE1 – NZ	E16 (Ub)	K501 (ALIX)	3.12	Charge-charge
OH – OD1	Y484 (LD)	D506 (ALIX)	2.72	Hydrogen Bond

E = glutamic acid, R = arginine, Y = tyrosine, D = aspartic acid, K = lysine

Figure legends

Figure 1. Consensus sequences and distribution of HIV-1 gag. (a) Aligned consensus
sequences of gag from HIV-1 type B wild-type (top), HIV-1 type C wild-type (middle), and HIV-
1 type C with tetrapeptide insertion (bottom) from the Los Alamos database. Sequence logos with
one-letter coded amino acid were generated using WebLogo 3. **(b-d)** Distribution of HIV-1 type
C with gag_{wt}, gag_{PYKE}, and gag_{PYQE} in patient cohorts from Sweden **(b)**, Germany **(c)** and
Ethiopia **(d)**.

Figure 2. *In silico* binding prediction between PYKE inserted within HIV-1 type C gag,
ALIX and ubiquitin. (a) Likelihood of ubiquitin binding to lysine residues in the vicinity of or
within the PYKE motif. K_{ins} represents the lysine residue within the PYKE motif. **(b)** All proteins
are presented as ribbon-like structures. The red ribbon represents gag and its PYKE insertion, the
blue ribbon ALIX and its gag-binding sites, and the green ribbon ubiquitin. Amino acids involved
in gag:ALIX binding are shown in one-letter format. Interactions between amino acids are shown
with black dotted lines. *In silico* docking was performed using Schrödinger software.

Figure 3. Co-localization of gag variants with ALIX. (a) Confocal images of HEK293T cells
co-transfected with the GFP-tagged ALIX and mCherry-tagged HIV-1C gag variants (wt,

PYKEi, & PYQEi) expression plasmids. Nuclei are visualized in blue, ALIX in green, and gag in

red. Bars, 20 μm. **(b)** Confocal images of HEK293T cells transfected with the GFP-tagged ALIX

and mCherry-tagged HIV-1C gag PYQEi and HA-tagged ubiquitin expression plasmids. Nuclei

are visualized in blue, ALIX in green, gag in red, and ubiquitin in magenta. Bar, 20 μm.

Figure 4. HIV-1C gag with PYxE insertion has increased binding with ALIX. (a-d) Protein-

peptide interaction using Microscale thermophoresis. The peptide sequences are given in the

individual graphs. **(e)** The equilibrium dissociation constant (Kd) of the individual interactions.

The smaller the Kd value the greater the binding affinity of the ligand for its target. **(f)** HEK293T

cells were co-transfected with GFP-tagged ALIX and gag variants from HIV-1 type B or C. At 24

hours post transfection, cells were lysed and subjected to GFP-pull down. Input and pull-down

samples were subjected to immunoblotting antibodies against GFP (ALIX), p55 (gag), and β-

actin. Data depicted is representative of at least two independent experiments.

Figure 5. PYxE insertion increases viral growth and drug sensitivity towards protease

inhibitor drug lopinavir (a) Gag-pol sequences were amplified from HIV-1 type C patient

isolates and cloned into HIV-1 molecular clone pNL4-3. **(b)** Recombinant viruses were

propagated in HEK293T cells and MOIs were determined in TZM-bl reporter cells. Viral growth

assay was performed in MT4 cells that were infected with the various recombinant viruses at an

MOI of 0.05. Supernatants were harvested on day 0, 3, 5, and 7. Viral load in supernatants were

measured by HIV-1 p24 analysis on the HIV COBAS 8000 platform. Data represents the mean ±

SD of triplicates. **(c)** TZM-bl reporter cells were infected with the individual recombinant HIV

viruses at an MOI of 0.05 and cultured in the presence of individual anti-retroviral drugs at

different dilutions. EC_{50} values for each virus with each drug were determined and EC_{50} fold

changes were calculated compared to pNL4-3 reference virus.

a.

b.

a.

b.

c.