Torsional turning motion of chromosomes as an accelerating force to align homolog

ous chromosomes during meiosis

Kazutaka Takao[1], Kazunori Takamiya[1], Da-Qiao Ding[3], Tokuko Haraguchi[3,4], Yasushi

Hiraoka[3,4], Hiraku Nishimori[1,2], Akinori Awazu[1,2].

[1] Department of Mathematical and Life Sciences, Hiroshima University, 1-3-1

Kagamiyama, Higashi-Hiroshima 739-8526, Japan

[2] Research Center for Mathematics on Chromatin Live Dynamics, Hiroshima University,

1-3-1 Kagamiyama, Higashi-Hiroshima 739-8526, Japan

[3] Advanced ICT Research Institute Kobe, National Institute of Information and

Communications Technology, 588-2 Iwaoka, Iwaoka-cho, Nishi-ku, Kobe 651-2492,

Japan

[4] Graduate School of Frontier Biosciences, Osaka University, 1-3 Yamadaoka, Suita

565-0871, Japan

Abstract

Homologous sets of parental chromosomes must pair during meiosis to produce

recombined sets of chromosomes for their progeny. This is accompanied by nuclear

oscillatory movements. This study aimed to elucidate the significance of these

movements with a model, wherein external force was applied to the oscillating nucleus

and via hydrodynamic interactions within the nucleus. Simulations revealed that a major

force for aligning homologous chromosomes is length-dependent sorting during

chromosomal torsional turning, which occur when the nucleus reverses the direction of

its movement.

Meiosis is an important process for eukaryotic organisms to generate inheritable haploid gametes from a diploid cell. During meiosis, homologous (paternal and maternal) chromosomes recombine; these recombined chromosomes are inherited by their progeny. For recombination of homologous chromosomes, side-by-side alignment of homologous loci along the chromosome is necessary [1]. During this alignment of homologous chromosomes, telomeres form a cluster beneath the nuclear envelope in various eukaryotic organisms [2].

Unicellular fission yeast *Schizosaccharomyces pombe* is a useful model system for studying the alignment of homologous chromosomes during meiosis [3, 4]. A meiotic cell of *S. pombe* contains 3 pairs of homologous chromosomes (chromosomes 1, 2, and 3 of 5.6, 4.6, and 3.5 Mb, respectively) [5]. During meiotic prophase, the nucleus elongates, oscillates between the cell poles, and shows drastic deformation, so-called "horsetail movement" [6], during which telomeres remain clustered at the spindle pole body (SPB) at the leading edge of the nuclear movement [6]. Telomere clustering and nuclear movements reportedly promote the alignment of homologous chromosomes [7,8,9].

To elucidate mechanisms underlying the alignment of homologous chromosomes, computer simulation strongly facilitates the examination and prediction of biological events under conditions that are never accomplished empirically. Various theoretical models of the pairing of homologous loci during meiosis have been proposed on the basis of telomere clustering [10, 11, 12, 13]. However, the role of the nuclear envelope, which confines the chromosomes, remains unknown in these models.

Studies are required to consider the role of the nuclear envelope in models of the meiotic nucleus of *S. pombe*, since its nucleus migrates and undergoes striking deformation during meiosis. During the horsetail movements, the nuclear envelope receives drag forces from the cytoplasm and from the nucleoplasm, while each chromosome receives drag forces only from the nucleoplasm. Thus, models of the meiotic nucleus of *S. pombe* would help describe the motion of the deformable nuclear envelope, direct collisions between the nuclear envelope and chromosomes, effects of the cytoplasm on the nuclear envelope, and effects of the nucleoplasm on chromosomes and the nuclear envelope, in addition to telomere clustering at the SPB. This study

aimed to propose the first model displaying the aforementioned nuclear membrane

dynamics and investigate the role the horsetail nuclear movements in the alignment of

homologous chromosomes via simulations of the presented model..

Model construction: A model of chromosomes confined within the nuclear envelope of

S. pombe cells (Fig. 1(a)) was developed using the following methods. We described

each chromosome as a chain and the nuclear envelope as a one-layered shell—both

comprising particles with an excluded volume (Fig. 1(b)). Each chromosome was

assumed to consist of particles with uniform radii since they were expected as almost

uniformly condensed chains during meiosis. A meiotic nucleus of *S. pombe* contains 3

homologous pairs of chromosomes; thus, the model contains 6 chains in total confined

in the shell. To account for the effects of the SPB on chromosomal movement, we

constructed a virtual sphere around the SPB ("the SPB sphere"), penetrating a limited

area of the shell (nuclear envelope) extending toward the space inside the shell (Fig.

1(b)). To simulate telomere movement, which are clustered near the SPB during the

horsetail movement, we assumed that ends of the chromosome chains move along the

surface of the SPB sphere inside the shell.

To simulate the horsetail movements, we assumed that the SPB sphere oscillates

periodically in the cell. Here, the drag force from the cytoplasm is exerted exclusively

on the nuclear envelope. Since the cytoplasm can be considered a static intracellular

fluid, we assumed that this drag force is proportional to the velocity of the nuclear

envelope. However, the drag forces from the nucleoplasm to the nuclear envelope and

chromosomes depend on the intranuclear velocity profile of the nucleoplasm, which is

primarily determined on the basis of velocity profiles of chromosomes and the nuclear

envelope. Thus, drag forces from the nucleoplasm could be described by the

hydrodynamic interaction among particles in the chains and the shell. Furthermore, we

assumed the volume of the nucleus tend to be sustained since the in and out flow of

nucleoplasm through the nuclear envelope were considered to be little during meiosis.

These methods allowed for the generation of an appropriate model involving the effects

indispensable for simulating the *S. pombe* meiotic nucleus.

Model implementation: We assumed that chromosomes 1, 2, and 3 consist of N_n

particles (n = 1, 2, or 3) and the nuclear envelope consists of *M* particles with radii = *r*.

The motion of the *i*-th particle is given as

95
$$m\ddot{\vec{x_i}} = -\gamma_i \dot{\vec{x_i}} - \vec{\nabla_i} U(\{\vec{x_i}\}) + \sum_j \vec{F_{ij}} \quad (1)$$

where $\vec{x_i} = (x_i, y_i, z_i)$ and m are the position and mass of the *i*-th particle,

respectively. γ_i indicates the coefficient of a drag force from the cytoplasm where

$\gamma_i = 6\pi r\eta$ if the *i*-th particle is on the nuclear envelope and $\gamma_i = 0$ if the *i*-th particle

is on the chromosome. Here, η was assumed as the viscosity of cytoplasm.

In the second term of equation (1), the potential of the system $U(\{\vec{x_i}\})$ is given as

follows:

102
$$U(\{\vec{x_i}\}) = U_{cs} + U_{ms} + U_e \qquad (2)$$

103   where U_{cs} is the contribution from the chain, U_{ms} is the contribution from the nuclear

104   envelope, and U_e *is* the potential of the excluded volume effects.

105   The potential contribution from the chromosome chain U_{cs} is given as follows:

106
$$U_{cs} = \sum_{i<j} \frac{k_c}{2} (|\vec{x_i} - \vec{x_j}| - 2r)^2 + \sum_{i=telo} \frac{k'_c}{2} (|\vec{x_i} - \vec{X_{SPB}}| - (R_s + r))^2 \qquad (3)$$

where the *i*-th particle is a part of chromosomes, *r* indicates the radii of particles, and

$\vec{X_{SPB}}$ and R_S indicate the position of the center and the radius of the SPB sphere. We

assumed each chromosome is a chain of particles connected in a line. Σ in the 1st term

indicates the sum for the *i*-th and *j*-th particles (*i* < *j*) that are adjacent and bonded along

the chromosome, and Σ in the 2nd term indicates the sum for the *i*-th particle at the

ends of chromosomes corresponding to the telomeres.

The potential contribution from the nuclear envelope U_{ms} is given as follows:

$$U_{ms} = \sum_{i<j,L_{i,j}<3r} \frac{k_m}{2} (|\vec{x_i} - \vec{x_j}| - L_{i,j})^4 + \sum_{L_{i,SPB}<R_s} \frac{k'_m}{2} (|\vec{x_i} - \overrightarrow{X_{SPB}}| - L_{i,SPB})^2$$

$$+ \frac{k_V}{2} \left(\sum_i \frac{\pi r^2}{3} |\vec{x_i} - \overrightarrow{x_{CM}}| - V \right)^2$$

$$+ \sum_i \theta(\sqrt{x_i^2 + y_i^2} - (R_c - r)) \frac{k_w}{2} (\sqrt{x_i^2 + y_i^2} - (R_c - r))^2$$

114   (4)

where the i-th particle is a part of the nuclear envelope. θ is Heaviside step function

and $L_{i,j}$ indicates the initial distance between the centers of the i-th and j-th particles

constructing the nuclear envelope ($L_{i,SPB}$, the initial distance between the centers of the

i-th particle and the SPB sphere). Here, the initial structure of the nuclear envelope is

indicated as a spherical shell with a thickness of one particle layer constructed by the

rings along with y-axis with various radii consisting of particles (Fig. 1(c)), and initial

position of the center of the SPB sphere is denoted as $\overrightarrow{X_{SPB}}(0) = (R_c, 0, 0)$ (R_C is the

radius of the cell as described below). Σ in the 1st term of U_{ms} indicates the sum for

the i-th and j-th particles ($i < j$) that are adjacent and bonded obeying $L_{i,j} < 3r$; Σ in

the 2nd term indicates the sum for the i-th particle obeying $L_{i,SPB} < R_s$; Σ in other

terms indicate sum for all particles constructing the nuclear envelope. Notably, we

assumed that the nuclear envelope can deform easily since it consists of lipids that can

exhibit the elasticity as well as the fluidity. Thus, we used an unharmonic potential in

the 1st term of U_{ms} which provides weaker tension for local small changes in relative

positions among particles in the nuclear envelope than the harmonic potential, inducing

larger global deformation of nuclear envelope.

The 2nd term of U_{ms} indicates the binding of the SPB to the nuclear envelope,

where particles constituting the nuclear envelope and confined in the SPB sphere were

connected tightly to the center of the SPB sphere. The 3rd term of U_{ms} provides the

force to sustain the volume of the cell nucleus where $\overrightarrow{x_{CM}}$ and V indicate the position

of the center of mass and the volume of the nucleus, respectively. Here, V is denoted by

$\sum_i \frac{\pi r^2}{3} |\vec{x_i} - \overrightarrow{x_{CM}}|$ with initial values of $\vec{x_i}$. The 4th term of U_{ms} indicates the

excluded volume effect of the cell wall, where the cellular shape was assumed to be a

parallel cylinder along the z-axis, with a radius R_C. Here, the central axis of this cylinder

is denoted as $(0,0,z)$, and θ indicates the Heaviside function.

U_e denotes the potential of the excluded volume effects among particles and the SPB

sphere as follows:

$$U_e = \sum_{i<j} \theta(2r - |\vec{x_i} - \vec{x_j}|)\frac{k_e}{2}(|\vec{x_i} - \vec{x_j}| - 2r)^2$$

$$+ \sum_i \theta((R_S + r) - |\vec{x_i} - \overrightarrow{X_{SPB}}|)\frac{k'_e}{2}(|\vec{x_i} - \overrightarrow{X_{SPB}}| - (R_S + r))^2$$

142   (5).

In the third term of equation (1), $\overrightarrow{F_{ij}}$ indicates the force of hydrodynamic effects

from the j-th particle to the i-th particle, providing a drag force to each particle from the

nucleoplasm. Herein, we employ the lubrication approximation [14, 15] because it can

describe hydrodynamic effects among polymers modeled by particle chains in the

simulation of the polymers suspension in shear flow suitably [15], which is denoted as

follows:

$$\overrightarrow{F_{ij}} = \begin{cases} \frac{3}{2}r^2\pi\eta \frac{(\dot{\vec{x_j}}-\dot{\vec{x_i}})\cdot\overrightarrow{n_{ij}}}{|\vec{x_j}-\vec{x_i}|-2r}\overrightarrow{n_{ij}} & (if\ |\vec{x_j}-\vec{x_i}|>2.001r) \\ \frac{3}{2}r^2\pi\eta \frac{(\dot{\vec{x_j}}-\dot{\vec{x_i}})\cdot\overrightarrow{n_{ij}}}{0.001r}\overrightarrow{n_{ij}} & (if\ |\vec{x_j}-\vec{x_i}|\leq2.001r) \end{cases}$$ (6)

where r and $\overrightarrow{n_{ij}}$ denoted the radius of the particle and a relative unit vector from the

j-th particle to the i-th particle $\overrightarrow{n_{ij}} = \frac{\vec{x_i}-\vec{x_j}}{|\vec{x_i}-\vec{x_j}|}$, respectively. Nucleoplasm viscosity was

assumed the same as the cytoplasm viscosity η in this model.

Parameters for simulations: In the simulations, we assumed a particle radius

$r = 0.084 \times 10^{-6}\ (m)$, where each particle models a local region of chromatin of ~100

kb. Here, chromosomes 1, 2, and 3 (5.6 Mb, 4.6 Mb, and 3.5 Mb, respectively) were

described by the chains consisting of 56, 46, and 34 particles ($N_1 = 56$, $N_2 = 46$, and N_3

= 34; we considered N_3 as an even number for convenience of analysis). The nuclear

envelope was constructed by using 994 particles ($M = 994$) where we assumed their radii

were the same as those of chromosome particles ($r = 0.084 \times 10^{-6}(m)$) because of the

convenience to simulate the hydrodynamic effects using Eq. (6). Here, the inner

diameter of the initial spherical structure of the nuclear envelope was assumed to be

$2.7 \times 10^{-6}(m)$, the initial distance between the center of the SPB sphere and that of the

nucleus was assumed to be $2.0 \times 10^{-6}(m)$, and the radius of the SPB sphere was

assumed $R_s = 10r$ (Fig. 1(b)). In this case, the diameter of the cross-section where

nucleus and the SPB sphere intersect was $\sim 1.0 \times 10^{-6}(m)$. The initial configurations

of the chromosomes were randomized in this spherical shell.

The viscosity of nucleoplasm and cytoplasm η was ~ 0.64 $(kg\ m^{-1}s^{-1})$ [12].

The elastic constants were assumed as appropriate large values for the excluded volume

effects of all particles and the connections among particles in chromosomes, among

particles in chromosomes and the SPB sphere, and among particles in the nuclear

envelope and SPB sphere, as $k_c = k_w = k_e = k'_e = 2.77 \times 10^8 (kg\ s^{-2})$, $k'_c =$

$6.22 \times 10^8 (kg\ s^{-2})$, and $k'_m = 8.12 \times 10^8 (kg\ s^{-2})$. We also assumed the elastic

constant to keep the connection of particles in the nuclear envelope and the nuclear

volume as appropriate values as $k_m = 4.77 \times 10^{20} (kg\ s^{-2}\ m^{-2})$ and $k_V = \frac{150}{\pi} k_m$

with which the present model reproduced a similar shape variation of the nuclear

envelope via horsetail movements. The inertia term of Eq. (1) cannot be eliminated in

order to employ the lubrication approximation although the equation of motion of such

microscopic system should obey the equation with overdamp limit. However, we

assumed m as $\sim m/6\pi r\eta \sim 10^{-3}$ sec ($m = 8.3 \times 10^{-10}(kg)$ was assumed in this

paper) that indicates the relaxation times of the particle velocities were negligibly

smaller than the period of horsetail movement $\sim 10^2$ sec. Therefore, we could neglect

the effects of inertia to simulate our focused phenomena. We confirmed that the

following results were qualitatively unaffected by small changes in these parameters

when k_m was chosen appropriate to exhibit sufficient shape variation of the nuclear

envelope where chromosomes can elongate. Thus, we consider these parameters

appropriate to simulate intracellular environment.

Results: To evaluate the effects of the horsetail movement, we simulated the following four cases of motions of the SPB sphere in the cylinder (cell) with radius $R_c = 2 \times 10^{-6} (m)$ as shown in Fig. 2: (a) 3-d 8-shaped oscillatory motions as $\overrightarrow{X_{SPB}} = (R_c - B(1 - \cos 4\pi wt), C\sin 4\pi wt, A\cos 2\pi wt)$, (b) 3-d O-shaped oscillatory motions as $\overrightarrow{X_{SPB}} = (R_c - B(1 - \cos 4\pi wt), C\sin 2\pi wt, A\cos 2\pi wt)$, (c) 1-d motions as $\overrightarrow{X_{SPB}} = (R_c, 0, A\cos 2\pi wt)$, and (d) 1-d constant velocity linear motion as $\overrightarrow{X_{SPB}} = (R_c, 0, \pi wAt)$. Herein, $A = 7.3 \times 10^{-6} (m)$ and $w = \frac{1}{3} \times 10^{-2} (s^{-1})$ $(= 12 (h^{-1}))$ since the SPB completes 30 oscillations in 2 ~3 h from end to end in spheroid-shaped *S. pombe* cells [9]. Furthermore, we assumed $B = 0.208 \times 10^{-6} (m)$, and $C = 0.67 \times 10^{-6} (m)$. The results were qualitatively independent of the detail values of *B* and *C*.

Chromosomal behavior in the migrating nucleus was simulated under the aforementioned 4 conditions. Simulations were repeated with eight initial states for each condition (Figs. 3, 4, and Supplementary movies 1 and 2). Figure 3 shows typical snapshots for t = 75, 3375, and 7950 (s) in an example of 8-shaped 3-d oscillatory motion. Elongation of the nuclear envelope (Supplementary movie 1) and switching of relative positions among chromosomes (Fig. 3, Supplementary movie 2) were observed during oscillatory motions of the SPB. The present results indicate the validity of our simulations.

To obtain a pictorial view of chromosomal movements, we plotted positions of the chromosomes as probability distributions along the x-y plane (Fig. 4). The plots were obtained from the 34 (= N_3) nearest particles from the SPB sphere ($N_3 / 2 = 17$ nearest particles from each end) along each of the 6 chromosomes (Fig. 4a). Figure 4b shows typical snapshots of the probability distributions of particles for t = 75, 3375, and 7950 (s) in the 3-d 8-shaped oscillatory motion. When t = 75, the probability distributions of particles of chromosome 1 and 3 exhibited two separated hills respectively, that indicated homologous pairs of chromosomes 1 and 3 were separated far away. On the other hand, the probability distributions of particles of all chromosomes exhibited only one hill respectively at different area when t = 7950. These facts indicated each homologous chromosome pair gathered temporally and non-homologous chromosomes

were separated from each other, that is, alignment of homologous chromosomes
occurred.

Figure 4 (c) – (e) show typical snapshots of the probability distributions of particles
at t = 7950 (s) when the SPB sphere exhibits a 3-d O-shaped oscillatory motion, a 1-d
oscillatory motion, and a linear motion. As shown in Fig. 4 (b) and (c), each
homologous pair of chromosomes gathered temporally and non-homologous
chromosomes were separated from each other when the SPB sphere exhibits 3-d
oscillatory motion involving 3D reversal (torsional turning) motion of chromosomes.

However, when the SPB sphere exhibits 1-d oscillatory motion involving no
torsional motion of chromosomes, some homologous chromosome pairs often stay
separated from each other (Fig. 4 (d); alignment of homologous chromosomes was
observed exceptionally only in one out of eight simulations). Furthermore, the assembly
of homologous chromosomes did not occur when the SPB sphere exhibits a linear
motion involving no chromosomal turning (Fig. 4 (e)). These results suggest that
chromosomal torsional turning motion is crucial for shuffling of chromosomes for
appropriate alignment of homologous chromosomes.

By the simulations of following virtual situations where chromosomes 1 and 2
became shorter as (i) 4.6 Mb and 4.0 Mb ($N_1 = 46$, $N_2 = 40$, and $N_3 = 34$), and (ii) 3.8
Mb and 3.6 Mb ($N_1 = 38$, $N_2 = 36$, and $N_3 = 34$), we found that the frequency of the
separation of non-homologous chromosomes decreased with the decrease in their
difference in chain lengths: the separation of non-homologous chromosomes was
observed in three out of eight simulations in both situations even when the SPB sphere
exhibits a 3-d 8-shaped oscillatory motion. We also found the separation of
non-homologous chromosomes occurred little if the elastic constant k_m among
particles in nuclear envelope was given so large as that the shape variation of the
nuclear envelope was not sufficient for the elongation of chromosomes via the SPB
movements. By these facts, non-homologous chromosomes are considered to separate
owing to differences in chain lengths among non-homologous chromosomes.

During the horsetail movement, the SPB movement reverses its direction
periodically. Chromosomes receive drag forces from the nucleoplasm at every turn of

the SPB, while no drag force is exerted on chromosomes and the nuclear envelope if

SPB migrates linearly with a constant velocity. During reversal of SPB, the region distal

from the SPB in longer chromosomes tends to be dragged by the slow (nearly stopped)

motion of the rear region of the bent nucleus, while that in the shorter one tends to

follow the SPB motion (Fig. 5). Thus, the relative positions of longer and shorter

chromosomes tend to separate gradually from each other owing to reversal of the

iterative direction of the SPB movement, resulting in the separation of non-homologous

chromosomes. Owing to this separation, homologous chromosomes are aligned and

stabilized during the periodic reciprocation of the SPB.

Discussion: Homologous chromosome pairing has received increasing attention in

studies in physics, and several models have been proposed to describe its underlying

mechanisms [10, 11, 12, 13, 16]. In the present simulations, separation of

non-homologous chromosomes was achieved when the SPB sphere shows 3-d

oscillations in the cell, but not when the SPB sphere exhibits only a linear motion. This

result is further substantiated by previous experimental evidence regarding *S. pombe*:

the efficiency of homologous recombination decreases drastically when the nucleus

does not exhibit the horsetail movement but just elongates [17]; the iterative reversals of

the direction of the SPB movement during the horsetail movement are essential for the

alignment of homologous chromosomes via separating non-homologous chromosomes

[7, 9]. Our simulation reveals that a major force for the alignment and sorting of

homologous/non-homologous chromosomes is the nucleoplasmic flow generated owing

to SPB reversal and deformation of the nuclear envelope, involving direct collision and

indirect hydrodynamic interactions between chromosomes and the nuclear envelope,

which were previously unclear. The present model is the first, to our knowledge, to

consider both direct collisions and indirect hydrodynamic interactions as essential

mechanisms for the alignment of homologous chromosomes.

This study was partly supported by the Platform Project for Support in Japan Agency

for Medical Research and Development (to AA); MEXT KAKENHI Grant Number JP

17K05614 (to AA), JP18H05528 (to TH) and JP18H05533 (to YH).

References

[1] D. Zickler, and N. Kleckner, Cold Spring Harb. Perspect. Biol. **7**, a016626 (2015).

[2] H. Scherthan, Nat. Rev. Mol. Cell Biol. **2**, 621-627 (2001).

[3] R. Egel, and D. Lankenau (ed.), *Recombination and Meiosis: Models, Means, and Evolution* (Springer, Berlin-Heidelberg, 2008).

[4] D.-Q. Ding, T. Haraguchi, and Y. Hiraoka, Y, FEBS J. **277**, 565-570 (2010).

[5] V. Wood et al., Nature **415**, 871-880 (2002).

[6] Y. Chikashige, D.-Q. Ding, H. Funabiki, T. Haraguchi, S. Mashiko, M. Yanagida, and Y. Hiraoka, Science **264**, 270-273 (1994).

[7] O. Niwa, M. Shimanuki, and F. Miki, EMBO J. **19**, 3831-3840 (2000).

[8] D.-Q. Ding, A. Yamamoto, T. Haraguchi, and Y. Hiraoka, Dev. Cell **6**, 329-341 (2004).

[9] M. R. Chacón, P. Delivani, and I. M. Tolić, Cell Reports **17**, 1632-1645 (2016).

[10] D. Dorninger, G. Karigl, and J. Loidl, J. Theor. Biol. **176**, 247–60 (1995).

[11] C. A. Penfold, P. E. Brown, N. D. Lawrence, and A. S. H. Goldman, PLoS Comput. Biol. **8**, e1002496, (2012).

[12] Y. T. Lin, D. Frömberg, W. Huang, P. Delivani, M. Chacón, I. M. Tolić, F. Jülicher, V. Zaburdaev, Phy. Rev. Lett. **115**, 208102 (2015).

[13] W. F. Marshall, and J. C. Fung, Phys. Biol. **13,** 026003 (2016).

[14] N. A. Frankel, and A. Acrivos, Chem. Eng. Sci. **22**, 847-85 (1967).

[15] S. Yamamoto, and T. Matsuoka, J. Chem. Phys. **102**, 2254-2260 (1995).

[16] K. Takamiya, K. Yamamoto, S. Isami, H. Nishimori, and A. Awazu, *Nonlinear Theory and Its Applications,* vol. **7**, issue 2, 66-75 (2016).

[17] A. Yamamoto, R. R. West, J. R. McIntosh, and Y. Hiraoka, J. Cell Biol. **145**, 1233-1249 (1999).

Figure legends

FIG. 1. (a) Illustration of the meiotic nucleus of *Schizosacchromyces pombe* containing

3 homologous pairs of chromosomes (Chr, 1, 2, and 3), the nuclear envelope (NE), the

spindle pole body (SPB), and microtubules (MT). (b) Illustration of the model

composed of particle populations. (c) Illustration of the model of an intracellular region

described as a cylindrical space, and the initial structure of the nuclear envelope

constructed by combining particle rings (along dashed circles) along the vertical y-axis

with various radii.

FIG. 2. Illustration of trajectories of the spindle pole body (SPB) sphere motions in the

322   cell (cylindrical space). (a) 3-d 8-shaped oscillatory motions, (b) 3-d O-shaped

323   oscillatory motions, (c) 1-d oscillatory motions, and (d) 1-d constant velocity linear

324   motion. Yellow particle indicates the SPB sphere; red and blue arrows indicate the

325   trajectory of downward and upward motion, respectively.

FIG. 3. Typical time course of chromosome dynamics in the case that the spindle pole
body (SPB) sphere exhibits 3-d 8-shaped oscillation. Snapshots of chromosome
configurations in the nucleus from two points of view (projected on y-z and x-z plane) at
time t = 75 s (after 0 oscillations), 3375 s (after 10 oscillations), and 7950 s (after 26
oscillations). The large sphere indicates the SPB sphere; red, black, and blue particle
chains indicate Chr. 1, Chr. 2, and Chr. 3, respectively. Blue arrows indicate the moving
directions of SPB.

FIG. 4. Probability distributions of particles in each of the chromosome chains near the
spindle pole body (SPB) sphere along the x-y plane. (a) The 34 ($= N_3$) nearest particles
from the SPB sphere along each of the 6 chromosomes (filled particles) were used to
estimate probability distributions. (b) Probability distributions of the 34 ($= N_3$) nearest
particles on each chromosome indicated as filled particles in (a). Curves represent
contour lines of probability distributions of particles in the 3-d 8-shaped oscillation of

the SPB sphere at t = 75, 3375, and 7950 s; red, black, and blue curves indicate Chr. 1,

Chr. 2, and Chr. 3, respectively. The intensity of the contour lines represents the

probability values of 0.3, 0.5, and 0.75 in an increasing order of darkness. (c) – (e)

Probability distributions at t = 7950 s in (c) 3-d O-shaped oscillation, (d) 1-d

oscillations, and (e) constant velocity linear motion of the SPB sphere. Colors of the

contour lines represent probability distributions of each chain as in (b).

FIG. 5. Illustration of velocities of the spindle pole body (SPB; white arrow), long and

short chromosomes (red and blue arrows, respectively), the nuclear envelope (yellow

arrows), and the nucleoplasmic flow (black arrows). When the SPB exhibits a constant

linear velocity motion, all regions of the nuclear envelope, the nucleoplasm, and

chromosomes relax to the same velocity as the SPB (Left). During the turn of the SPB,

the regions of the nuclear envelope, the nucleoplasm, and chromosomes at the nuclear

front end (near the SPB) follow the motion of the SPB, while those at the nuclear rear

end are almost stopped (Right). In the latter case, longer chromosomes tend to be

dragged by the slow motion of the nuclear rear region, while shorter chromosomes tend

to follow the motion of the SPB.