1   Taxonomy, ecology and distribution of *Juniperus oxycedrus* L. group in the Mediterranean

2   Region using morphometric, phytochemical and bioclimatic approaches

Taxonomy, ecology and distribution of *Juniperus oxycedrus* L.

Ana Cano Ortiz[1], Carmelo M. Musarella[1,2,*], José C. Piñar Fuentes[1], Carlos J. Pinto Gomes[3],

Giovanni Spampinato[2], Eusebio Cano[1]

[1]Department of Animal and Plant Biology and Ecology, Section of Botany, University

of Jaén, Jaén, Spain

[2]Department of AGRARIA, "Mediterranea" University of Reggio Calabria, Reggio

Calabria, Italy

[3]Department of Landscape, Environment and Planning, Institute for Mediterranean

Agrarian and Environmental Sciences (ICAAM), School of Science and Technology,

University of Évora, Évora, Portugal

* Corresponding author

E-mail: carmelo.musarella@unirc.it

Abstract

The ecology, taxonomy and distribution of the *Juniperus oxycedrus* L. group of taxa are studied. From an ecological aspect, this work proposes a new ombroedaphoxeric index to explain the presence of populations of *Juniperus* in ombrotypes that are not the optimum for these taxa. The controversy among various authors on the taxonomy of the *oxycedrus* group with regard to *J. oxycedrus* subsp. *badia* and *J. oxycedrus* subsp. *lagunae* is clarified. The phytochemical differences in essential oils are addressed, and their similarities analysed; greater similarities are observed between *oxycedrus* and *badia* (H. Gay) Debeaux, and between *navicularis* Grand. and *macrocarpa* (Sm.) Ball. The phytochemical, molecular and distribution differences allow *J. macrocarpa* and *J. navicularis* to be maintained as species.

Introduction

The prickly juniper species (*Juniperus oxycedrus* L.) is widely distributed throughout the Mediterranean area, and has three subspecies on the Iberian Peninsula: *oxycedrus, badia* and *macrocarpa* [1]. The subspecies *macrocarpa* and *oxycedrus* extend as far as the Balearic Islands, Corsica, Sardinia and the Italian Peninsula [2], whereas the subsp. *oxycedrus* is found as far as Croatia and Slovenia [3]. At one time the *J. oxycedrus* group included *Juniperus navicularis* (syn.: *Juniperus oxycedrus* L. subsp. *transtagana* Franco) and *Juniperus deltoides* R.P. Adams [syn.: *Juniperus oxycedrus* L. subsp. *deltoides* (R.P.Adams) N. G. Passal.], which was described as a new species by [4] for Greece. Subsequently, this same author [5] reported the distribution of *J. deltoides* in Italy, Croatia, Greece and Turkey –coexisting in Turkey with *Juniperus polycarpos* K. Koch–, and established phytochemical differences with *J. oxycedrus* due to its lower

alpha-pinene content and higher limonene content. These data have also been confirmed

by [6].

All the taxa in the *J. oxycedrus* group grow in xeric environments, generally in

inaccessible places such as limestone or siliceous screes, although isolated individuals

may appear in *Quercus rotundifolia* Lam. woodlands. The phytocoenoses of *Juniperus*

are of considerable ecological interest due to the presence of companion endemics in

these plant communities, which serves as the justification for their study. They form

small vegetation islands that act as species reservoirs as they are not used for either

agriculture or livestock farming and thus have not been destroyed by human action. In

these phytocoenoses it is frequent to find endemic species with different degrees of

distribution on the peninsula, such as *Echinospartum ibericum* Rivas Mart., Sánchez

Mata & Sancho, *Adenocarpus argyrophyllus* (Rivas Goday) Caball., *Digitalis purpurea*

L. subsp. *mariana* (Boiss.) Rivas Goday, *Sideritis lacaitae* Font Quer, *Coincya*

longirostra (Boiss) Greuter & Burdet, *Cytisus scoparius* (L.) Link subsp. *bourgaei*

(Boiss.) Riv.-Mart., *Cytisus striatus* (Hill) Rothm. subsp. *eriocarpus* (Boiss. & Reut.)

Rivas-Mart., *Genista polyanthos* R. Roem. ex Willk., *Dianthus crassipes* R. de Roemer,

Dianthus lusitanus Brot., *Digitalis thapsi* L., *Digitalis purpurea* L. subsp. *heywoodii* P.

Silva & M. Silva, *D. purpurea* L. subsp. *mariana* (Boiss) Rivas Goday, *Securinega*

tinctoria (L.) Rothm., *Lavandula stoechas* L. subsp. *luisieri* (Rozeira) Rozeira,

Lavandula stoechas L. subsp. *sampaiana* Rozeira, *Genista hirsuta* Vahl, *Thymus*

mastichina L., *Thymus grantensis* Boiss. subsp. *micranthus* (Willk.) O. Bolòs & Vigo,

Thymus zygis Loefl. ex L. subsp. *gracilis* (Boiss.) R. Morales, *Antirrhinum graniticum*

Roth. subsp. *onubensis* (Fernández Casas) Valdés. The territories studied are Sites of

Community Interest (SCI) due to their presence on the vertical walls of habitats such as

8210 "Calcareous rocky slopes with chasmophytic vegetation", and 8220 "Siliceous

rocky slopes with chasmophytic vegetation", which include many endemic plant

associations [7] and explain the need to conserve these areas. However, in less steeply

sloping rocky areas, the dominant species is *Juniperus oxycedrus* subsp. *badia*, which

characterises habitat 5210 "Arborescent matorral with *Juniperus spp.*". These zones can

therefore be classified as hotspots of special interest for conservation. All these

associations are included in the Habitats Directive, which justifies the ecological

importance of these areas and the need to study them for their subsequent conservation

[8].

The areas dominated by *Juniperus* species are currently undergoing a process of

expansion due to the increase in areas of scree, which are becoming more widespread

each year due to deforestation and forest fire. This phenomenon causes an increase in

edaphoxerophilous zones and a decrease in climatophilous zones, thus increasing the

potential areas that can act as a refuge for endemic species [9].

The aim of this work is to clarify the taxonomy, ecology and distribution of the taxa in

the *J. oxycedrus* group, which form the communities included in the Habitats Directive

such as the "Arborescent matorral with *Juniperus spp.*" (5210).

Materials and Methods

We studied the *J. oxycedrus* group and conducted herborisation and phytosociological

sampling campaigns in the field to determine the habitat in which these taxa occur. We

examined the differences in the ecology, distribution and taxonomy of the taxa in the *J.*

oxycedrus group by analysing their morphological, ecological and phytochemical

differences. With these characters, we studied the resemblance between the taxa using a

similarity analysis based on the data provided by [10], [11], and [12]. A bioclimatic

study was carried out to explain the presence of *Juniperus* species in different rocky

substrates.

To understand the presence of communities of *Juniperus* in territories dominated by

species in the genus *Quercus*, we applied Thornthwaite's, $ETP_{monthly} = 16(10.T/I)^a$, to

calculate the potential evapotranspiration and 0.2ETP by [13]. With these data we used

the new ombroedaphoxeric index Ioex proposed by [14] which, in a comparative

analysis with the ombrothermic index I_o proposed by [15], justifies the presence of

microwoodlands of *Juniperus* and *Pinus*.

Territories behave differently in response to the general climate, the type of substrate

and the topography of the terrain. For this reason areas on rocky crests –although they

may be located in rainy environments and surrounded by climactic forests– behave

differently from the territories around them. In these circumstances, islands evolve

which may contain edaphoseries, minoriseries and permaseries [16]. All plant

communities growing on rocky crests and steeply sloping areas with extreme gradients,

among others, are very significantly influenced by the soil, which conditions their

existence. All territories have a particular type of substrate and an orography that

determines whether they have a greater or lesser capacity to retain water. In ideal

situations with good soil texture and structure and with no slopes, the water retention

capacity (RC) can be assumed to be maximum (100%). Otherwise, losses occur due to

runoff and drainage, causing the RC to vary. Water is also lost through

evapotranspiration (ETP). However, as plants have the capacity to self-regulate their

losses, it can be accepted that the residual evapotranspiration e = 0.2ETP. There are

therefore two parameters (**e** and **RC**) implicated in the development of a vegetation,

which is essentially conditioned by rainfall. The ombroclimatic index I_o therefore does

not explain the presence of plant communities influenced by the substrate, and for this

reason, we use the Ombroedaphoxeric Index [14] to explain the presence of

communities of *Juniperus* and *Pinus* for territories with a thermotype ranging from the

thermo to the supramediterranean.

$I_{oex} = P_p\text{-}e/T_p \text{ x RC}$

P_p = mean annual precipitation. T_p = mean annual temperature [15]. e = residual

evapotranspiration whose value is 0.2 ETP [13]. RC = retention capacity in parts per

unit, whose values may be 0.25, 0.50, 0.75 and 1.

For the authorship of the taxa we have followed [2], [17], [18], [19], [20], [21], [22],

[23], [24].

129   Results

130   Ecology and distribution

Juniperus oxycedrus L. has its distribution in the Mediterranean region, eastern

Portugal, Morocco and even extends as far as northern Iran [1]. According to this

author, this species has three clearly differentiated subspecies. The subspecies

macrocarpa (Sm.) Ball. is typical of dunes and coastal sand flats, and may occasionally

occupy rocky areas. The communities characterized by this taxon on the Iberian

Peninsula are described and included in the alliance *Juniperion turbinatae* Rivas-

Martínez 1975 *corr.* 1987, along with other plant communities presided by *Juniperus*

navicularis Gand. [synonym of *Juniperus oxycedrus* L. subsp. *transtagana* Franco in

Feddes Repert. Spec. Nov. Regni Veg. 68:166 (1963)] and *Juniperus phoenicea* L.

subsp. *turbinata* (Guss.) Nyman, also typical of psamophilous environments and dunes

in coastal zones [25]. The subspecies *macrocarpa* is distributed in Iberian and Italian

territories and in the Balearic Islands, Corsica, Sardinia, Sicily (western

Mediterranean); this taxon does not reach the eastern Mediterranean and is substituted

in Greece by *Juniperus drupacea* Labill..

J. oxycedrus L. subsp. *oxycedrus* and *J. oxycedrus* L. subsp. *badia* (H. Gay) Debeaux

are found on the Iberian Peninsula on hard acid and basic substrates. The area of

distribution of *J. oxycedrus* extends as far as Italy [2], Croatia and Slovenia [3]. It is

generally found within *Quercus rotundifolia* Lam. and *Quercus ilex* L. woodlands, and

also frequently forms plant communities on screes; as both the subsp. *oxycedrus* and

the subsp. *badia* have their optimum development on skeletic-rocky substrates and in

semiarid-dry bioclimatic environments (Fig 1).

Fig 1. Optimum areas for the development of *Juniperus oxycedrus* group.

Optimum ombrotypes: semiarid, dry.

There are also some doubts as to the presence of *J. oxycedrus* subsp. *badia* on the

African continent. Some authors such as [26] do not recognise this taxon in northern

Africa, although [27] reports *J. oxycedrus* f. *badia* H. Gay in North Africa, and the

subsp. *oxycedrus* and subsp. *macrocarpa*.

Bolòs & Vigo [28] include the var. *laguna* Pau *ex* Bolòs et Vigo in *J. oxycedrus* subsp.

oxycedrus, as it has the same characters as the subspecies *badia*. Recently, based on the

work of [20], [29] formulated the new combination *Juniperus oxycedrus* L. subsp.

lagunae (Pau *ex* C. Vicioso) Rivas Mart. This all serves to highlight the complexity of

this taxon, whose distribution area is insufficiently known. However, its presence is

very evident in the centre and south of the Iberian Peninsula, where it grows in

formations with a broad extension, generally on screes and biotopes with shallow soils

where holm oaks (*Quercus rotundifolia*) cease to be dominant or simply cannot exist

due to absence of the necessary ecological and/or soil conditions for these taxa to

develop [30] (Fig 2).

Fig 2. Distribution of taxa of *Juniperus oxycedrus* group in the Mediterranean

bassin. • *J. oxycedrus* subsp. *oxycedrus*, ○ *J. oxycedrus* subsp. *badia*, ▲ *J. navicularis*,

△ *J. macrocarpa*, ● *J. polycarpos*, □ *J. deltoides*, ◊ *J. drupacea*.

Table 1 shows the values of Pp, Tp and Io according to the criterion established by

Rivas-Martínez. The formula of Thornthwaite $ETP_{monthly} = 16(10.T/I)^a$ is applied to

obtain the value of ETP, where T is the mean monthly temperature, I is the annual heat

index and a parameter that depends on the values taken by I.

Table 1. Comparative value of indices Io and I$_{oex}$ in some localities in the southern

Iberian Peninsula.

Weather Station	Pp	Tp	Io	Ombrotype	ETP	e	Ioex1	Ioex2	Ioex3	Ombroclimatic behavior of the locality
Almadén-Minas (CR)	625.20	194.40	3.20	dry	808.54	161.70	0.59	1.19	1.78	semiarid
Cabezas Rubias (H)	993.40	177.60	5.60	subhumid	702.14	140.42	1.20	2.40	3.60	dry
Aracena (H)	1025.80	175.20	5.90	subhumid	703.46	140.69	1.26	2.52	3.78	dry
Santiago Pontones (J)	1148.70	164.40	7.00	subhumid	675.23	135.04	1.54	3.08	4.62	dry
Vadillo Castril (J)	1182.20	140.40	8.40	humid	488.88	97.72	1.93	3.86	5.79	subhumid
Grazalema (Ca)	1962.20	183.60	11.00	humid	726.22	145.24	2.47	4.94	7.42	subhumid
Montoro (Co)	522.40	210.00	2.50	dry	903.15	180.63	0.40	0.81	1.22	arid
Pozoblanco (Co)	514.40	193.20	2.70	dry	805.45	161.09	0.45	0.91	1.37	arid
Villanueva del Arzobispo (J)	698.20	196.80	3.50	dry	915.70	183.14	0.65	1.30	1.96	semiarid

Applying the formula I$_{oex}$ for the assumptions that RC is 0.25, 0.50, 0.75, we obtain

three values, of which the most representative is I$_{oex2}$. Table 1 shows the equivalence

values so that although the territorial bioclimate allows the existence of climactic

forests, in wild areas with RC = 50% the humid ombrotype becomes dry or subhumid

depending whether the value of RC =25% or 50%. The subhumid becomes dry and the

dry becomes semiarid or arid. Therefore, in areas with Io > 8 we obtain I_{oex2} values of

3.86 and 4.94, which is equivalent to subhumid. This explains the fact that in rocky

areas there is an edaphoxerophilous community of *Quercus faginea* s.l. or *Abies pinsapo*

Boiss., as occurs in Grazalema (Cadiz); or that a value of I_{oex1} = 2.47 is obtained in the

case that RC = 25%. In this situation, there is a presence of an edaphoxerophilous

community of *Quercus rotundifolia* in Cazorla (Jaén) and in Grazalema (Cadiz). When

the underlying ombrotype is subhumid, the equivalence value of I_{oex2} corresponds to

dry; and a starting situation with a dry I_o gives semiarid and even arid values of I_{oex2} if

we start from a horizon that is lower than dry. This does not allow development of tree

species of *Quercus*, but does allow development of the genera *Juniperus* and *Pinus*. The

value of I_{oex} is affected by climate change, as evidenced by Del Río et al. [31], who

report a heterogeneous trend in terms of annual rainfall redistribution, with a decline in

most of the mountainous areas of Grazalema, Ronda, Cazorla, Segura, Sierra Nevada

and a large part of the Sierra Morena. However, these authors have detected an increase

in rainfall on the Andalusian coast and particularly in Almería. This affects the forest

stands, and in conjunction with human activity [8], favours a redistribution of the

current forests, with a decline in *Quercus* woodlands and an increase in the micro-

woodlands of *Juniperus*.

Taxonomy

As specified by [10] and [12] there is a clear phytochemical differentiation between the

three subspecies of *Juniperus oxycedrus*. Subsequently [5] clarifies the phytochemical

differences between *J. oxycedrus* and *J. deltoides*, a species described by this author [4]

in Greek territories. This taxon has a low content in alpha-pinene and a high content in

limonene, which, among other morphological differences, justifies the rank of species

for this taxon. There are also substantial differences in the essential oils of *Juniperus*

navicularis Gand. compared to the rest [11] .

Alpha-pinene is common to the four taxa and p-cymene to *J. oxycedrus* subsp.

oxycedrus, *J. oxycedrus* subsp. *macrocarpa* and *J. oxycedrus* subsp. *navicularis* (Table

1). This is justification for including them all in the *Juniperus oxycedrus* group.

Moreover, there are certain morphological and ecological similarities between *J.*

oxycedrus subsp. *oxycedrus* and *J. oxycedrus* subsp. *badia* and between *J. oxycedrus*

subsp. *macrocarpa* and *J. navicularis*, according to the presence of certain oils. The

component myrcene is exclusive to subspecies *oxycedrus*, whereas the exclusive

compounds in subspecies *badia* are: germacrene D, alpha-camphelenal, beta

bourbenene and nine sesquiterpenes. The subspspecies *macrocarpa* does not have any

of its own essential oils as it shares ganma-terpinene and terpinen 4-ol with *J.*

navicularis, while the exclusive chemical compounds in *J. navicularis* are alpha-

phellandrene, alpha-terpinene, terpinolene, alpha terpinol and ten sesquiterpenes.

The Pearson correlation analysis (Tables 2 and 3, Fig 3) gives values of r = 1 or near 1

for group G2 (*J. oxycedrus* subsp. *macrocarpa* and *J. navicularis*). In the case of group

G1 (*J. oxycedrus* subsp. *oxycedrus* and *J. oxycedrus* subsp. *badia*) the values of rare

close or equal to 1; however the relation between both groups is low owing to the

phytochemical differences between them.

Table 2. Essential oil content in four species of *Juniperus* from [10, 11].

	J. oxycedrus subsp. *oxycedrus*	*J. oxycedrus* subsp. *badia*	*J. oxycedrus* subsp. *macrocarpa*	*J. navicularis*
alpha-pinene	34	0.1	0.1	30.8
limonene	16.25	0	0	0
beta-pinene	0.1	0	0	3.3
myrcene	0.1	0	0	0

p-cymene	0.1	0	0.1	3
beta-phellandrene	0.1	0	0	0
manoyl oxide	0.1	0.1	0	0
germacrene D	0	0.1	0	0
alpha-campholenal	0	0.1	0	0
beta-bourbenene	0	0.1	0	0
Sesquiterpenes /only for 1 species	0	0.1	0	0
ganma-terpinene	0	0	0.1	0.5
terpinen 4-ol	0	0	0.1	0.9
alpha-phellandrene	0	0	0	11.1
limonene + beta-phellandrene	0	0	0	27.2
alpha-terpinene	0	0	0	0.5
terpinolene	0	0	0	3.4
alpha-terpinol	0	0	0	1.1
Sesquiterpenes /only for 1 species	0	0	0	0.1

234 **Table 3. r- Pearson.** The values in bold are different from 0 with a level of significance of alpha=0.05.
235

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19
1	1	0,633	0,543	0,633	0,532	0,633	0,049	-0,576	-0,576	-0,576	-0,576	0,424	0,470	0,519	0,519	0,519	0,519	0,519	0,519
2	0,633	1	-0,306	**1,000**	-0,318	**1,000**	0,577	-0,333	-0,333	-0,333	-0,333	-0,420	-0,382	-0,333	-0,333	-0,333	-0,333	-0,333	-0,333
3	0,543	-0,306	1	-0,306	**1,000**	-0,306	-0,565	-0,347	-0,347	-0,347	-0,347	**0,977**	**0,992**	**1,000**	**1,000**	**1,000**	**1,000**	**1,000**	**1,000**
4	0,633	**1,000**	-0,306	1	-0,318	**1,000**	0,577	-0,333	-0,333	-0,333	-0,333	-0,420	-0,382	-0,333	-0,333	-0,333	-0,333	-0,333	-0,333
5	0,532	-0,318	**1,000**	-0,318	1	-0,318	-0,590	-0,363	-0,363	-0,363	-0,363	**0,983**	**0,995**	**0,999**	**0,999**	**0,999**	**0,999**	**0,999**	**0,999**
6	0,633	**1,000**	-0,306	**1,000**	-0,318	1	0,577	-0,333	-0,333	-0,333	-0,333	-0,420	-0,382	-0,333	-0,333	-0,333	-0,333	-0,333	-0,333
7	0,049	0,577	-0,565	0,577	-0,590	0,577	1	0,577	0,577	0,577	0,577	-0,728	-0,662	-0,577	-0,577	-0,577	-0,577	-0,577	-0,577
8	-0,576	-0,333	-0,347	-0,333	-0,363	-0,333	0,577	1	**1,000**	**1,000**	**1,000**	-0,420	-0,382	-0,333	-0,333	-0,333	-0,333	-0,333	-0,333
9	-0,576	-0,333	-0,347	-0,333	-0,363	-0,333	0,577	**1,000**	1	**1,000**	**1,000**	-0,420	-0,382	-0,333	-0,333	-0,333	-0,333	-0,333	-0,333
10	-0,576	-0,333	-0,347	-0,333	-0,363	-0,333	0,577	**1,000**	**1,000**	1	**1,000**	-0,420	-0,382	-0,333	-0,333	-0,333	-0,333	-0,333	-0,333
11	-0,576	-0,333	-0,347	-0,333	-0,363	-0,333	0,577	**1,000**	**1,000**	**1,000**	1	-0,420	-0,382	-0,333	-0,333	-0,333	-0,333	-0,333	-0,333
12	0,424	-0,420	**0,977**	-0,420	**0,983**	-0,420	-0,728	-0,420	-0,420	-0,420	-0,420	1	**0,996**	**0,980**	**0,980**	**0,980**	**0,980**	**0,980**	**0,980**
13	0,470	-0,382	**0,992**	-0,382	**0,995**	-0,382	-0,662	-0,382	-0,382	-0,382	-0,382	**0,996**	1	**0,994**	**0,994**	**0,994**	**0,994**	**0,994**	**0,994**
14	0,519	-0,333	**1,000**	-0,333	**0,999**	-0,333	-0,577	-0,333	-0,333	-0,333	-0,333	**0,980**	**0,994**	1	**1,000**	**1,000**	**1,000**	**1,000**	**1,000**
15	0,519	-0,333	**1,000**	-0,333	**0,999**	-0,333	-0,577	-0,333	-0,333	-0,333	-0,333	**0,980**	**0,994**	**1,000**	1	**1,000**	**1,000**	**1,000**	**1,000**
16	0,519	-0,333	**1,000**	-0,333	**0,999**	-0,333	-0,577	-0,333	-0,333	-0,333	-0,333	**0,980**	**0,994**	**1,000**	**1,000**	1	**1,000**	**1,000**	**1,000**
17	0,519	-0,333	**1,000**	-0,333	**0,999**	-0,333	-0,577	-0,333	-0,333	-0,333	-0,333	**0,980**	**0,994**	**1,000**	**1,000**	**1,000**	1	**1,000**	**1,000**
18	0,519	-0,333	**1,000**	-0,333	**0,999**	-0,333	-0,577	-0,333	-0,333	-0,333	-0,333	**0,980**	**0,994**	**1,000**	**1,000**	**1,000**	**1,000**	1	**1,000**
19	0,519	-0,333	**1,000**	-0,333	**0,999**	-0,333	-0,577	-0,333	-0,333	-0,333	-0,333	**0,980**	**0,994**	**1,000**	**1,000**	**1,000**	**1,000**	**1,000**	1

Types of oils studied: 1. Alpha-pinene, 2. Limonene, 3. Beta-pinene, 4. Myrcene, 5. P-cymene, 6. Beta-phellandrene, 7. Monoyl oxide, 8. Germacrene D, 9. Alpha-campholenal, 10. Beta-bourbenene, 11. Sesquiterpenes exclusives, 12. Gamma-terpinene, 13. Terpinen 4-ol, 14 Alpha-phellandrene, 15. Limonene + beta-phellandrene, 16. Alpha-terpinene, 17. Terpinolene, 18. Alpha-terpinol, 19. Sesquiterpenes.

236
237 **Fig 3. Similarity analysis between the taxa *Juniperus oxycedrus* subsp. *oxycedrus*,**
238 **subsp. *badia*, *J. oxycedrus* subsp. *macrocarpa* and *J. navicularis*.**
239

Discussion

Linnaeus in the first part of his work *Speics Plantarum* did not describe *Juniperus*

oxycedrus L., where he merely noted the existing synonymy. It was Clusius who later

described this species and gave specific localities for Spain, which were subsequently

included by [32].

The main differences between *Juniperus oxycedrus* subsp. *oxycedrus* and *Juniperus*

oxycedrus subsp. *badia* are based on their physiognomy and the size of their mature

fruits [1]. Whereas the subspecies *oxycedrus* tends to occur as a shrub, the subspecies

badia (H. Gay) Debeaux is a tree of considerable height and with a pyramidal form.

The size of the mature galbuli in the subspecies *oxycedrus* does not generally exceed 1

cm, while in the subspecies *badia* it is over 1 cm. The leaves of *oxycedrus* have a width

of 1-1.5 mm while those of subsp. *badia* are 1.2-2 mm. Coincidentally these subspecies

are frequently found coexisting in similar biotopes, which has led to frequent confusion

by several authors. The subsp. *macrocarpa* (Sm.) Ball. has galbuli of 1.2-1.5 cm. with a

purplish chestnut colour when ripe; this is an erect tree, rarely prostrate, growing up to

3 m. In contrast *Juniperus navicularis* Gand. has a similar ecology to subsp.

macrocarpa but grows to a maximum height of 2 m; its leaves are 1-1.5 mm wide, and

its galbuli are between 0.7-1 cm (Table 4).

258
259 **Table 4. Morphometric comparison in the taxa of *Juniperus oxycedrus* group from [1].**

	J. oxycedrus subsp. *oxycedrus*	*J. oxycedrus* subsp. *badia*	*J. oxycedrus* subsp. *macrocarpa*	*J. navicularis*
leaf length (mm)	8-15(25) mm	12-20 mm	20-25 mm	4-12 mm
leaf width (mm)	1-1.5 mm	1.2-2 mm	2-2.5 mm	1-1.5 mm
fruit (gabuli)	8-10 mm	10-13 mm	12-15 mm	7-10 mm

tree habit	scrub < 4 m	tree < 15 m	scrub < 3 m	scrub < 2 m

Theoretically, the authorship of *Juniperus oxycedrus* L. should be attributed to Clusius,

who described it: *"Nusquam autem majorem vidisse memini, quam supra Segoviam et*

Guadarrama, itinere Madritiano, ubi magnarum arborum interdum altitudinem". He

also says of the fruit *"fructum initio viridem... postremo, cum maturuit punicea colorís,*

Juniperi fructu multo majorem ut interdum avellanam magnitudine aequet" [33]. This

description is specifically included in the work of [19]. Clusius' descrtiption, in which

he states literally that it has a "thick pruinose red fruit", corresponds to *Juniperus*

oxycedrus L. subsp. *badia* (H. Gay) Debeaux [Basión. *Juniperus oxycedrus* var. *badia*

H. Gay in Assoc. Franç. Avancem. Sci. Compt. Rend. 1889:501 (1889)], included in [1].

The lack of typification of the specimen described by Clusius is justification for these

facts, and supports the valid acceptance of *Juniperus oxycedrus* L. subsp. *badia* (H.

Gay) Debeaux (ICN article 41). Due to the high number of morphological and

ecological differences and in essential oils between subsp. *macrocarpa*, subsp.

oxycedrus and subsp. *badia,* and to its similarity with *J. navicularis*, the subspecies

macrocarpa can be proposed to be subordinated to *J. navicularis* as *J. navicularis* Gand.

subsp. *macrocarpa* (Sm.) Ball; however, the molecular study by [34] establishes major

molecular differences between the four taxa *J. oxycedrus* subsp. *oxycedrus, J. oxycedrus*

subsp. *badia, J. oxycedrus* subsp. *macrocarpa* and *J. navicularis*, and based on this he

raises them to the rank of species. The phytochemical differences between groups G1

and G2 are very clear, so the species rank could be accepted for the four taxa. However,

in the case of their distribution, the populations of *J. oxycedrus* subsp. *macrocarpa* and

J. navicularis are separated despite having similar ecologies, while this is not the case

of *J. oxycedrus* subsp. *oxycedrus* and subsp. *badia.* Therefore, it is logical to accept *J.*

macrocarpa Sm. and *J. navicularis* Gand. as species and to maintain *J. oxycedrus* L.

subsp. *oxycedrus* and *J. oxycedrus* L. subsp. *badia* (H. Gay) Debeaux with the rank of

subspecies.

Finally, considering the phytosociological point of view, in vegetation studies the

subspecies *badia* (H. Gay) Debeaux and the subspecies *lagunae* (Pau ex Vicioso) Rivas-

Martínez have been used indistinctly as a result of accumulated and persistent errors

[30]; the name *Juniperus oxycedrus* L. subsp. *lagunae* (Pau ex Vicioso) Rivas-Martínez

is invalid and based on the iconography of Laguna by Pau [*Juniperus oxycedrus* L.

subsp. *lagunae* (Pau ex C.Vicioso) Rivas Mart. - Itinera Geobot. 15(2): 703 (2002);

nom. inval.], as the name of the basionym is not validly published (ICN articulo 7). The

correct name is therefore *Juniperus oxycedrus* L. subsp. *badia* (H. Gay) Debeaux. Based

on these facts, the various authors of plant associations must rectify their names.

297   Conclusions

In this work, we update the state of knowledge on the ecology, taxonomy and

distribution of the taxa of the *Juniperus oxycedrus* group, based on our study and on

research by various authors, which has occasionally been the source of controversy. The

similarity study based on the quantity, type of essential oils, molecular analyses and

different bioclimatic distribution justify maintaining the rank of species for *J.*

macrocarpa Sm. and *J. navicularis* Gand.. Instead, the distribution and co-existence of

the two remaining taxa, in spite of their phytochemical and molecular differences, do

not warrant the rank of species, and we thus maintain their rank as subspecies: *J.*

oxycedrus L. subsp. *oxycedrus* and *J. oxycedrus* L. subsp. *badia* (H. Gay) Debeaux.

This study, in accordance to others [6, 35, 36, e.g.] also confirm that for a good

discrimination among species of groups with difficult interpretation, morphometric,

phytochemical and bioclimatic approaches are very useful to clarify the belonging rank

to the taxa. All these results and conclusions are according to [37].

Acknowledgments

313   Ms Pru Brooke-Turner (M.A. Cantab.), a native English speaker specialized in scientific

314   texts, has translated this article. This manuscript has been released as a Pre-Print at:

315   Biorxiv 459651; https://doi.org/10.1101/459651 2018.

Author Contributions

Conceptualization: Eusebio Cano.

Data curation: Eusebio Cano; Ana Cano Ortiz; Carmelo M. Musarella; José C. Piñar

Fuentes; Carlos J. Pinto Gomes; Giovanni Spampinato.

Formal analysis: Ana Cano Ortiz; Carmelo M. Musarella; José C. Piñar Fuentes.

Investigation: Eusebio Cano; Ana Cano Ortiz; Carmelo M. Musarella; José C. Piñar

Fuentes; Carlos J. Pinto Gomes; Giovanni Spampinato.

Methodology: Eusebio Cano; Ana Cano Ortiz.

Resources: Eusebio Cano; Ana Cano Ortiz; Carmelo M. Musarella; Carlos J. Pinto

Gomes; Giovanni Spampinato.

Supervision: Eusebio Cano; Carmelo M. Musarella.

Validation: Eusebio Cano.

Visualization: Eusebio Cano; Ana Cano Ortiz.

Writing ± original draft: Eusebio Cano; Ana Cano Ortiz; Carmelo M. Musarella; José

C. Piñar Fuentes; Carlos J. Pinto Gomes; Giovanni Spampinato.

Writing ± review & editing: Eusebio Cano; Ana Cano Ortiz; Carmelo M. Musarella.

References

1. Amaral Franco J. *Juniperus*. In: Flora Ibérica I. Madrid: Real Jardin Botánico; 1986. pp.181-188.

2. Pignatti S. *Juniperus*. Flora d'Italia 1. Bologna: Edagricole; 1982.

3. Brus R, Idzjtic´ M, Jarni K. Morphologic variation in northern marginal *Juniperus oxycedrus* L. subsp. *oxycedus* population in Istria. Plant Biosyst. 2016; 150(2): 274-284. https://doi.org/10.1080/11263504.2014.984790

4. Adams RP. *Juniperus deltoides*, a new species, and nomenclatural notes on *Juniperus polycarpos* and *Juniperus turcomanica* (Cupressaceae). Phytologia 2004; 86(2): 49-53.

5. Adams RP. Morphological comparison and key to *Juniperus deltoides* and *J. oxycedrus*. Phytologia. 2014; 96(2): 58-62.

6. Roma-Marzio F, Najar B, Alessandri J, Pistelli L, Peruzzi L. Taxonomy of prickly juniper (*Juniperus oxycedrus* group): a phytochemical-morphometric combined approach at the contact zone of two cryptospecies. Phytochemistry. 2017; 141:48-60. doi: 10.1016/j.phytochem.2017.05.008.

7. Cano E, Melendo M, Valle F. The plant communities of the *Asplenietea trichomanis* in the SW Iberian Península. Folia Geobot. 1997; 32: 361-376.

8. Cano Ortiz A, Piñar JC, Pinto Gomes CJ, Musarella CM, Cano E. Expansion of the *Juniperus* genus due to anthropic activity. In: Weber RP, editor. Old-Growth Forest and Coniferous Forests. Nova Science Publishers; 2015. pp. 55-65.

9. Musarella CM, Cano-Ortiz A, Piñar Fuentes JC, del Río S, Pinto Gomes CJ, Spampinato G, et al. Analysis vegetation of *Juniperus oxycedrus* L. subsp. *badia* (H. Gay) Debeaux microforests in the Iberian Peninsula: increasing their

knowledge for a better restoration. Fl. Medit. 2017; 27: 54-55. doi:
10.7320/FlMedit27.005.

10. Adams RP, Joaquin Altarejos J, Fernandez C, Camacho A. The leaf essential oils
and taxonomy of *Juniperus oxycedrus* L. subsp. *oxycedrus,* subsp. *badia* (H.
Gay) Debeaux, and subsp. *macrocarpa* (Sibth. & Sm.) Ball. J. Essent. Oil Res.
1999; 11(2): 167-172.

11. Velasco Negueruela A, Pérez Alonso MJ, Palá-Paúl J, Íñigo A, López G. Leaf
essential oils analysis of *Juniperus navicularis* Gandoger. Bot Complut. 2002;
26:85-91.

12. Salido S, Altarejos J, Nogueras M, Sánchez A, Pannecouque C,Witvrouw M, De
Clercq E. Chemical studies of essential oils of *Juniperus oxycedrus* ssp. *badia.* J
Ethnopharmacol. 2002; 81(1):129-34.

13. Montero Burgos JL, González Rebollar JL. Diagramas bioclimáticos. Madrid.
ICONA, 1983.

14. Cano E, Musarella CM, Cano-Ortiz A, Piñar Fuentes JC, Rodríguez Torres A,
Del Rio González S, Pinto Gomes C, Quinto-Canas R, Spampinato G.
Geobotanical Study of the Microforests of *Juniperus oxycedrus* subsp. *badia* in
the Central and Southern Iberian Peninsula. Preprints 2018; doi:
10.20944/preprints201810.0362.v1.

15. Rivas-Martínez S, Loidi J. 1999. Bioclimatology of the Iberian Peninsula. Itinera
Geobot. 1999; 13: 41-47.

16. Rivas-Martínez S et al. Mapa se series, geoseries y geopermaseries de
vegetación de España. Parte II. Itinera Geobot. 2011; 18 (1): 5-424.

17. Castroviejo, S. (coord. gen.). 1986-2012. Flora iberica 1-8, 10-15, 17-18, 21.
Real Jardín Botánico, CSIC, Madrid.

18. Tutin T, Heywood VH, Burges DA, Valentine DH, Walter SM, Webb DA. Flora

Europaea. Vol I al V. Cambrige at the University Pres. (eds.). 1964-1980.

19. Valdés B, Talavera S, Fernández Galiano E. Flora Vascular de Andalucía

Occidental. Vol. 1, 2 y 3. Barcelona. Ketres Editora; 1987.

20. Vicioso C. Notas sobre la flora española. Madrid. Anales Jar. Bot. Madrid. 1946;

6(2): 5-89.

21. Laguna M. Flora Forestal Española de los Árboles, Arbustos y Matas. Parte

primera. 1883; 1-374. http://bdh-rd.bne.es/viewer.vm?id=0000042028&page=1

22. Laguna M. Flora Forestal Española de los Árboles, Arbustos y Matas. Parte

segunda. 1890; 375-839. http://bdh-

rd.bne.es/viewer.vm?id=0000042028&page=1

23. Fiori A. Nuova Flora Analitica d´Italia. New York: Library New York Botanical

Garden, 1923-1925; I: 944.

24. Blanca G, Cabezudo B, Cueto M, Fernández López C, Morales Torres C. Flora

Vascular de Andalucía Oriental. Vol. I: *Selaginellaceae-Ceratophyllaceae*.

Sevilla: Consejería de Medio Ambiente Junta de Andalucía (eds); 2009.

25. Minissale P, Sciandrello S. A relic wood of *Juniperus turbinata* Guss.

(Cupressaceae) in Sicily: Structural and ecological features, conservation

perspectives. Plant Biosyst. 2013; 147(1): 145-157.

https://doi.org/10.1080/11263504.2012.743931

26. Valdés B, Rejdali M, Achhal El Kadmiri A, Jury JL, Monserrat JM. Checklist of

vascular plants of N. Morocco with identification keys, Vol I y II. Madrid. CSIC;

2002.

27. Maire R. 1952 Flore de l´Afrique du Nort. Paris. Ed. Paul Lechevalier, 1952. 1-

266 .

28. Bolòs O, Vigo J. Flora dels Països Catalans. Barcelona. ed. Barcino; 1984.

29. Rivas-Martínez S, Díaz TE, Fernández González F, Izco J, Loidi J, Lousa M, et
al. Vascular plant communities of Spain and Portugal. Itinera Geobot. 2002;
15(2): 433-922.

30. Cano E, Rodríguez Torres A, Pinto Gomes CJ, García Fuentes A, Torres JA,
Salazar C, et al. Analysis of the *Juniperus oxycedrus* L. in the centre and south
of the Iberian Peninsula (Spain and Portugal). Acta Bot. Gallica. 2007; 154 (1):
79-99.

31. Del Río S, Cano-Ortiz A, Herrero L, Penas A. Recent trends in mean maximum
and minimum air temperatures over Spain (1961–2006). Theor Appl Climatol.
2012; 109: 605–626 DOI 10.1007/s00704-012-0593-2

32. Linnaeus C. Species plantarum, exhibentes plantas rite cognitas, ad genera
relatas, cum differentiis specificis, nominibus trivialibus, synonymis selectis,
locis natalibus, secundum systema sexuale digestas. Holmiae, Impensis
Laurentii Salvii. [L. Salvius, Stockholm.], 1 May 1753.

33. Clusio C. Rariorum aliquot stirpium per Hispanias observatarum historia.
Antwerpen: Ex officina Christophori Plantini. Biblioteca del Real Jardín
Botánico de Madrid, 1576.

34. Adams RP. Systematics of *Juniperus* section *Juniperus* based on leaf essential
oils and random amplified polymorphic DNAs (RAPDs). Biochem Syst Ecol.
2000; 28(6): 515-528. https://doi.org/10.1016/S0305-1978(99)00089-7

35. Cano E, Musarella CM, Cano-Ortiz A, Piñar Fuentes JC, Spampinato G, Pinto
Gomes CJ. Morphometric analysis and bioclimatic distribution of *Glebionis*
coronaria s.l. (Asteraceae) in the Mediterranean area. PhytoKeys 2017; 81: 103-
126. https://doi.org/10.3897/phytokeys.81.11995

36. Panuccio MR, Fazio A, Musarella CM, Mendoza-Fernández AJ, Mota JF,

Spampinato G. Seed germination and antioxidant pattern in *Lavandula multifida*

(Lamiaceae): A   comparison between core and peripheral populations.

Plant   Biosyst.   2018;   152(3):   398-406.

https://doi.org/10.1080/11263504.2017.1297333

37. Cano Ortiz A, Musarella CM, Piñar Fuentes JC, Pinto Gomes CJ, Spampinato G,

Cano E. Taxonomy, ecology and distribution of *Juniperus oxycedrus* L. group in

the Mediterranean Region using morphometric, phytochemical and bioclimatic

approaches. bioRxiv 459651; doi: https://doi.org/10.1101/459651.

Map Legend

Ombrothermic index

- Arid - semirarid
- Lower dry
- Upper dry
- Lower subhumid
- Upper subhumid
- Humid - hyperhumid

| 0 | 375 | 750 | 1.500 | 2.250 | 3.000 |

Kilómetros