Characterization of Pacific oyster (*Crassostrea gigas*) proteomic response to
natural environmental differences

Running head: Oyster proteomic response to environment

Yaamini R. Venkataraman[1], Emma Timmins-Schiffman[2], Micah J. Horwith[3], Alexander T. Lowe[4],
Brook Nunn[2], Brent Vadopalas[1], Laura H. Spencer[1], Steven B. Roberts*[1]
[1]School of Aquatic and Fishery Sciences, University of Washington, Seattle, Washington 98105,
United States
[2]Department of Genome Sciences, University of Washington, Seattle, Washington 98105,
United States
[3]Washington Department of Natural Resources, Olympia, Washington 98504, United States
[4]Department of Biology, University of Washington, Seattle, Washington 98105, United States

Abstract

Global climate change is rapidly altering coastal marine ecosystems important for food
production. A comprehensive understanding of how organisms will respond to these
complex environmental changes can come only from observing and studying species
within their natural environment. To this end, the effects of environmental drivers — pH,
dissolved oxygen content, salinity, and temperature — on *Crassostrea gigas* physiology
were evaluated in an outplant experiment. Sibling juvenile oysters were outplanted to
eelgrass and unvegetated habitat at five different estuarine sites within the Acidification
Nearshore Monitoring Network in Washington State, USA to evaluate how regional
environmental drivers influenced molecular physiology. We tested effects environmental
conditions at outplant sites and habitat to determine if macrophyte presence and diurnal
cycling buffered pH conditions and changed the oysters' expressed proteome. A novel,
two-step, gel-free proteomic approach was used to identify differences in protein
abundance in *C. gigas* ctenidia tissue after a 29 day outplant by 1) identifying proteins in
a data independent acquisition survey step and 2) comparing relative quantities of
targeted environmental response proteins using selected reaction monitoring. While
there was no difference in protein abundance detected between habitats or within Puget
Sound, *C. gigas* outplanted at Willapa Bay had significantly higher abundances of
antioxidant enzymes and carbohydrate metabolism proteins. Environmental factors at
Willapa Bay, such as higher average temperature, could have driven this protein
abundance pattern. These findings generate a suite of new hypotheses for lab and field
experiments to compare the effects of regional conditions on physiological responses of
marine invertebrates.

Keywords: proteomics, oysters, in situ measurements, estuarine systems, antioxidant
enzymes, carbohydrate metabolism

Introduction

Global climate change will influence estuarine dynamics and impact the organisms that inhabit these environments. Estuaries are already variable across spatial and temporal scales in terms of phytoplankton production (Pennock and Sharp 1986), nutrient availability, (Paerl et al. 2014), heavy metal contamination (Liu et al. 2015), salinity (Banas et al. 2004), and carbonate chemistry (Pelletier et al. 2018; Feely et al. 2010; Ruesink et al. 2015; Baumann and Smith 2018). Since climate change will affect these parameters, it is important to consider how estuarine organisms will respond.

Proteomics, or the study of protein abundance and expression, can be used to shed light on physiological changes on a molecular level. Proteins direct all major cellular functions, thus examining protein abundance provides direct evidence of an organism's physiological response to the estuarine environment (Tomanek 2014). The proteome is dynamic, as it must rapidly respond to perturbation, providing mechanistic information that standard gene expression and mRNA quantification studies cannot (Veldhoen, Ikonomou, and Helbing 2012; Flores-Nunes, Gomes, et al. 2015). As a result of the proteome's dynamic nature, proteins analyzed at the time of collection represent an organism's response to the environment in near real-time. Discovery-based proteomic methods can unbiasedly elucidate responses to environmental drivers (Flores-Nunes, Gomes, et al. 2015). Several studies have connected protein abundances with changes in laboratory-simulated environmental conditions, identifying key proteins and mechanisms involved in specific environmental responses (Timmins-Schiffman et al. 2014; Tomanek 2014; Papakostas et al. 2012; Meng et al. 2017; Parker et al. 2011; Dineshram et al. 2016). While these studies provide insight into organismal

adaptation and physiology, laboratory studies alone cannot fully encapsulate the effects

of multiple environmental drivers within an ecosystem context (Riebesell and Gattuso

2014).

Although challenging, *in situ* field studies provide a necessary biological realism

when considering variable environments (Cornwall and Hurd 2016; Slattery et al. 2012).

Such experiments can be leveraged to study the effects of multiple environmental

drivers on organismal physiology and to incorporate realistic variability, as opposed to

examining the effect of a single stressor on an organism (Riebesell and Gattuso 2014).

Through transcriptomics Chapman et al. (2011) demonstrated the power of an *in situ*

experimental design for examining the impacts of regional environmental conditions on

Eastern oyster (*Crassostrea virginica*) physiology. Transcript signatures from *C.*

virginica sampled from various locations in southeastern United States revealed that

temperature, pH, salinity, dissolved oxygen and pollutant load at each location impacted

gene expression. Furthermore, they were able to disentangle the interactions of these

environmental factors on gene expression. RNA and protein abundances can be

influenced by several environmental factors, and *in situ* studies can determine which

drivers will be more important to consider for organismal physiology.

Marine invertebrates have proven to be informative bioindicators in proteomic

studies to examine the effects of *in situ* conditions on organismal physiological

responses to environmental change. When marine invertebrates have been exposed to

varying environmental conditions, proteomics has demonstrated changes in cellular

defense, immune responses, and genome function (Veldhoen, Ikonomou, and Helbing

2012). Changes in protein abundance in bivalves like blue mussels (*Mytilus edulis spp.*)

and the Pacific oyster (*Crassostrea gigas*) have been used to develop biomarkers for

environmental contaminants (Beyer et al. 2017; Slattery et al. 2012). Proteomic

responses to natural environmental drivers have also been evaluated in bivalves. For

example, shotgun proteomic analysis of *M. edulis* ctenidia from Baltic Sea microcosms

revealed that low salinity conditions impact abundance of cytoskeleton proteins and

signalling and intracellular membrane trafficking pathways (Campos et al. 2016). With a

growing wealth of genomic information for these species, and the importance of this

information for commercial aquaculture, understanding how these species fare under

different environmental conditions is critical.

Pacific oyster (*Crassostrea gigas*) rearing in estuarine environments in

Washington State, USA provides a perfect system to examine the effect of *in situ*

environmental conditions on the expressed proteome. *C. gigas* are extensively farmed

in two different estuarine systems that show substantial regional variation: Puget Sound

and Willapa Bay. Puget Sound is a complex estuarine system with interconnected sub-

basins, each with different freshwater inputs, residence times, and stratification levels

(Feely et al. 2010; Pelletier et al. 2018; Bianucci et al. 2018). Willapa Bay is a large

shallow estuary on the Pacific coast that exchanges approximately half its water volume

with the Pacific Ocean at each tide (Banas et al. 2004, 2007). Seasonality and location

within Puget Sound dictates temperature, dissolved oxygen, salinity, and pH conditions,

while Willapa Bay conditions are influenced by diurnal fluctuations and proximity to

either the ocean or rivers draining into the bay (Pelletier et al. 2018; Feely et al. 2010;

Banas et al. 2007; Ruesink et al. 2015). Both Puget Sound and Willapa Bay also host

eelgrass beds (*Zostera spp.*) that affect environmental conditions, such as oxygen

concentrations, on diurnal time scales and may provide refuge from ocean acidification

or rapid change in temperature through photosynthetic activity. Understanding how

different aquaculture grow-out locations and habitats will affect the oyster's ability to

persist through environmental change is crucial for the industry and the ecosystem.

The purpose of this study was to use proteomic techniques to uncover the

impacts of environmental drivers on Pacific oyster physiological outcomes in estuarine

environments in Washington State, USA. Naturally existing environmental variation was

harnessed by outplanting *C. gigas* in different locations within Puget Sound and Willapa

Bay, and habitat effects were taken into consideration by placing oysters in eelgrass

and unvegetated habitats. Gel-free proteomic methods were used to examine the

effects of outplant conditions on relative quantities of all expressed proteins in a series

of *in situ* experiments in order to identify differentially abundant proteins. We predicted

that differences in environmental drivers at each outplant location and within outplant

habitats would yield unique protein abundance patterns. Oysters at outplant locations

with higher temperatures, lower dissolved oxygen content, lower salinity, or lower pH

may have higher abundances of proteins related to environmental response. Eelgrass

beds were expected to ameliorate stressful conditions, resulting in lower abundances of

environmental stress response proteins than oysters in unvegetated habitats.

Methods

Shellfish deployment

Sibling *C. gigas* (average shell length = 27.2 cm) were outplanted for 29 days

starting June 19, 2016 at five locations: Case Inlet (CI), Fidalgo Bay (FB), Port Gamble

Bay (PG), Skokomish River Delta (SK), and Willapa Bay (WB) in Washington State,

USA (Table 1; Figure S1). These sites were selected for differences in environmental

parameters, as well as for the presence of unvegetated areas and eelgrass beds within

each site. All sites were part of the Acidification Nearshore Monitoring Network, a

network of sensors placed in various Washington locations to monitor marine chemistry

(ANeMoNe; Washington Department of Natural Resources). At each site, fifteen

juvenile oysters were placed in bags directly onto the substrate at a tidal height of -1.5

MLLW, both inside and outside of eelgrass beds (n=15 per habitat type), for a total of

thirty outplanted oysters per site. Oysters were housed in exclusion cages to prevent

predation. Custom-built Durafet-based sensors (Honeywell) were used to monitor pH.

Commercially-available MiniDOT loggers (Precision Measurement Engineering) were

used to measure dissolved oxygen, and Odyssey loggers were used to measure

conductivity. All sensors recorded temperature measurements, and all sensors logged

at 10-minute intervals across the outplant period, with the exception of SK, where

sensors were installed 21 days into the outplant period.

Juvenile oysters remained at each site for a 29-day exposure period. Oyster shell

length was measured before and after the outplant period. Because the ctenidia is the

primary site where oysters interact with the environment, ctenidia samples were

dissected at the end of the outplant and held on dry ice until storage at -80ºC (Beyer et

al. 2017; Meng et al. 2017; Meistertzheim et al. 2007).

Environmental data was treated as follows. Conductivity observations were

removed when less than 0, which occurs when the instrument is dry. Remaining

observations were converted to salinity measurements using the *swSCTp* function in

the *oce* package in R (Kelley and Richards 2018; R Core Team 2018), with temperature

at 25ºC and pressure at 10 dbar. For dissolved oxygen, pH, and salinity datasets, data

were removed when collected by probes 1) during low tide or 2) when tidal depth was

less than one foot to remove readings where the probes may have been exposed.

These values were retained for temperature datasets. Outliers were screened using the

Tukey method for temperature, dissolved oxygen, pH, and salinity datasets (Hoaglin,

Iglewicz, and Tukey 1986). Uniform outplant tidal heights were checked using *prop.test*

in R (R Core Team 2018). R Scripts are available in the supplementary Github

repository (S2).

Protein Discovery

To identify select protein targets for characterization across locations and

environmental conditions, a subset of tissue samples (2 from each site) were analyzed

with data independent acquisition (DIA) mass spectrometry analysis (Table S3).

Protein Quantification

Tissue samples were homogenized in a solution of 50 mM NH_4HCO_3 with 6M

Urea (500µl). Tissues were then sonicated 3 times (Fisher Scientific Sonic

Dismembrator Model 100) for 10 seconds each and cooled between sonications in a

bath of ethanol and dry ice. Protein quantities were measured with the ThermoScientific

BCA Protein Assay Kit microplate assay with a limited quantity of sonicated sample (11

µL). The protein concentration was measured via spectroscopy at 540 nm in a

Labsystems (Waltham, MA) Multiskan MCC/340 and accompanying Ascent Software

Version 2.6. Protein concentrations were calculated based on a standard curve with

BSA (Pierce) per manufacturer's instructions.

Protein Digestion

Protein digestion followed the protocol outlined in Timmins-Schiffman et al.

(2013). To each sample of 30 µg protein,1.5 M Tris pH 8.8 buffer (6.6 µL) and 200 mM

TCEP (2.5 µL) were added. After solvent additions, each sample's pH was verified to be

basic (pH ≥ 8), and placed on a 37ºC heating block for one hour. Iodoacetamide (200

mM, 20 µL) was then added to each sample to maximize digestion enzyme access to

protein cleavage sites. Samples were covered with aluminum foil to incubate in the dark

for 1 hour at room temperature. Afterwards, diothiolthreitol (200 mM, 20 µL) was added

and samples were incubated at room temperature for one hour. Lysyl Endopeptidase

(Wako Chemicals) was then added to each sample in a 1 µg enzyme:30 µg oyster

protein ratio, followed by one hour of incubation at room temperature. Urea was diluted

with NH_4HCO_3 (25 mM, 800 µL) and HPLC grade methanol (200 µL). Trypsin

(Promega) was added to each sample in a 1 µg trypsin: 30 µg oyster protein ratio for

overnight digestion at room temperature.

Peptide Isolation

After overnight incubation, samples were evaporated to near dryness at 4ºC with

a speedvac (CentriVap ® Refrigerated Centrifugal Concentrator Model 7310021).

Samples were then reconstituted in 100 µL of a 5% Acetonitrile and 0.1%

Trifluoroacetic Acid (Solvent B) to isolate peptides. If samples were not at pH ≤ 2, 10-20

µL aliquots of 10% Formic Acid were added until this pH was achieved.

Before desalting peptide samples, Macrospin C18 columns (The Nest Group)

were prepared by adding 200 µL of a 60% Acetonitrile with 0.1% Trifluoroacetic Acid

(Solvent A). The columns were spun for three minutes at 2000 rpm, and flow-through

liquid from the column was discarded. The spinning and discarding process was

completed a total of four times.To wash columns, 200 µL of Solvent B was added to

each column. The columns were once again spun for three minutes at 2000 rpm and

liquid was discarded afterwards; the solvent addition, spinning, and discarding process

was completed a total of three times.

To bind peptides to the columns, digested peptides were added to prepared

columns, then the columns were spun at 3000 rpm for three minutes. The filtrate was

pipetted back onto the column and spun again at 3000 rpm for three minutes. Solvent B

(200 µL) was added to each column three separate times, then the column was spun for

three minutes at 3000 rpm to wash salts off the column.

Peptides were eluted with two additions of 100 µL Solvent A to each column.

Columns were spun at 3000 rpm for three minutes and the peptide fraction (filtrate) was

reserved. Samples were placed in a speed vacuum at 4ºC until they were nearly dry

(approximately two hours) to dry peptides. Peptides were reconstituted with 60 µL of 3%

Acetonitrile + 0.1% Formic Acid, and stored at -80ºC.

Peptide Retention Time Calibration (PRTC; Pierce) is a mix of known peptides in

known quantities used as an internal standard to ensure consistency of peptides

detected and measured throughout a mass spectrometry run. The stock solution of

PRTC was diluted to 0.2 pmol/µL PRTC using 3% Acetonitrile with 0.1% Formic Acid. In

a clean centrifuge tube, 6 µg of oyster protein and 0.376 pmol of PRTC were mixed

together. Sample volume was brought up to 15 µL using a 3% acetonitrile and 0.1%

formic acid solution. A quality control solution was also prepared (1 µL PRTC + BSA:3

µL 3% Acetonitrile and 0.1% Formic Acid solution).

Data Independent Acquisition Mass Spectrometry

Peptides were analyzed on an Orbitrap Fusion Lumos mass spectrometer

(Thermo Scientific) for Data Independent Acquisition Mass Spectrometry (DIA). DIA

analyses were completed as a comprehensive, non-random analytical method for

detecting all peptide ions present within a sample. The detection of peptides was then

leveraged to develop a targeted proteomics assay for quantification (see Selected

Reaction Monitoring Assay). A 30 cm analytical column and 3 cm pre-column were

packed in-house with 3 µm C18 beads (Dr. Maisch). Samples were run in a randomized

order. A blank injection followed each sample, with procedural blanks run at the very

end. Every injection was 3 µL, which included 1 µg of oyster protein and 0.0752 pmol of

PRTC. Peptides were analyzed in MS1 over the *m/z* range of 450-950 with 12 *m/z* wide

windows with 5 *m/z* overlaps (Egertson et al. 2013). MS1 resolution was 60,000 and

AGC target was 400,000 with a three second cycle time. The MS2 loop count was set to

20 and MS2 data was collected with a resolution of 15,000 on charge state of 2 with an

AGC target of 50,000. No dynamic exclusion was used (Table S4).

Peptide-Centric Proteomic Analyses

Unknown peptide spectra from mass spectrometry samples were matched with

known peptides using Peptide-Centric Analysis in the PECAN software (Ting et al.

2015). Raw mass spectrometry files were converted to mzML files, then demultiplexed

using the MSConvert command line interface (S5; Chambers et al. 2012). The *C. gigas*

proteome was digested with in silico tryptic digest using Protein Digestion Simulator

(S6) . All known peptides from the mzML files were identified in an comparison to the

digested *C. gigas* proteome (S6).

The PECAN-generated spectral library (.blib) file was used to detect peptides of interest

in raw DIA files in Skyline (MacLean et al. 2010). Skyline identified peptides using

chromatogram peak picking, where ions that elute at the same time and mass are

detected as a peptide (S7). All PRTC peptides and approximately 100 different oyster

proteins and their peptide transitions were manually checked for retention time and

peak area ratio consistency to determine a Skyline auto peak picker error rate (24.3% ±

25%, range: 0% to 100%).

Proteins had to satisfy 4 criteria to be considered appropriate targets for the

study. 1) To select proteins for a targeted MS assay, protein data was first evaluated in

Skyline to ensure there was no missing data for any peptide or sample. 2) Peaks with

significant interference were also not considered. 3) Proteins needed at least two

peptides with three transitions per peptide to qualify as a potential target for

259 downstream assays. 4) Proteins with annotated functions related to oxidative stress,

260 hypoxia, heat shock, immune resistance, shell formation growth and cellular

261 maintenance were considered for targets to address hypotheses of response to in situ

262 environmental differences and eelgrass vs. bare habitat. From this list, fifteen proteins

263 (41 peptides and 123 transitions total), with various environmental response and

264 general maintenance functions, were selected as targets (Table 2).

265 *Selected Reaction Monitoring Assay*

266  Following the protein discovery phase (DIA), proteins were isolated as described

267 above from an additional five randomly selected samples per site and habitat

268 combination (for a total of 5 oysters per group) and analyzed with Selected Reaction

269 Monitoring (SRM). Samples were prepared as described for DIA, except tissue samples

270 were homogenized in 100 µL, and peptide samples were evaporated at 25ºC after

271 peptide isolation.

272  Proteins of interest identified from the DIA analysis were used as targets in a

273 SRM assay following the workflow and informatic pipeline of Timmins-Schiffman et al.

274 2016; 2017. Target peptide transitions were monitored using SRM on a Thermo TSQ

275 Vantage. SRM data were collected during a gradient of 2-60% acetonitrile over 40

276 minutes. All samples were run in technical duplicates in a randomized order with a 1 µg

277 oyster peptide and 0.0752 pmol PRTC injection. A quality control injection and blank

278 injection were run after every five sample injections, and PRTC peptides were

279 monitored throughout the experiment.

 Target Peptide Specificity

To ensure SRM assay specificity to oyster peptides of interest, oyster peptides

were diluted in a background matrix of similar complexity (Pacific geoduck — *Panopea*

generosa — peptides), then analyzed using the oyster SRM assay. An oyster-specific

SRM target would decrease in abundance with a decreasing abundance of oyster

peptides in a mixture. Non-specific peptides — more likely to be found in background

matrix of similar complexity — or peptides susceptible to interference would not

correlate with oyster peptide abundance, and therefore, would be uninformative. Five *C.*

gigas samples used for SRM were randomly selected and pooled in equal quantities. A

ten-sample oyster:geoduck dilution series was prepared and run using the same

methods as other SRM samples (Table S8).

Target Analysis

Raw SRM files, a background *C. gigas* proteome, and the PECAN spectral

library file from DIA were used to create a Skyline document (S9, S10). Correct

transition peaks were selected based on predicted retention times from DIA results by

comparing the relative retention times between identical PRTC peptides in the DIA and

SRM datasets (R^2 = 0.99431). Based on peptide specificity analyses, heat shock protein

70 B2 and one constituent peptide of

glucose-6-phosphate 1-dehydrogenase were removed from analyses (S11).

Coefficients of variation were calculated between technical replicates for each

peptide transition. Any sample missing data for more than 50% of peptide transitions

was deemed poor quality for downstream analyses and excluded. Abundance data was

normalized using total ion current (TIC) values from the mass spectrometer.

Consistency between technical replicates was verified in remaining samples using a

non-metric multidimensional scaling plot (NMDS) with TIC-normalized data and a

euclidean dissimilarity matrix, and technical replicates were averaged after verification

with the NMDS. Analysis of Variance and post-hoc Tukey's Honest Significant

Difference tests were used to determine significant differences in peptide abundance

among sites for average abundances. The Benjamini-Hochberg method with a false

discovery rate (FDR) of 10% was used to correct for multiple comparisons.

Results

Conditions at Outplant Locations

Over the course of the outplant, Willapa Bay (WB) had the highest average

temperatures (Table 3, Figure S12). There were no large differences in mean pH

between locations (Table 3). Oysters at Case Inlet spent more time at low tide than

those at the other four sites ($\chi2 = 25.29$, df = 4, p = 4.408e-05).

Data Independent Acquisition Mass Spectrometry

Out of 39,816 predicted proteins in the *C. gigas* FASTA proteome, 9,047 proteins

were detected in *C. gigas* across five sites and two habitats using DIA (Skyline auto

peak picker error rate 24.3% ± 25% , range: 0% to 100%). Proteins detected included,

but were not limited to, those annotated from processes such as responses to hypoxia

and oxidative stress, removal of superoxide radicals, protein folding, muscle organ

development and negative regulation of apoptosis.

 Selected Reaction Monitoring Assay

Differential abundance of protein targets was evaluated at the peptide level after

combining technical replicates (Figure S16, Figure S17). Proteins were considered

differentially abundant if at least one monitored peptide was differentially abundant.

There was no significant difference in SRM peptides between unvegetated and eelgrass

habitats across sites (Figure S18). Abundance data from both habitats were pooled for

downstream analyses comparing protein abundances among sites.

Peroxiredoxin-5 (PRX), thioredoxin reductase 3 (TrxR), catalase (EC 1.11.1.6)

(CAT), NAD(P) transhydrogenase (NADPt), glucose-6-phosphate 1-dehydrogenase

(G6PD), protein phosphatase 1B (PP1), and glycogen phosphorylase (GPH) had

differentially abundant peptides between Willapa Bay and the other four locations (Table

S19; ANOVA and Tukey's Honest Significant Difference, FDR = 0.1). These proteins

are either involved in oxidative stress response or related to carbohydrate metabolism

(Table 4).

All differentially abundant proteins were detected at higher levels in the WB

oysters than in oysters from the other four PS locations (CI, FB, PG, and SR),

regardless of protein function (Figure 1). No differences in protein abundance were

detected among the PS sites (Table S19; ANOVA and Tukey's Honest Significant

Difference).

Discussion

Among sibling Pacific oysters placed at five sites in Washington State, we found

higher abundances of antioxidant enzymes and carbohydrate metabolism proteins in

oysters outplanted at Willapa Bay (WB). Significant differences in protein abundances

were detected between WB and all four PS sites; no differences were observed within

PS. Outplanted individuals at WB had higher abundances of antioxidant enzymes and

proteins involved in carbohydrate metabolism. Increased antioxidant activity and

changes in carbohydrate metabolism have been consistently documented as responses

to extreme temperatures, hyposalinity, low pH and air exposure (Tomanek 2014; Zhang

et al. 2015). Higher average temperatures may have driven differential protein

abundances (Table 3, Figures S12).

To our knowledge, this is the first time combined use of DIA and SRM analytical

methods have been applied in an ecological context. Previously, both gel-based

(Campos et al. 2016; Tomanek 2014; Zhang et al. 2015; Parker et al. 2011) and gel-free

(Timmins-Schiffman et al. 2014; Timmins-Schiffman et al. 2017; Campos et al. 2016;

Dineshram et al. 2016; Müller et al. 2018) proteomic methods have been used in lab

settings to evaluate the effects of environmental conditions on organismal physiology,

with gel-based methods having a limited dynamic range. Similar gel-free methods has

also been used to examine physiology of other non-model marine species (Kültz et al.

2015; Plumel et al. 2013). Our methods allowed for increased acquisition of proteomic

data for a non-model organism. We detected over 9,000 proteins — far more than

previous studies have detected. The application of a novel, two step, gel-free proteomic

approach broadened the scope of proteins detected in the DIA phase, thus revealing

more possibilities for SRM target design.The SRM assay allowed for increased

detection when protein abundance was relatively low.

 Antioxidant enzymes

Higher antioxidant enzyme abundances can be a direct response to an increase

in reactive oxygen species (ROS) that arise from changes in oxygen conditions (Limón-

Pacheco and Gonsebatt 2009; Zhang et al. 2015). During electron transport, oysters

can produce ROS that induce oxidative stress if not neutralized. (Abele et al. 2007;

Limón-Pacheco and Gonsebatt 2009). Peroxiredoxin-5 (PRX), thioredoxin reductase

(TrxR), and catalase (CAT) scavenge these ROS and degrade them before they cause

cellular harm (Tomanek 2014; Sussarellu et al. 2012; Limón-Pacheco and Gonsebatt

2009; Flores-Nunes, Mattos, et al. 2015; Sussarellu et al. 2010), while NAD(P)

transhydrogenase (NADPt) maintains the cellular redox state. Higher abundances of

antioxidant enzymes in WB oysters suggest the need for ROS scavenging to

acclimatize to local conditions.

The shallow bathymetry of WB may have also contributed to elevated ROS and

the increased need for antioxidant enzymes. At high tide, shallow waters would warm

much faster, leading to the observed higher temperatures at WB (Table 1). Warmer

waters at low tide may prompt oysters to spend more times with their shells closed,

producing higher levels of ROS that would need to be neutralized. Lack of oxygen and

higher pCO_2 while the oyster is closed would lead to lower hemolymph pH, causing

higher levels of ROS. Oysters would respond through increased abundance of

antioxidant enzymes (PRX, CAT, TrxR, and NADPt).

ROS are produced in response to many environmental changes, thus biomarkers

of ROS scavenging are difficult to link to a single environmental difference in a variable

and complex setting. In particular, reduction of ROS species was found to be a common

response to both increased temperatures and aerial exposure in another experiment

with *C. gigas* (Zhang et al. 2015). Because ROS mediation is a conserved response to

several stressors, it is possible that environmental parameters we did not measure (e.g.,

contaminants, microbiota abundance, trace metals) could explain the observed variation

in antioxidant enzyme abundance. Future work at these locations should take these

variables into account.

Carbohydrate metabolism

Higher abundance of carbohydrate metabolism proteins at WB — protein

phosphatase 1B (PP1), glucose-6-phosphate 1-dehydrogenase (G6PD), and glycogen

phosphorylase (GPH) — indicate increased glycogen catabolism at this location (Table

3). Increased glycogen catabolism (PP1 and GPH) has been reported with respect to

environmental drivers (Fuhrmann et al. 2018; Papakostas et al. 2012; Bacca et al. 2005;

Serrano et al. 1993; David et al. 2005). For example, enhanced GPH activity was found

in *M. galloprovincialis* responding to thermal stress to provide more energy during high

temperature acclimation (Serrano et al. 1993). *C. gigas* outplanted at WB could be

using the same process to acclimatize to higher average temperatures. Warmer WB

temperatures could have also sped up basal metabolism, necessitating igher

abundances of PP1 and GPH to catabolize glycogen stores at a faster rate (Wu et al.

1998; She et al. 2015; Bacca et al. 2005). Increasing the amount of energy available to

bivalves by catabolizing glycogen could explain higher abundance of PP1 and GPH in

WB oysters.

Increased glycogen catabolism at WB may also be explained by decreased rates

of food capture due to less time spent filter feeding or less food availability at this

location. With closed shells, *C. gigas* are unable to filter-feed and would need to

catabolize glycogen for energy, facilitated by increased abundances of PP1 and GPH.

The WB oysters also had lower total fatty acid content when compared to oysters

outplanted at PS locations, indicative of less access to food (less food availability or less

time with valves open for filtration) or poorer quality food (Lowe et al. *in prep*). Whether

due to more time spent with their shells closed or decreased caloric intake, the WB

oysters had to rely on energy stores of glycogen more than oysters throughout Puget

Sound.

Higher G6PD abundance in WB oysters indicate that oysters at WB were still

metabolically active. G6PD catalyzes the oxidative portion of the pentose phosphate

pathway, and products from this pathway are often precursors for nucleic and aromatic

amino acids ((L., n.d.)). Additionally, G6PD activity generates NADPH and can indirectly

ameliorate effects of ROS ((L., n.d.)), as demonstrated in *C. gigas*: after exposure to

linear alklybenzenes and other pollutants, *C. gigas* ctenidia expressed higher G6PD

activity to regulate the redox environment (Flores-Nunes, Mattos, et al. 2015). Increased

abundance of G6PD in WB not only could have maintained transcription and translation

processes, but also levels of cellular and metabolic activity by indirectly dealing with

ROS. Increased G6PD abundance, therefore, provides a link between carbohydrate

metabolism and oxidative stress response in oysters outplanted at Willapa Bay.

Proteomic Responses in Puget Sound and Between Habitats

Due to the observed differences across environmental parameters between

Willapa Bay and Puget Sound locations, similar abundances for proteins involved in

various environmental responses may be evidence of physiological plasticity. One

particular protein that we expected to be differentially abundant was heat shock 70 kDA

protein 12A (HSP 12A), since Willapa Bay had higher average temperatures (Table 3).

Higher abundances of heat shock proteins (HSPs) are generally induced when

organisms are exposed to thermal stress (Hamdoun, Cheney, and Cherr 2003; Zhang

et al. 2015; Meistertzheim et al. 2007). In our experiment, average temperatures *C.*

gigas experienced were lower than the threshold to induce elevated HSP expression

(Figure 1; Table 2) (Hamdoun, Cheney, and Cherr 2003). While HSP12A is involved in

a general response to environmental drivers, differentially expressed antioxidant

enzymes and carbohydrate metabolism proteins could be more indicative of molecular

processes required for acclimatization.

The lack of differential abundance for protein targets — both among Puget

Sound sites and between unvegetated and eelgrass habitats — was unexpected. These

similar proteomic profiles could be due to three factors. First, it is possible that a

different suite of environmental response proteins in the SRM assay could have yielded

a different view of acclimatization to the various outplant sites; however, the targets we

chose have proven to yield insight into a range of environmental responses in previous

studies. Second, outplant duration could have been too short to capture varied

physiological response within Puget Sound, or oysters could have also acclimatized to

conditions in their outplant locations Finally, is that it is possible that the proteomic

response was not different because the environmental conditions that would elicit up- or

down-regulation of monitored proteins were similar across these five locations (Table 3).

While eelgrass beds are touted as a potential refuge for organisms from the negative

effects of environmental drivers like pH, they can also drive more extreme carbonate

chemistry conditions (Pacella et al. 2018). It is possible that the refuge effects and

extreme effects cancelled each other out over the course of our outplant, effectively

making the eelgrass beds just like the unvegetated sites for outplanted oysters. If this is

the case, it would explain why there was no detected difference in protein abundance

between unvegetated and eelgrass habitats.

Conclusion

The differential protein abundance observed between sibling oysters placed for

29 days in Willapa Bay or Puget Sound indicates that environmental factors at Willapa

Bay lend themselves to increased antioxidant enzyme and carbohydrate metabolism

protein abundance. Because the observed proteomic responses can be triggered by a

variety of drivers, from a field trial it is difficult to discern if high temperatures or other

environmental factors at WB led to these responses. However, the observed responses

generate a suite of new hypotheses for lab and field experiments comparing the effects

of environmental conditions on physiological responses of marine invertebrates. This is

the first to link regional environmental conditions to physiological responses in Puget

Sound, and one of the first to compare responses between Puget Sound and Willapa

Bay. Linking environmental drivers to other performance metrics, such as growth and

fatty acid composition, is crucial for contextualizing these protein abundance findings

(Lowe et al. *in prep*).

Understanding the difference between these two estuaries is important for the

persistence of oyster reefs and aquaculture in the face of climate change. Since

aquaculture species are outplanted in the same location for many months to years, the

lack of differential protein abundance among Puget Sound outplants could mean that

environmental response should not be heavily considered when selecting grow-out

locations in this region. Future experiments with a longer outplant duration and multiple

sampling points may provide a clearer connection between environmental drivers,

protein expression, and performance, as well as assist with developing a practical assay

for elucidating stress responses in the field.

 Figures and Tables

489 **Table 1.** Latitude and longitude of *C. gigas* outplants. Oysters were placed at five

490 locations sites: Case Inlet (CI), Fidalgo Bay (FB), Port Gamble Bay (PG), Skokomish

491 River Delta (SK), and Willapa Bay (WB).

Location	Latitude	Longitude
CI	47.3579367	-122.7957627
FB	48.481691	-122.58353
PG	47.842676	-122.583832
SK	47.35523	-123.1572
WB	46.4944789	-124.0261356

Table 2. Proteins used as targets for a Selected Reaction Monitoring Assay (SRM).

Targets were identified based on differential abundance and stress-related annotations.

At least two peptides and six transitions were included in the assy for each protein. The

protein Catalase had two isoforms under separate proteome IDs —

CHOYP_CATA.1.3|m.11120 and CHOYP_CATA.3.3|m.21642 — and target peptides

were chosen from both IDs. A total of four peptides and twelve associated transitions

were used as SRM targets.

Protein	Proteome ID	Number of Peptides	Number of Transitions
3-ketoacyl-CoA thiolase	CHOYP_ACAA2. 1.1|m.30666	3	9

Peroxiredoxin-5	CHOYP_BRAFLD RAFT_119799.1. 1|m.23765	2	6
Thioredoxin reductase 3	CHOYP_BRAFLD RAFT_122807.1. 1|m.3729	3	9
Protein phosphatase 1B	CHOYP_BRAFLD RAFT_275870.1. 1|m.12895	3	9
Carbonic anyhdrase 2	CHOYP_CAH2.1. 1|m.42306	3	9
Catalase 1	CHOYP_CATA.1. 3|m.11120	3	9
Catalase 2	CHOYP_CATA.3. 3|m.21642	1	3
Glucose-6-phosphate 1-dehydrogenase	CHOYP_G6PD.2. 2|m.46923	3	9
Heat shock 70 kDa protein	CHOYP_HS12A. 25.33|m.60352	2	6

Heat shock protein 70 B2	CHOYP_HSP74.1.1|m.13095	2	6
NAD(P) transhydrogenase	CHOYP_LOC100633041.1.1|m.35428	2	6
Glycogen phosphorylase	CHOYP_LOC100883864.1.1|m.41791	3	9
Multidrug resistance-associated protein	CHOYP_MRP1.5.10|m.34368	2	6
Protein disulfide-isomerase	CHOYP_PDIA1.1.1|m.5297	3	9
Protein disulfide-isomerase	CHOYP_PDIA3.1.1|m.60223	3	9
Puromysin-sensitive amirase	CHOYP_PSA.1.1|m.27259	3	9

501 **Table 3.** Environmental conditions (mean ± standard deviation) at outplant locations

502 (Case Inlet (CI), Fidalgo Bay (FB), Port Gamble Bay (PG), Skokomish River Delta (SK),

503 Willapa Bay (WB).

Metric	CI	FB	PG	SK	WB
Mean Temperature (ºC)	16.1 ± 1.7	14.8 ± 1.8	15.1 ± 2.9	15.1± 2.2	18.0 ± 1.3
Mean Salinity (PSU)	24.47 ± 1.35	30.20 ± 0.38	23.25 ± 1.56	13.43 ± 1.04	27.28 ± 0.73
Mean Dissolved Oxygen Content (mg/L)	8.27 ± 1.91	9.63 ± 4.59	11.70 ± 3.70	9.99 ± 4.58	8.76 ± 1.78
Mean pH	7.63 ± 0.19	7.54 ± 0.23	7.33 ± 0.25	7.37 ± 0.24	7.56 ± 0.18

505   **Table 4.** Functions of differentially abundant proteins

Protein	Function	Citations
Peroxiredoxin-5 (PRX)	Scavenges ROS located in mitochondria using cysteine residuals to reduce substrates	Tomanek 2014; Sussarellu et al. 2012
Thioredoxin reductase	Thiol-specific antioxidant	Limón-Pacheco and

(TrxR)	reducing agent	Gonsebatt 2009
Catalase (CAT)	Degrades hydrogen peroxide into water and oxygen	Flores-Nunes, Mattos, et al. 2015; Sussarellu et al. 2012
NAD(P) transhydrogenase (NADPt)	Involved in maintenance of cellular redox state	Sussarellu et al. 2010, 2012
Glucose-6-phosphate 1-dehydrogenase (G6PD)	Catalyzes first step in the pentose phosphate pathway and controls rate of pathway's oxidative portion	Livingstone., 1982
Protein phosphatase 1B (PP1)	Regulation of glygcogen metabolism by dephosphorylating glycogen synthase	Wu et al. 1998
Glycogen phosphorylase (GPH)	Key glycogen catabolism enzyme. Removes glucose from glycogen molecules to aid in glycogen breakdown	David et al. 2005; Oyster and Gigas 1993; She et al. 2015

507

WB   CI   SK   FB   PG

Figure 1. Average protein abundance by constituent peptides across experimental sites

from SRM. Peptide abundance data at each site was averaged, then log transformed.

Proteins were considered differentially abundant if at least one constituent peptide was

significantly different (indicated by an asterisk). There were no significant differences in

protein abundance among the Puget Sound locations, or between unvegetated and

eelgrass habitats.

Acknowledgements

This work was funded by the Washington State Department of Natural

Resources Interagency Agreement 16-241 and Washington Sea Grant. Brittany Taylor

(Taylor Shellfish) provided *C. gigas* for the outplant experiment. The University of

Washington Proteomic Resource and Priska von Haller assisted with mass

spectrometry. Jarrett Egertson (Department of Genome Sciences) helped design the

DIA acquisition. Austin Keller (Department of Genome Sciences) adapted MSConvert to

correctly demultiplex and convert DIA data files. Sean Bennet (School of Aquatic and

Fishery Sciences), and Han-Yin Yang and Brian Searle (Department of Genome

Sciences) assisted with running Pecan.

References

Abele, Eoris, Eva Philip, Paula M. Gonzalez, and Susana Puntarulo. 2007. "Marine Invertebrate
Mitochondria and Oxidative Stress." *Frontiers in Bioscience: A Journal and Virtual Library* 12
(January): 933–46.
Bacca, H., A. Huvet, C. Fabioux, J-Y Daniel, M. Delaporte, S. Pouvreau, A. Van Wormhoudt,
and J. Moal. 2005. "Molecular Cloning and Seasonal Expression of Oyster Glycogen
Phosphorylase and Glycogen Synthase Genes." *Comparative Biochemistry and Physiology.*
Part B, Biochemistry & Molecular Biology 140 (4): 635–46.
Banas, N. S., B. M. Hickey, P. MacCready, and J. A. Newton. 2004. "Dynamics of Willapa Bay,
Washington: A Highly Unsteady, Partially Mixed Estuary." *Journal of Physical Oceanography* 34
(11). American Meteorological Society: 2413–27.
Banas, N. S., B. M. Hickey, J. A. Newton, and J. L. Ruesink. 2007. "Tidal Exchange, Bivalve
Grazing, and Patterns of Primary Production in Willapa Bay, Washington, USA." *Marine Ecology*
Progress Series 341. Inter-Research Science Center: 123–39.
Beyer, Jonny, Norman W. Green, Steven Brooks, Ian J. Allan, Anders Ruus, Tânia Gomes,
Inger Lise N. Bråte, and Merete Schøyen. 2017. "Blue Mussels (Mytilus Edulis Spp.) as Sentinel
Organisms in Coastal Pollution Monitoring: A Review." *Marine Environmental Research* 130
(September): 338–65.
Bianucci, Laura, Wen Long, Tarang Khangaonkar, Gregory Pelletier, Anise Ahmed, Teizeen
Mohamedali, Mindy Roberts, and Cristiana Figueroa-Kaminsky. 2018. "Sensitivity of the
Regional Ocean Acidification and Carbonate System in Puget Sound to Ocean and Freshwater
Inputs." *Elem Sci Anth* 6 (1): 22.
Campos, Alexandre, Gabriela Danielsson, Ana Paula Farinha, Jacob Kuruvilla, Per Warholm,
and Susana Cristobal. 2016. "Shotgun Proteomics to Unravel Marine Mussel (Mytilus Edulis)
Response to Long-Term Exposure to Low Salinity and Propranolol in a Baltic Sea Microcosm."
Journal of Proteomics 137 (March): 97–106.
Chambers, Matthew C., Brendan Maclean, Robert Burke, Dario Amodei, Daniel L. Ruderman,
Steffen Neumann, Laurent Gatto, et al. 2012. "A Cross-Platform Toolkit for Mass Spectrometry
and Proteomics." *Nature Biotechnology* 30 (10): 918–20.
Chapman, Robert W., Annalaura Mancia, Marion Beal, Artur Veloso, Charles Rathburn, Anne
Blair, A. F. Holland, et al. 2011. "The Transcriptomic Responses of the Eastern Oyster,
Crassostrea Virginica, to Environmental Conditions." *Molecular Ecology* 20 (7): 1431–49.
Cheney, D. P., B. F. MacDonald, and R. A. Elston. 2000. "Summer Mortality of Pacific Oysters,
Crassostrea Gigas (Thunberg): Initial Findings on Multiple Environmental Stressors in Puget
Sound, Washington, 1998." *Journal of Shellfish Research* 19 (1). [S.l. :National Shellfisheries
Association,: 6.
Cornwall, Christopher E., and Catriona L. Hurd. 2016. "Experimental Design in Ocean
Acidification Research: Problems and Solutions." *ICES Journal of Marine Science: Journal Du*
Conseil 73 (3). Oxford University Press: 572–81.
Dan Kelley and Clark Richards (2018). oce: Analysis of Oceanographic Data. R package
version 0.9-23. https://CRAN.R-project.org/package=oce
David, Elise, Arnaud Tanguy, Karine Pichavant, and Dario Moraga. 2005. "Response of the
Pacific Oyster Crassostrea Gigas to Hypoxia Exposure under Experimental Conditions." *The*

FEBS Journal 272 (21): 5635–52.

Egertson, Jarrett D., Andreas Kuehn, Gennifer E. Merrihew, Nicholas W. Bateman, Brendan X.
MacLean, Ying S. Ting, Jesse D. Canterbury, et al. 2013. "Multiplexed MS/MS for Improved
Data-Independent Acquisition." *Nature Methods* 10 (8): 744–46.

Feely, Richard A., Simone R. Alin, Jan Newton, Christopher L. Sabine, Mark Warner, Allan
Devol, Christopher Krembs, and Carol Maloy. 2010. "The Combined Effects of Ocean
Acidification, Mixing, and Respiration on pH and Carbonate Saturation in an Urbanized
Estuary." *Estuarine, Coastal and Shelf Science* 88 (4): 442–49.

Flores-Nunes, Fabrício, Tânia Gomes, Rui Company, Roberta R. M. Moraes, Silvio T. Sasaki,
Satie Taniguchi, Márcia C. Bicego, Cláudio M. R. Melo, Afonso C. D. Bainy, and Maria J.
Bebianno. 2015. "Changes in Protein Expression of Pacific Oyster Crassostrea Gigas Exposed
in Situ to Urban Sewage." *Environmental Science and Pollution Research International* 22 (22):
17267–79.

Flores-Nunes, Fabrício, Jacó J. Mattos, Flávia L. Zacchi, Miguel A. S. Serrano, Clei E. Piazza,
Silvio T. Sasaki, Satie Taniguchi, Márcia C. Bicego, Cláudio M. R. Melo, and Afonso C. D.
Bainy. 2015. "Effect of Linear Alkylbenzene Mixtures and Sanitary Sewage in Biochemical and
Molecular Responses in Pacific Oyster Crassostrea Gigas." *Environmental Science and
Pollution Research* 22 (22): 17386–96.

Fuhrmann, Marine, Lizenn Delisle, Bruno Petton, Charlotte Corporeau, and Fabrice Pernet.
2018. "Metabolism of the Pacific oyster,Crassostrea Gigas, Is Influenced by Salinity and
Modulates Survival to the Ostreid Herpesvirus OsHV-1." *Biology Open* 7 (2).
https://doi.org/10.1242/bio.028134.

Hamdoun, Amro M., Daniel P. Cheney, and Gary N. Cherr. 2003. "Phenotypic Plasticity of
HSP70 and HSP70 Gene Expression in the Pacific Oyster (Crassostrea Gigas): Implications for
Thermal Limits and Induction of Thermal Tolerance." *The Biological Bulletin* 205 (2): 160–69.

Hoaglin, David C., Boris Iglewicz, and John W. Tukey. 1986. "Performance of Some Resistant
Rules for Outlier Labeling." *Journal of the American Statistical Association* 81 (396). Taylor &
Francis: 991–99.

Kültz, Dietmar, Johnathon Li, Xuezhen Zhang, Fernando Villarreal, Tuan Pham, and Darlene
Paguio. 2015. "Population-Specific Plasma Proteomes of Marine and Freshwater Three-Spined
Sticklebacks (Gasterosteus Aculeatus)." *Proteomics* 15 (23-24): 3980–92.

Limón-Pacheco, Jorge, and María E. Gonsebatt. 2009. "The Role of Antioxidants and
Antioxidant-Related Enzymes in Protective Responses to Environmentally Induced Oxidative
Stress." *Mutation Research* 674 (1-2): 137–47.

Liu, Zhiyong, Shaoming Pan, Zhuyou Sun, Renfeng Ma, Lanhua Chen, Yanlong Wang, and
Shuao Wang. 2015. "Heavy Metal Spatial Variability and Historical Changes in the Yangtze
River Estuary and North Jiangsu Tidal Flat." *Marine Pollution Bulletin* 98 (1-2): 115–29.

L., Mussel Mytilus Edulis. n.d. "INDUCTION OF ENZYMES AS A MECHANISM FOR THE
SEASONAL CONTROL OF METABOLISM IN MARINE INVERTEBRATES: GLUCOSE-6-
PHOSPHATE DEHYDROGENASES FROM THE MANTLE AND HEPATOPANCREAS OF THE
COMMON."

Lockwood, Brent L., and George N. Somero. 2011. "Transcriptomic Responses to Salinity
Stress in Invasive and Native Blue Mussels (genus Mytilus)." *Molecular Ecology* 20 (3): 517–29.

Lowe, Alexander T. et al. "Growth and fatty acid composition differences in outplanted Pacific

oysters" *in prep.*

MacLean, Brendan, Daniela M. Tomazela, Nicholas Shulman, Matthew Chambers, Gregory L.
Finney, Barbara Frewen, Randall Kern, David L. Tabb, Daniel C. Liebler, and Michael J.
MacCoss. 2010. "Skyline: An Open Source Document Editor for Creating and Analyzing
Targeted Proteomics Experiments." *Bioinformatics* 26 (7): 966–68.

Meistertzheim, Anne-Leila, Arnaud Tanguy, Dario Moraga, and Marie-Thérèse Thébault. 2007.
"Identification of Differentially Expressed Genes of the Pacific Oyster Crassostrea Gigas
Exposed to Prolonged Thermal Stress." *The FEBS Journal* 274 (24): 6392–6402.

Meng, Jie, Wenxiong Wang, Li Li, Qi Yin, and Guofan Zhang. 2017. "Cadmium Effects on DNA
and Protein Metabolism in Oyster (Crassostrea Gigas) Revealed by Proteomic Analyses."
Scientific Reports 7 (1): 11716.

Oyster, O. F. The, and Crassostrea Gigas. 1993. "PURIFICATION AND PROPERTIES OF
GLYCOGEN PHOSPHORYLASE FROM THE ADDUCTOR MUSCLE." *Planning Perspectives:*
PP 481: 486.

Pacella, Stephen R., Cheryl A. Brown, George G. Waldbusser, Rochelle G. Labiosa, and Burke
Hales. 2018. "Seagrass Habitat Metabolism Increases Short-Term Extremes and Long-Term
Offset of CO2under Future Ocean Acidification." *Proceedings of the National Academy of*
Sciences of the United States of America, April. https://doi.org/10.1073/pnas.1703445115.

Paerl, Hans W., Nathan S. Hall, Benjamin L. Peierls, and Karen L. Rossignol. 2014. "Evolving
Paradigms and Challenges in Estuarine and Coastal Eutrophication Dynamics in a Culturally
and Climatically Stressed World." *Estuaries and Coasts* 37 (2): 243–58.

Papakostas, Spiros, Anti Vasemägi, Juha-Pekka Vähä, Mikael Himberg, Lauri Peil, and Craig R.
Primmer. 2012. "A Proteomics Approach Reveals Divergent Molecular Responses to Salinity in
Populations of European Whitefish (Coregonus Lavaretus)." *Molecular Ecology* 21 (14): 3516–
30.

Parker, Laura, Pauline Ross, David Raftos, Emma Thompson, and Wayne O'Connor. 2011.
"The Proteomic Response of Larvae of the Sydney Rock Oyster, Saccostrea Glomerata to
Elevated P CO2." *The Australian Zoologist* 35 (4). Royal Zoological Society of NSW: 1011–23.

Pelletier, Gregory, Mindy Roberts, Mya Keyzers, and Simone R. Alin. 2018. "Seasonal Variation
in Aragonite Saturation in Surface Waters of Puget Sound – a Pilot Study."
https://doi.org/10.1525/elementa.270.

Pennock, Jonathan R., and Jonathan H. Sharp. 1986. "Phytoplankton Production in the
Delaware Estuary: Temporal and Spatial Variability." *Marine Ecology Progress Series* 34 (1/2).
Inter-Research Science Center: 143–55.

Plumel, Marine I., Thierry Wasselin, Virginie Plot, Jean-Marc Strub, Alain Van Dorsselaer,
Christine Carapito, Jean-Yves Georges, and Fabrice Bertile. 2013. "Mass Spectrometry-Based
Sequencing and SRM-Based Quantitation of Two Novel Vitellogenin Isoforms in the
Leatherback Sea Turtle (Dermochelys Coriacea)." *Journal of Proteome Research* 12 (9): 4122–
35.

R Core Team (2018). R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Riebesell, Ulf, and Jean-Pierre Gattuso. 2014. "Lessons Learned from Ocean Acidification
Research." *Nature Climate Change* 5 (December). Nature Publishing Group, a division of
Macmillan Publishers Limited. All Rights Reserved.: 12.

Ruesink, J. L., A. Sarich, and A. C. Trimble. 2018. "Similar Oyster Reproduction across
Estuarine Regions Differing in Carbonate Chemistry." *ICES Journal of Marine Science: Journal*
Du Conseil 75 (1). Oxford University Press: 340–50.
Serrano, Fuencisla San Juan, Pilar Suarez Alonso, Marta Lombraña Tascon, and L. Oscar
Garcia Martin. 1993. "Effect of Temperature on Glycogen Phosphorylase Activity of Mytilus
Galloprovincialis Mantle Tissue." In *Actas Del IV Congreso Nacional de Acuiculture, Centro de*
Investigaciones Marinas, Pontevedra (Spain), 425–30.
She, Zhicai, Li Li, Haigang Qi, Kai Song, Huayong Que, and Guofan Zhang. 2015. "Candidate
Gene Polymorphisms and Their Association with Glycogen Content in the Pacific Oyster
Crassostrea Gigas." *PloS One* 10 (5): e0124401.
Silbiger, Nyssa J., and Cascade J. B. Sorte. 2018. "Biophysical Feedbacks Mediate Carbonate
Chemistry in Coastal Ecosystems across Spatiotemporal Gradients." *Scientific Reports* 8 (1):
796.
Slattery, Marc, Sridevi Ankisetty, Jone Corrales, K. Erica Marsh-Hunkin, Deborah J. Gochfeld,
Kristine L. Willett, and John M. Rimoldi. 2012. "Marine Proteomics: A Critical Assessment of an
Emerging Technology." *Journal of Natural Products* 75 (10): 1833–77.
Sussarellu, Rossana, Caroline Fabioux, Miguel Camacho Sanchez, Nelly Le Goïc, Christophe
Lambert, Philippe Soudant, and Dario Moraga. 2012. "Molecular and Cellular Response to
Short-Term Oxygen Variations in the Pacific Oyster Crassostrea Gigas." *Journal of*
Experimental Marine Biology and Ecology 412 (January): 87–95.
Sussarellu, Rossana, Caroline Fabioux, Gilles Le Moullac, Elodie Fleury, and Dario Moraga.
2010. "Transcriptomic Response of the Pacific Oyster Crassostrea Gigas to Hypoxia." *Marine*
Genomics 3 (3-4): 133–43.
Timmins-Schiffman, Emma B., Grace A. Crandall, Brent Vadopalas, Michael E. Riffle, Brook L.
Nunn, and Steven B. Roberts. 2016. "Integrating Discovery-Driven Proteomics and Selected
Reaction Monitoring to Develop a Non-Invasive Assay for Geoduck Reproductive Maturation."
bioRxiv. https://doi.org/10.1101/094615.
———. 2017. "Integrating Discovery-Driven Proteomics and Selected Reaction Monitoring To
Develop a Noninvasive Assay for Geoduck Reproductive Maturation." *Journal of Proteome*
Research 16 (9): 3298–3309.
Timmins-Schiffman, Emma, William D. Coffey, Wilber Hua, Brook L. Nunn, Gary H. Dickinson,
and Steven B. Roberts. 2014. "Shotgun Proteomics Reveals Physiological Response to Ocean
Acidification in Crassostrea Gigas." *BMC Genomics* 15 (November): 951.
Ting, Ying S., Jarrett D. Egertson, Samuel H. Payne, Sangtae Kim, Brendan MacLean, Lukas
Käll, Ruedi Aebersold, Richard D. Smith, William Stafford Noble, and Michael J. MacCoss.
2015. "Peptide-Centric Proteome Analysis: An Alternative Strategy for the Analysis of Tandem
Mass Spectrometry Data." *Molecular & Cellular Proteomics: MCP* 14 (9): 2301–7.
Tomanek, Lars. 2014. "Proteomics to Study Adaptations in Marine Organisms to Environmental
Stress." *Journal of Proteomics* 105 (June): 92–106.
Veldhoen, Nik, Michael G. Ikonomou, and Caren C. Helbing. 2012. "Molecular Profiling of
Marine Fauna: Integration of Omics with Environmental Assessment of the World's Oceans."
Ecotoxicology and Environmental Safety 76 (2): 23–38.
Wu, J., J. Liu, I. Thompson, C. J. Oliver, S. Shenolikar, and D. L. Brautigan. 1998. "A Conserved
Domain for Glycogen Binding in Protein Phosphatase-1 Targeting Subunits." *FEBS Letters* 439

(1-2): 185–91.

Zhang, Yang, Jin Sun, Huawei Mu, Jun Li, Yuehuan Zhang, Fengjiao Xu, Zhiming Xiang, Pei-

Yuan Qian, Jian-Wen Qiu, and Ziniu Yu. 2015. "Proteomic Basis of Stress Responses in the

Gills of the Pacific Oyster Crassostrea Gigas." *Journal of Proteome Research* 14 (1): 304–17.