Accurate sequence variant genotyping in cattle using variation-aware genome graphs

Danang Crysnanto[1, *], Christine Wurmser[2], Hubert Pausch[1]

[1]Animal Genomics, ETH Zurich, Zurich, Switzerland
[2]Chair of Animal Breeding, TU München, Freising, Germany

[*]Corresponding author: danang.crysnanto@usys.ethz.ch

Abstract

Background: The genotyping of sequence variants typically involves as a first step the alignment of sequencing reads to a linear reference genome. Because a linear reference genome represents only a small fraction of sequence variation within a species, reference allele bias may occur at highly polymorphic or diverged regions of the genome. Graph-based methods facilitate to compare sequencing reads to a variation-aware genome graph that incorporates non-redundant DNA sequences that segregate within a species. We compared sensitivity and accuracy of graph-based sequence variant genotyping using the *Graphtyper* software to two widely used methods, i.e., *GATK* and *SAMtools*, that rely on linear reference genomes using whole-genomes sequencing data of 49 Original Braunvieh cattle.

Results: We discovered 21,140,196, 20,262,913 and 20,668,459 polymorphic sites using *GATK*, *Graphtyper,* and *SAMtools*, respectively. Comparisons between sequence variant and microarray-derived genotypes showed that *Graphtyper* outperformed both *GATK* and *SAMtools* in terms of genotype concordance, non-reference sensitivity, and non-reference discrepancy. The sequence variant genotypes that were obtained using *Graphtyper* had the lowest number of mendelian inconsistencies for both SNPs and indels in nine sire-son pairs with sequence data. Genotype phasing and imputation using the *Beagle* software improved the quality of the sequence variant genotypes for all tools evaluated particularly for animals that have been sequenced at low coverage. Following imputation, the concordance between sequence- and microarray-derived genotypes was almost identical for the three methods evaluated, i.e., 99.39, 99.29, and 99.52 % for *GATK*, *SAMtools*, and *Graphtyper*, respectively. *Graphtyper* required considerably more computing resources than *SAMtools* but it required less than *GATK*.

Conclusions: Sequence variant genotyping using *Graphtyper* is accurate, sensitive and computationally feasible in cattle. Graph-based methods enable sequence variant genotyping from variation-aware reference genomes that may incorporate cohort-specific sequence variants which is not possible with the current implementations of state-of-the-art methods that rely on linear reference genomes.

Keywords: Sequence variant genotyping, Genome graph, Variation-aware graph, cattle, Whole-genome sequencing

Introduction

The sequencing of important ancestors of many cattle breeds revealed millions of sequence variants that are polymorphic in dairy and beef populations [1–4]. In order to compile an exhaustive catalog of polymorphic sites that segregate in *Bos taurus*, the 1000 bull genomes consortium was established [5]. The 1000 bull genomes project imputation reference panel facilitates to infer sequence variant genotypes for large cohorts of genotyped animals thus enabling genomic investigations at nucleotide resolution [5–8].

Sequence variant discovery and genotyping typically involves two steps that are carried out successively [9–12]: first, raw sequencing data are generated, filtered to remove bases with low sequencing quality, and aligned towards a linear reference genome using, e.g., *Bowtie* [13] or the Burrows-Wheeler Alignment (*BWA)* software [14]. Second, the aligned reads are compared to the nucleotide sequence of a reference genome in order to discover and genotype polymorphic sites using, e.g., *SAMtools* [15] or the *Genome Analysis Toolkit (GATK)* [16–18]. The sensitivity and specificity of sequence variant discovery is higher using multi-sample than single-sample approaches particularly when the sequencing depth is low [19–23]. However, the genotyping of sequence variants from multiple samples simultaneously is a computationally demanding task, particularly when the number of sequenced individuals is high [18]. The multi-sample sequence variant genotyping approach that has been implemented in the *SAMtools* software has to be restarted for the entire cohort once new samples are added. In contrast, the joint genotyping approach that has been implemented in *GATK* separates the discovery of polymorphic sites within samples from the estimation of genotype likelihoods across samples. Once new samples are added to an existing cohort, only the latter needs to be performed for the entire cohort, thus enabling computationally efficient parallelization of sequence variant genotyping in a large number of samples.

Ideally, the accuracy and sensitivity of sequence variant discovery is assessed using high-confidence sequence variants as a truth set e.g., Genome in a Bottle, GIAB [24]. For species where such a resource is not available, the accuracy of sequence variant genotyping may be evaluated by comparing sequence variant to microarray-derived genotypes. Due to the ascertainment bias of the microarray-derived SNPs, this type of comparison may overestimate the accuracy of sequence variant discovery [25, 26].

Sequence variant genotyping approaches that rely on alignments to a linear reference genome have limitations to variant discovery, because a monoploid reference sequence does not reflect variation within a species. As a result, read alignments may be erroneous particularly at genomic regions that differ substantially between the sequenced individual and the reference sequence, thus introducing reference allele bias, flawed genotypes, and false-positive variant discovery around indels [27–29]. Aligning reads to population- or breed-specific reference genomes may overcome most of these problems [30–32]. However, considering multiple (population-specific) linear reference genomes with distinct genomic coordinates complicates the biological interpretation and annotation of sequence variant genotypes across populations [33].

Genome graph-based methods consider non-linear reference sequences for variant discovery [33–37]. A variation-aware genome graph may incorporate distinct (population-specific) reference sequences and known sequence variants. Recently, the *Graphtyper* software has been developed in order to facilitate sequence variant discovery from a genome graph that has been constructed and iteratively augmented using variation of the sequenced cohort [34]. So far, sequence variant genotyping using variation-aware genome graphs has not been evaluated in cattle.

In this study, we compare sequence variant discovery and genotyping from a variation-aware genome graph using *Graphtyper* to two state-of-the-art methods (*GATK*, *SAMtools*) that rely on linear reference genomes in 49 Original Braunvieh cattle. We compare sequence variant to microarray-derived genotypes in order to assess sensitivity and accuracy of sequence variant genotyping for each of the three methods evaluated.

Results

Following quality control (removal of adapter sequences and low-quality bases), we aligned more than 13 billion paired-end reads (2 x 125 and 2 x 150 basepairs) from 49 Original Braunvieh cattle to the UMD3.1 assembly of the bovine genome. On average, 98.44% (91.06-99.59%) of the reads mapped to the reference genome. 4.26% (2.0-10.91%) of the mapped reads were flagged as duplicates and not considered for further analyses. The average sequencing depth per animal was 12.75 and it ranged from 6.00 to 37.78. Although the re-alignment of sequencing reads around indels is not required when sequence variants are genotyped using the latest version of *GATK* (v 4), it is recommended to improve the genotyping of indels using *SAMtools*. To ensure an unbiased comparison of the three

methods evaluated, we realigned the reads around indels and used the re-aligned BAM files as input for all tools (**Figure 1**). All sequencing read data were deposited at the European Nucleotide Archive (ENA) (http://www.ebi.ac.uk/ena) under primary accession PRJEB28191.

Figure 1 Schematic representation of three sequence variant discovery and genotyping methods evaluated.

Sequence variant discovery and genotyping

Polymorphic sites (SNPs, short insertions and deletions) were discovered and genotyped in the 49 animals using either *GATK* (version 4), *Graphtyper* (version 1.3) or *SAMtools* (version 1.8). All software programs were run using well-established recommendations and workflow descriptions for variant discovery (**Figure 1**, also see Material and Methods). Only autosomal sequence variants were considered to evaluate the accuracy and sensitivity of sequence variant genotyping.

The number of polymorphic sites discovered was 21,140,196, 20,262,913, and 20,668,459 using *GATK, Graphtyper* and *SAMtools*, respectively (**Table 1)**. The vast majority (85.11-89.42%) of the detected variants were biallelic SNPs. Of the 18,594,182, 18,120,724 and 17,592,038 SNPs detected using *GATK*, *Graphtyper* and *SAMtools*, respectively, 7.46%, 8.31% and 5.02% were novel, i.e., they were not among 102,091,847 polymorphic sites of the most recent version (150) of the Bovine dbSNP database. The transition to transversion

(Ti/Tv) ratio of the detected SNPs was 2.09, 2.07 and 2.05 using *GATK*, *Graphtyper* and *SAMtools*, respectively. Using *GATK* revealed four times more multiallelic SNPs (246,220) than either *SAMtools* or *Graphtyper*.

The number of indels identified was 2,478,489, 2,044,585, and 3,076,421, using *GATK*, *Graphtyper*, and *SAMtools*, respectively and 26.78%, 29.15%, and 41.75% of them were novel. *SAMtools* revealed the highest number and proportion (14.9%) of indels. Between 12 and 14% of the detected indels were multiallelic. While *Graphtyper* and *GATK* identified more (12%) deletions than insertions, the ratio was almost equal using *SAMtools*.

On average, each Original Braunvieh cattle carried between 7 and 8 million variants that differed from the UMD3.1 reference genome. Of those, between 2.4 and 2.6 million SNPs were homozygous for the alternate allele, between 3.8 and 4.7 million SNPs were heterozygous and between 0.7 and 1 million were indels (**Table 2**).

An intersection of 15,901,526 biallelic SNPs was common to all sequence-variant discovery tools evaluated (**Figure 2A**), i.e., between 85.51 and 90.39% of the detected SNPs of each tool, and 466,029 (2.93%, Ti/Tv: 1.81) of them were novel, i.e., they were not present in dbSNP 150. The Ti/Tv-ratio of the common SNPs was 2.22. *SAMtools* had the largest number of SNPs in common with the other two tools (90.39%). The number of private SNPs, i.e., SNPs that were detected by one but not the other tools was highest for *GATK* and lowest for *Graphtyper*.

The number of biallelic indels (**Figure 2B**) that was common to all tools evaluated was 1,299,467 and 98,931 (13.13%) of those were novel, i.e., they were not present in dbSNP 150. The intersection of the three tools was considerably smaller for indels than SNPs. *Graphtyper* had the highest proportion of indels in common with the other tools (74.11%). *SAMtools* discovered the highest number (2,704,413) of biallelic indels and most of them (41.26%) were not detected using either *GATK* or *Graphtyper*. *GATK* (21.2 %) and *Graphtyper* (12.38%) discovered less private indels than *SAMtools*.

Table 1 Numbers of different types of autosomal sequence variants detected in 49 Original Braunvieh cattle using three sequence variant genotyping methods.

The number of variants is presented for the three tools evaluated before (Full) and after (Filtered) applying recommended filters to identify and exclude low quality variants.

	Full			Filtered		
	GATK	*Graphtyper*	*SAMtools*	*GATK*	*Graphtyper*	*SAMtools*
Variants	21,140,196	20,262,913	20,668,459	19,761,679	17,679,155	18,871,549
SNPs	18,594,182	18,120,724	17,592,038	17,248,593	15,777,446	16,272,917
Not in dbSNP	1,387,781	1,505,586	882,575	867,838	564,326	570,901
Biallelic	18,347,962	18,053,396	17,528,249	17,111,806	15,730,153	16,218,714
Multi-allelic	246,220	67,328	63,789	136,787	47,293	54,203
Ti/Tv ratio	2.09	2.07	2.05	2.17	2.18	2.16
Indels	2,478,489	2,044,585	3,076,421	2,445,766	1,826,808	2,598,632
Not in dbSNP	663,831	596,137	1,279,162	639,219	456,752	979,291
Biallelic	2,166,352	1,753,391	2,704,413	2,133,840	1,571,195	2,310,386
Multi-allelic	312,137	291,194	372,008	311,926	255,613	288,246
Insertion/Deletion	0.88	0.88	1	0.88	0.88	0.99
Complex variation	67,525	97,604	0	67,320	74,901	0

Table 2 Average number of autosomal variants identified per animal using three sequence variant genotyping methods.

The number of variants is presented for the three tools evaluated before (Full) and after (Filtered) applying recommended filters to identify and exclude low quality variants.

	Full			Filtered		
	GATK	*Graphtyper*	*SAMtools*	*GATK*	*Graphtyper*	*SAMtools*
Total biallelic SNPs	6,324,455	7,384,058	6,617,948	6,105,674	6,533,711	6,564,229
Heterozygous	3,890,351	4,758,297	4,187,882	3,744,336	4,074,011	4,147,033
Homozygous ALT	2,434,104	2,625,761	2,430,066	2,361,338	2,459,700	2,417,196
Ti/Tv	2.17	2.13	2.11	2.20	2.14	2.13
Total biallelic indels	693,697	767,261	1,007,420	691,765	697,637	960,218
Heterozygous	390,495	441,172	616,981	388,622	391,856	593,417
Homozygous ALT	303,202	326,089	390,439	303,143	305,781	366,801
Singletons	49,166	23,406	32,810	41,408	17,999	32,398

Figure 2 Number of biallelic SNPs (A) and indels (B) identified in 49 Original Braunvieh cattle using three sequence variant genotyping methods.

Blue horizontal bars represent the total number of sites discovered for each method. Vertical bars indicate private and common variants detected by the methods evaluated.

Sequence variant genotyping using *Graphtyper* is accurate

Genotype concordance, non-reference sensitivity and non-reference discrepancy were calculated between the microarray-derived genotypes at 732,970 autosomal SNPs from the BovineHD Bead chip and the sequence variant genotypes that were obtained using either *GATK*, *Graphtyper*, or *SAMtools* (**Table 3**). All metrics were calculated either before or after applying the algorithm implemented in the *Beagle* software for haplotype phasing and imputation.

The proportion of missing non-reference sites was 1.90%, 0.56%, and 0.47% using *GATK*, *Graphtyper*, and *SAMtools*, respectively. The genotype concordance between the sequence- and microarray-derived genotypes was higher using *Graphtyper* (97.72%) than either *SAMtools* (97.68%) or *GATK* (95.99%). The genotype concordance was higher for homozygous than heterozygous sites, particularly in animals that have been sequenced at low depth (**Additional file 1 Figure S1**). In order to take the variable proportions of missing genotypes in the sequence variants into account, we calculated non-reference sensitivity and non-reference discrepancy. The non-reference sensitivity is calculated for all positions where microarray-derived genotypes are either heterozygous or homozygous for the alternate allele, thus reflecting the sensitivity to discover variant sites. The non-reference discrepancy reflects the genotype concordance of all but the concordant reference sites. Non-reference sensitivity was almost identical using *Graphtyper* (98.26%) and *SAMtools* (98.21%). However, non-reference sensitivity was clearly lower using *GATK* (93.81%). Non-reference discrepancy was lower using *Graphtyper* (3.53%) than either *SAMtools* (3.6%) or *GATK* (6.35%).

Next, we analysed the proportion of opposing homozygous genotypes for SNPs and indels in nine sire-son pairs that were included among the sequenced animals (**Table 4**). We observed that SNPs that were discovered using either *Graphtyper* or *SAMtools* had almost a similar proportion of genotypes with mendelian inconsistencies (0.33% and 0.36%). The proportion of genotypes with mendelian errors was two times higher using *GATK* (0.73%). The proportion of opposing homozygous genotypes for indels was lower using *Graphtyper* (0.54%) than either *GATK* (0.98%) or *SAMtools* (0.66%).

Table 3 Comparisons between microarray-derived and sequence variant genotypes.

Genotype concordance, non-reference sensitivity and non-reference discrepancy (in percentage) was calculated between the genotypes from the BovineHD SNP Bead chip and sequence –derived genotypes for 49 Original Braunvieh cattle considering either the raw or imputed (imp) sequence variant genotypes before (full) and after (filtered) recommended quality control was applied on the variants. Bold type indicates the best value that was observed for each parameter.

	Genotype concordance				Non-reference sensitivity				Non-reference discrepancy			
	full		filtered		full		filtered		full		filtered	
	raw	imp	raw	imp	raw	imp	raw	imp	raw	imp	raw	imp
GATK	95.99	99.32	96.02	99.39	93.81	**99.36**	93.67	**99.15**	6.35	1.05	6.30	0.95
Graphtyper	**97.71**	**99.46**	**97.75**	**99.52**	**98.26**	99.35	**97.91**	99.00	**3.53**	**0.83**	**3.47**	**0.73**
SAMtools	97.68	99.24	97.70	99.29	98.21	99.35	97.53	98.67	3.60	1.17	3.56	1.09

Table 4 Proportions of opposing homozygous genotypes observed in nine sire-son pairs.

The ratio (in percentage) was calculated using autosomal sequence variants considering either the raw or imputed (imp) sequence variant genotypes before (full) and after (filtered) variants with low quality were removed from the data. Bold type indicates the best value that was observed for a given parameter.

	SNPs				indels			
	full		filtered		full		filtered	
	raw	imp	raw	imp	raw	imp	raw	imp
Bovine HD SNP array								**0.001**
GATK	0.73	0.15	0.72	0.13	0.98	0.24	0.99	0.21
Graphtyper	0.36	**0.11**	0.36	**0.11**	**0.54**	**0.13**	**0.54**	**0.13**
SAMtools	**0.33**	0.28	**0.32**	0.25	0.67	0.54	0.61	0.57

Figure 3 Accuracy and sensitivity of sequence variant genotyping at different sequencing depth.

Genotype concordance, non-reference sensitivity and non-reference discrepancy were calculated for 49 Original Braunvieh cattle considering either raw (red) or imputed (blue) sequence variant genotypes.

Beagle genotype refinement improved the genotype quality

Using well-established and recommended filtering parameters (see Material & Methods), we excluded 1,378,517 (6.52%, Ti/Tv 1.24), 2,583,758 (12.75%, Ti/Tv 1.47) and 1,796,910 (8.69%, Ti/Tv 1.36) variants due to low mapping or genotyping quality from the *GATK*, *Graphtyper*, and *SAMtools* datasets, respectively. The genotype concordance between sequence- and microarray-derived genotypes was slightly higher in the filtered than raw genotypes, but the non-reference sensitivity was less in the filtered than raw genotypes

indicating that the filtering step also removed some true variant sites from the raw data (**Table 3 & 4**). We used the *Beagle* software to refine the primary genotype calls and infer missing genotypes in the raw and filtered datasets. Following imputation, the quality of the sequence variant genotypes increased for all tools evaluated particularly in cattle where the sequencing coverage was less than 10-fold (**Figure 3**). The largest increase in the concordance metrics was observed for the sequence variants obtained using *GATK* (**Table 3 & 4**). Following imputation, the variants identified using *Graphtyper* had the highest quality in eight out of ten metrics evaluated.

Computing requirements

The multi-sample sequence variant genotyping pipelines that were implemented using either *GATK* or *SAMtools* were run separately for each chromosome in single-threading mode. The *SAMtools mpileup* module took between 3.07 and 11.4 hours and it required between 0.12 and 0.25 gigabytes (GB) peak random-access memory (RAM) per chromosome. To genotype 20,668,459 sequence variants in 49 animals, *SAMtools mpileup* required 192 CPU hours (**Figure 4**).

For *GATK*, we submitted 1421 parallel jobs of the *HaplotypeCaller* module (*i.e.*, one job for each animal and chromosome) that required between 3.9 and 12.3 GB RAM and between 0.36 and 11 CPU hours to complete. To process 29 chromosomes in 49 samples, the *HaploytpeCaller* module required a total of 2428 CPU hours. Subsequently, we ran the *GATK GenomicsDBImport* module which required between 7.98 and 20.88 GB RAM and between 2.81 and 19.31 CPU hours per chromosome. *GATK Joint Genotyping* required between 4.33 and 17.32 GB of RAM and between 1.81 and 14.01 hours per chromosome. To genotype 21,140,196 polymorphic sequence variants in 49 animals, the *GATK* pipeline required a total of 2792 CPU hours (**Figure 4**).

The *Graphtyper* pipeline including the construction of the variation graph and the genotyping of sequence variants was run in parallel for 2538 non-overlapping segments of 1 million basepairs as recommended by Eggertson *et al.* [34]. The peak RAM required by *Graphtyper* was between 1 and 3 GB per segment. Twelve segments for which *Graphtyper* either ran out of memory or did not finish within the allocated time were subdivided into smaller segments of 10 Kb and subsequently re-run (**Additional file 2 Table S1**). The genotyping of 20,262,913 polymorphic sites in 49 animals using our implementation of the *Graphtyper* pipeline took a total of 1066 CPU hours (**Figure 4**).

Figure 4 Computing time required to genotype all autosomal sequence variants in 49 Original Braunvieh cattle.

The runtime of *GATK* and *Graphtyper* is shown for the different steps (see **Figure 1** for more details).

The computing resources required by *SAMtools* and *GATK* increased linearly with chromosome length. The computing time required to genotype sequence variants was highly heterogeneous along the genome using *Graphtyper*. The average CPU time per 1 Mb segment was 0.42 hours, but it varied in between 0.196 and 10.11 hours. We suspected that flaws in the reference genome might increase the complexity of the variation-aware graph and that the construction of the graph might benefit from an improved assembly. To test this hypothesis, we re-mapped the sequencing reads to the recently released new bovine reference genome (ARS-UCD1.2, https://www.ncbi.nlm.nih.gov/assembly/GCF_002263795.1) and repeated the graph-based sequence variant discovery. We indeed observed a decrease in the computing time required to genotype polymorphic sites (particularly at chromosomes 12, 27 and 29, **Figure 5**) and a more uniform runtime along the genome possibly indicating that graph-based variant discovery in cattle will be faster and more accurate from highly contiguous reference sequences.

Figure 5 Sequence variant genotyping on chromosome 12 using Graphtyper.

Computing time required (A) and number of variants discovered (B) for bovine chromosome 12 using *Graphtyper*. Each dot represents an interval of 1 million basepairs. Blue and red colour represents values for the UMD3.1 and ARS-UCD1.2 versions of the bovine assembly.

Discussion

In this study, we used either *GATK, Graphtyper* or *SAMtools* to discover and genotype polymorphic sequence variants in whole genome sequencing data of 49 Original Braunvieh cattle that have been sequenced at between 6 and 38-fold genome coverage. While *SAMtools* and *GATK* discover variants from a linear reference genome, *Graphtyper* locally re-aligns reads to a variation-aware reference graph that incorporates cohort-specific sequence variants [34]. Our graph-based variant discovery pipeline that has been implemented using the *Graphtyper* software used the existing bovine reference sequence to construct the genome graph. The graph was subsequently augmented with variants that

were detected from linear alignments of 49 Original Braunvieh cattle [34]. The use of more sophisticated genome graph-based approaches that have been developed very recently facilitates to map raw sequencing reads directly against a genome graph without the need to first align reads towards a linear reference [36]. While genome graph-based variant discovery has been explored recently in mammalian-sized genomes [29, 33, 34, 37], our work is the first to apply graph-based sequence variant genotyping in cattle.

In order to evaluate graph-based variant discovery in cattle, we compared accuracy and sensitivity of *Graphtyper* to *SAMtools* and *GATK*, i.e., two state-of-the-art methods on linear reference genomes that have been evaluated thoroughly in many species including cattle [2, 19]. Our evaluation of the software tools may suffer from ascertainment bias because we relied on SNPs that have been included in the BovineHD SNP array, i.e., they are located predominantly at well accessible regions of the genome [25, 26, 38]. Thus, the global accuracy and sensitivity of sequence variant discovery might be overestimated in our study. However, this ascertainment bias is unlikely to affect the relative performance of the methods evaluated. In 49 Original Braunvieh cattle, sequence variant genotyping was more accurate using *Graphtyper* than either *GATK* or *SAMtools.* However, the difference in accuracy was small between the three tools particularly in samples that were sequenced at high coverage. Irrespective of the methods evaluated, we observed heterozygous under-calling in animals that had been sequenced at low coverage, i.e., heterozygous variants were erroneously genotyped as homozygous due to an insufficient number of sequencing reads supporting the heterozygous genotype [9, 39–41]. In agreement with previous studies [2, 5], *Beagle* imputation improved genotype concordance and reduced heterozygous under-calling particularly in cattle that had been sequenced at low coverage. After the imputation step, the genotype concordance, non-reference sensitivity, and non-reference discrepancy of the three tools was almost identical, indicating that genotyping sequence variants from samples with medium coverage is possible at high accuracy using either of the three tools evaluated and subsequent *Beagle* imputation. While this has been concluded previously for *SAMtools* and *GATK* [2, 19], our findings demonstrate that the genotype likelihoods estimated from the *Graphtyper* software are also compatible with and benefit from the imputation algorithm implemented in the *Beagle* software. However, integrating the phasing and imputation of missing genotypes directly in a graph-based variant genotyping approach would simplify sequence variant genotyping from variation-aware graphs [33, 42, 43].

Using *Graphtyper* for variant genotyping and *Beagle* for genotype refinement enabled us to genotype sequence variants in 49 Original Braunvieh cattle at a genotypic concordance of 99.52%, i.e., higher than previously achieved using either *GATK* or *SAMtools* to genotype variants in cattle that had been sequenced at similar genome coverage [2–5, 19, 44],

indicating that graph-based variant discovery improves sequence variant genotyping. The accuracy and sensitivity of graph-based variant discovery may be higher when known variants are incorporated in the initial construction of the variation graph [34]. Indeed, we observed a slight increase in genotype concordance (**Additional file 3 Figure S2**) when sequence variants were genotyped from a variation-aware genome-graph that incorporated variants that are listed in bovine dbSNP 150.

Using microarray-derived genotypes as a truth set does not allow for assessing accuracy and sensitivity of indel discovery because variants other than SNPs are currently not routinely genotyped with commercially available microarrays. The proportion of opposing homozygous genotypes between parent-offspring pairs may be a useful diagnostic metric to detect sequencing artefacts or flawed genotypes at indels [45]. Our results show that genotypes at indels are more accurate using *Graphtyper* than either *SAMtools* or *GATK* because *Graphtyper* produced less opposing homozygous genotypes at indels in nine sire-son pairs than the other methods. These findings are in line with Eggertsson *et al* [34], who showed that the mapping of the sequencing reads against a variation-aware graph could improve read alignment nearby indels, enabling highly accurate sequence variant genotyping also for variants other than SNPs. Recently, Garrison *et al.* [36] showed that graph-based variant discovery may also mitigate reference allele bias. An assessment of reference allele bias was, however, not possible in our study because the sequencing depth was too low for most samples.

In our study, *Graphtyper* required less computing time than *GATK* to genotype sequence variants for 49 cattle. *SAMtools* required the least computing resources likely because the implemented *mpileup* algorithm readily produces genotypes from the aligned reads without performing the computationally intensive local re-alignment of the reads. However, with an increasing number of samples, the multi-sample variant genotyping implementation of *GATK* seems to be more efficient than *SAMtools mpileup* because variant discovery within samples can be separated from the joint genotyping across samples [18]. A highly parallelized graph-based variant discovery pipeline also offers a computationally efficient and scalable framework for variant discovery in thousands of samples [34]. However, the computing time for graph-based variant genotyping might be high in genomic regions where the nucleotide diversity is high or the assembly is flawed [37, 46]. In our study, the algorithm implemented in the *Graphtyper* software failed to finish within the allocated time for twelve 1 Mb segments including a segment on chromosome 12 that contains a large segmental duplication [47–49] possibly because many mis-mapped reads increased graph complexity. Our results show that *Graphtyper* may also produce genotypes for problematic segments when they are split and processed in smaller bits. Moreover, most of these problems disappeared when we

considered the latest assembly of the bovine genome, possibly corroborating that more complete and contiguous genome assemblies may facilitate more reliable genotyping from variation-aware graphs [25, 50].

Conclusion

Genome graphs facilitate sequence variant discovery from non-linear reference genomes. Sequence variant genotyping from a variation-aware graph is possible in cattle using *Graphtyper*. Sequence variant genotyping at both SNP and indels is more accurate and sensitive using *Graphtyper* than either *SAMtools* or *GATK*. The proportion of mendelian inconsistencies at both SNP and indels is low using *Graphtyper* indicating that sequence variant genotyping from a variation-aware genome graph facilitates accurate variant discovery at different types of genetic variation. Considering highly-informative variation-aware genome graphs that have been constructed from multiple breed-specific de-novo assemblies and high-confidence sequence variants facilitates more accurate, sensitive and unbiased sequence variant genotyping in cattle.

Methods

Animal Selection

We selected 49 Original Braunvieh bulls that were either frequently used in artificial insemination or explained a large fraction of the genetic diversity of the breeding population. Semen straws of the bulls were purchased from an artificial insemination center and DNA was prepared following standard DNA extraction protocols.

Sequencing data pre-processing

All samples were sequenced at either Illumina HiSeq 2500 (30 animals) or Illumina HiSeq 4000 (19 animals) instruments using 125 or 150 basepair paired-end sequencing libraries with insert sizes between 400 and 450 basepairs. Quality control (removal of adapter sequences and bases with low quality) of the raw sequencing data was carried out using the *fastp* software (version 0.19.4) with default parameters [51]. The filtered reads were mapped to the UMD3.1 version of the bovine reference genome [52] using *BWA mem* (version 0.7.12) [14] with option -M to mark shorter split hits as secondary alignments, otherwise the default parameters were applied. Optical and PCR duplicates were marked using

Samblaster (version 0.1.24) [53]. The output of *Samblaster* was converted into BAM format using *SAMtools view* (version 1.3) [15] and subsequently coordinate sorted using *Sambamba* (vesion 0.6.6) [54]. We used the *GATK* (version 3.8) *RealignerTargetCreator* and *IndelRealigner* modules to realign reads around indels. The realigned BAM files served as an input for *GATK BQSR* (base quality score recalibration) using 732,970 and 102,091,847 positions from the Illumina BovineHD SNP chip and Bovine dbSNP version 150, respectively, as known variants. The *mosdepth* software (version 0.2.2) [55] was used to extract the number of reads covering a genomic position.

Sequence variant discovery

We followed the best practice guidelines recommended for variant discovery and genotyping using *GATK* (version 4.0.6) [16, 17, 23]. First, genotype likelihoods were calculated separately for each sequenced animal using *GATK HaplotypeCaller* [18], resulting in files in gVCF (genomic Variant Call Format) format for each sample [56]. The gVCF files from 49 samples were consolidated using *GATK GenomicsDBImport*. Subsequently, *GATK CombineGVCFs* was applied to genotype polymorphic sequence variants for all samples simultaneously.

Graphtyper (version 1.3) was run in multi-sample mode as recommended in Eggertsson *et al.* [34]. Because the original implementation of *Graphtyper* is limited to the analysis of the human chromosome complement, we cloned the *Graphtyper GitHub* repository (https://github.com/DecodeGenetics/graphtyper), modified the source code to allow analyzing the cattle chromosome complement and compiled the program from the modified source code. The *Graphtyper* workflow consisted of four steps that were executed successively. First, sequence variants were identified from the read alignments that were produced using *BWA mem* (see above). Next, these cohort-specific variants were used to augment the UMD3.1 reference genome and construct the variation-aware genome graph. Subsequently, the sequencing reads were locally re-aligned against the variation-aware graph. A clean variation graph was produced by removing unobserved haplotypes paths from the raw graph. In the final step, genotypes were identified from the re-aligned reads in the clean graph. The *Graphtyper* pipeline was run in segments of 1 million base-pairs and whenever the program failed to genotype variants for a particular segment either because it run out of memory or exceeded the allocated runtime of twelve hours, the interval was subdivided into smaller segments (10 kb).

Our implementation of *SAMtools mpileup* (version 1.8) [57] was run in multi-sample mode to calculate genotype likelihoods from the aligned reads for all samples simultaneously. The parameters *-E* and *–t* were used to recalculate (and apply) base alignment quality and produce per-sample genotype annotations, respectively. Next, the estimated genotype likelihoods were converted into genotypes using *BCFtools call* using the *-v* and *-m* flags to output variant sites only and permit that sites may have more than two alternative alleles, respectively.

We implemented all pipelines using *Snakemake* (version 5.2.0) [58]. The scripts for the pipelines are available via *GitHub* repository (https://github.com/danangcrysnanto/Graph-genotyping-paper-pipelines).

Sequence variant filtering and genotype refinement

The *GATK VariantFiltration* module was used to parse and filter the raw VCF files. Quality control on the raw sequencing variants and genotypes was applied according to well established guidelines that were recommended for each variant caller. Variants that were identified using *GATK* were retained if they met the following criteria: QualByDepth (QD) > 2.0, FisherStrand < 60, RMSMappingQuality (MQ) > 40, MappingQualityRankSumTest (MQRankSum) > 12.5, ReadPosRankSumTest (ReadPosRankSum) > -8.0 (SNPs) and QD > 2.0, FS < 200, ReadPosRankSum > 20 (indels). For the variants identified using SAMtools, the thresholds that have been applied by the 1000 bull genomes project [5] were considered to remove variants with indication of low quality. Variants were retained if they met the following criteria: QUAL > 20, MQ > 30, ReadDepth (DP) > 10, DP < median(DP) + 3*mean(DP). Moreover, SNPs were removed from the data if they had the same positions as the starting position of an indels. The output of Graphtyper was filtered to only include variants that met the criteria recommended by Eggertsson *et al.* (2017): ABHet < 0.0 | ABHet > 0.33, ABHom < 0.0 | ABHom > 0.97, MaxAASR > 0.4, and MQ > 30.

We used *Beagle* (version 4.1) [59] to improve the raw sequence variant genotype quality and impute missing genotypes. The genotype likelihood (*gl*) mode of *Beagle* was applied to infer missing and modify existing genotypes based on the phred-scaled likelihood (*PL*) of all other genotypes.

Evaluation of sequence variant genotyping

To ensure consistent variant representation across the different sequence variant genotyping methods evaluated, we applied the *vt normalize* software (version 0.5) [60]. Normalized variants are parsimonious (i.e., represented by as few nucleotides as possible and left aligned [60]). The number of variants detected and Ti/Tv ratios were calculated using *vt peek* [60] and *BCFtools stats* [57]. The intersection of variants that were common to the tools evaluated was calculated and visualized using *BCFtools isec* [57] and the UpSet R package [61], respectively.

Mendelian inconsistencies were calculated as the proportion of variants showing opposing homozygous genotypes in nine parent-offspring pairs that were included among the 49 sequenced animals. For this comparison, we considered only sites where the genotypes of both sire and son were not missing.

The sequenced animals of this study were also genotyped using the Illumina BovineHD Bead Chip comprising 777,962 SNPs. In order to assess the quality of sequence variant genotyping, the genotypes detected by the different variant calling methods were compared to the microarray-derived variants in terms of genotype concordance, non-reference sensitivity and non-reference discrepancy [23, 38], see **Additional file 4 Figure S3** for more details on the metrics.

Computing environment and statistical analysis

All computations were performed on the ETH Zurich Leonhard Open Cluster with access to multiple nodes equipped with 18 cores Intel Xeon E5-2697v4 processors (base frequency rated at 2.3 GHz) and 128 GB of random-access memory. Unless otherwise stated, the R (version 3.3.3) software environment [62] was used for statistical analyses and ggplot2 (version 3.0.0) [63] was used for data visualisation.

Declarations

Animal ethics statement

Not applicable

Author contributions

Analyzed data: CD HP; Wrote the software pipelines: CD; Generated sequencing data: CW; Wrote the paper: CD HP; Read and approved the final version of the paper: CD CW HP

Acknowledgements

We thank Braunvieh Schweiz for providing pedigree and genotype data of Original Braunvieh cattle. Semen samples of the sequenced bulls were provided by Swissgenetics. Sequencing of the Original Braunvieh bulls was partially funded from the Federal Office for Agriculture (FOAG), Bern, Switzerland. The sequence data were partly generated at the Functional Genomics Center Zurich.

Availability of supporting data

The scripts for three pipelines are available via *GitHub* repository (https://github.com/danangcrysnanto/Graph-genotyping-paper-pipelines). Sequencing read data of 49 Original Braunvieh bulls are available at the European Nucleotide Archive (ENA) (http://www.ebi.ac.uk/ena) under primary accession PRJEB28191.

References

1. Hoff JL, Decker JE, Schnabel RD, Taylor JF. Candidate lethal haplotypes and causal mutations in Angus cattle. BMC Genomics. 2017;18. doi:10.1186/s12864-017-4196-2.

2. Jansen S, Aigner B, Pausch H, Wysocki M, Eck S, Benet-Pagès A, et al. Assessment of the genomic variation in a cattle population by re-sequencing of key animals at low to medium coverage. BMC Genomics. 2013;14:446. doi:10.1186/1471-2164-14-446.

3. Stothard P, Liao X, Arantes AS, De Pauw M, Coros C, Plastow GS, et al. A large and diverse collection of bovine genome sequences from the Canadian Cattle Genome Project. Gigascience. 2015. doi:10.1186/s13742-015-0090-5.

4. Boussaha M, Michot P, Letaief R, Hozé C, Fritz S, Grohs C, et al. Construction of a large collection of small genome variations in French dairy and beef breeds using whole-genome sequences. Genet Sel Evol. 2016;48:87. doi:10.1186/s12711-016-0268-z.

5. Daetwyler HD, Capitan A, Pausch H, Stothard P, Van Binsbergen R, Brøndum RF, et al. Whole-genome sequencing of 234 bulls facilitates mapping of monogenic and complex traits in cattle. Nat Genet. 2014;46:858–65. doi:10.1038/ng.3034.

6. Pausch H, Emmerling R, Gredler-Grandl B, Fries R, Daetwyler HD, Goddard ME. Meta-analysis of sequence-based association studies across three cattle breeds reveals 25 QTL for fat and protein percentages in milk at nucleotide resolution. BMC Genomics. 2017;18:853. doi:10.1186/s12864-017-4263-8.

7. Bouwman AC, Daetwyler HD, Chamberlain AJ, Ponce CH, Sargolzaei M, Schenkel FS, et al. Meta-analysis of genome-wide association studies for cattle stature identifies common genes that regulate body size in mammals. Nat Genet. 2018;50:362–7. doi:10.1038/s41588-018-0056-5.

8. Raymond B, Bouwman AC, Schrooten C, Houwing-Duistermaat J, Veerkamp RF. Utility of whole-genome sequence data for across-breed genomic prediction. Genet Sel Evol. 2018;50:27. doi:10.1186/s12711-018-0396-8.

9. Nielsen R, Paul JS, Albrechtsen A, Song YS. Genotype and SNP calling from next-generation sequencing data. Nat Rev Genet. 2011;12:443–51. doi:10.1038/nrg2986.

10. Guo Y, Ye F, Sheng Q, Clark T, Samuels DC. Three-stage quality control strategies for DNA re-sequencing data. Brief Bioinform. 2013;15:879–89. doi:10.1093/bib/bbt069.

11. Goodwin S, McPherson JD, McCombie WR. Coming of age: ten years of next-generation sequencing technologies. Nat Rev Genet. 2016;17:333–51. doi:10.1038/nrg.2016.49.

12. Pfeifer SP. From next-generation resequencing reads to a high-quality variant data set. Heredity. 2017;118:111–24. doi:10.1038/hdy.2016.102.

13. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods. 2012;9:357–9. doi:10.1038/nmeth.1923.

14. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform. Bioinformatics. 2009;25:1754–60. doi:10.1093/bioinformatics/btp324.

15. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence Alignment/Map format and SAMtools. Bioinformatics. 2009;25:2078–9. doi:10.1093/bioinformatics/btp352.

16. McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, et al. The Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data. Genome Res. 2010;20:1297–303. doi:10.1101/gr.107524.110.

17. Van der Auwera GA, Carneiro MO, Hartl C, Poplin R, del Angel G, Levy-Moonshine A, et al. From fastQ data to high-confidence variant calls: The genome analysis toolkit best practices pipeline. Curr Protoc Bioinforma. 2013; SUPL.43:11.10.1-11.10.33. doi:10.1002/0471250953.bi1110s43.

18. Poplin R, Ruano-Rubio V, Depristo MA, Fennell TJ, Carneiro MO, Van Der Auwera GA, et al. Scaling accurate genetic variant discovery to tens of thousands of samples. bioRxiv. 2017. doi:10.1101/201178.

19. Baes CF, Dolezal MA, Koltes JE, Bapst B, Fritz-Waters E, Jansen S, et al. Evaluation of variant identification methods for whole genome sequencing data in dairy cattle. BMC Genomics. 2014;15:948. doi:10.1186/1471-2164-15-948.

20. Liu X, Han S, Wang Z, Gelernter J, Yang B-Z. Variant Callers for Next-Generation Sequencing Data: A Comparison Study. PLoS One. 2013;8:e75619. doi:10.1371/journal.pone.0075619.

21. Cheng AY, Teo YY, Ong RTH. Assessing single nucleotide variant detection and genotype calling on whole-genome sequenced individuals. Bioinformatics. 2014;30:1707–13. doi:10.1093/bioinformatics/btu067.

22. Kumar P, Al-Shafai M, Al Muftah W, Chalhoub N, Elsaid MF, Aleem A, et al. Evaluation of SNP calling using single and multiple-sample calling algorithms by validation against array base

genotyping and Mendelian inheritance. BMC Res Notes. 2014;7:747. doi:10.1186/1756-0500-7-747.

23. DePristo MA, Banks E, Poplin R, Garimella K V., Maguire JR, Hartl C, et al. A framework for variation discovery and genotyping using next-generation DNA sequencing data. Nat Genet. 2011;43:491–8. doi:10.1038/ng.806.

24. Zook JM, Catoe D, McDaniel J, Vang L, Spies N, Sidow A, et al. Extensive sequencing of seven human genomes to characterize benchmark reference materials. Sci Data. 2016;3:160025. doi:10.1038/sdata.2016.25.

25. Li H, Wren J. Toward better understanding of artifacts in variant calling from high-coverage samples. Bioinformatics. 2014;30:2843–51. doi:10.1093/bioinformatics/btu356.

26. Malomane DK, Reimer C, Weigend S, Weigend A, Sharifi AR, Simianer H. Efficiency of different strategies to mitigate ascertainment bias when using SNP panels in diversity studies. BMC Genomics. 2018;19:22. doi:10.1186/s12864-017-4416-9.

27. Degner JF, Marioni JC, Pai AA, Pickrell JK, Nkadori E, Gilad Y, et al. Effect of read-mapping biases on detecting allele-specific expression from RNA-sequencing data. Bioinformatics. 2009;25:3207–12. doi:10.1093/bioinformatics/btp579.

28. Brandt DYC, Aguiar VRC, Bitarello BD, Nunes K, Goudet J, Meyer D. Mapping Bias Overestimates Reference Allele Frequencies at the HLA Genes in the 1000 Genomes Project Phase I Data. G3;Genes|Genomes|Genetics. 2015;5:931–41. doi:10.1534/g3.114.015784.

29. Dilthey A, Cox C, Iqbal Z, Nelson MR, McVean G. Improved genome inference in the MHC using a population reference graph. Nat Genet. 2015;47:682–8. doi:10.1038/ng.3257.

30. Fakhro KA, Staudt MR, Ramstetter MD, Robay A, Malek JA, Badii R, et al. The Qatar genome: a population-specific tool for precision medicine in the Middle East. Hum Genome Var. 2016;3:16016. doi:10.1016/j.ejr.2016.03.001.

31. Shi L, Guo Y, Dong C, Huddleston J, Yang H, Han X, et al. Long-read sequencing and de novo assembly of a Chinese genome. Nat Commun. 2016;7:12065. doi:10.1038/ncomms12065.

32. Ameur A, Che H, Martin M, Bunikis I, Dahlberg J, Höijer I, et al. De Novo Assembly of Two Swedish Genomes Reveals Missing Segments from the Human GRCh38 Reference and Improves Variant Calling of Population-Scale Sequencing Data. Genes. 2018;9:486. doi:10.3390/genes9100486.

33. Rakocevic G, Semenyuk V, Spencer J, Browning J, Johnson I, Arsenijevic V, et al. Fast and Accurate Genomic Analyses using Genome Graphs. bioRxiv. 2018. doi:10.1101/194530.

34. Eggertsson HP, Jonsson H, Kristmundsdottir S, Hjartarson E, Kehr B, Masson G, et al. Graphtyper enables population-scale genotyping using pangenome graphs. Nat Genet. 2017;49:1654–60. doi:10.1038/ng.3964.

35. Novak AM, Hickey G, Garrison E, Blum S, Connelly A, Dilthey A, et al. Genome Graphs. bioRxiv. 2017. doi:10.1101/101378.

36. Garrison E, Sirén J, Novak AM, Hickey G, Eizenga JM, Dawson ET, et al. Variation graph toolkit improves read mapping by representing genetic variation in the reference. Nat Biotechnol. 2018;36:875–81. doi:10.1038/nbt.4227.

37. Sibbesen JA, Maretty L, Krogh A. Accurate genotyping across variant classes and lengths using variant graphs. Nat Genet. 2018;50:1054–9. doi:10.1038/s41588-018-0145-5.

38. Linderman MD, Brandt T, Edelmann L, Jabado O, Kasai Y, Kornreich R, et al. Analytical validation of whole exome and whole genome sequencing for clinical applications. BMC Med Genomics. 2014;7:20. doi:10.1186/1755-8794-7-20.

39. Sims D, Sudbery I, Ilott NE, Heger A, Ponting CP. Sequencing depth and coverage: key considerations in genomic analyses. Nat Rev Genet. 2014;15:121–32. doi:10.1038/nrg3642.

40. Fragoso CA, Heffelfinger C, Zhao H, Dellaporta SL. Imputing genotypes in biallelic populations from low-coverage sequence data. Genetics. 2016;202:487–95. doi:10.1534/genetics.115.182071.

41. Bilton TP, McEwan JC, Clarke SM, Brauning R, van Stijn TC, Rowe SJ, et al. Linkage disequilibrium estimation in low coverage high-throughput sequencing data. Genetics. 2018;209:389–400. doi:10.1534/genetics.118.300831.

42. Sirén J, Garrison E, Novak AM, Paten B, Durbin R. Haplotype-aware graph indexes. arXiv. 2018. doi:10.4230/LIPIcs.WABI.2018.4.

43. Novak AM, Garrison E, Paten B. A graph extension of the positional Burrows–Wheeler transform and its applications. Algorithms Mol Biol. 2017;12:18. doi:10.1186/s13015-017-0109-9.

44. Stafuzza NB, Zerlotini A, Lobo FP, Yamagishi MEB, Chud TCS, Caetano AR, et al. Single nucleotide variants and InDels identified from whole-genome re-sequencing of Guzerat, Gyr, Girolando and Holstein cattle breeds. PLoS One. 2017;12:e0173954. doi:10.1371/journal.pone.0173954.

45. Patel ZH, Kottyan LC, Lazaro S, Williams MS, Ledbetter DH, Tromp hbGerard, et al. The struggle

to find reliable results in exome sequencing data: filtering out Mendelian errors. Front Genet. 2014;5:16. doi:10.3389/fgene.2014.00016.

46. Koren S, Harhay GP, Smith TPL, Bono JL, Harhay DM, Mcvey SD, et al. Reducing assembly complexity of microbial genomes with single-molecule sequencing. Genome Biol. 2013;14:R101. doi:10.1186/gb-2013-14-9-r101.

47. Pausch H, MacLeod IM, Fries R, Emmerling R, Bowman PJ, Daetwyler HD, et al. Evaluation of the accuracy of imputed sequence variant genotypes and their utility for causal variant detection in cattle. Genet Sel Evol. 2017;49:24. doi:10.1186/s12711-017-0301-x.

48. Liu GE, Ventura M, Cellamare A, Chen L, Cheng Z, Zhu B, et al. Analysis of recent segmental duplications in the bovine genome. BMC Genomics. 2009;10:571. doi:10.1186/1471-2164-10-571.

49. Bickhart DM, Hou Y, Schroeder SG, Alkan C, Cardone MF, Matukumalli LK, et al. Copy number variation of individual cattle genomes using next-generation sequencing. Genome Res. 2012;22:778–90. doi:10.1101/gr.133967.111.

50. Guo Y, Dai Y, Yu H, Zhao S, Samuels DC, Shyr Y. Improvements and impacts of GRCh38 human reference on high throughput sequencing data analysis. Genomics. 2017;109:83–90. doi:10.1016/j.ygeno.2017.01.005.

51. Chen S, Zhou Y, Chen Y, Gu J. Fastp: An ultra-fast all-in-one FASTQ preprocessor. Bioinformatics. 2018;34:i884–90. doi:10.1093/bioinformatics/bty560.

52. Zimin A V, Delcher AL, Florea L, Kelley DR, Schatz MC, Puiu D, et al. A whole-genome assembly of the domestic cow, Bos taurus. Genome Biol. 2009;10:R42. doi:10.1186/gb-2009-10-4-r42.

53. Faust GG, Hall IM. SAMBLASTER: Fast duplicate marking and structural variant read extraction. Bioinformatics. 2014;30:2503–5. doi:10.1093/bioinformatics/btu314.

54. Tarasov A, Vilella AJ, Cuppen E, Nijman IJ, Prins P. Sambamba: fast processing of NGS alignment formats. Bioinformatics. 2015;31:2032–4. doi:10.1093/bioinformatics/btv098.

55. Pedersen BS, Quinlan AR. Mosdepth: Quick coverage calculation for genomes and exomes. Bioinformatics. 2018;34:867–8. doi:10.1093/bioinformatics/btx699.

56. Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, et al. The variant call format and VCFtools. Bioinformatics. 2011;27:2156–8. doi:10.1093/bioinformatics/btr330.

57. Li H. A statistical framework for SNP calling, mutation discovery, association mapping and population genetical parameter estimation from sequencing data. Bioinformatics. 2011;27:2987–93. doi:10.1093/bioinformatics/btr509.

58. Köster J, Rahmann S. Snakemake--a scalable bioinformatics workflow engine. Bioinformatics. 2012;28:2520–2. doi:10.1093/bioinformatics/bts480.

59. Browning BL, Browning SR. Genotype Imputation with Millions of Reference Samples. Am J Hum Genet. 2016;98:116–26. doi:10.1016/j.ajhg.2015.11.020.

60. Tan A, Abecasis GR, Kang HM. Unified representation of genetic variants. Bioinformatics. 2015;31:2202–4. doi:10.1093/bioinformatics/btv112.

61. Lex A, Gehlenborg N, Strobelt H, Vuillemot R, Pfister H. UpSet: Visualization of intersecting sets. IEEE Trans Vis Comput Graph. 2014;20:1983–92. doi:10.1109/TVCG.2014.2346248.

62. R Core Team. R: A Language and Environment for Statistical Computing. 2017. https://www.r-project.org.

63. Wickham H. Elegant Graphics for Data Analysis. Springer-Verlag New York; 2016.

Additional files

Additional file 1 Figure S1

File format: pdf

Description: Concordance of heterozygous and alternate homozygous genotypes. (a) The average concordance across 49 Original Braunvieh cattle and the concordance for different sequencing depths based on the (b) raw and (c) imputed datasets.

Additional file 2 Table S1

File format: xlsx

Description: Twelve 1 Mb regions where *Graphtyper* initially failed to genotype sequence variants because the algorithm either ran out of memory or exceeded the allocated runtime (12 hours). *Graphtyper* eventually produced genotypes for the sequence variants when these regions were re-run in 10 kb segments.

Additional file 3 Figure S2

File format: tiff

Description: Accuracy and sensitivity of sequence variant genotyping on bovine chromosome 25 from a variation-aware genome graph that incorporated 2,143,417 dbSNP variants as prior known variants.

Additional file 4 Figure S3

File format: tiff

Description: Properties of the different metrics used for the evaluation of sequence variant genotyping accuracy. The metrics were calculated using the sum of the red cells as numerator and the cells within the green frame as denominator.