Real-time projections of epidemic transmission and estimation of vaccination impact during an Ebola virus disease outbreak in Northeastern Democratic Republic of Congo

Lee Worden[1], Rae Wannier[1,2], Nicole A. Hoff[3], Kamy Musene[3], Bernice Selo[4], Mathias Mossoko[4], Emile Okitolonda-Wemakoy[5], Jean Jacques Muyembe-Tamfum[6], George W. Rutherford[2], Thomas M. Lietman[1,2], Anne W. Rimoin[3], Travis C. Porco[1,2], J. Daniel Kelly[1,2*]

1 F. I. Proctor Foundation, University of California, San Francisco (UCSF), San Francisco, CA, USA
2 School of Medicine, UCSF, San Francisco, CA, USA
3 School of Public Health, University of California, Los Angeles, Los Angeles, CA, USA
4 Ministry of Health, Kinshasa, Democratic Republic of Congo
5 School of Public Health, University of Kinshasa, Kinshasa, Democratic Republic of Congo
6 National Institute of Biomedical Research, Kinshasa, Democratic Republic of Congo

* Corresponding author: Dan.Kelly@ucsf.edu

Abstract

Background: As of February 25, 2019, 875 cases of Ebola virus disease (EVD) were reported in North Kivu and Ituri Provinces, Democratic Republic of Congo. Since the beginning of October, the outbreak has largely shifted into regions in which active armed conflict has occurred, and in which EVD cases and their contacts have been difficult for health workers to reach. We used available data on the current outbreak with case-count time series from prior outbreaks to project the short-term and long-term course of the outbreak.

Methods: For short- and long-term projections, we modeled Ebola virus transmission using a stochastic branching process that assumes gradually quenching transmission rates estimated from past EVD outbreaks, with outbreak trajectories conditioned on agreement with the course of the current outbreak, and with multiple levels of vaccination coverage. We used two regression models to estimate similar projection periods. Short- and long-term projections were estimated using negative binomial autoregression and Theil-Sen regression, respectively. We also used Gott's rule to estimate a baseline minimum-information projection. We then constructed an ensemble of forecasts to be compared and recorded for future evaluation against final outcomes. From August 20, 2018 to February 25, 2019, short-term model projections were validated against known case counts.

Results: During validation of short-term projections, from one week to four weeks, we found models consistently scored higher on shorter-term forecasts. Based on case counts as of February 25, the stochastic model projected a median case count of 933 cases by February 18 (95% prediction interval: 872–1054) and 955 cases by March 4 (95% prediction interval: 874–1105), while the auto-regression model projects median case counts of 889 (95% prediction interval: 876–933) and 898 (95% prediction interval: 877–983) cases for those dates, respectively. Projected median final counts range from 953 to 1,749. Although the outbreak is already larger than all past Ebola outbreaks other than the 2013–2016 outbreak of over 26,000 cases, our models do not project that it is likely to grow to that scale. The stochastic model estimates that vaccination coverage in this outbreak is lower than reported in its trial setting in Sierra Leone.

Conclusions: Our projections are concentrated in a range up to about 300 cases beyond those already reported. While a catastrophic outbreak is not projected, it is not ruled out, and prevention and vigilance are warranted. Prospective validation of our models in real time allowed us to generate more accurate short-term forecasts, and this process may prove useful for future real-time short-term forecasting. We estimate that transmission rates are higher than would be seen under target levels of 62% coverage due to contact tracing and vaccination, and this model estimate may offer a surrogate indicator for the outbreak response challenges.

Author summary

As of February 25, 2019, 875 cases of Ebola virus disease (EVD) were reported in North Kivu and Ituri Provinces, Democratic Republic of Congo. Since the beginning of October 2018, the outbreak has largely shifted into regions in which active armed conflict has been reported, and in which EVD cases and their contacts have been difficult for health workers to reach. We used an ensemble of models to estimate EVD transmission rates and to forecast the short- and long-term course of the outbreak. Our models project that a final size of roughly up to 300 additional cases is most likely, and estimate that transmission rates are higher than would be seen under optimal levels of contact tracing and vaccination. While a catastrophic outbreak is not projected, is it not ruled out, and prevention and vigilance are warranted.

Introduction 1

On August 1, 2018, the World Health Organization (WHO) announced a new outbreak 2 of Ebola virus disease (EVD) in North Kivu Province, Democratic Republic of Congo 3 (DRC) [1]. Epidemiological investigations traced EVD cases back to the week of April 4 30 and identified the initial epicenter to be Mabalako. This region of DRC (Fig 1) has 5 over five million inhabitants, some of whom suffer from armed conflict, humanitarian 6 crisis, and displacement to the bordering countries of Uganda and Rwanda [2]. Since 7 the outbreak began, its magnitude has surpassed all other EVD outbreaks in DRC, 8 becoming the second largest in history to date. As of February 25, 2019, 875 EVD cases 9 had been reported (65 probable and 810 confirmed), and the Ministry of Health of DRC, 10 the World Health Organization, and other organizations were responding to the Ebola 11 outbreak [3]. 12

This is the first EVD outbreak to occur in an armed conflict zone, and this new 13 context has created challenges to the outbreak response [4]. Uptake of traditional 14 control measures such as contact tracing and safe burials has varied with the 15 community, even with the rapid deployment and widespread availability of experimental 16 vaccines and therapeutics [3]. Our understanding of the impact of scientific advances on 17 EVD transmission has been evolving; there has been an underlying, untested 18

Fig 1. Map of health districts in which confirmed and/or probable cases have occurred in Northeastern DRC, with zones in which active conflict has occurred identified by color. Data sourced from Référentiel Géographique Commun [7], made available under the Open Data Commons Open Database License.

assumption that vaccination may reduce epidemic transmission [5]. Mathematical modeling of simulated EVD epidemics suggested that vaccination coverage as low as 40% in the general population and 95% in healthcare workers (HCWs) may avert another epidemic similar to size of the West African one [6]. Studies have yet to estimate the levels of vaccination coverage associated with epidemic decline. An even greater potential problem in difficult-to-access outbreak areas is ascertainment of vaccination coverage among contacts and contacts of contacts. Mathematical models are needed that can produce relatively accurate forecasts in the setting of such unknown but important outbreak response metrics.

During an Ebola outbreak, real-time forecasting has the potential to support decision-making and allocation of resources, but highly accurate forecasts have proven difficult for Ebola [8,9] as well as other diseases [10–13]. Highly accurate forecasts of small, noisy outbreaks may be a fundamentally elusive ideal [14]. Previous work has found that probabilistic forecasts can have relatively high accuracy within a few weeks, though they become less useful as time horizons grow longer [15]. Thus, short-term

forecasts may provide useful information for response organizations. 34

In this paper, we applied a suite of independent methods to estimate short- and 35
long-term projections of future EVD case counts in northeastern DRC in real time. 36
Given the unique context of the current EVD outbreak, we validated short-term 37
projections before generating projections on February 25. One of our models also 38
estimated vaccination coverage. It is our hope that these projections offer insight into 39
the outbreak trajectory and vaccination coverage, particularly in the setting of armed 40
conflict. 41

Methods 42

We used four independent methods to generate projections of future case counts: (1) a 43
stochastic simulation model, (2) a negative binomial auto-regression model, (3) a 44
Theil-Sen regression model, and (4) a final size projection based on Gott's rule. The 45
stochastic simulation model was calibrated to time-dependent transmission rates 46
measured from past EVD outbreaks and constrained to the observed partial trajectory 47
of the current outbreak, extending a model used in previous work [16]. This model 48
generated short- and long-term projections. The negative binomial auto-regression 49
model produced short-term projections based on case counts from the outbreak to date 50
and prior outbreaks. The Theil-Sen regression model predicted final outbreak size based 51
on prior outbreaks. We used a probability mass function derived from Gott's rule as a 52
minimum-information projection of final outbreak size, using only the assumption that 53
the proportion of the outbreak observed so far is entirely unknown. From these, we 54
assembled an ensemble of projections to be compared and recorded for future evaluation 55
against final outcomes. 56

Data sources 57

Data on the current outbreak were collected from the WHO website in real time as 58
updated information was published [3]. A cumulative case count of probable and 59
confirmed cases was extracted to be consistent with the best knowledge at the time. 60
Snapshots of the table of case counts were kept as of multiple dates (Fig S1 in S1 61
Supporting Information), to be used in retrospective scoring of model projections 62

against subsequently known counts. Although the epidemic was officially reported on $_{63}$
August 1 as a cluster of cases occurring in June and July, seven sporadic early cases $_{64}$
from April and May were subsequently linked to the current outbreak and were added $_{65}$
to later case totals [3]. This additional knowledge was added retrospectively to the time $_{66}$
series of cumulative case counts only for predictions made for days on or after $_{67}$
September 15, when these cases were officially linked to the current outbreak. $_{68}$

Stochastic model $_{69}$

We modeled Ebola virus (EBOV) transmission using a stochastic branching process $_{70}$
model, parameterized by transmission rates estimated from the dynamics of prior EVD $_{71}$
outbreaks and conditioned on agreement with reported case counts from the 2018 EVD $_{72}$
outbreak to date. We incorporated high and low estimates of vaccination coverage into $_{73}$
this model. We used this model to generate a set of probabilistic projections of the size $_{74}$
of simulated outbreaks in the current setting. This model is similar to one described in $_{75}$
previous work [16], with the addition of a smoothing step allowing for a continuum of $_{76}$
transmission rates interpolated between those estimated from prior outbreaks. $_{77}$

On the assumption that past outbreaks provide a basis for projection of the current $_{78}$
outbreak, we used estimates of transmission rates from past EVD outbreaks to $_{79}$
parameterize simulations of the current outbreak. To estimate the reproduction number $_{80}$
R in past outbreaks as a function of the number of days from the beginning of the $_{81}$
outbreak, we included reported cases by date from fourteen prior outbreaks (Table S1 in $_{82}$
S1 Supporting Information), [17–30]. To reflect the Ebola response system in DRC $_{83}$
during what is now its tenth outbreak, the first historical outbreak reported in each $_{84}$
country was excluded (e.g. the 1976 outbreak in Yambuko, DRC), as there is a $_{85}$
difference in the Ebola response system as well as community sensitization to EVD $_{86}$
following a country's first outbreak. We used the Wallinga-Teunis technique to estimate $_{87}$
R for each case and therefore for each reporting date in these outbreaks [31]. The serial $_{88}$
interval distribution used for this estimation was a gamma distribution with a mean of $_{89}$
14.5 days and a standard deviation of 5 days, with intervals rounded to the nearest $_{90}$
whole number of days, consistent with the understanding that the serial interval of EVD $_{91}$
cases ranges from 3 to 36 days with mean 14 to 15 days. $_{92}$

Transmission rates estimated by day in these outbreaks tend to decline from initially high to eventually low values, though they may display substantial fluctuations. This "quenching" of transmission may be driven by formal interventions such as quarantine, or by informal changes in individuals' behavior in response to the disease or by depletion of uninfected contacts of infective individuals, or by other causes. We incorporated this pattern into our model by estimating an initial reproduction number R_{initial} and quenching rate τ for each outbreak by fitting an exponentially quenched curve to the outbreak's estimates of R by day d (Fig S2 in S1 Supporting Information), and used these pairs of parameters, one from each past outbreak, to construct a joint distribution of initial reproduction numbers and quenching rates for outbreak simulation.

We simulated EBOV transmission using a stochastic branching process model in which the number of secondary cases caused by any given primary case is drawn from a negative binomial distribution, whose mean is the reproduction number R as a function of day of the outbreak, and variance is controlled by a dispersion parameter k [32, 33]. All transmission events were assumed to be independent. The interval between date of detection of each primary case and that of each of its secondary cases is assumed gamma distributed with mean 14.5 days and standard deviation 5 days, rounded to the nearest whole number of days, as above.

We used the $(R_{\text{initial}}, \tau)$ pairs estimated from prior outbreaks to provide R values for simulation. R_{initial} values were sampled uniformly from the range of values estimated from past outbreaks. We applied a linear regression to the values of R_{initial} and $\log(\tau)$ estimated for prior outbreaks and used the resulting regression line to assign a mean τ to each R, used with the residual variance of $\log(\tau)$ as a distribution from which to sample τ values for simulation given R_{initial}. Note that the range of fast and slow quenching scenarios modeled in this way is not limited to the exact combinations estimated from past outbreaks, but extends over a continuous distribution that includes those values. The parameters R_{initial} and τ sampled in this way, together with three values of the dispersion parameter k, 0.3, 0.5, and 0.7, consistent with transmission heterogeneity observed in past Ebola outbreaks, were used to generate simulated outbreaks.

This model generated randomly varying simulated outbreaks with a range of case counts per day. The outbreak was assumed to begin with a single case. The simulation was run multiple times, each instance producing a proposed epidemic trajectory,

generated by the above branching process with the given parameters R_{initial}, τ, and k, and these were then filtered by discarding all proposed trajectories except those whose cumulative case counts matched known counts of the current 2018 EVD outbreak on known dates. In earlier, smaller data sets we filtered against all reported case counts, while in later, more complete data sets we used a thinned series of case counts for filtering, for computational tractability, by selecting five case counts evenly spaced in the data set plus the final case count (Fig S1 in S1 Supporting Information). The filtration required an exact match of the first target value, and at subsequent target dates accepted epidemics within a number of cases more or less than each recorded value. On the earlier data sets in which the beginning dates of the epidemic were unknown, the first target value was allowed to match on any day, and subsequent target dates were assigned relative to that day.

In order to produce model outbreak trajectories consistent with the case counts reported more recently in the outbreak, it was necessary to make the filtering step of the model more tolerant to variation in counts in order to accommodate the rapidly rising count. This was because higher transmission rates in late September and early October were necessary to generate case counts of that size than were consistent with the earlier counts.

Thus this model embodies a set of assumptions that transmission rates are overall gradually declining from the start of the outbreak to its end, though possibly with a high level of variability in transmission rate between cases and between simulations. When the tolerance of the filter on case counts is small, quenching of transmission through time must closely track case counts, while when tolerance is high, fluctuations in the rate of generation of new cases can reflect a pattern of ongoing quenching of transmission more loosely and on the long term, while being more insensitive to short-term up and down fluctuations in transmission rates reflected in the true case counts.

We varied the tolerance as the data set became more complete to maintain a roughly fixed rate of generation of filtered trajectories. As larger tolerances became necessary, in data sets from after October, we introduced one further distinction: while it is possible for cumulative case counts to decrease as inaccurately classified cases are removed from the counts, due to the precision of the labeling of cases probable and confirmed we

expect this to happen rarely, so we limit the tolerance of matching to only at most 15 ₁₅₇ cases below the reported count regardless of the tolerance of counts above the reported ₁₅₈ count. This limit on underestimates was applied only to analysis of data sets from later ₁₅₉ than October 13, to preserve unaltered the projection methods we reported in a ₁₆₀ preprint of this paper [34]. Filtering tolerances were as follows: when using the August ₁₆₁ 20 data set, we allowed a tolerance of 4 cases more or less than each target count; for ₁₆₂ August 27 and September 5, 6 cases; for September 15, 10 cases; for October 7, 12 cases; ₁₆₃ for October 13, 17 cases; for November 1, 41 cases; for November 20, 55 cases; for ₁₆₄ January 6, 2019, 75 cases; and for February 25, 150 cases. This one-step particle ₁₆₅ filtering technique produced an ensemble of model outbreaks, filtered on agreement with ₁₆₆ the recorded trajectory of the outbreak to date. This filtered ensemble was then used to ₁₆₇ generate projections of the outbreak in the short term (one week, two weeks, four weeks, ₁₆₈ and eight weeks) and long term (final size) [35]. ₁₆₉

To model vaccination coverage with respect to total transmission (unreported and ₁₇₀ reported), we multiplied the estimate of vaccine effectiveness by low and high estimates ₁₇₁ of reported cases. In a ring vaccination study at the end of the West Africa outbreak, ₁₇₂ the overall estimated recombinant vesicular stomatitis virus (rVSV)-vectored vaccine ₁₇₃ efficacy was 100% and vaccine effectiveness was 64.6% in protecting all contacts and ₁₇₄ contacts of contacts from EVD in the randomized clusters, including unvaccinated ₁₇₅ cluster members [36]. We used estimates of vaccine effectiveness in our stochastic model. ₁₇₆ The ring vaccination study found the vaccine to be effective against cases with onset ₁₇₇ dates 10 days or more from the date of vaccine administration, so we modeled the ₁₇₈ vaccination program as a proportionate reduction in the number of new cases with ₁₇₉ onsets 10 days or more after the program start date. ₁₈₀

We used past estimates of the proportion of unreported cases to estimate the ₁₈₁ proportion of exposed individuals not covered by the vaccination process. Based on a ₁₈₂ Sierra Leonean study from the 2013–2016 outbreak [37], we estimated the proportion of ₁₈₃ reported cases (out of all known cases) in DRC to range from a low of 68% to a high of ₁₈₄ 96%, as extremes of the best known range of estimates, to be evaluated on their fit to ₁₈₅ data. Given these low and high estimates of reported cases and the estimate of vaccine ₁₈₆ effectiveness, a low estimate of vaccination program coverage was 44% (68% × 64.6%) ₁₈₇ and a high estimate of vaccination program coverage was 62% (96% × 64.6%). ₁₈₈

Vaccination was included in the simulation using a start date of August 14, 2018, estimated from dates available from situation reports [3].

To generate forecasts from each data set, 320 to 330 outbreaks were collected that passed the step of filtering on approximate agreement with DRC case counts. The simulated outbreaks that were retained after filtering were continued until they generated no further cases. Rare outlying simulated outbreaks that exceeded 300,000 cases were capped at the first value reached above that number, to limit unnecessary computation. We used this ensemble to derive a distribution of final outbreak sizes, and of cumulative counts at specific forecasting dates. Projection distributions were constructed using kernel density estimation with leave-one-out cross-validation to determine bandwidth, using a log-normal kernel for final sizes, due to the extended tail of the values, and a normal kernel for all other estimates. We calculated median values and 95% prediction intervals using the 2.5th and 97.5th percentiles of simulated outbreak size.

The frequencies of occurrence of the three vaccination coverage scenarios modeled (zero, low, and high coverage) among the simulated outbreaks accepted by the stochastic model's filtering step were used to estimate likelihoods of the three scenarios. The frequencies of $(R_{\mathrm{initial}}, \tau)$ pairs selected by the filtering process were similarly recorded as an estimate of the likelihood of those transmission rate parameters.

We provide a detailed report of the parameters, simulations, and performance of the stochastic model in the Supplemental Material. This analysis was conducted using R 3.4.2 (R Foundation for Statistical Computing, Vienna, Austria).

Auto-Regression model

A negative binomial autoregressive model was chosen through a validation process to forecast additional new case counts at time points one week, two weeks, four weeks, and eight weeks from the current date. To adjust for disparities in the frequency of case reporting in historic outbreaks, the data were weighted by the inverse square root of the number of observations contributed to the model. Models considered included parameters for historic cumulative case counts (probable and confirmed) at different time points, logs of historic case counts, ratio of historic case counts to try and capture

the trend of the epidemic curve, log(time), and an offset for current case total. When ²¹⁹ historic case counts for specific dates were missing, each missing case count was linearly ²²⁰ interpolated from the two nearest case counts, allowing the model to remain agnostic ²²¹ about the current trend of the epidemic. After model fitting and validation, the final ²²² model chosen was a log-link regression for additional cases on the number of new cases ²²³ identified in the previous two and four weeks and the ratio of these two case counts. ²²⁴

This analysis was conducted using R 3.4.2 (R Foundation for Statistical Computing, ²²⁵ Vienna, Austria). ²²⁶

Theil-Sen regression model ²²⁷

We conducted a simple regression forecast based solely on outbreaks of size 10 or ²²⁸ greater, based on prior outbreaks [17–30]. Nonparametric Theil-Sen regression (R ²²⁹ package `mblm` [38]) was used to project the final outbreak size based on values of the ²³⁰ outbreak size at a specific earlier time. All time series were aligned on the day they ²³¹ reached 10 cases. Finally, we reported the median and 95% central coverage intervals for ²³² the prediction distribution, conditional on the predicted value being no smaller than the ²³³ observed value for each day. More details can be found elsewhere [16]. ²³⁴

This analysis was conducted using R 3.4.2 (R Foundation for Statistical Computing, ²³⁵ Vienna, Austria). ²³⁶

Gott's rule model ²³⁷

Gott's rule assumes we have no special knowledge of our position on the epidemic ²³⁸ curve [39]. Given Y_0 cases reported, assuming a non-informative uniform prior for the ²³⁹ portion α of the epidemic observed to date, the corresponding probability density ²⁴⁰ function for the final size $Y = Y_0/\alpha$ is Y_0/y^2, $Y_0 \leq y$. We constructed a probability ²⁴¹ mass function by assigning all probability density to the whole number of days given by ²⁴² the integer part of each value. We used this probability mass function as a projection of ²⁴³ the final outbreak size. ²⁴⁴

This analysis was conducted using R 3.4.2 (R Foundation for Statistical Computing, ²⁴⁵ Vienna, Austria). ²⁴⁶

Scoring

Models' performance on short-term projections was scored using the natural logarithm   248
of the probability assigned to subsequently known reported case counts.   249

The stochastic and negative binomial auto-regression models were scored based on   250
projections at multiple time points, where available: 251

- Case count 1 week after the last available case count   252

- Case count 2 weeks after the last available case count   253

- Case count 4 weeks after the last available case count   254

- Case count 8 weeks after the last available case count   255

For each projection, we generated an assignment of probability to possible values of   256
multiple quantities. As Ebola situation reports were released before February 25, we   257
generated short-term projections in real time with stochastic and negative binomial   258
auto-regression models. We scored these projections and calibrated the models during   259
the outbreak. Then we used these calibrated models for our projections on February 25.   260
The Theil-Sen and Gott's rule models were not calibrated as the outbreak's final size is   261
not known. Final outbreak size projections generated by the stochastic, Theil-Sen, and   262
Gott's rule models were recorded for future evaluation of their performance.   263

Evaluation of a catastrophic outbreak   264

Because the question has been raised of whether the current outbreak might grow to the   265
scale of the catastrophic West Africa outbreak, we evaluated each model's projected   266
probability of exceeding three large outbreak scenarios of 1,000 cases, 10,000 cases, or   267
the 28,616 cases reported in the West Africa outbreak [30]. Each of these three   268
thresholds is more than double the size of the previous second largest outbreak (n=425,   269
Uganda, 2000) [22].   270

Results   271

When we started performing our short-term forecasts on August 20, 2018, there were 102   272
reported EVD cases in North Kivu and Ituri provinces, DRC. We used our stochastic   273

and auto-regression models to project one-week, two-week, four-week, and eight-week 274
forecasts of outbreak size. As time lapsed, we compared predicted and known outbreak 275
sizes and found a higher probability of accurate forecasts at one week than at eight 276
weeks (Figs 2, 3). Log-likelihood scores typically declined as projection times extended 277
further into the future. These lower-scoring longer-term projections tended to include 278
wider prediction intervals, reflecting less certain outcomes in which less probability was 279
assigned to any one value. The epidemic curve accelerated in early October, and 280
stochastic model forecasts occurring just before that change were especially low scoring, 281
as they failed to anticipate the coming rise in case counts. Subsequent projections took 282
into account the reported acceleration and their performance recovered. 283

Fig 2. Prospective validation of probabilistic projections of auto-regression
and stochastic models, by comparing projected cumulative case counts from past data
to known case counts. Vertical lines indicate known case counts (height not to scale).

After our model validation process was completed, we used the stochastic and 284
auto-regression models to project one-week, two-week, four-week, and eight-week 285
outcomes (Figs 4, 5). We used the Gott's rule and Theil-Sen regression models together 286
with the stochastic model to project final outbreak sizes (Figs 6, 7). 287

Fig 3.   Log-likelihood scores of stochastic and auto-regression projections in prospective validation.

As of February 25, the total reported count of probable and confirmed EVD cases ₂₈₈ was 875 cases. With the stochastic model, the four-week projection of median outbreak ₂₈₉ size was 955 cases (95% prediction interval: 874–1105). With the auto-regression model, ₂₉₀ the four-week projection of median outbreak size was 898 cases (95% prediction interval: ₂₉₁ 877–983). ₂₉₂

Median final outbreak size projected by the stochastic model was 990 cases (95% ₂₉₃ prediction interval: 860–1295). With the Gott's rule model, the median final outbreak ₂₉₄ size was 1749 cases (95% prediction interval: 921–17498). Median final outbreak size ₂₉₅ projected by the regression model was 953 cases (95% prediction interval: 882–1136). ₂₉₆

Fig 4.   Short-term probabilistic projections of case counts based on reported counts as of February 25, 2019. Stochastic and auto-regression models were used. Vertical scale of probabilities, on right side of plot, applies to all dates.

The possibility of a final outbreak size exceeding 1,000 cases was projected to be ₂₉₇

Fig 5. Medians and prediction intervals from short-term projections of case counts based on reported counts as of February 25, 2019. Solid dots indicate median count.

Fig 6. Probabilistic projections of final case counts based on reported counts as of February 25, 2019. Stochastic, Theil-Sen regression, and Gott's rule models were used.

Fig 7. Medians and prediction intervals from projections of final case counts based on reported counts as of February 25, 2019. Solid dots indicate median size. Inset provides detail at smaller numbers.

quite likely by all the models, with the Gott's rule forecast being particularly pessimistic, as it was in all cases. The probability of a very large outbreak (10,000 or more cases) was calculated to be below 8% for Gott's rule and negligibly small for the other models. A final outbreak size of a catastrophic outbreak larger than the West Africa outbreak (28,616 or more cases) was projected to have probability less than 3% by Gott's rule and negligibly small for the other models (Table S2 in S1 Supporting Information).

We generated short- and long-term projections from earlier snapshots of the current outbreak case counts with each model, for the purpose of both scoring and forecasting. These additional results can be found in the Supplemental Material.

Probability of vaccination coverage scenarios

The stochastic model estimated likelihoods of the three scenarios of zero, low, and high vaccination coverage, based on how often models using the different coverage assumptions were able to pass the particle filtering step. In the estimate based on data through February 25, vaccination coverage was estimated substantially more likely than low or high coverage, as was also true in most earlier estimates. The lower vaccination coverage scenario was estimated more likely than the higher one. Higher vaccination coverage scenarios were estimated more likely in estimates made before October, at which time the outbreak epicenter shifted into the conflict zone and situation reports described an increase in case counts (Fig S5 in S1 Supporting Information). Assuming that other transmission factors were held constant, our result may provide evidence that that geographical shift in transmission contributed to a decreased likelihood of a high vaccination scenario.

Discussion

As of February 25, 2019, with 875 cases reported to date, most of our model projections were concentrated in a range up to about 300 additional cases overall, and even the Gott's rule forecast, whose prediction intervals are especially wide, placed low probability on outcomes on the scale of the West African outbreak. The current outbreak, however, has so far behaved unpredictably, possibly due to complex social circumstances, and could continue to be unpredictable and render our projections

inaccurate. EVD has never before been introduced into a conflict zone with such ₃₂₇ political instability, potential disease mobility and community impenetrability [2]. In ₃₂₈ October, WHO reported that up to 80% of contacts were not being traced [3]. At ₃₂₉ present, the most reliable data source of EBOV transmission has been the weekly case ₃₃₀ counts that can be found in the WHO situation reports. Despite such situations of data ₃₃₁ scarcity and new outbreak circumstances, our models generated relatively accurate ₃₃₂ short-term projections of outbreak size in the months of January and February 2019, ₃₃₃ suggesting that short-term projections made in real time can be useful in decision ₃₃₄ making and resource allocation. ₃₃₅

Another outbreak response challenge has been ascertaining the level of vaccination ₃₃₆ coverage that has occurred during the ring vaccination program. For the ring ₃₃₇ vaccination program to be theoretically effective at reducing transmission, an uncertain ₃₃₈ proportion of contacts and contacts of contacts need to receive the vaccine [6]. We ₃₃₉ estimated that transmission rates are higher than would be expected under target levels ₃₄₀ of 62% coverage due to contact tracing and vaccination. Under our stochastic model's ₃₄₁ assumptions, the elevated transmission rates, relative to those of past outbreaks, needed ₃₄₂ to reproduce this outbreak's epidemic trajectory are highly unlikely to be randomly ₃₄₃ observed with high vaccination coverage. This model estimate may serve as a loose ₃₄₄ proxy for the outbreak response challenges. Thus, outbreak response teams may need to ₃₄₅ consider other control and care strategies to end the outbreak. Under current ₃₄₆ circumstances, mass vaccination in regions of high prevalence or areas where the ₃₄₇ outbreak has newly arrived may be a more effective strategy to reduce EBOV ₃₄₈ transmission than ring vaccination. Aggressive supportive care, experimental ₃₄₉ therapeutics and high-quality facilities (*e.g.* air-conditioned, individualized) have also ₃₅₀ been described as part of the outbreak response. These interventions have the potential ₃₅₁ to improve health-seeking behaviors and possibly reduce EBOV transmission in ₃₅₂ communities that are resistant to control efforts [15], but further studies are needed. ₃₅₃

The mathematical models we adopted for this project near the beginning of the ₃₅₄ outbreak were developed to predict the course of prior outbreaks, and did not consider ₃₅₅ circumstances unique to the current outbreak. To improve the accuracy of short-term ₃₅₆ projections, we used the short-term forecasting performance of the mathematical models ₃₅₇ during the current outbreak to calibrate them and improve their performance on this ₃₅₈

outbreak. This may be a useful pattern for short-term forecasting of ongoing disease   359
outbreaks in real time.   360

The performance of our short-term prediction models can be assessed on the   361
outbreak to date. The relatively simple auto-regression model we used performed more   362
consistently on the range of partial data sets used for scoring than the more complex   363
stochastic simulation model did. The stochastic model has tended to produce tighter   364
prediction distributions that are prone to extreme failure when they get it wrong, while   365
the auto-regression model's predictions are more tolerant of unpredictable outcomes.   366
Conversely, the stochastic model outperforms the auto-regression when it gets the   367
prediction right. It should also be noted that because the stochastic simulation model is   368
based on mechanistic knowledge of the transmission process generating the outbreak   369
while the regression model is a purely statistical inference from past outbreaks, it may   370
be that if conditions emerge that are substantially different from past outbreak   371
conditions, the mechanistic model may produce sensible predictions where a purely   372
statistical model fails. Because the other models included produce only final outbreak   373
size projections, they can not be fully evaluated before the outbreak has ended.   374
However, we can note that our early projections of final size (Figs S9, S10 in S1   375
Supporting Information) fell below the counts that have been observed as of February.   376
In other words, our forecasts based on early reports and on an assumption that past   377
outbreaks can be used to forecast the present one were more optimistic than warranted   378
by subsequent events. Events to come may shed light on whether this outbreak is   379
qualitatively different than the past ones we have used to construct forecasts, perhaps   380
due to the impact of conflict conditions on the outbreak.   381

There are limitations to our projections. Projection distributions are right-skewed,   382
with long tails (and we therefore report the median instead of the mean). We were   383
unable to include all the 23 observed EVD outbreaks with a case count greater than ten   384
cases in our estimates due to data availability. Our regression models are based entirely   385
on past outbreaks of Ebola virus disease (measured and reported in different ways), and   386
cannot account for the improved control measures and vaccination in the way that a   387
mechanistic model does. We included as much real-time information in our models as   388
possible, but situations such as the introduction of EVD into a zone of armed conflict   389
and the recent introduction of vaccination are not reflected by the suite of past   390

outbreaks. The stochastic model used estimated vaccination effectiveness, reported cases, and timing of onset dates affected by vaccination from studies from West Africa, not DRC, and did not include vaccination of healthcare workers. Our forecasts do not account for possible unreported cases or changes in reporting over time; such gaps in reporting can not be ruled out, though given the intense efforts at case finding that started in July and carried forward, we think it unlikely that there were large changes in reporting. Furthermore, as the outbreak moved into areas affected by violent conflict as the outbreak continued, we think it likely that case reporting, if anything, decreased over time, and therefore underreporting would not explain the apparent increase in transmission from June to October. While it would be desirable to use the vaccination coverage estimates to estimate the number of cases prevented by the vaccination program, our models were not designed to produce a testable estimate of such an effect.

Gott's rule [39], presented initially as reflecting a "Copernican" principle with respect to time, assumed that ones observation of the age of a phenomenon occurred at a random time during its lifetime. Gott applied this principle in a variety of settings, including the lifetime of Broadway shows, the Berlin Wall, and of the human race [39,40]. Critics suggest that Gott's rule gives misleading predictions when the phenomenon in question exhibits known time scales, such as in prediction of human lifetimes [40,41]. Gott's rule has not been validated for epidemics, and in this case, is based on an assumption that the current case count in an epidemic occurs at a random point in its progression. For an easily detected but uncontrollable supercritical epidemic, we would expect forecasts based on the first reports to underestimate the final size. For an easily controlled epidemic, forecasts would more likely be too pessimistic, since control measures would (by assumption) swiftly end the epidemic after recognition. Our application of the rule is intended to serve as a null hypothesis-like comparator for the other methods.

Our models do not account for spatial heterogeneity in transmission, which may be relevant to the course of this outbreak, in which an increase in transmission appears to coincide with a shift in location. The stochastic model may be sensitive to the assumption of an exponentially decaying curve for transmission rate by day, and alternative assumptions might lead to a different distribution of forecast outcomes. Indeed, all of our projections are conditional on model assumptions being met. If

unpredictable events were to change patterns of EBOV transmission in ways not seen in past outbreaks, assumptions used for model projections could be violated and our results could change. For example, exceedingly improbable events such as a catastrophic outbreak (more than 10,000 cases or approaching the size of the 2013–2016 West Africa outbreak) might become more probable. A strength of our approach was the use of multiple methods to estimate the outbreak size, including both mechanistic modeling and purely statistical approaches.

Before October, there was limited EBOV transmission in active, armed conflict zones. When more reported EVD cases occurred in Beni, it was unclear how the context would affect secondary EBOV transmission. The October data suggests that EBOV transmission increased there. We believe the increased rate of reported EVD cases corresponded to the shift of EBOV transmission into conflict zones. It may be that a model that explicitly distinguishes transmission rates in these zones from those in other areas would model the dynamics underlying these cases more precisely and produce more accurate projections.

Many of the reported EVD cases in this outbreak have been identified as new transmission chains. While a reduced ability to detect and vaccinate contacts in difficult-to-access communities has been anecdotally reported, our stochastic model provided empirical evidence of outbreak challenges, suggesting that transmission rates were consistent with lower than target levels of vaccine coverage. beyond those already reported. While a catastrophic outbreak of 10,000 or more cases is not projected as probable, vigilance is warranted. New circumstances—such as epidemic spread to Uganda—call for newly validated projections, whenever possible, even in the short term.

References

1. World Health Organization Regional Office for Africa. Health topics: Ebola virus disease; 2018.
   http://www.afro.who.int/health-topics/ebola-virus-disease.

2. Wikipedia contributors. North Kivu; 2018. Available from: https://en.wikipedia.org/w/index.php?title=North_Kivu&oldid=845077998.

3. World Health Organization. Ebola situation reports: Democratic Republic of the ⁠452
Congo; 2018. `http://www.who.int/ebola/situation-reports/drc-2018/en/`. ⁠453

4. Damon IK, Rollin PE, Choi MJ, Arthur RR, Redfield RR. New Tools in the ⁠454
Ebola Arsenal. New England Journal of Medicine. 2018;. ⁠455

5. Barry A, Ahuka-Mundeke S, Ahmed YA, Allarangar Y, Anoko J, Archer BN, ⁠456
et al. Outbreak of Ebola virus disease in the Democratic Republic of the Congo, ⁠457
April–May, 2018: an epidemiological study. The Lancet. 2018;392(10143):213–221. ⁠458

6. Bodine EN, Cook C, Shorten M. The potential impact of a prophylactic vaccine ⁠459
for Ebola in Sierra Leone. Mathematical Biosciences & Engineering. ⁠460
2018;15(1551-0018_2018_2_337):337. doi:10.3934/mbe.2018015. ⁠461

7. Référentiel Géographique Commun. Democratic Republic of Congo and ⁠462
neighboring countries Health Boundaries; 2018. `https://data.humdata.org/` ⁠463
`dataset/democratic-republic-of-congo-health-boundaries`. ⁠464

8. Butler D. Models overestimate Ebola cases. Nature. 2014;515(7525):18. ⁠465
doi:10.1038/515018a. ⁠466

9. Shaman J, Yang W, Kandula S. Inference and forecast of the current West ⁠467
African Ebola outbreak in Guinea, Sierra Leone and Liberia. PLoS Curr. 2014;6. ⁠468
doi:10.1371/currents.outbreaks.3408774290b1a0f2dd7cae877c8b8ff6. ⁠469

10. Reis J, Yamana T, Kandula S, Shaman J. Superensemble forecast of respiratory ⁠470
syncytial virus outbreaks at national, regional, and state levels in the United ⁠471
States. Epidemics. 2018;S1755-4365(17):30174–3. ⁠472
doi:10.1016/j.epidem.2018.07.001. ⁠473

11. Yamana T, Kandula S, Shaman J. Superensemble forecasts of dengue outbreaks. ⁠474
J R Soc Interface. 2016;13(123). ⁠475

12. Reich N, Lauer S, Sakrejda K, Iamsirithaworn S, Hinjoy S, Suangtho P, et al. ⁠476
Challenges in Real-Time Prediction of Infectious Disease: A Case Study of ⁠477
Dengue in Thailand. PLoS Negl Trop Dis. 2016;10(6):e0004761. ⁠478
doi:10.1371/journal.pntd.0004761. ⁠479

13. Graham M, Suk J, Takahashi S, Metcalf C, Jimenez A, Prikazsky V, et al. Challenges and Opportunities in Disease Forecasting in Outbreak Settings: A Case Study of Measles in Lola Prefecture, Guinea. Am J Trop Med Hyg. 2018;98(5):1489–1497. doi:10.4269/ajtmh.17-0218.

14. Li M, Dushoff J, Bolker B. Fitting mechanistic epidemic models to data: A comparison of simple Markov chain Monte Carlo approaches. Stat Methods Med Res. 2018;27(7):1956–1967. doi:10.1177/0962280217747054.

15. Funk S, Camacho A, Kucharski AJ, Lowe R, Eggo RM, Edmunds WJ. Assessing the performance of real-time epidemic forecasts: A case study of the 2013-16 Ebola epidemic. bioRxiv. 2018;doi:10.1101/177451.

16. Kelly JD, Worden L, Wannier SR, Hoff NA, Mukadi P, Sinai C, et al. Projections of Ebola outbreak size and duration with and without vaccine use in Équateur, Democratic Republic of Congo, as of May 27, 2018. PLOS ONE. 2019;14(3):e0213190. doi:10.1371/journal.pone.0213190.

17. Ebola haemorrhagic fever in Sudan, 1976. Report of a WHO/International Study Team. Bull World Health Organ. 1978;56(2):247–70.

18. Breman JG, Heymann DL, Lloyd G, McCormick JB, Miatudila M, Murphy FA, et al. Discovery and Description of Ebola Zaire Virus in 1976 and Relevance to the West African Epidemic During 2013-2016. J Infect Dis. 2016;doi:10.1093/infdis/jiw207.

19. Baron RC, McCormick JB, Zubeir OA. Ebola virus disease in southern Sudan: hospital dissemination and intrafamilial spread. Bull World Health Organ. 1983;61(6):997–1003.

20. Georges AJ, Leroy EM, Renaut AA, Benissan CT, Nabias RJ, Ngoc MT, et al. Ebola hemorrhagic fever outbreaks in Gabon, 1994-1997: epidemiologic and health control issues. J Infect Dis. 1999;179 Suppl 1:S65–75. doi:10.1086/514290.

21. Khan AS, Tshioko FK, Heymann DL, Le Guenno B, Nabeth P, Kerstiëns B, et al. The reemergence of Ebola hemorrhagic fever, Democratic Republic of the Congo,

1995. Commission de Lutte contre les Epidémies à Kikwit. J Infect Dis. 1999;179   508
Suppl 1:S76–86. doi:10.1086/514306.   509

22. CDC and Ministry of Health: T Oyok, Odonga C, Mulwani E, Abur J, Kaducu F,   510
Akech M, et al. Outbreak of Ebola Hemorrhagic Fever — Uganda, August   511
2000–January 2001. MMWR: Morbidity and Mortality Weekly Report.   512
2001;50(5):73–77.   513

23. Outbreak(s) of Ebola haemorrhagic fever, Congo and Gabon, October 2001-July   514
2002. Wkly Epidemiol Rec. 2003;78(26):223–8.   515

24. Boumandouki P, Formenty P, Epelboin A, Campbell P, Atsangandoko C,   516
Allarangar Y, et al. Prise en charge des malades et des défunts lors de l'épidémie   517
de fièvre hémorragique due au virus Ebola d'octobre à décembre 2003. Bull Soc   518
Pathol Exot. 2005;98(3):218–223.   519

25. World Health Organization. Weekly epidemiological record. Abonnement annuel.   520
2005;43(80):369–376.   521

26. Nkoghe D, Kone ML, Yada A, Leroy E. A limited outbreak of Ebola   522
haemorrhagic fever in Etoumbi, Republic of Congo, 2005. Trans R Soc Trop Med   523
Hyg. 2011;105(8):466–72. doi:10.1016/j.trstmh.2011.04.011.   524

27. Rosello A, Mossoko M, Flasche S, Van Hoek AJ, Mbala P, Camacho A, et al.   525
Ebola virus disease in the Democratic Republic of the Congo, 1976-2014. Elife.   526
2015;4. doi:10.7554/eLife.09015.   527

28. MacNeil A, Farnon EC, Morgan OW, Gould P, Boehmer TK, Blaney DD, et al.   528
Filovirus outbreak detection and surveillance: lessons from Bundibugyo. J Infect   529
Dis. 2011;204 Suppl 3:S761–7. doi:10.1093/infdis/jir294.   530

29. Uganda: Ebola Situation Report. Bull World Health Organ. 2013;.   531

30. Centers for Disease Control and Prevention. Number of Cases and Deaths in   532
Guinea, Liberia, and Sierra Leone during the 2014-2016 West Africa Ebola   533
Outbreak. 2017;.   534

31. Wallinga J, Teunis P. Different epidemic curves for severe acute respiratory syndrome reveal similar impacts of control measures. Am J Epidemiol. 2004;160(6):509–16. doi:10.1093/aje/kwh255.

32. Blumberg S, Lloyd-Smith JO. Comparing methods for estimating R_0 from the size distribution of subcritical transmission chains. Epidemics. 2013;5(3):131–45. doi:10.1016/j.epidem.2013.05.002.

33. Lloyd-Smith JO, Schreiber SJ, Kopp PE, Getz WM. Superspreading and the effect of individual variation on disease emergence. Nature. 2005;438(7066):355–9. doi:10.1038/nature04153.

34. Worden L, Wannier R, Hoff NA, Musene K, Selo B, Mossoko M, et al. Real-time projections of epidemic transmission and estimation of vaccination impact during an Ebola virus disease outbreak in the Eastern region of the Democratic Republic of Congo. bioRxiv. 2018;doi:10.1101/461285.

35. Dalziel B, Lau M, Tiffany A, et al. Unreported cases in the 2014–2016 Ebola epidemic: Spatiotemporal variation, and implications for estimating transmission. PLoS Negl Trop Dis. 2018;12(e0006161).

36. Henao-Restrepo AM, Camacho A, Longini IM, Watson CH, Edmunds WJ, Egger M, et al. Efficacy and effectiveness of an rVSV-vectored vaccine in preventing Ebola virus disease: final results from the Guinea ring vaccination, open-label, cluster-randomised trial (Ebola Ça Suffit!). Lancet. 2017;389(10068):505–518. doi:10.1016/S0140-6736(16)32621-6.

37. Dalziel BD, Lau MSY, Tiffany A, McClelland A, Zelner J, Bliss JR, et al. Unreported cases in the 2014-2016 Ebola epidemic: Spatiotemporal variation, and implications for estimating transmission. PLoS Negl Trop Dis. 2018;12(1):e0006161. doi:10.1371/journal.pntd.0006161.

38. Komsta L. mblm: Median-Based Linear Models; 2019. Available from: https://CRAN.R-project.org/package=mblm.

39. Gott III JR. Implications of the Copernican principle for our future prospects. Nature. 1993;363:315–319.

40. Caves CM. Predicting future duration from present age: A critical assessment. 564
Contemporary Physics. 2000;41(3):143–153. 565

41. Nelson K. How and how not to make predictions with temporal Copernicanism. 566
Synthese. 2009;166(1):91–111. 567

S1 Supporting Information. Appendix containing supplemental figures 568
and tables. 569

Supporting information

Table S1 summarizes the past outbreaks used as data to inform our models.   571

Table S1. Table of past outbreaks by year and country.

Time Period	Country	Reported Count	Time Series Count	Regression?	Stochastic?	Auto-Regression?
Aug–Sep 1976	DRC*	318	262	Yes	No	Yes
Jun–Nov 1976	Sudan	284	284	Yes	No	Yes
Aug–Sep 1979	Sudan	34	34	Yes	Yes	Yes
Dec 1994–Feb 1995	Gabon	52	49	Yes	No	Yes
May–Jul 1995	DRC	315	317	Yes	Yes	Yes
Jan–Apr 1996	Gabon	37	29	Yes	Yes	Yes
Jul 1996–Mar 1997	Gabon	60	–	No	No	No
Oct 2000–Jan 2001	Uganda	425	436	Yes	No	Yes
Oct 2001–Jul 2002	Gabon, Republic of the Congo	124	124	Yes	Yes	Yes
Dec 2002–Mar 2003	Republic of the Congo	143	–	No	No	No
Nov–Dec 2003	Republic of the Congo	35	35	Yes	Yes	Yes
Apr–Jun 2004	Sudan	17	17	Yes	Yes	Yes
Apr–May 2005	DRC	12	12	Yes	Yes	Yes
Aug–Nov 2007	DRC	264	264	Yes	Yes	Yes
Dec 2007–Jan 2008	Uganda	131	127	Yes	Yes	Yes
Dec 2008–Feb 2009	DRC	32	32	Yes	Yes	Yes
Jun–Aug 2012	Uganda	24	24	Yes	Yes	Yes
Jun–Nov 2012	DRC	52	52	Yes	Yes	Yes
Aug–Nov 2014	DRC	66	62	Yes	Yes	Yes
Jul–Oct 2014	Nigeria (offshoot of West African outbreak)	20	–	No	No	No
Jan 2014–Jun 2016	Guinea, Liberia, Sierra Leone	28,616	21,422	Yes	No	Yes
Apr–Jun 2018	DRC	53	53	Yes	Yes	Yes

Official reported case counts for each epidemic are given, including suspected cases ("Reported Count"). Case counts for the time series data included in the models include only probable and confirmed cases ("Time Series Count"). Case counts for historic outbreaks were pulled from publicly available literature [17–30]. Lastly, each historic outbreak's inclusion in the regression, stochastic, and auto-regression models is enumerated.
*Democratic Republic of Congo (formerly Zaïre)

We retained snapshots of the set of available case counts at multiple time points, for   572

use in scoring of retrospective model projections against known subsequent counts   573

(Fig S1). In later data sets, due to the larger number of data points, a subset of the case   574

counts was selected for use in the stochastic model's particle filtering step, as noted in   575

the figure.   576

Fig S1. Reported case counts in current outbreak by date, in multiple snapshots of available data. Where not otherwise noted, all case counts shown were used in the stochastic model's particle filtering step.

Stochastic Model

Epidemic curves reported for past Ebola outbreaks were used to estimate time series of

effective reproduction number (R) by day, which were then fit to an exponential

quenching curve (Fig S2). The quenching parameter estimates the rate of change in R

per day from $R_{initial}$ that results from outbreak control efforts, behavioral changes in

response to the outbreak, and potential local depletion of susceptibles. The parameters

$R_{initial}$ and τ estimated by that curve fitting on past epidemics were then used to create

a distribution from which values were sampled to parametrize the stochastic simulation   584
(Fig S3).   585

Fig S2. Estimates of reproduction number R by day in past Ebola outbreaks.
Thin curves are exponentially quenched curves $R = R_{\mathrm{initial}} e^{-\tau d}$ fit to each series of R
estimates.

Fig S3. Distribution of transmission rates sampled for simulation. Black
dots are pairs R_{initial} and quenching rate τ estimated from past Ebola outbreaks, and
blue cloud is the continuous distribution from which pairs are sampled for simulation.

The R_{initial} and τ parameters driving simulated outbreaks that were successful in   586
passing the particle filtering step tended to cluster in particular locations within the   587
assumed distribution (Fig S4). In some cases, distinct ranges of R_{initial} and/or τ were   588
selected in conjunction with the different vaccine coverage scenarios. The frequencies of   589
selection of the three vaccine coverage scenarios were used to estimate the likelihood of   590
each scenario (Fig S5).   591

For each data set, simulations were generated until between 320 and 330 simulated   592
outbreaks were obtained after filtering based on approximate agreement with reported   593
case counts from the current outbreak.   594

The simulations passing the particle filtering step, representing a distribution of parameter values and vaccine scenarios, were continued beyond the particle filtering points to generate a spreading set of projections of case counts at later dates (Fig S6), which was smoothed to create probabilistic projections of future case counts at the desired dates.

Projections

We have recorded the projections generated by our models from older data sets to assess the development of the projections as the outbreak has progressed, in Figs S7, S8, S9, and S10.

Table S2 summarizes the medians and 95% prediction intervals produced by each model on the most recent data set included, and their probabilities of large outcomes.

Table S2. Table of medians and prediction intervals of model projections.

Forecast as of	Forecast	Model	Lower	Median	Upper	Size 1,000	Size 10,000	Size 28,616
2019-02-25	2019-03-04	Auto-Regression	876	883	908	0.00002	0.00000	0.00000
2019-02-25	2019-03-04	Stochastic	868	920	1028	0.11077	0.00000	0.00000
2019-02-25	2019-03-11	Auto-Regression	876	889	933	0.00204	0.00000	0.00000
2019-02-25	2019-03-11	Stochastic	872	933	1054	0.16451	0.00000	0.00000
2019-02-25	2019-03-25	Auto-Regression	877	898	983	0.03327	0.00000	0.00000
2019-02-25	2019-03-25	Stochastic	874	955	1105	0.26913	0.00000	0.00000
2019-02-25	2019-04-22	Auto-Regression	877	915	1095	0.16126	0.00000	0.00000
2019-02-25	2019-04-22	Stochastic	874	976	1180	0.39925	0.00000	0.00000
2019-02-25	final	Gott's Rule	921	1749	17498	0.86625	0.07875	0.02183
2019-02-25	final	Stochastic	860	990	1295	0.46620	0.00000	0.00000
2019-02-25	final	Theil-Sen Regression	882	953	1136	0.29805	0.00000	0.00000

Fig S4. Transmission rates selected by the particle filtering process, by vaccine coverage scenario, for successive snapshots of available case count data. As in previous figure, black dots for R_{initial}, τ pairs estimated for past outbreaks (for comparison), and colors illustrate the density of R_{initial}, τ pairs selected by filtering simulated outbreaks, classified by level of vaccine coverage.

Fig S5. Likelihood estimates of vaccine coverage scenarios consistent with model transmission rates, estimated by number of simulated outbreaks under the three scenarios accepted by the stochastic model's filtering step, in which simulated outbreaks must match reported case counts. Note that each bar of this plot is an estimate based on the entire time series up to the date indicated, that is, they are cumulative estimates of the overall vaccine coverage, not successive estimates of short-term coverage.

Fig S6. Cumulative case counts by date projected by individual realizations of the stochastic model, by vaccine coverage scenario, using successive snapshots of available case count data. The vertical axis is cut off at the upper limit of the 95% prediction interval for outbreak sizes, for readability.

Fig S7. Short-term projections based on multiple data sets. Probability scale (marked on right) is consistent across dates within each subplot.

Fig S8. **Medians and prediction intervals from short-term projections** based on multiple data sets.

Fig S9. Final outbreak size projections based on multiple data sets.

Fig S10. Medians and prediction intervals from final outbreak size projections based on multiple data sets, with inset for detail in smaller numbers.