Title: *C. albicans* white and opaque cells exhibit distinct spectra of organ colonization in standard mouse models of infection

Author/Affiliations: Julie Takagi[1*], Sheena D. Singh-Babak[1], Matthew B. Lohse[1], Chiraj K. Dalal[1#] & Alexander D. Johnson[1,2#]

[1]Department of Microbiology and Immunology, University of California, San Francisco, 600 16 Street, San Francisco, CA 94158, USA
[2]Department of Biochemistry and Biophysics, University of California, San Francisco, 600 16th Street, San Francisco, CA 94158, USA

*Current Address:
Department of Biology, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, MA 02139, USA

#Correspondence: ajohnson@cgl.ucsf.edu
                 chirajdalal.ucsf@gmail.com

Abstract

Candida albicans can thrive in diverse niches of its mammalian hosts; it is a

normal resident of the GI tract and mucosal surfaces but it can also enter the

bloodstream and colonize internal organs causing serious disease. The ability of *C.*

albicans to thrive in these different host environments has been attributed, at least

in part, to its ability to assume different morphological forms. In this work, we

examine one such morphological change known as white-opaque switching. White

cells are the default state of *C. albicans*, and most animal studies have been carried

out exclusively with white cells. Here, we compared the proliferation of white and

opaque cells in two murine models of infection and also monitored, using specially

constructed strains, switching between the two states in the host. We found that

white cells outcompeted opaque cells in many niches; however, in some organs

(specifically, the heart and spleen), opaque cells competed favorably with white

cells. In environments where the introduced white cells outcompeted the introduced

opaque cells, opaque cells often switched to white cells. We did not observe white

cells switching to opaque cells in any of the niches we examined.

Introduction

Candida albicans is a source of serious fungal infections in the United

States; it is the fourth most commonly cultured microbe from blood (Pfaller and

Diekema, 2007). Unlike other major fungal pathogens, *C. albicans* is also a part of

the normal human microbiome, particularly in the gastrointestinal tract, skin and

mucosal surfaces (Odds, 1988). As a pathogen, *C. albicans* poses health risks,

particularly for people whose immune system is suppressed, who have surgeries,

implanted medical devices, or who have been treated with long courses of

antibiotics. The attributable mortality from *C. albicans* bloodstream infections in

adults is at least 15% and has been reported to be as high as 40% (Zaoutis et al.,

2005).

While some human fungal pathogens exist primarily as budding yeast cells

(for example, *Cryptococcus neoformans* (Buchanan, 1998)) or filamentous hyphal

structures (for example, *Aspergillus* spp. (Baker and Bennett, 2007)), *C. albicans*

alternates between these and other morphologies, often in response to specific

environmental cues. In this paper, we explore the switch *C. albicans* makes from its

normal, round-to-oval yeast cell morphology (white) to an elongated, mating

competent cell type termed opaque (for reviews see Lohse and Johnson, 2009; Soll

et al., 2009). White-opaque switching is unusual in that it is a heritable change, one

that occurs without a change in the DNA sequence of the genome. In standard

laboratory media, switching occurs rarely, approximately once every 10^4 cell

divisions (Slutsky et al., 1987; Soll et al., 1993); therefore, each cell type is

maintained across many cycles of cell division. Approximately one-sixth of all genes

(~6000 total) are differentially regulated (at least two-fold) between the two cell

types (Lan et al., 2002; Tuch et al., 2010). These expression differences involve

many types of genes, including those responsible for the ability of opaque cells (but

not white cells) to mate. These differentially regulated genes also encode a variety

of metabolic enzymes, suggesting that white and opaque cells are each

'metabolically specialized' to thrive in specific niches (reviewed in Huang, 2012;

Lohse and Johnson, 2009; Morschhäuser, 2010; Soll, 2014, 1992, 2004).

Supporting this hypothesis, environmental factors such as temperature, carbon

source, and oxygen can significantly alter the switching rate between cell-types

(Ene et al., 2016; Huang et al., 2009, 2010; Rikkerink et al., 1988). Another

difference between white and opaque cells lies in their recognition by macrophages

and neutrophils, indicating that the white-opaque switch may play a role in evading

the host innate immune system response and in colonizing internal environments

(Geiger et al., 2004; Kolotila and Diamond, 1990; Kvaal et al., 1999, 1997; Lohse

and Johnson, 2008; Sasse et al., 2013). These observations suggest the

hypothesis that white-opaque switching evolved during *C. albicans*' long association

with its warm-blooded host and plays a crucial role in the pathogen-host

relationship. Consistent with this hypothesis, deletion of the master regulator of

white-opaque switching (Wor1) impairs the ability of *C. albicans* to survive in a

mouse gastrointestinal model of colonization (Pande et al., 2013). Although not all

clinically isolated strains of *C. albicans* readily undergo white-opaque switching,

many do so. Even "non-switching" clinical strains can be converted to switching

strains by simple genetic manipulation, indicating that the mechanisms underlying

white-opaque switching are deeply conserved. Also consistent with this idea is the

observation that two pathogenic fungi closely related to *C. albicans* (*Candida*

dubliniensis and *Candida tropicalis*) also undergo white-opaque switching (Porman

et al., 2013; Pujol et al., 2004).

To systematically investigate the behavior of white and opaque cells *in vivo*,

genetically identical strains of white and opaque cells were tagged with spectrally

distinct fluorophores and co-inoculated into rodent models of infection and

colonization. We found, using the standard tail-vein injection model (Odds et al.,

2000), that both white and opaque cells are capable of disseminating from the

bloodstream into multiple organs; in some organs, white cells significantly

outcompete opaque cells, but in others, they are roughly equivalent. In a gut model

of colonization (Rosenbach et al., 2010), long-term colonization by opaque cells is

dependent on switching to the white state.

Results:

Fluorescent markers enable unbiased comparison of white and opaque cells
in vivo

In order to develop a quantitative, reproducible approach to observe the

relative colonization of white and opaque cells *in vivo*, we constructed fluorescently

tagged white and opaque strains of *C. albicans*. These strains were matched

genetically, except for a single locus where the Tef2 protein was C-terminally

tagged with either GFP or mCherry, two spectrally distinct fluorophores. These

strains were introduced together (in 50:50 mixes) to mice via tail-vein injection or

oral gavage; after the mice were sacrificed, *C. albicans* cells were harvested and

immediately plated at 25°C. After seven days of growth, we scored colonies as

white or opaque to determine the state of each cell at the point the mice were

sacrificed. (At 25°C on plates, very little switching occurs, so the appearance of a

given colony reflected the identity of the single cell that produced it.) Using a

fluorescent stereomicroscope, we also determined whether each colony expressed

GFP or mCherry. In this way, we were able to determine whether a given cell was

injected as a white or opaque cell and whether it remained as the same cell type or

underwent a switch to the other cell type. This approach offers several advantages

when compared with previous studies: 1) it does not require the use of auxotrophies

or drug resistance markers that have been shown to affect *C. albicans* behavior *in*

vivo (Bain et al., 2001; Lay et al., 1998)*,* 2) it does not require the use of strains that

are artificially "locked" in one cell-type or another, for example by deleting or

overexpressing the master regulator Wor1 (Huang et al., 2006; Pande et al., 2013;

Zordan et al., 2006), and 3) it does not require any downstream processing, such

as qPCR, which can introduce errors through amplification. Finally, because white

and opaque cells were injected together, this approach enabled us to directly

compare colonization, proliferation, and switching of white and opaque cells *in vivo*.

To test whether the fluorophores introduced any bias *in vivo*, the tail veins of

four Balb/C mice (18-20g) were injected with 2.2×10^6 white cells, half of which were

tagged with GFP and the other half with mCherry. The mice were sacrificed after 24

hours and the fungal burden in five organs (the kidney, liver, spleen, heart, and

brain) was analyzed (Figure 1a). Across all five organs, the distributions of mCherry

and GFP-tagged colonies were not significantly different (p-values all >0.05,

Wilcoxon matched-pairs signed rank test, Figure 1, Dataset 1), indicating that the

fluorophores themselves introduce little or no bias *in vivo*.

Opaque cells can disseminate from the bloodstream into many organs

To compare the relative colonization of white and opaque cells in a systemic

model of infection, white-mCherry and opaque-GFP cells were co-injected (at a

50:50 ratio) into the tail vein of four mice each at two inocula: 8×10^5 and 1.6×10^6

cells (Figure 2a). After 24 hours, the mice were euthanized and their organs

harvested and processed. Both white and opaque cells were present in all five

organs (as indicated by the colony counts) and were at a higher density in the mice

with the higher inoculum (Figure 2, Dataset 2). A comparison of the white and

opaque cells across the five organs showed different distributions depending on the

organ (Figure 2, Dataset 2). The most pronounced difference was observed in the

kidney, where white cells greatly outnumbered opaque cells in every mouse

examined (p-value = 0.0078, Wilcoxon matched-pairs signed rank test, Figure 2b,

Dataset 2). Of the white cells recovered from the kidney, 70% had been injected as

white cells; the remaining 30% had been initially injected as opaque cells but had

switched to white cells. The situation in the heart and spleen was different; here,

opaque cells outnumbered white cells (p-value 0.0156, Wilcoxon matched-pairs

signed rank test, Figure 2d, 2e Dataset 2). In the heart, virtually none of the cells

had switched in either direction; in the spleen, a small fraction of the white cells

recovered had been injected as opaque cells. Finally, in the liver and brain, both

white and opaque cells were recovered with a slight preference for white cells

(Figure 2c, 2f). Because of significant mouse-to-mouse variation (particularly in the

liver), we do not feel confident that this preference is biologically meaningful even

though it is statistically significant (p-values =0.0078, Wilcoxon matched-pairs

signed rank test, Figure 2c, 2f, Dataset 2). Rather, we view the colonization of white

and opaque cells in these organs as roughly equivalent. We do note that, of the

white cells recovered from the brain, a significant fraction had been initially injected

as opaque cells.

As described above, our observations document a significant level of

opaque-to-white switching that occurred *in vivo*. We do not know when, during the

course of the infection, this switching occurred; however, we suspect it happened

after the injected cells entered the different organs. Had the opaque cells switched

in the bloodstream immediately following injection, we would have expected to

observe similar opaque-to-white switching patterns across all organs tested, and

this was not the case. Finally, we note that we did not observe any opaque-

mCherry colonies (in any organ, in any mouse, and at any inoculum, >1000 colony-

forming units visually inspected), indicating that stable switching from white cells to

opaque cells must be rare or non-existent in the mouse models we examined.

To test whether the colonization by opaque cells was dependent on co-

injection with white cells, we injected the tail veins of four additional mice with a

50:50 mixture of opaque-mCherry and opaque-GFP inoculum ($1.6*10^6$ cells) and

measured colonization in the kidney, spleen, liver, brain, and heart (Figure S1,

Dataset 2). We observed opaque cell colonization of all five organs (Figure S1,

Dataset 2), indicating that a large white cell population was not needed to "aid" the

opaque cells in colonization.

Taken together, these results show that opaque cells are capable of

efficiently colonizing and proliferating in numerous niches *in vivo*, and in some

cases (heart and spleen) somewhat better than white cells. However, opaque cells

are severely outcompeted by white cells in the kidney, the organ that is typically

used to assess systemic fungal infection in murine models.

C. albicans cells that persist for long times during systemic infection are
predominantly white
166
In the previous experiments, we used a relatively large inocula of *C. albicans*

and sacrificed the mice after 24 hours. In the next set of experiments, we compared

white and opaque cells in a longer-term infection model initiated with a lower

inoculum. This model is the standard systemic model of fungal infection (Noble et

al., 2010; Odds et al., 2000; Perez et al., 2013). Seven mice were infected with

$5*10^5$ cells containing an approximately equal number of white-mCherry and

opaque-GFP cells. Mice were sacrificed upon signs of sickness (occurring between

7 and 14 days after injection) and fungal burden was assessed across the five

organs as described above (Figure 3a). In this experiment, *C. albicans* cells were

recovered reproducibly only from the kidney; either the *Candida* cells were cleared

from the other organs or the inoculum was sufficiently low that *Candida* did not

efficiently disseminate to the other organs.

Consistent with the previous experiments, cells recovered from the kidney

were overwhelmingly white. In several mice, a significant portion of the white cells

isolated from the kidney had been initially injected as opaque cells indicating

significant opaque-to-white switching *in vivo* (0-58% of cells, depending on the

mouse, Figure 3b, 3c, Dataset 3). We also performed an experiment with an

inoculum comprised only of opaque cells, half of which were marked with GFP and

half of which were marked with mCherry ($6.2 * 10^5$ cells). Both populations were

recovered from the kidney, and most recovered cells had switched from opaque-to-

white (Figure S2). We did observe an unanticipated outcome of this experiment.

Rather than an equal ratio of the two strains (as was present in the inoculum), the

majority of cells in each individual mouse expressed either GFP or mCherry. We

speculate that this observation reflects the stochastic nature of white-opaque

switching: there would be a strong bias towards colonization by whichever

fluorescent strain switched first.

Consistent with this interpretation is an additional control experiment: mice

injected with an inoculum containing an equal number of white-GFP and white-

mCherry cells ($5*10^5$ cells total) resulted in approximately equal numbers of GFP

and mCherry-expressing white cells in the kidneys of three of four mice (Figure S3).

Opaque cells switch to white cells in a commensal model of infection
198
To investigate how white and opaque cells compare in GI tract colonization,

we utilized the standard intestinal colonization model (Pande et al., 2013; Perez et

al., 2013; Rosenbach et al., 2010). Here, we introduced an inoculum comprised of

an equal ratio of white and opaque cells by oral gavage ($5*10^7$ cells total per

mouse, n=8 mice, Figure 4a). Colonization was monitored by collecting fecal pellets

at various time points post-inoculation. The opaque cells switched to white cells

within 24 hours and remained white until the mice were sacrificed after 17 days

(Figure 4b). All cells injected as white cells remained as white cells throughout the

experiment; we did not observe any instance of white-to-opaque switching (>3500

colony-forming units observed). We note that these results mirror, at least

superficially, those observed in the kidney following tail-vein injection infections

described above. Moreover, they are consistent with results from the oropharyngeal

candidiasis model, where introduced opaque cells were either cleared from the

oropharynx or switched rapidly to white cells (Solis et al., 2018).

The standard intestinal colonization model employed above involves a 4-day

antibiotic treatment of the mice preceding the *C. albicans* inoculation. To compare

white and opaque cells in the GI tract in the presence of an intact mouse

microbiota, we inoculated non-antibiotic treated mice with an equal number of white

and opaque cells via oral gavage ($1*10^8$ cells total per mouse, n=4 mice, Figure 4c)

and then tracked white and opaque cells during the infection through fecal pellets.

As expected, we found that significantly fewer *C. albicans* cells colonized the GI

tract when an intact microbiota was present (Figure 4d, ~100-1000 times fewer than

in the standard model described above and in Figure 4a) and that the few cells that

did colonize the GI tract were almost exclusively white cells that had been injected

as white cells. We did not observe long-term survival of any cells that were

introduced as opaque cells. These results indicate that opaque cells are out-

competed by white cells in the murine gut colonization model and that this trend is

more pronounced in the untreated mice than in the antibiotic-treated mice.

Discussion

White and opaque cells are distinct, heritable cell types of *C. albicans*

produced from the same genome. White cells and opaque cells differ in many

characteristics and are believed to be specialized for different niches in their

mammalian hosts (reviewed in Huang, 2012; Lohse and Johnson, 2009;

Morschhäuser, 2010; Soll, 2014, 1992, 2004). We used variations on two murine

models of *C. albicans* infection combined with isogenic fluorescent reporter strains

to directly compare the fate of white and opaque cells *in vivo* and to also monitor

switching between the two cell types in the host.

Many published experiments have monitored the fate of *C. albicans* white

cells in murine models of infection, and our results with white cells are entirely

consistent with this large body of data. While most of the experiments in the

literature have utilized SC5314-derived strains, which are a/α and therefore remain

locked as white cells, we used mating type **a** derivatives of SC5314, which have the

ability to undergo white-opaque switching. Despite this difference, white **a** cells, in

our experiments, behaved similarly to white a/α cells in the published literature. In a

sense, we treated white cells as a control to understand the fate of genetically

identical opaque cells that were co-injected with them. Our results show that during

systemic infection initiated by tail-vein injection, opaque cells can disseminate into

the bloodstream and colonize all five of the organs we monitored. However, the

pattern of colonization across organs differs significantly from that of white cells. For

example, when injected in equal numbers, white cells greatly predominated in the

kidney with a substantial fraction of the white cells having initially been injected as

opaque cells and undergone opaque-to-white switching. In contrast, opaque cells

outnumbered white cells in the heart and spleen. Roughly equivalent colonization of

the two cell types was observed in the liver and brain.

In the mouse GI model, white cells out-numbered opaque cells at every time

point in every mouse. All of the cells we examined that had been introduced as

opaque cells and subsequently recovered in the GI model had switched to the white

cell type, further supporting the advantage of white cells for this model.

Finally, we observed evidence of wide-spread opaque-to-white switching in

both murine models. In the GI model, extensive opaque-to-white switching occurred

early (day 1) in the colonization process. In the systemic model, we recovered white

cells that had initially been injected as opaque cells from the kidney, liver, spleen,

brain and heart (Figure 2, Figure S1). Although we do not know exactly when these

cells switched, the results suggest that it generally occurred after opaque cells had

colonized the organ in question (see results section). If this model is correct, then

switching rates must differ from organ to organ ranging from little or none occurring

in the heart to extensive switching in the brain and kidney. This conclusion is

consistent with a number of *in vitro* studies showing that both switching rates and

relative growth rates of white and opaque cells differ across media conditions (Ene

et al., 2016).

Since its discovery in 1987 by Soll and colleagues, white-opaque switching

has been extensively studied by numerous laboratories. However, the selective

pressure that has preserved white-opaque switching across a clade of species (*C.*

albicans, C. dubliniensis, and *C. tropicalis*) representing 50 million years of

evolutionary time is not understood. White and opaque cells differ in their

appearance, their mating ability, their recognition by macrophages, and their growth

on different laboratory media. Yet, the extensive gene expression changes between

the two cell types suggest that these phenotypic differences constitute only a

subset of the total range of variation. In this paper, we have documented several

additional phenotypic differences; specifically, differences in organ colonization. It

also appears that opaque-to-white switching occurs at significantly different rates in

different organs. These observations strongly support the hypothesis that white and

opaque cells are specialized to "match" different niches in the host. It is now

possible to test this hypothesis explicitly by analyzing colonization by white and

opaque cells in which different "opaque-specific" and "white-specific" genes have

been deleted.

Acknowledgements

We thank the Preclinical Therapeutic Core Facility, and especially Donghui

Wang for help with tail-vein injections and oral gavage, respectively. We thank the

Li Lab for use of the Leica stereomicroscope and members of the Johnson lab for

discussion. This study was supported by NIH grant R01AI049187 (A.D.J.).

Materials and Methods

Plasmid construction

Plasmids for GFP or mCherry tagging of *TEF2* were constructed as follows. The last 500 bp of the *TEF2* open reading frame (ORF) (excluding the stop codon) were PCR amplified with a 5' SphI site and a 3' linker for either GFP or mCherry. The 500bp immediately 3' of the *TEF2* ORF was PCR amplified between NotI and AatII sites. *C. albicans* optimized GFP (Cormack et al., 1997) was PCR amplified with a 5' 3x Gly linker and a 3' XhoI site. *C. albicans* optimized mCherry (Lohse and Johnson, 2016) was PCR amplified with a 5' GRRIPGLIN linker and a 3' XhoI site. We then merged the two *TEF2* fragments and either GFP or mCherry into one fragment during a second round of PCR. This resulted in a SphI-end of TEF2 ORF-GFP-XhoI-NotI-TEF2 3' flank-AatII fragment and a SphI-end of *TEF2* ORF-mCherry-XhoI-NotI-TEF2 3' flank-AatII fragment that were digested with SphI and AatII and ligated into pUC19 (NEB) to form pMBL183 and pMBL184. Following sequencing, the plasmids were then digested with XhoI and NotI and ligated with the similarly digested SAT1 flipper cassette from pSFS2A (Reuss et al., 2004) to form pMBL187 and pMBL188. Both plasmids were digested with SphI and AatII prior to transformation into *C. albicans.*

C. albicans **Strain construction**

MLY612/613 are derived from the switching-capable AHY135 background (Hernday et al., 2013), a strain where the *HIS1* and *LEU2* markers were added back to RZY47 (Zordan et al., 2006). AHY135 was transformed with linearized pMBL187 or pMBL188 and selected for growth on Yeast Extract Peptone Dextrose

(YEPD) supplemented with 200 μg/ml nourseothricin (clonNAT, WERNER

BioAgents, Jena, Germany). Insertion was verified by colony PCR against the 5'

and 3' flanks. The *SAT1* marker was then recycled by growth in Yeast Extract

Peptone (YEP) media lacking dextrose that was supplemented with 2% Maltose for

6-24 hours at 30°C. Cells were then plated on YEPD supplemented with 25 μg/ml

nourseothricin and grown for 24 hours at 30°C. Loss of the *SAT1* cassette was then

verified by a second round of colony PCR against the 5' and 3' flanks. The strain

genotypes are listed in detail in Table S1.

MLY612/613 were grown on synthetic complete media supplemented with

2% glucose and 100 μg/ml uridine (SD+AA+Uri) at room temperature

(approximately 22°C) and examined for opaque sectors, which were then isolated

to make opaque glycerol stocks (MLY629/642).

Gastrointestinal Tract Colonization Model (Commensal)
330
The procedure used was essentially as described previously (Perez et al.,

2013, Rosenbach et al., 2010; White et al., 2007). In brief, female Swiss Webster

mice (6 weeks old, 18–20 g) were housed two per cage and treated with antibiotics

(tetracycline [1 mg/ml]), streptomycin [2 mg/ml], and gentamycin [0.1 mg/ml]) added

to their drinking water throughout the experiment, beginning 4 days before

inoculation. In one experiment, as indicated, mice were not treated with antibiotics

prior to inoculation or for the duration of the experiment.

Prior to inoculation, *C. albicans* strains were grown for 18 h at 25°C in

SD+AA+Uri liquid medium, washed twice with PBS, and counted in a

hemocytometer. Mice were orally inoculated, with either $5*10^7$ or $1*10^8$ *C. albicans*

cells (in a 0.1 or 0.2 ml volume) as indicated, by gavage using a feeding needle.

Colonization was monitored through the collection of fecal pellets (produced at the

time of collection) at various days post-inoculation and at the end of the experiment

when the mice were sacrificed.

Fecal pellets were weighed and suspended in sterile PBS. The fecal

homogenates were serially diluted and 100 µl was plated onto SD medium

containing ampicillin (50 mg/ml) and gentamycin (15 mg/ml). The antibiotics were

used to prevent the growth of any contaminating bacteria. Colony forming units

were determined per gram of feces. All mice were sacrificed by CO_2 euthanasia

followed by cervical dislocation.

A summary of mouse experiments is tabulated in Table S2.

Systemic Infection Model

Female BALB/c mice (6 weeks old, 18–20 g) were infected with the *C.*

albicans inoculum via tail vein injection. Saturated *C. albicans* cultures were grown

at 25°C for 18 hours in SD+AA+Uri liquid medium. Cells were washed twice with

sterile saline, counted in a hemocytometer and the specified inoculum was injected

via tail vein in a 0.1 ml volume. Signs of infection were monitored twice per day

throughout the experimental time course. In some experiments, mice were

sacrificed after 24 hours. In other experiments, mice were sacrificed upon signs of

illness (including inability to ambulate, staggered gait, hypothermia or significant

weight loss). All mice were sacrificed by CO_2 euthanasia followed by cervical

dislocation. A summary of mouse experiments is tabulated in Table S2.

Organ Processing and Visualization

After dissection, organs were harvested, flash frozen, and placed in a -80° C freezer. Organs were thawed on ice, homogenized in 1 ml PBS, serially diluted in PBS, and plated onto SD medium containing ampicillin (50 mg/ml) and gentamycin (15 mg/ml). The antibiotics were used to prevent the growth of any contaminating bacteria. Colonies were visualized for fluorescence using a Leica stereomicroscope. Colony forming units were determined per gram of organ.

Data Analysis

The colony forming units per organ are tabulated in Datasets 1-4. The raw data were normalized to the inoculum such that the inoculum was always 50:50 (this is why some of the numbers are not whole numbers). These normalized colony counts were plotted in all figures and were used to determine whether or not there were statistical differences between different distributions. First, we tested whether or not there were statistical differences between cells that were injected as white and cells that were injected as opaque. In other words, irrespective of whether the cell types switched to another cell-type after they were injected, we tested whether or not there was a difference in their initial colonization. These two distributions (normalized colony counts of cell that were injected white and normalized colony counts of cells that were injected opaque) were compared with a non-parametric test, the Wilcoxon matched-pairs signed rank test, to generate p-values and determine significance. Next, we tested whether or not there were statistical differences between the stability of the cell types. In other words, we compared two distributions (normalized colony counts of cells that were injected white and

387   remained white and normalized colony counts of cells that were injected opaque

388   and remained opaque) using the same non-parametric statistical test (Wilcoxon

389   matched-pairs signed rank test) used above. The statistical analysis was conducted

390   independently for each experiment and for each organ with GraphPad Prism 7.

391   The raw data, normalized data and p-values are reported in Datasets 1-4.

Figure Legends

Figure 1: The use of fluorescent reporters does not influence *C. albicans*

activity *in vivo*. (a) Using a flowchart, the experimental setup, cell type, and

potential fluorescence phenotypes for each strain are tabulated. In this case, white

cells expressing GFP and mCherry were co-injected into the tail-veins of 4 mice.

Five organs, the kidney, liver, heart, spleen and brain, were processed to measure

white and opaque cell colonization as well as white-opaque switching. The

mechanistic interpretation of each phenotype; in other words, whether or not it

indicates cell-type switching, is also indicated. The colony-forming units of white

cells expressing GFP (yellow) and the colony-forming units of white cells

expressing mCherry (red) are plotted for each mouse for the (b) kidney, (c) liver, (d)

heart, (e) spleen and (f) brain. (g) The mean percentage (from n=4 mice) of total

cells that are white cells expressing GFP (yellow) and white cells expressing

mCherry (red) are plotted per organ as a horizontal bar graph. The raw data for this

experiment is available in Dataset 1.

Figure 2: Opaque cells can colonize multiple organs but white cells are

preferred in the kidney. (a) Using a flowchart, the experimental setup, cell type,

and potential fluorescence phenotypes for each strain are tabulated. In this case,

white cells expressing mCherry and opaque cells expressing GFP were co-injected

into the tail-veins of 8 mice. Five organs, the kidney, liver, heart, spleen and brain,

were processed to measure white and opaque cell colonization as well as white-

opaque switching. The mechanistic interpretation of each phenotype; in other

words, whether or not it indicates cell-type switching, is also indicated. The colony-

forming units of cells that remained white (i.e. white cells expressing mCherry, red),

that remained opaque (i.e. opaque cells expressing GFP, blue), and that switched

from opaque-to-white (i.e. white cells expressing GFP, dashed pink) are plotted for

each mouse for the (b) kidney, (c) liver, (d) heart, (e) spleen and (f) brain. The left

side of each horizontal bar graph refers to cells that were white at the end of the

experiment while the right side of each horizontal bar graph refers to cells that were

opaque at the end of the experiment. The four mice that received the lower

inoculums are indicated in each panel. (g) The mean percentage of total cells that

remained white (i.e. white cells expressing mCherry, red), that remained opaque

(i.e. opaque cells expressing GFP, blue), and that switched from opaque-to-white

(i.e. white cells expressing GFP, dashed pink) are plotted per organ as a horizontal

bar graph. The left side of the horizontal bar graph refers to cells that were white at

the end of the experiment while the right side refers to cells that were opaque at the

end of the experiment. The raw data for this experiment is available in Dataset 2.

Figure 3: White cells are preferred in the kidney in long-term infections. (a)

Using a flowchart, the experimental setup, cell type, and potential fluorescence

phenotypes for each strain are tabulated. In this case, white cells expressing

mCherry and opaque cells expressing GFP were co-injected into the tail-veins of 7

mice. Once the mice became ill, the kidney was processed to measure white and

opaque cell colonization as well as white-opaque switching. The mechanistic

interpretation of each phenotype; in other words, whether or not it indicates cell-

type switching, is also indicated. (b) The number of colony-forming units of cells that

remained white (i.e. white cells expressing mCherry, red), that remained opaque

(i.e. opaque cells expressing GFP, blue), and that switched from opaque-to-white

(i.e. white cells expressing GFP, dashed pink) are plotted per mouse as a horizontal

bar graph. The left side of the horizontal bar graph refers to cells that were white at

the end of the experiment while the right side refers to cells that were opaque at the

end of the experiment. (c) The percentage of total cells that remained white (i.e.

white cells expressing mCherry, red), that remained opaque (i.e. opaque cells

expressing GFP, blue), and that switched from opaque-to-white (i.e. white cells

expressing GFP, dashed pink) are plotted for each mouse as a horizontal bar

graph. The left side of the horizontal bar graph refers to cells that were white at the

end of the experiment while the right side refers to cells that were opaque at the

end of the experiment. The raw data for this experiment is available in Dataset 3.

Figure 4: White cells are preferred in a commensal model of infection. (a)

Using a flowchart, the experimental setup, cell type, and potential fluorescence

phenotypes for each strain are tabulated. In this case, white cells expressing

mCherry and opaque cells expressing GFP were inoculated into 8 mice via oral

gavage. Fecal pellets were processed to measure white and opaque cell

colonization as well as white-opaque switching. The mechanistic interpretation of

each phenotype; in other words, whether or not it indicates cell-type switching, is

also indicated. (b) The colony-forming units of white cells expressing GFP (meaning

they underwent opaque-to-white switching, blue circles) and the colony-forming

units of white cells expressing mCherry (red squares) are plotted for each mouse

over the course of the 17-day experiment. The raw data from this figure is available

in Dataset 4. (c) Using a flowchart, the experimental setup, cell type, and potential

fluorescence phenotypes for each strain are tabulated. In this case, white cells

expressing mCherry and opaque cells expressing GFP were inoculated into 4 mice

via oral gavage. Unlike the experiment in panels a-b, these mice were not treated

with antibiotics prior to the inoculation. Fecal pellets were processed to measure

white and opaque cell colonization as well as white-opaque switching. The

mechanistic interpretation of each phenotype; in other words, whether or not it

indicates cell-type switching, is also indicated. (d) The colony-forming units of white

cells expressing GFP (meaning they underwent opaque-to-white switching, blue

circles) and the colony-forming units of white cells expressing mCherry (red

squares) are plotted for each mouse over the course of the seven day experiment.

The raw data from this Figure is available in Dataset 4.

Figure S1, related to Figure 2: Opaque cells can colonize organs without

being co-injected with white cells. (a) Using a flowchart, the experimental setup,

cell type, and potential fluorescence phenotypes for each strain are tabulated. In

this case, opaque cells expressing mCherry and opaque cells expressing GFP were

co-injected into the tail-veins of 4 mice. Five organs, the kidney, liver, heart, spleen

and brain, were processed to measure white and opaque cell colonization as well

as white-opaque switching. The mechanistic interpretation of each phenotype; in

other words, whether or not it indicates cell-type switching, is also indicated. The

colony-forming units of cells that remained opaque (i.e. opaque cells expressing

mCherry (light blue) or GFP (blue)) and of cells that switched from opaque-to-white

(i.e. white cells expressing GFP (dashed pink) or mCherry (dashed orange)) are

plotted for each mouse for the (b) kidney, (c) liver, (d) heart, (e) spleen and (f) brain.

The left side of each horizontal bar graph refers to cells that were white at the end

of the experiment while the right side of each horizontal bar graph refers to cells

that were opaque at the end of the experiment. (g) The mean percentage of total

cells that remained opaque (i.e. opaque cells expressing mCherry (light blue) or

GFP (blue)) or that switched from opaque-to-white (i.e. white cells expressing GFP

(dashed pink) or mCherry (dashed orange)) are plotted per organ as a horizontal

bar graph. The left side of the horizontal bar graph refers to cells that were white at

the end of the experiment while the right side of the horizontal bar graph refers to

cells that were opaque at the end of the experiment. The raw data for this

experiment is available in Dataset 2.

Figure S2, related to Figure 3: Opaque cells switch to white cells in the

kidney. (a) Using a flowchart, the experimental setup, cell type, and potential

fluorescence phenotypes for each strain are tabulated. In this case, opaque cells

expressing mCherry and opaque cells expressing GFP were co-injected into the

tail-veins of 8 mice. Upon the onset of illness, the kidney was processed to

measure opaque cell colonization as well as opaque-to-white switching. The

mechanistic interpretation of each phenotype; in other words, whether or not it

indicates cell-type switching, is also indicated. (b) The colony-forming units of cells

that remained opaque (i.e. opaque cells expressing mCherry (light blue) or GFP

(blue)) and of cells that switched from opaque-to-white (i.e. white cells expressing

GFP (dashed pink) or mCherry (dashed orange)) are plotted per mouse as a

horizontal bar graph. The left side of the horizontal bar graph refers to cells that

were white at the end of the experiment while the right side of the horizontal bar

graph refers to cells that were opaque at the end of the experiment. (c) The

percentage of total cells that remained opaque (i.e. opaque cells expressing

mCherry (light blue) or GFP (blue)) or that switched from opaque-to-white (i.e. white

cells expressing GFP (dashed pink) or mCherry (dashed orange)) are plotted for

each mouse as a horizontal bar graph. The left side of the horizontal bar graph

refers to cells that were white at the end of the experiment while the right side of the

horizontal bar graph refers to cells that were opaque at the end of the experiment.

The raw data for this experiment is available in Dataset 3.

Figure S3, related to Figure 3: White cells are stable in the kidney. (a) Using a

flowchart, the experimental setup, cell type, and potential fluorescence phenotypes

for each strain are tabulated. In this case, white cells expressing mCherry and white

cells expressing GFP were co-injected into the tail-veins of 4 mice. Upon the onset

of illness, the kidney was processed to measure white cell colonization as well as

white-opaque switching. The mechanistic interpretation of each phenotype; in other

words, whether or not it indicates cell-type switching, is also indicated. (b) The

colony-forming units of white cells expressing GFP (yellow) and the colony-forming

units of white cells expressing mCherry (red) are plotted per mouse as a horizontal

bar graph. The left side of the horizontal bar graph refers to cells that were white at

the end of the experiment while the right side of the horizontal bar graph refers to

cells that were opaque at the end of the experiment. (c) The percentage of total

cells that remained white (i.e. white cells expressing mCherry (red) or GFP (yellow))

are plotted for each mouse as a horizontal bar graph. The left side of the horizontal

bar graph refers to cells that were white at the end of the experiment while the right

side of the horizontal bar graph refers to cells that were opaque at the end of the

experiment. The raw data for this experiment is available in Dataset 3.

Table S1: List of *C. albicans* isolates used in this paper

Table S2: Summary of all *in vivo* experiments reported in this paper

Dataset 1: Raw data, normalized data, and statistical analysis of data plotted

in Figure 1.

Dataset 2: Raw data, normalized data, and statistical analysis of data plotted

in Figures 2 and S1.

Dataset 3: Raw data, normalized data, and statistical analysis of data plotted

in Figures 3, S2 and S3.

Dataset 4: Raw data, normalized data, and statistical analysis of data plotted

in Figure 4.

References

Bain, J.M., Stubberfield, C., and Gow, N.A.R. (2001). Ura-status-dependent adhesion of *Candida albicans* mutants. FEMS Microbiol. Lett. *204*, 323–328.

Baker, S., and Bennett, J. (2007). An Overview of the Genus *Aspergillus*. pp. 3–13.

Buchanan, K. (1998). What Makes *Cryptococcus neoformans* a Pathogen? Emerg. Infect. Dis. *4*, 71–83.

Cormack, B.P., Bertram, G., Egerton, M., Gow, N.A., Falkow, S., and Brown, A.J. (1997). Yeast-enhanced green fluorescent protein (yEGFP): a reporter of gene expression in C*andida albicans*. Microbiology *143 (Pt 2)*, 303–311.

Ene, I. V., Lohse, M.B., Vladu, A. V., Morschhäuser, J., Johnson, A.D., and Bennett, R.J. (2016). Phenotypic Profiling Reveals that *Candida albicans* Opaque Cells Represent a Metabolically Specialized Cell State Compared to Default White Cells. MBio *7*, e01269-16.

Geiger, J., Wessels, D., Lockhart, S.R., and Soll, D.R. (2004). Release of a Potent Polymorphonuclear Leukocyte Chemoattractant Is Regulated by White-Opaque Switching in *Candida albicans*. Infect. Immun. *72*, 667–677.

Hernday, A.D., Lohse, M.B., Fordyce, P.M., Nobile, C.J., DeRisi, J.D., and Johnson, A.D. (2013). Structure of the Transcriptional Network Controlling White-Opaque Switching in *Candida albicans*. Mol. Microbiol. *90*, 22–35.

Huang, G. (2012). Regulation of phenotypic transitions in the fungal pathogen *Candida albicans*. Virulence *3*, 251–261.

Huang, G., Wang, H., Chou, S., Nie, X., Chen, J., and Liu, H. (2006). Bistable expression of WOR1, a master regulator of white-opaque switching in *Candida albicans*. Proc. Natl. Acad. Sci. *103*, 12813–12818.

Huang, G., Srikantha, T., Sahni, N., Yi, S., and Soll, D.R. (2009). CO_2 regulates white-to-opaque switching in *Candida albicans*. Curr. Biol. *19*, 330–334.

Huang, G., Yi, S., Sahni, N., Daniels, K.J., Srikantha, T., and Soll, D.R. (2010). N-acetylglucosamine induces white to opaque switching, a mating prerequisite in *Candida albicans*. PLoS Pathog. *6*, e1000806.

Kolotila, M.P., and Diamond, R.D. (1990). Effects of neutrophils and in vitro oxidants on survival and phenotypic switching of *Candida albicans* WO-1. Infect. Immun. *58*, 1174–1179.

Kvaal, C., Lachke, S.A., Srikantha, T., Daniels, K., McCoy, J., and Soll, D.R. (1999). Misexpression of the opaque-phase-specific gene *PEP1* (*SAP1*) in the white phase of *Candida albicans* confers increased virulence in a mouse model of cutaneous infection. Infect. Immun. *67*, 6652–6662.

Kvaal, C.A., Srikantha, T., and Soll, D.R. (1997). Misexpression of the white-phase-specific gene *WH11* in the opaque phase of *Candida albicans* affects switching and virulence. Infect. Immun. *65*, 4468–4475.

Lan, C.-Y., Newport, G., Murillo, L.A., Jones, T., Scherer, S., Davis, R.W., and Agabian, N. (2002). Metabolic specialization associated with phenotypic switching in *Candida albicans*. Proc. Natl. Acad. Sci. U. S. A. *99*, 14907–14912.

Lay, J., Henry, L.K., Clifford, J., Koltin, Y., Bulawa, C.E., and Becker, J.M. (1998). Altered expression of selectable marker URA3 in gene-disrupted *Candida albicans* strains complicates interpretation of virulence studies. Infect. Immun. *66*, 5301–5306.

Lohse, M.B., and Johnson, A.D. (2008). Differential phagocytosis of white versus opaque

Candida albicans by *Drosophila* and mouse phagocytes. PLoS One *3*, e1473.

Lohse, M.B., and Johnson, A.D. (2009). White-opaque switching in *Candida albicans*. Curr. Opin. Microbiol. *12*, 650–654.

Lohse, M.B., and Johnson, A.D. (2016). Identification and Characterization of Wor4, a New Transcriptional Regulator of White-Opaque Switching. G3 (Bethesda). *6*, 721–729.

Morschhäuser, J. (2010). Regulation of white-opaque switching in *Candida albicans*. Med. Microbiol. Immunol. *199*, 165–172.

Noble, S.M., French, S., Kohn, L.A., Chen, V., and Johnson, A.D. (2010). Systematic screens of a *Candida albicans* homozygous deletion library decouple morphogenetic switching and pathogenicity. Nat. Genet. *42*, 590–598.

Odds, F.C. (1988). Candida and candidosis (London: Bailliere Tindall).

Odds, F.C., Van Nuffel, L., and Gow, N.A. (2000). Survival in experimental *Candida albicans* infections depends on inoculum growth conditions as well as animal host. Microbiology *146 (Pt 8*, 1881–1889.

Pande, K., Chen, C., and Noble, S.M. (2013). Passage through the mammalian gut triggers a phenotypic switch that promotes *Candida albicans* commensalism. Nat. Genet. *45*, 1088–1091.

Perez, J.C., Kumamoto, C.A., Johnson, A.D. (2013). *Candida albicans* Commensalism and Pathogenicity Are Intertwined Traits Directed by a Tightly Knit Transcriptional Regulatory Circuit. PLoS Biol *11*, e1001510.

Pfaller, M.A., and Diekema, D.J. (2007). Epidemiology of Invasive Candidiasis: a Persistent Public Health Problem. Clin. Microbiol. Rev. *20*, 133–163.

Porman, A.M., Hirakawa, M.P., Jones, S.K., Wang, N., and Bennett, R.J. (2013). MTL-independent phenotypic switching in *Candida tropicalis* and a dual role for Wor1 in regulating switching and filamentation. PLoS Genet. *9*, e1003369.

Pujol, C., Daniels, K.J., Lockhart, S.R., Srikantha, T., Radke, J.B., Geiger, J., and Soll, D.R. (2004). The closely related species *Candida albicans* and *Candida dubliniensis* can mate. Eukaryot. Cell *3*, 1015–1027.

Reuss, O., Vik, A., Kolter, R., and Morschhäuser, J. (2004). The *SAT1* flipper, an optimized tool for gene disruption in *Candida albicans*. Gene *341*, 119–127.

Rikkerink, E.H., Magee, B.B., and Magee, P.T. (1988). Opaque-white phenotype transition: a programmed morphological transition in *Candida albicans*. J. Bacteriol. *170*, 895–899.

Rosenbach, A., Dignard, D., Pierce, J. V, Whiteway, M., and Kumamoto, C.A. (2010). Adaptations of *Candida albicans* for growth in the mammalian intestinal tract. Eukaryot Cell *9*, 1075–1086.

Sasse, C., Hasenberg, M., Weyler, M., Gunzer, M., and Morschhauser, J. (2013). White-Opaque Switching of *Candida albicans* Allows Immune Evasion in an Environment-Dependent Fashion. Eukaryot. Cell *12*, 50–58.

Slutsky, B., Staebell, M., Anderson, J., Risen, L., Pfaller, M., and Soll, D.R. (1987). "White-opaque transition";: a second high-frequency switching system in *Candida albicans*. J. Bacteriol. *169*, 189–197.

Solis, N. V, Park, Y.-N., Swidergall, M., Daniels, K.J., Filler, S.G., and Soll, D.R. (2018). *Candida albicans* White-Opaque Switching Influences Virulence but Not Mating during Oropharyngeal Candidiasis. Infect. Immun. *86*, e00774-e00777.

Soll, D. (2014). The role of phenotypic switching in the basic biology and pathogenesis of *Candida albicans*. J. Oral Microbiol. *6*, 22993-23005.

Soll, D.R. (1992). High-frequency switching in *Candida albicans*. Clin. Microbiol. Rev. *5*,
183–203.
Soll, D.R. (2004). Mating-type locus homozygosis, phenotypic switching and mating: a
unique sequence of dependencies in *Candida albicans*. BioEssays *26*, 10–20.
Soll, D.R., Morrow, B., and Srikantha, T. (1993). High-frequency phenotypic switching in
Candida albicans. Trends Genet. *9*, 61–65.
Soll, D.R., Anderson, J., Mihalik, R., Soll, D., Anderson, J., Soll, D., Aparicio, O.,
Billington, B., Gottschling, D., Arkowitz, R., et al. (2009). Why does *Candida albicans*
switch? FEMS Yeast Res. *9*, 973–989.
Tuch, B.B., Mitrovich, Q.M., Homann, O.R., Hernday, A.D., Monighetti, C.K., De La
Vega, F.M., and Johnson, A.D. (2010). The transcriptomes of two heritable cell types
illuminate the circuit governing their differentiation. PLoS Genet. *6*, e1001070.
Zaoutis, T.E., Argon, J., Chu, J., Berlin, J.A., Walsh, T.J., and Feudtner, C. (2005). The
Epidemiology and Attributable Outcomes of Candidemia in Adults and Children
Hospitalized in the United States: A Propensity Analysis. Clin. Infect. Dis. *41*, 1232–1239.
Zordan, R.E., Galgoczy, D.J., and Johnson, A.D. (2006). Epigenetic properties of white-
opaque switching in *Candida albicans* are based on a self-sustaining transcriptional
feedback loop. Proc. Natl. Acad. Sci. *103*, 12807–12812.

Takagi et al. Figure 1

Takagi et al. Figure 2

A.

Strains
White-mCherry
Opaque-GFP

→

***in vivo* Model**
Systemic Infection
7 mice
5×10^5 CFU

→

Sample Collection
Extended assay,
kidneys harvested
upon onset of
sickness

→

Quantification Interpretation
(colony phenotype, marker)
White, mCherry: Entered White, Stayed White
White, GFP: Entered Opaque, Switched to White
Opaque, mCherry: Entered White, Switched to Opaque
Opaque, GFP: Entered Opaque, Stayed Opaque

■ Remained White (mCherry)   ■ Remained Opaque (GFP)   ▨ Injected as Opaque, Switched to White (GFP)

B.

White Cells — Colony Forming Units — Opaque Cells

x-axis: 1.2×10^5, 6.0×10^4, 0, 6.0×10^4, 1.2×10^5

Mouse 1, Mouse 2, Mouse 3, Mouse 4, Mouse 5, Mouse 6, Mouse 7

C.

White Cells — % of Cells — Opaque Cells

x-axis: 100%, 50%, 0%, 50%, 100%

Mouse 1, Mouse 2, Mouse 3, Mouse 4, Mouse 5, Mouse 6, Mouse 7

Takagi et al. Figure 3

A.

| Strains | *in vivo* Model | Sample Collection | Quantification Interpretation (colony phenotype, marker) |

Strains
White-mCherry
Opaque-GFP

in vivo Model
GI Tract
Colonization
8 mice
5×10^7 CFU

Sample Collection
17 day assay,
feces collected
during experiment

Quantification Interpretation
(colony phenotype, marker)
White, mCherry: Entered White, Stayed White
White, GFP: Entered Opaque, Switched to White
Opaque, mCherry: Entered White, Switched to Opaque
Opaque, GFP: Entered Opaque, Stayed Opaque

■ Remained White (mCherry) ● Injected as Opaque, Switched to White (GFP)

B. With Antibiotics

CFU/g vs Time (days)

D. Without Antibiotics

CFU/g vs Time (days)

C.

Strains
White-mCherry
Opaque-GFP

in vivo Model
antibiotic-free GI
Tract Colonization
4 mice
1×10^8 CFU

Sample Collection
7 day assay,
feces collected
during experiment

Quantification Interpretation
(colony phenotype, marker)
White, mCherry: Entered White, Stayed White
White, GFP: Entered Opaque, Switched to White
Opaque, mCherry: Entered White, Switched to Opaque
Opaque, GFP: Entered Opaque, Stayed Opaque

Takagi et al. Figure 4

Takagi et al. Figure S1

Takagi et al. Figure S2

Takagi et al. Figure S3

Strain Number	Common Name	Strain Genotype
MLY612	White-GFP	WHITE MATa/a C.m. *LEU2/leu2Δ* C.d.*HIS1/his1Δ URA3/ura3Δ*::imm[434] *IRO1/iro1Δ*::imm[434] *TEF2-GFP/TEF2*
MLY613	White-mCherry	WHITE MATa/a C.m. *LEU2/leu2Δ* C.d.*HIS1/his1Δ URA3/ura3Δ*::imm[434] *IRO1/iro1Δ*::imm[434] *TEF2-mCherry/TEF2*
MLY629	Opaque-GFP	OPAQUE MATa/a C.m. *LEU2/leu2Δ* C.d.*HIS1/his1Δ URA3/ura3Δ*::imm[434] *IRO1/iro1Δ*::imm[434] *TEF2-GFP/TEF2*
MLY642	Opaque-mCherry	OPAQUE MATa/a C.m. *LEU2/leu2Δ* C.d.*HIS1/his1Δ URA3/ura3Δ*::imm[434] *IRO1/iro1Δ*::imm[434] *TEF2-mCherry/TEF2*

Table S1: *C. albicans* Strains

Inoculum Strain 1	Inoculum Strain 2	Infection Type (Figure #)	Total Inoculum	Duration	Number of Mice	Samples Harvested
White-GFP	White-mCherry	Systemic (Figure 1)	2.2×10^6	24 hours	4	5 Organs
White-mCherry	Opaque-GFP	Systemic (Figure 2)	8.0×10^5, 1.6×10^6	24 hours	8 (4 Each Inoculum)	5 Organs
Opaque-GFP	Opaque-mCherry	Systemic (Figure S1)	1.6×10^6	24 hours	4	5 Organs
White-mCherry	Opaque-GFP	Systemic (Figure 3)	5.0×10^5	Variable (Onset of Sickness)	7	Kidneys
Opaque-GFP	Opaque-mCherry	Systemic (Figure S2)	6.2×10^5	Variable (Onset of Sickness)	8	Kidneys
White-GFP	White-mCherry	Systemic (Figure S3)	5.0×10^5	Variable (Onset of Sickness)	4	Kidneys
White-mCherry	Opaque-GFP	Commensal (Figure 4B)	5.0×10^7	17 days	8	Fecal Pellets
White-mCherry	Opaque-GFP	Commensal, No Antibiotic (Figure 4D)	1.0×10^8	7 days	4	Fecal Pellets

Table S2: Summary of *in vivo* experiments