Loss of cytoplasmic incompatibility and minimal fecundity effects explain relatively low *Wolbachia* frequencies in *Drosophila mauritiana*

Megan K. Meany[1], William R. Conner[1], Sophia V. Richter[1], Jessica A. Bailey[1], Michael Turelli[2] and Brandon S. Cooper[1]¶

[1]Division of Biological Sciences, University of Montana, Missoula, MT USA
[2]Department of Evolution and Ecology, University of California, Davis, CA USA

Correspondence:
¶Division of Biological Sciences
University of Montana
32 Campus Drive, ISB
Missoula, MT 59812, USA
Email: brandon.cooper@umontana.edu

RUNNING TITLE: *w*Mau *Wolbachia* in *Drosophila mauritiana*

KEYWORDS

host-microbe interactions, introgression, maternal transmission, mitochondria, spatial spread, WO phage

ABSTRACT

Maternally transmitted *Wolbachia* bacteria infect about half of all insect species. Many *Wolbachia* cause cytoplasmic incompatibility (CI), reduced egg hatch when uninfected females mate with infected males. Although CI produces a frequency-dependent fitness advantage that leads to high equilibrium *Wolbachia* frequencies, it does not aid *Wolbachia* spread from low frequencies. Indeed, the fitness advantages that produce initial *Wolbachia* spread and maintain non-CI *Wolbachia* remain elusive. *w*Mau *Wolbachia* infecting *Drosophila mauritiana* do not cause CI, despite being very similar to CI-causing *w*No from *D. simulans* (0.068% sequence divergence over 682,494 bp), suggesting recent CI loss. Using draft *w*Mau genomes, we identify a deletion in a CI-associated gene, consistent with theory predicting that selection within host lineages does not act to increase or maintain CI. In the laboratory, *w*Mau shows near-perfect maternal transmission; but we find no significant effect on host fecundity, in contrast to published data. Intermediate *w*Mau frequencies on the island Mauritius are consistent with a balance between unidentified small, positive fitness effects and imperfect maternal transmission. Our phylogenomic analyses suggest that group-B *Wolbachia*, including *w*Mau and *w*Pip, diverged from group-A *Wolbachia*, such as *w*Mel and *w*Ri, 6–46 million years ago, more recently than previously estimated.

INTRODUCTION

Maternally transmitted *Wolbachia* infect about half of all species throughout all major insect

orders (Werren and Windsor 2000; Zug and Hammerstein 2012; Weinert et al. 2015), as well as

other arthropods (Jeyaprakash and Hoy 2000; Hilgenboecker et al. 2008) and nematodes (Taylor

et al. 2013). Host species may acquire *Wolbachia* from common ancestors, from sister species

via hybridization and introgression, or horizontally (O'Neill et al. 1992; Rousset and Solignac

1995; Huigens et al. 2004; Baldo et al. 2008; Raychoudhury et al. 2009; Gerth and Bleidorn

2016; Schuler et al. 2016; Turelli et al. 2018). *Wolbachia* often manipulate host reproduction,

inducing cytoplasmic incompatibility (CI) and male killing in *Drosophila* (Laven 1951; Yen and

Barr 1971; Hoffmann et al. 1986; Hoffmann and Turelli 1997; Hurst and Jiggins 2000). CI

reduces egg hatch when *Wolbachia*-uninfected females mate with infected males. Three

parameters usefully approximate the frequency dynamics and equilibria of CI-causing *Wolbachia*

that do not distort sex ratios: the relative hatch rate of uninfected eggs fertilized by infected

males (H), the fitness of infected females relative to uninfected females (F), and the proportion

of uninfected ova produced by infected females (μ) (Caspari and Watson 1959; Hoffmann et al.

1990). To spread deterministically from low frequencies, *Wolbachia* must produce $F(1 - \mu) > 1$,

irrespective of CI. Once they become sufficiently common, CI-causing infections, such as *w*Ri-

like *Wolbachia* in *Drosophila simulans* and several other *Drosophila* (Turelli et al. 2018), spread

to high equilibrium frequencies, dominated by a balance between CI and imperfect maternal

transmission (Turelli and Hoffmann 1995; Kreisner et al. 2016). In contrast, non-CI-causing

Wolbachia, such as *w*Au in *D. simulans* (Hoffmann et al. 1996), typically persist at lower

frequencies, presumably maintained by a balance between positive *Wolbachia* effects on host

fitness and imperfect maternal transmission (Hoffmann and Turelli 1997; Kreisner et al. 2013).

When $H < F(1 - \mu) < 1$, "bistable" dynamics result, producing stable equilibria at 0 and at a

higher frequency denoted p_s, where $0.50 < p_s \leq 1$ (Turelli and Hoffmann 1995). Bistability

explains the pattern and (slow) rate of spread of *w*Mel transinfected into *Aedes aegypti* to

suppress the spread of dengue, Zika and other human diseases (Hoffmann et al. 2011; Barton and

Turelli 2011; Turelli and Barton 2017; Schmidt et al. 2017).

In contrast to the bistability observed with *w*Mel transinfections, natural *Wolbachia*

infections seem to spread via "Fisherian" dynamics with $F(1 - \mu) > 1$ (Fisher 1937; Kriesner et

al. 2013; Hamm et al. 2014). Several *Wolbachia* effects could generate $F(1 - \mu) > 1$, but we do

not yet know which ones actually do. For example, *w*Ri has evolved to increase *D. simulans*

fecundity in only a few decades (Weeks et al. 2007), *w*Mel seems to enhance *D. melanogaster*

fitness in high and low iron environments (Brownlie et al. 2009), and several *Wolbachia*

including *w*Mel protect their *Drosophila* hosts from RNA viruses (Hedges et al. 2008; Teixeira

et al. 2008; Martinez et al. 2014). However, it remains unknown which if any these potential

fitness benefits underlie *Wolbachia* spread in nature. For instance, *w*Mel seems to have little

effect on viral abundance in wild-caught *D. melanogaster* (Webster et al. 2015; Shi et al. 2018).

D. mauritiana, D. simulans and *D. sechellia* comprise the *D. simulans* clade within the nine-

species *D. melanogaster* subgroup of *Drosophila*. The *D. simulans* clade diverged from *D.*

melanogaster approximately three million years ago (mya), with the island endemics *D. sechellia*

(Seychelles archipelago) and *D. mauritiana* (Mauritius) thought to originate in only the last few

hundred thousand years (Lachaise et al. 1986; Ballard 2000a; Dean and Ballard 2004;

McDermott and Kliman 2008; Garrigan et al. 2012; Brand et al. 2013; Garrigan et al. 2014). *D.*

simulans is widely distributed around the globe, but has never been collected on Mauritius

(David et al. 1989; Legrand et al. 2011). However, evidence of mitochondrial and nuclear

introgression supports interisland migration and hybridization between these species (Ballard

2000a; Nunes et al. 2010; Garrigan et al. 2012), which could allow introgressive *Wolbachia*

transfer (Rousset and Solignac 1995).

D. mauritiana is infected with *Wolbachia* denoted *w*Mau, likely acquired via introgression

from other *D. simulans*-clade hosts (Rousset and Solignac 1995). *Wolbachia* variant *w*Mau may

also infect *D. simulans* (denoted *w*Ma in *D. simulans*) in Madagascar and elsewhere in Africa

and the South Pacific (Ballard 2000a; Ballard 2004). *w*Mau does not cause CI in *D. mauritiana*

or when transinfected into *D. simulans* (Giordano et al. 1995). Yet it is very closely related to

*w*No strains that do cause CI in *D. simulans* (Merçot et al. 1995; Rousset and Solignac 1995;

James and Ballard 2000, 2002). (Also, *D. simulans* seems to be a "permissive" host for CI, as

evidenced by the fact that *w*Mel, which causes little CI in its native host, *D. melanogaster*,

causes intense CI in *D. simulans* [Poinsot et al. 1998].) Fast et al. (2011) reported that a *w*Mau

variant increased *D. mauritiana* fecundity four-fold. This fecundity effect occurred in concert

with *w*Mau-induced alternations of programmed cell death in the germarium and of germline

stem cell mitosis, possibly providing insight into the mechanisms underlying increased egg

production (Fast et al. 2011). However, the generality of this finding across *w*Mau variants and

host genetic backgrounds remains unknown.

Here, we assess the genetic and phenotypic basis of *w*Mau frequencies in *D. mauritiana* on

Mauritius by combining analysis of *w*Mau draft genomes with analysis of *w*Mau transmission in

the laboratory and *w*Mau effects on host fecundity and egg hatch. We identify a single mutation

that disrupts a locus associated with CI. The loss of CI in *w*Mau is consistent with theory

demonstrating that selection within host species does not act to increase or maintain the level of

CI (Prout 1994; Turelli 1994; Haygood and Turelli 2009), but instead acts to increase $F(1 - \mu)$,

the product of *Wolbachia* effects on host fitness and maternal transmission efficiency (Turelli

1994). The loss of CI helps explain the intermediate *w*Mau frequencies on Mauritius, reported by

us and Giordano et al. (1995). We find no *w*Mau effects on host fecundity, and theoretical

analyses show that even a two-fold fecundity increase cannot be reconciled with the observed

intermediate population frequencies, unless maternal *w*Mau transmission is exceptionally

unreliable in the field. Finally, we present theoretical analyses illustrating that the persistence of

two distinct classes of mtDNA haplotypes among *Wolbachia*-uninfected *D. mauritiana* is

unexpected under a simple null model. Together, our results contribute to understanding the

genomic and phenotypic basis of global *Wolbachia* persistence, which is relevant to improving

Wolbachia-based biocontrol of human diseases (Ritchie 2018).

MATERIALS AND METHODS

***Drosophila* Husbandry and Stocks**

The *D. mauritiana* isofemale lines used in this study ($N = 32$) were sampled from Mauritius in

2006 by Margarita Womack and kindly provided to us by Prof. Daniel Matute from the

University of North Carolina, Chapel Hill. We also obtained four *D. simulans* stocks (lines 196,

297, 298, and 299) from the National *Drosophila* Species Stock Center that were sampled from

Madagascar. Stocks were maintained on modified version of the standard Bloomington-cornmeal

medium (Bloomington Stock Center, Bloomington, IN) and were kept at 25°C, 12 light:12 dark

photoperiod prior to the start of our experiments.

Determining *Wolbachia* infection status and comparing infection frequencies

One to two generations prior to our experiments DNA was extracted from each isofemale line

using a standard 'squish' buffer protocol (Gloor et al. 1993), and infection status was determined

using a polymerase chain reaction (PCR) assay (Simpliamp ThermoCycler, Applied Biosystems,

Singapore). We amplified the *Wolbachia*-specific *wsp* gene (Forward: 5'-

TGGTCCAATAAGTGATGAAGAAAC-3'; Reverse: 5'-AAAAATTAAACGCTACTCCA-3';

Braig et al. 1998) and a nuclear control region of the *2L* chromosome (Forward: 5'-

TGCAGCTATGGTCGTTGACA-3'; Reverse: 5'-ACGAGACAATAATATGTGGTGCTG-3';

designed here). PCR products were visualized using 1% agarose gels that included a molecular-

weight ladder. Assuming a binomial distribution, we estimated exact 95% binomial confidence

intervals for the infection frequencies on Mauritius. Using Fisher's Exact Test, we tested for

temporal differences in *w*Mau frequencies by comparing our frequency estimate to a previous

estimate (Giordano et al. 1995). All analyses were performed using R version 3.5.1 (R Team

2015).

We used quantitative PCR (qPCR) (MX3000P, Agilent Technologies, Germany) to confirm

that tetracycline-treated flies were cleared of *w*Mau. DNA was extracted from *D. mauritiana*

flies after four generations of tetracycline treatment (1-2 generations prior to completing our

experiments), as described below. Our qPCR used a PowerUp™ SYBR™ Green Master Mix

(Applied Biosystems™, California, USA) and amplified *Wolbachia*-specific *wsp* (Forward: 5'-

CATTGGTGTTGGTGTTGGTG-3'; Reverse: 5'-ACCGAAATAACGAGCTCCAG-3') and

Rpl32 as a nuclear control (Forward: 5'-CCGCTTCAAGGGACAGTATC-3'; Reverse: 5'-

CAATCTCCTTGCGCTTCTTG-3';

Newton and Sheehan 2014).

***Wolbachia* DNA extraction, library preparation, and sequencing**

We sequenced *w*Mau-infected *R9*, *R29*, and *R60 D. mauritiana* genotypes. Tissue samples for

genomic DNA were extracted using a modified CTAB Genomic DNA Extraction protocol. DNA

quantity was tested on an Implen Nanodrop (Implen, München, Germany) and total DNA was

quantified by Qubit Fluorometric Quantitation (Invitrogen, Carlsbad, California, USA). DNA

was cleaned using Agencourt AMPure XP beads (Beckman Coulter, Inc., Brea, CA, U.S.A),

following manufacturers' instructions, and eluted in 50 µl 1× TE Buffer for shearing. DNA was

sheared using a Covaris E220 Focused Ultrasonicator (Covaris Inc., Woburn, MA) to a target

size of 400 bp. We prepared libraries using NEBNext® Ultra™ II DNA Library Prep with

Sample Purification Beads (New England BioLabs, Ipswich, Massachusetts). Final fragment

sizes and concentrations were confirmed using a TapeStation 2200 system (Agilent, Santa Clara,

California). We indexed samples using NEBNext® Multiplex Oligos for Illumina® (Index

Primers Set 3 & Index Primers Set 4), and 10 µl of each sample was shipped to Novogene

(Sacramento, CA) for sequencing using Illumina HiSeq 4000 (San Diego, CA), generating

paired-end, 150 bp reads.

***Wolbachia* assembly**

We obtained published reads ($N = 6$) from Garrigan et al. (2014), and assembled these genomes

along with the *R9*, *R29*, and *R60* genomes that we sequenced. Reads were trimmed using Sickle

v. 1.33 (Joshi and Fass 2011) and assembled using ABySS v. 2.0.2 (Jackman et al. 2017). *K*

values of 41, 51, 61, and 71 were used, and scaffolds with the best nucleotide BLAST matches to

known *Wolbachia* sequences with E-values less than 10^{-10} were extracted as the draft *Wolbachia*

assemblies. We deemed samples infected if the largest *Wolbachia* assembly was at least 1

million bases and uninfected if the largest assembly was fewer than 100,000 bases. No samples

produced *Wolbachia* assemblies between 100,000 and 1 million bases. Of the six sets of

published reads we analyzed (Garrigan et al. 2014), only lines *R31* and *R41* were *w*Mau-infected.

We also screened the living copies of these lines for *wsp* using PCR, and both were infected,

supporting reliable *w*Mau transmission in the lab since these lines were sampled in nature.

To assess the quality of our draft assemblies, we used BUSCO v. 3.0.0 to search for

homologs of the near-universal, single-copy genes in the BUSCO proteobacteria database

(Simao et al. 2015). As a control, we performed the same search using the reference genomes for

*w*Ri (Klasson et al. 2009), *w*Au (Sutton et al. 2014), *w*Mel (Wu et al. 2004), *w*Ha (Ellegaard et

al. 2013), and *w*No (Ellegaard et al. 2013).

***Wolbachia* gene extraction and phylogenetics**

To determine phylogenetic relationships and estimate divergence times, we obtained the public

Wolbachia group-B genomes of: *w*AlbB that infects *Aedes albopictus* (Mavingui et al. 2012),

*w*Pip_Pel that infects *Culex pipiens* (Klasson et al. 2008), *w*Pip_Mol that infects *Culex molestus*

(Pinto et al. 2013), *w*No that infects *Drosophila simulans* (Ellegaard et al. 2013), and *w*VitB that

infects *Nasonia vitripennis* (Kent et al. 2011); in addition to group-A genomes of: *w*Mel that

infects *D. melanogaster* (Wu et al. 2004), *w*Suz that infects *D. suzukii* (Siozios et al. 2013), four

Wolbachia that infect *Nomada* bees (*w*NFe, *w*NPa, *w*NLeu, and *w*NFa; Gerth and Bleidorn

2016), and three *Wolbachia* that infect *D. simulans* (*w*Ri, *w*Au and *w*Ha; Klasson *et al.* 2009;

Sutton *et al.* 2014; Ellegaard *et al.* 2013). The previously published genomes and the five *w*Mau-

infected *D. mauritiana* genomes were annotated with Prokka v. 1.11, which identifies homologs

to known bacterial genes (Seemann 2014). To avoid pseudogenes and paralogs, we used only

genes present in a single copy, and with no alignment gaps, in all of the genome sequences.

Genes were identified as single copy if they uniquely matched a bacterial reference gene

identified by Prokka v. 1.11. By requiring all homologs to have identical length in all of the draft

Wolbachia genomes, we removed all loci with indels. 143 genes, a total of 113,943 bp, met these

criteria when comparing all of these genomes. However, when our analysis was restricted to the

five *w*Mau genomes, our criteria were met by 694 genes, totaling 704,613 bp. Including *w*No

with the five *w*Mau genomes reduced our set to 671 genes with 682,494 bp. We calculated the

percent differences for the three codon positions within *w*Mau and between *w*Mau and *w*No.

We estimated a Bayesian phylogram of the *Wolbachia* sequences with RevBayes 1.0.8

under the GTR + Γ model, partitioning by codon position (Höhna et al. 2016). Four independent

runs were performed, which all agreed.

We estimated a chronogram from the *Wolbachia* sequences using the absolute chronogram

procedure implemented in Turelli et al. (2018). Briefly, we generated a relative relaxed-clock

chronogram with the GTR + Γ model with the root age fixed to 1 and the data partitioned by

codon position. The relaxed clock branch rate prior was Γ(2,2). We used substitution-rate

estimates of Γ(7,7) × 6.87×10^{-9} substitutions/3rd position site/year to transform the relative

chronogram into an absolute chronogram. This rate estimate was chosen so that the upper and

lower credible intervals matched the posterior distribution estimated by Richardson et al. (2012),

assuming 10 generations/year, normalized by their median estimate of 6.87×10^{-9} substitutions/3rd

position site/year. Although our relaxed-clock analyses allow for variation in substitution rates

across branches, our conversion to absolute time depends on the unverified assumption that the

median substitution rate estimated by Richardson et al. (2012) for *w*Mel is relevant across these

broadly diverged *Wolbachia*. (To assess the robustness of our conclusions to model assumptions,

we also performed a strict-clock analysis and a relaxed-clock analysis with branch-rate prior

$\Gamma(7,7)$.) For each analysis, four independent runs were performed, which all agreed. Our analyses

all support wNo as sister to wMau.

We also estimated a relative chronogram for the host species using the procedure

implemented in Turelli et al. (2018). Our host phylogeny was based on the same 20 nuclear

genes used in Turelli et al. (2018): *aconitase, aldolase, bicoid, ebony, enolase, esc, g6pdh, glyp,*

glys, ninaE, pepck, pgi, pgm, pic, ptc, tpi, transaldolase, white, wingless and *yellow*.

Analysis of *Wolbachia* and mitochondrial genomes

We looked for copy number variation (CNV) between wMau and its closest relative, wNo across

the whole wNo genome. Reads from the five infected wMau lines were aligned to the wNo

reference (Ellegaard et al. 2013) with bwa 0.7.12 (Li and Durbin 2009). We calculated the

normalized read depth for each alignment over sliding 1,000-bp windows by dividing the

average depth in the window by the average depth over the entire wNo genome. The results were

plotted and visually inspected for putative copy number variants (CNVs). The locations of CNVs

were specifically identified with ControlFREEC v. 11.5 (Boeva et al. 2012), using a ploidy of

one and a window size of 1,000. We calculated P-values for each identified CNV with the

Wilcoxon Rank Sum and the Kolmogorov-Smirnov tests implemented in ControlFREEC.

We used BLAST to search for pairs of CI-factor (*cif*) homologs in wMau and wNo genomes

that are associated with CI (Beckmann and Fallon 2013; Beckmann et al. 2017; LePage et al.

2017; Lindsey et al. 2018; Beckmann et al. 2019). (We adopt Beckmann et al. (2019)'s

nomenclature that assigns names to loci based on their predicted enzymatic function, with

superscripts denoting the focal *Wolbachia* strain.) These include predicted CI-inducing

deubiquitylase (*cid*) wPip_0282-wPip_0283 (*cidA-cidB*$^{w\text{Pip}}$) and CI-inducing nuclease (*cin*)

wPip_0294-wPip_0295 (*cinA-cinB*$^{w\text{Pip}}$) pairs that induce toxicity and rescue when expressed/co-

expressed in *Saccharomyces cerevisiae* (Beckmann et al. 2017 and Beckmann et al. 2019);

WD0631-WD632 (*cidA-cidB*$^{w\text{Mel}}$) that recapitulate CI when transgenically expressed in *D.*

melanogaster (LePage *et al*. 2017); and wNo_RS01055 and wNo_RS01050 that have been

identified as a Type III *cifA-cifB* pair in the wNo genome (LePage et al. 2017; Lindsey et al.

2018). wNo_RS01055 and wNo_RS01050 are highly diverged from *cidA-cidB*$^{w\text{Mel}}$ and *cidA-*

cidB$^{w\text{Pip}}$ homologs and from *cinA-cinB*$^{w\text{Pip}}$; however, this wNo pair is more similar to *cinA-*

cinB[wPip] in terms of protein domains, lacking a ubiquitin-like protease domain (Lindsey et al.

2018). We refer to these loci as *cinA-cinB*[wNo].

We found only homologs of the *cinA-cinB*[wNo] pair in *w*Mau genomes, which we extracted

from our draft *w*Mau assemblies and aligned with MAFFT v. 7 (Katoh and Standley 2013). We

compared *cinA-cinB*[wNo] to the *w*Mau homologs to identify single nucleotide variants (SNVs)

among our *w*Mau assemblies.

D. mauritiana carry either the *ma*I mitochondrial haplotype, associated with *w*Mau

infections, or the *ma*II haplotype (Rousset and Solignac 1995; Ballard 2000a; James and Ballard

2000). To determine the mitochondrial haplotype of each *D. mauritiana* line, we assembled the

mitochondrial genomes by down-sampling the reads by a factor of 100, then assembling with

ABySS 2.0.2 using a *K* value of 71 for our data (150 bp reads) and 35 for the published data (76

bp reads) (Garrigan et al. 2014). Down-sampling reads prevents the nuclear genome from

assembling but does not inhibit assembly of the mitochondrial genome, which has much higher

coverage. We deemed the mitochondrial assembly complete if all 13 protein-coding genes were

present on the same contig and in the same order as in *D. melanogaster*. If the first attempt did

not produce a complete mitochondrial assembly, we adjusted the down-sampling fraction until a

complete assembly was produced for each line.

Annotated reference mitochondrial sequences for the *D. mauritiana* mitochondrial

haplotypes *ma*I and *ma*II were obtained from Ballard et al. (2000b), and the 13 protein-coding

genes were extracted from our assemblies using BLAST and aligned to these references. The

*ma*I and *ma*II reference sequences differ at 343 nucleotides over these protein-coding regions.

We identified our lines as carrying the *ma*I haplotype if they differed by fewer than five

nucleotides from the *ma*I reference and as *ma*II if they differed by fewer than five nucleotides

from the *ma*II reference. None of our assemblies differed from both references at five or more

nucleotides.

***w*Mau phenotypic analyses**

Previous analyses have demonstrated that *w*Mau does not cause CI (Giordano et al. 1995). To

check the generality of this result, we reciprocally crossed *w*Mau-infected *R31 D. mauritiana*

with uninfected *R4* and measured egg hatch. Flies were reared under controlled conditions at

25°C for multiple generations leading up to the experiment. We paired 1–2-day-old virgin

females with 1–2-day-old males in a vial containing spoons with cornmeal media and yeast

paste. After 24 hr, pairs were transferred to new spoons, and this process was repeated for five

days. Eggs on each spoon were given 24 hr at 25°C to hatch after flies were removed. To test for

CI, we used nonparametric Wilcoxon tests to compare egg hatch between reciprocal crosses that

produced at least 10 eggs. All experiments were carried out at 25°C with a 12 light:12 dark

photoperiod.

To determine if wMau generally enhances *D. mauritiana* fecundity, we assessed the

fecundity of two wMau-infected isofemale lines from Mauritius (*R31* and *R41*); we also

reciprocally introgressed wMau from each of these lines to assess host effects. To do this we

crossed *R31* females with *R41* males and backcrossed F_1 females to *R41* males—this was

repeated for four generations to generate the reciprocally introgressed *R41^{R31}* genotypes (wMau

variant denoted by superscripts). A similar approach was taken to generate *R31^{R41}* genotypes.

This approach has previously revealed *D. teissieri*-host effects on wTei-induced CI (Cooper et al.

2017). To assay fecundity, we reciprocally crossed each genotype (*R31, R41, R31^{R41}, R31^{R41}*) to

uninfected line *R4* to generate paired infected- and uninfected-F_1 females with similar genetic

backgrounds. The wMau-infected and uninfected F_1 females were collected as virgins and placed

in holding vials. We paired 3–7-day-old females individually with an uninfected-*R4* male (to

stimulate oviposition) in vials containing a small spoon filled with standard cornmeal medium

and coated with a thin layer of yeast paste. We allowed females to lay eggs for 24 hours, after

which pairs were transferred to new vials. This was repeated for five days. At the end of each 24-

hr period, spoons were frozen until counted. All experiments were carried out at 25°C with a 12

light:12 dark photoperiod.

We also measured egg lay of wMau-infected (*R31*) and tetracycline-cleared uninfected

(*R31-tet*) genotypes over 24 days, on apple-agar plates, to more closely mimic the methods of

Fast et al. (2011). We fed flies 0.03% tetracycline concentrated medium for four generations to

generate the *R31-tet* genotype. We screened F_1 and F_2 individuals for wMau using PCR, and we

then fed flies tetracycline food for two additional generations. In the fourth generation, we

assessed wMau titer using qPCR to confirm that each genotype was cleared of wMau infection.

We reconstituted the gut microbiome by rearing *R31-tet* flies on food where *R31* males had fed

and defecated for 48 hours. Flies were given at least three more generations to avoid detrimental

effects of tetracycline treatment on mitochondrial function (Ballard and Melvin 2007). We then

paired individual 6–7-day-old virgin *R31* ($N = 30$) and *R31-tet* ($N = 30$) females in bottles on

yeasted apple-juice agar plates with an *R4* male to stimulate oviposition. Pairs were placed on

new egg-lay plates every 24 hrs. After two weeks, we added one or two additional *R4* males to

each bottle to replace any dead males and to ensure that females were not sperm limited as they

aged.

We used nonparametric Wilcoxon tests to assess *w*Mau effects on host fecundity. We then

estimated the fitness parameter *F* in the standard discrete-generation model of CI (Hoffmann et

al. 1990; Turelli 1994). We used the '*pwr.t2n.test*' function in the '*pwr*' library in R to assess the

power of our data to detect increases to *F*. Pairs that laid fewer than 10 eggs across each

experiment were excluded from analyses, but our results are robust to this threshold.

To estimate the fidelity of maternal transmission, *R31* and *R41* females were reared at 25°C

for several generations prior to our experiment. In the experimental generation, 3-5 day old

inseminated females were placed individually in vials that also contained two males. These *R31*

($N = 17$) and *R41* ($N = 19$) sublines were allowed to lay eggs for one week. In the following

generation we screened F1 offspring for *w*Mau infection using PCR as described above.

RESULTS

Wolbachia **infection status**

Out of 32 *D. mauritiana* lines that we analyzed, 11 were infected with *w*Mau *Wolbachia*

(infection frequency = 0.34; binomial confidence interval: 0.19, 0.53). In contrast, none of the *D.*

simulans stocks ($N = 4$) sampled from Madagascar were infected, precluding our ability to

directly compare *w*Mau and *w*Ma. Our new *w*Mau frequency estimate is not statistically different

from a previous estimate (Giordano et al. 1995: infection frequency, 0.46; binomial confidence

interval, (0.34, 0.58); Fisher's Exact Test, $P = 0.293$), based largely on assaying a heterogenous

collecton of stocks in various laboratories. These relatively low infection frequencies are

consistent with theoretical expectations given that *w*Mau does not cause CI (Giordano et al.

1995; our data reported below). The intermediate *w*Mau frequencies on Mauritius suggest that

*w*Mau persists at a balance between positive effects on host fitness and imperfect maternal

transmission. Quantitative predictions, based on the idealized model of Hoffmann and Turelli

(1997), are discussed below. The maintenance of *w*Mau is potentially analogous to the

persistence of other non-CI-causing *Wolbachia*, specifically *w*Au in some Australian populations

of *D. simulans* (Hoffmann et al. 1996; Kriesner et al. 2013) and *w*Suz in *D. suzukii* and *w*Spc in

D. subpulchrella (Hamm et al. 2014; Conner et al. 2017; Turelli et al. 2018; but see Cattel et al.

2018).

Draft *w*Mau genome assemblies and comparison to *w*No

The five *w*Mau draft genomes we assembled were of very similar quality (Supplemental Table

1). N50 values ranged from 60,027 to 63,676 base pairs, and our assemblies varied in total length

from 1,266,004 bases to 1,303,156 bases (Supplemental Table 1). Our BUSCO search found

exactly the same genes in each draft assembly, and the presence/absence of genes in our *w*Mau

assemblies was comparable to those in the complete genomes used as controls (Supplemental

Table 2). In comparing our five *w*Mau draft genomes over 694 single-copy, equal-length loci

comprising 704,613 bp, we found only one SNP. Four sequences (*R9*, *R31, R41* and *R60*) are

identical at all 704,613 bp. *R29* differs from them at a single nucleotide, a nonsynonymous

substitution in a locus which Prokka v. 1.11 annotates as "bifunctional DNA-directed RNA

polymerase subunit beta/beta."

Comparing these five *w*Mau sequences to the *w*No reference (Ellegaard et al. 2013) over

671 genes with 682,494 bp, they differ by 0.068% overall, with equivalent divergence at all three

codon positions (0.067%, 0.061%, and 0.076%, respectively).

***Wolbachia* phylogenetics**

As expected from the sequence comparisons, our group-B phylogram places *w*Mau sister to *w*No

(Figure 1A). This is consistent with previous analyses using fewer loci that placed *w*Mau (or

*w*Ma in *D. simulans*) sister to *w*No (James and Ballard 2000; Zabalou et al. 2008; Toomey et al.

2013). Our chronogram (Figure 1B) estimates the 95% credible interval for the split between the

group-B versus group-A *Wolbachia* strains as 6 to 36 mya (point estimate, 16 mya). Reducing

the variance on the substitution-rate-variation prior by using $\Gamma(7,7)$ rather than $\Gamma(2,2)$, changes

the credible interval for the A-B split to 8 to 46 mya (point estimate, 21 mya). In contrast, a strict

clock analysis produces a credible interval of 12 to 64 mya (point estimate, 31 mya). These

estimates are roughly comparable to an earlier result based on a general approximation for the

synonymous substitution rate in bacteria (Ochman and Wilson 1987) and data from only the *ftsZ*

locus (59–67 mya, Werren et al. 1995). However, our estimates are much lower than an

alternative estimate based on comparative genomics (217 mya, Gerth and Bleidorn 2016). We

discuss this discrepancy below.

The observed divergence between *w*No and *w*Mau is consistent across all three codon

positions, similar to other recent *Wolbachia* splits like that between *w*Ri and *w*Suz (Turelli et al.

2018). Conversely, observed divergence at each codon position generally varies across the

chronogram, leading to inflation of the *w*No-*w*Mau (181,476 years; credible interval = 27,711 to

701,205 years; Figure 1B) and *w*Ri-*w*Suz (16,214; credible interval = 1,254 to 70,584)

divergence point estimates; the latter is about 1.6 times as large as the value in Turelli et al.

(2018). (Nevertheless, the confidence intervals of our and Turelli et al. (2018)'s *w*Ri-*w*Suz

divergence estimates overlap.) To obtain an alternative estimate of *w*No-*w*Mau divergence, we

estimated divergence time using the observed third-position pairwise divergence (0.077%, or

0.039% from tip to MRCA) and Richardson et al. (2012)'s estimate of the "short-term

evolutionary rate" of *Wolbachia* third-position divergence within *w*Mel. This approach produces

a point estimate of 57,000 years, with a credible interval of 30,000 to 135,000 years for the *w*No-

*w*Mau split. In Cooper et al. (2019), we address in detail how a constant substitution-rate ratio

among codon positions across the tree, assumed by the model, affects these estimates.

Figure 1. A) An estimated phylogram for various group-A (red) and group-B (blue) *Wolbachia*
strains. All nodes have Bayesian posterior probabilities of 1. The phylogram shows significant
variation in the substitution rates across branches, with long branches separating the A and B
clades. **B)** An estimated chronogram for the same strains, with estimated divergence times and
their confidence intervals at each node. To obtain these estimates, we generated a relative
relaxed-clock chronogram with the GTR + Γ model with the root age fixed to 1, the data
partitioned by codon position, and with a Γ(2,2) branch rate prior. We used substitution-rate
estimates of $\Gamma(7,7) \times 6.87 \times 10^{-9}$ substitutions/3rd position site/year to transform the relative
chronogram into an absolute chronogram.

Analysis of *Wolbachia* and mitochondrial genomes

We looked for CNVs in *w*Mau relative to sister *w*No by plotting normalized read depth along the

*w*No genome. There were no differences in copy number among the *w*Mau variants, but

compared to *w*No, ControlFREEC identified four regions deleted from all *w*Mau that were

significant according to the Wilcoxon Rank Sum and Kolmogorov-Smirnov tests (Figure 2 and

Supplemental Table 3). These deleted regions of the *w*Mau genomes include many genes,

including many phage-related loci, providing interesting candidates for future work (listed in

Supplementary Table 4).

Figure 2. All *w*Mau variants share four large deletions, relative to sister *w*No. Top panel) The
normalized read depth for *w*Mau *R60* plotted across the four focal regions of the *w*No reference
genome; 10 kb of sequence surrounding regions are plotted on either side of each region. Bottom
panel) The normalized read depth of *w*Mau *R60* plotted across the whole *w*No reference genome.
Regions that do not contain statistically significant CNVs are plotted in dark blue, and regions
with significant CNVs are plotted in red. All *w*Mau variants share the same CNVs, relative to
*w*No.

To test the hypothesis that *cif* loci are disrupted, we searched for pairs of loci known to be

associated with CI and found homologs to the *cinA-cinB*[*w*No] pair in each of our draft assemblies,

but we did not find homologs to the *cidA-cidB*[*w*Mel], *cidA-cidB*[*w*Pip], or to the *cinA-cinB*[*w*Pip] pairs.

There were no variable sites in *cinA-cinB*[*w*No] homologs among our five *w*Mau assemblies.

Relative to *w*No, all *w*Mau variants share a one base pair deletion at base 1133 out of 2091

(amino acid 378) in the *cinB*[*w*No] homolog. This frameshift introduces over 10 stop codons, with

the first at amino acid 388, potentially making this predicted CI-causing-toxin protein

nonfunctional. We also identified a nonsynonymous substitution in amino acid 264 of the

cinB[*w*No] homolog (*w*No codon ACA, Thr; *w*Mau codon AUA, Ile) and two SNVs in the region

homologous to *cinA*[*w*No]: a synonymous substitution in amino acid 365 (*w*No codon GUC, *w*Mau

codon GUU) and a nonsynonymous substitution in amino acid 397 (*w*No codon GCU, amino

acid Ala; *w*Mau codon GAU, amino acid Asp). Disruption of CI is consistent with theoretical

analyses showing that selection within a host species does not act directly on the level of CI

(Prout 1994; Turelli 1994; Haygood and Turelli 2009). Future functional analyses will determine

whether disruption of regions homologous to *cinA-cinB*[*w*No] underlie the lack of *w*Mau CI.

Of the *D. mauritiana* lines tested ($N = 9$), one line (uninfected-*R44*) carries the *ma*II

mitochondrial haplotype, while the other eight carry *ma*I. Rousset and Solignac (1995) reported a

similar *ma*II frequency, with 3 of 26 lines sampled in 1985 carrying *ma*II. The *ma*I and *ma*II

references differ by 343 SNVs across the proteome, and *R44* differs from the *ma*II reference by 4

SNVs in the proteome. Four of our *ma*I lines (*R23*, *R29*, *R32*, and *R39*) are identical to the *ma*I

reference, while three (*R31*, *R41*, and *R60*) have one SNV and one (*R9*) has two SNVs relative to

*ma*I reference. One SNV is shared between *R9* and *R60*, but the other three SNVs are unique.

Our results agree with past analyses that found *w*Mau is perfectly associated with the *ma*I

mitochondrial haplotype (Rousset and Solignac 1995; Ballard 2000a; James and Ballard 2000).

The presence of *ma*II among the uninfected is interesting. In contrast to *ma*I, which is associated

with introgression with *D. simulans* (Ballard 2000a; James and Ballard 2000), *ma*II appears as an

outgroup on the mtDNA phylogeny of the *D. simulans* clade and is not associated with

Wolbachia (Ballard 2000b, Fig. 5; James and Ballard 2000). Whether or not *Wolbachia* cause CI,

if they are maintained by selection-imperfect-transmission balance, we expect all uninfected flies

to eventually carry the mtDNA associated with infected mothers (Turelli et al. 1992). We present

a mathematical analysis of the persistence of *ma*II below.

Analysis of *w*Mau phenotypes

In agreement with Giordano et al. (1995), we found no difference between the egg hatch of

uninfected females crossed to *w*Mau-infected males (0.34 ± 0.23 SD, $N = 25$) and the reciprocal

cross (0.29 ± 0.28 SD, N = 24), indicating no CI. In contrast to Fast et al. (2011), we find no

evidence that *w*Mau affects *D. mauritiana* fecundity (Supplemental Table 5 and Figure 3),

regardless of host genetic backgrounds. Across both experiments assessing *w*Mau fecundity

effects in their natural backgrounds (*R31* and *R41*), we counted 27,221 eggs and found no

difference in the number of eggs laid by infected (mean = 238.20, SD = 52.35, N = 60) versus

uninfected (mean = 226.82, SD = 67.21, N = 57) females over the five days of egg lay (Wilcoxon

test, W = 1540.5, P = 0.357); and across both experiments that assessed *w*Mau fecundity effects

in novel host backgrounds (*R31^{R41}* and *R41^{R31}*), we counted 30,358 eggs and found no difference

in the number of eggs laid by infected (mean = 253.30, SD = 51.99, N = 60) versus uninfected

(mean = 252.67, SD = 63.53, N = 60) females over five days (Wilcoxon test, W = 1869.5, P =

0.719). [The mean number of eggs laid over five days, standard deviation (SD), sample size (*N*),

and *P*-values from Wilcoxon tests are presented in Supplemental Table 5 for all pairs.]

Figure 3. *w*Mau infections do not influence *D. mauritiana* fecundity, regardless of host genomic
background. **A)** Violin plots of the number of eggs laid by *D. mauritiana* females over five days
when infected with their natural *w*Mau variant (*R31I* and *R41I*), when infected with a novel
*w*Mau variant (*R31^{R41}I* and *R41^{R31}I*), and when uninfected (*R31U, R41U, R31^{R41}U*, and
R41^{R31}U). Large black dots are medians, and small black dots are the eggs laid by each replicate
over five days. **B)** The daily egg lay of these same infected (solid circles) and uninfected (open
circles) *R31* (aqua), *R41* (red)*, R31^{R41}* (gray), and *R41^{R31}* (dark blue) genotypes is reported. Large
circles are means of all replicates, and small circles are the raw data. Only days where females
laid at least one egg are plotted. Cytoplasm sources are denoted by superscripts for the
reciprocally introgressed strains.

We sought to determine if *w*Mau fecundity effects depend on host age with a separate

experiment that assessed egg lay over 24 days on apple-agar plates, similar to Fast et al. (2011).

Across all ages, we counted 9,459 eggs and found no difference in the number of eggs laid by

infected (mean = 156.29, SD = 138.04, N = 28) versus uninfected (mean = 187.70, SD = 168.28,

N = 27) females (Wilcoxon test, W = 409, P = 0.608) (Figure 4). While our point estimates

indicate that *w*Mau does not increase host fecundity, egg lay was generally lower and more

variable on agar plates relative to our analyses of egg lay on spoons described above.

Figure 4. The mean number of eggs laid by infected *R31* (*R31I*, large aqua dots) and uninfected
R31-tet (large tan dots) genotypes are similar. Egg counts for each replicate are also plotted
(small dots). Violin plots show egg lay across all ages for each genotype; large black circles are
medians, and small black circles are the number of eggs laid by each replicate.

With these data, we estimated the fitness parameter F in the standard discrete-generation

model of CI (Hoffmann et al. 1990; Turelli 1994). Taking the ratio of replicate mean fecundity

observed for wMau-infected females to the replicate mean fecundity of uninfected females in

naturally sampled $R31$ and $R41$ $D. mauritiana$ backgrounds, we estimated $F = 1.05$ (95% BC$_a$

interval: 0.96, 1.16). Following reciprocal introgression of wMau and host backgrounds (i.e., the

$R31^{R41}$ and $R41^{R31}$ genotypes), we estimated $F = 1.0$ (95% BC$_a$ interval: 0.93, 1.09). Finally,

across all 24 days of our age-effects experiment, we estimated $F = 0.83$, 95% BC$_a$ interval: 0.52,

1.32) for $R31$, which overlaps with our estimate of F for $R31$ in our initial experiment (Table 1).

BC$_a$ confidence intervals were calculated using the two-sample acceleration constant given by

equation 15.36 of Efron and Tibshirani (1993). (Estimates of F and the associated BC$_a$

confidence intervals are reported in Table 1 for each genotype and condition.) Consistent with

our other analyses, we find little evidence that wMau significantly increases fecundity. However,

our data do not have much statistical power to detect values of F on the order of 1.05, which may

suffice to produce $F(1 - \mu) > 1$ and deterministic spread of wMau from low frequencies. We

present our power calculations in Figure 1B of the Supplementary Information.

Table 1. Estimates of the relative fitness parameter F indicate that wMau fecundity effects are likely to be minimal.

wMau variant/age class	F	95% BC$_a$ interval
$R31$	0.988	(0.862, 1.137)
$R41$	1.107	(0.995, 1.265)
$R31^{R41}$	1.012	(0.911, 1.122)
$R41^{R31}$	0.992	(0.884, 1.143)
$R31$(across age)	0.833	(0.515, 1.323)

Finally, we assessed the fidelity of wMau maternal transmission under standard laboratory

conditions. We excluded sublines that produced fewer than 8 F1 offspring. In all cases, $R31$ ($N =$

17) sublines produced offspring that were all infected, indicating perfect maternal transmission.

In contrast, one $R41$ subline produced one uninfected individual out of a total of 18 F1 offspring

produced; all other $R41$ sublines meeting our criteria ($N = 15$) produced only infected F1

offspring, resulting in nearly perfect maternal transmission across all $R41$ sublines ($\mu = 0.0039$).

Mathematical analyses of *Wolbachia* frequencies and mtDNA polymorphism

If *Wolbachia* do not cause CI (or any other reproductive manipulation), their dynamics can be approximated by a discrete-generation haploid selection model. Following Hoffmann and Turelli (1997), we assume that the relative fecundity of *Wolbachia*-infected females is F, but a fraction μ of their ova do not carry *Wolbachia*. Given our ignorance of the nature of *Wolbachia*'s favorable effects, the F represents an approximation for all fitness benefits. If $F(1 - \mu) > 1$, the equilibrium *Wolbachia* frequency among adults is

$$\hat{p} = 1 - \frac{\mu F}{F - 1}. \tag{1}$$

Imperfect maternal transmisson has been documented for field-collected *D. simulans* infected with *w*Ri (Hoffmann and Turelli 1988; Turelli and Hoffmann 1995; Carrington et al. 2011), *D. melanogaster* infected with *w*Mel (Hoffmann et al. 1998) and *D. suzukii* infected with *w*Suz (Hamm et al. 2014). The estimates range from about 0.01 to 0.1. Given that we have documented imperfect maternal transmission of *w*Mau in the laboratory, we expect (more) imperfect transmission in nature (Turelli and Hoffmann 1995; Carrington et al. 2011). In order for the equilibrium *Wolbachia* frequency to be below 0.5, approximation (1) requires that the relative fecundity of infected females satisfies

$$F < \frac{1}{1 - 2\mu}. \tag{2}$$

Thus, even for μ as large a 0.15, which greatly exceeds our laboratory estimates for *w*Mau and essentially all estimates of maternal transmission failure from nature, *Wolbachia* can increase fitness by at most 43% and produce an equilibrium frequency below 0.5 (Supplemental Figure 1A). Conversely, (1) implies that a doubling of relative fecundity by *Wolbachia* would produce an equilibrium frequency $1 - 2\mu$. If $\mu \leq 0.25$, consistent with all available data, the predicted equilibrium implied by a *Wolbachia*-induced fitness doubling significantly exceeds the observed frequency of *w*Mau. Hence, a four-fold fecundity effect, as described by Fast et al. (2011), is inconsistent with the frequency of *w*Mau in natural populations of *D. mauritiana*. Field estimates of μ for *D. mauritiana* will provide better theory-based bounds on *w*Mau fitness effects that

would be consistent with wMau tending to increase when rare on Mauritius, i.e., conditions for

$F(1 - \mu) > 1$.

Our theoretical analysis, addressing the plausibility of a four-fold fitness increase caused by

wMau, assumes that the observed frequency of wMau approximates selection-transmission

equilibrium, as described by (1). With only two frequency estimates (one from a heterogeneous

collection of laboratory stocks), we do not know that the current low frequency is temporarlly

stable. Also, we do not know that the mutations we detect in $cinA$-$cinB^{wNo}$ homologs are

responsible for the lack of wMau CI. One alternative is that $D.\ mauritiana$ has evolved to

suppress CI (for host suppression of male killing, see Hornet et al. 2006 and Vanthournout and

Hendrickx 2016). Host suppression of CI is expected (Turelli 1994), and it may explain the low

CI caused by wMel in $D.\ melanogaster$ (Hoffmann and Turelli 1997). However, the fact that

wMau does not produce CI in $D.\ simulans$, a host that allows wMel and other strains to induce

strong CI even though little CI is produced in their native hosts, argues against host suppression

as the explanation for the lack of CI caused by wMau in $D.\ mauritiana$. Nevertheless, the loss of

CI from wMau may be quite recent; and wMau may be on its way to elimination in $D.$

$mauritiana$. If so, our equilibrium analysis is irrelevant – but this gradual-loss scenario is equally

inconsistent with the four-fold fecundity effect proposed by Fast et al. (2011).

As noted by Turelli et al. (1992), if $Wolbachia$ is introduced into a population along with a

diagnostic mtDNA haplotype that has no effect on fitness, imperfect $Wolbachia$ maternal

transmission implies that all infected and uninfected individuals will eventually carry the

$Wolbachia$-associated mtDNA, because all will have had $Wolbachia$-infected maternal ancestors.

We conjectured that a stable mtDNA polymorphism might be maintained if $Wolbachia$-

associated mtDNA introduced by introgression is deleterious in its new nuclear background. We

refute our conjecture in Appendix 1. We show that the condition for $Wolbachia$ to increase when

rare, $F(1 - \mu) > 1$, ensures that the native mtDNA will be completely displaced by the

$Wolbachia$-associated mtDNA, even if it lowers host fitness once separated from $Wolbachia$.

How fast is the mtDNA turnover, among $Wolbachia$-uninfected individuals, as a new

$Wolbachia$ invades? This is easiest to analyze when the mtDNA introduced with $Wolbachia$ has

no effect on fitness, so that the relative fitness of $Wolbachia$-infected versus uninfected

individuals is F, irrespective of the mtDNA haplotype of the uninfected individuals. As shown in

Appendix 1, the frequency of the ancestral mtDNA haplotype among uninfected individuals,

denoted r_t, declines as

$\qquad r_{t+1} = r_t/[F(1 - \mu)].$ (3)

Assuming $r_0 = 1$, recursion (3) implies that even if $F(1 - \mu)$ is only 1.01, the frequency of the

ancestral mtDNA haplotype should fall below 10^{-4} after 1000 generations. A much more rapid

mtDNA turnover was seen as the CI-causing wRi swept northward through California

populations of *D. simulans* (Turelli et al. 1992; Turelli and Hoffmann 1995). Thus, it is

theoretically unexpected, under this simple model, that mtDNA haplotype *ma*II, which seems to

be ancestral in *D. mauritiana* (Rousset and Solignac 1995; Ballard 2000a), persists among

Wolbachia-uninfected *D. mauritiana*, given that all sampled *Wolbachia*-infected individuals

carry *ma*I. However, spatial variation in fitnesses is one possible explanation for this

polymorphism (Gliddon and Strobeck 1975), which has persisted since at least 1985.

DISCUSSION

wMau is sister to wNo and diverged from group-A *Wolbachia* less than 100 mya

Our phylogenetic analyses place wMau sister to wNo, in agreement with past analyses using

fewer data (James and Ballard 2000; Zabalou et al. 2008; Toomey et al. 2013). The relationships

we infer agree with those from recently published phylograms (Gerth and Bleidorn 2016;

Lindsey et al. 2018) (Figure 1A).

Depending on the prior used for substitution-rate variation, we estimate that wMau and other

group-B *Wolbachia* diverged from group-A strains about 6–46 mya. This is roughly consistent

with a prior estimate using only *ftsZ* (58–67 mya, Werren et al. 1995), but is inconsistent with a

recent estimate using 179,763 bases across 252 loci (76–460 mya, Gerth and Bleidorn 2016).

There are several reasons why we question the Gerth and Bleidorn (2016) calibration. First,

Gerth and Bleidorn (2016)'s chronogram placed wNo sister to all other group-B *Wolbachia*, in

disagreement with their own phylogram (Gerth and Bleidorn 2016, Figure 3). In contrast, our

phylogram and that of Lindsey et al. (2018) support wAlbB splitting from all other strains at this

node. Second, the Gerth and Bleidorn (2016) calibration estimated the split between wRi that

infects *D. simulans* and wSuz that infects *D. suzukii* at 900,000 years. This estimate is more than

an order of magnitude higher than ours (16,214 years) and nearly two orders of magnitude higher

than the 11,000 year estimate of Turelli et al. (2018) who found 0.014% third position

divergence between wRi and wSuz (i.e., 0.007% along each branch) over 506,307 bases.

Raychoudhury et al. (2009) and Richardson et al. (2012) both estimated a rate of about 7×10^{-9}

substitutions/3$^{\mathrm{rd}}$ position site/year between *Wolbachia* in *Nasonia* wasps and within wMel,

respectively. An estimate of 900,000 years requires a rate about 100 times slower, 7.8×10^{-11}

substitutions/3$^{\mathrm{rd}}$ position site/year, which seems implausible. Finally, using data kindly provided

by Michael Gerth, additional analyses indicate that the third-position rates required for the

Wolbachia divergence times estimated by Gerth and Bleidorn (2016) between *Nomada flava* and

N. leucophthalma (1.72×10^{-10}), *N. flava* and *N. panzeri* (3.78×10^{-10}) (their calibration point),

and *N. flava* and *N. ferruginata* (4.14×10^{-10}) are each more than 10 times slower than those

estimated by Raychoudhury et al. (2009) and Richardson et al. (2012), which seems unlikely.

Our analyses suggest that the A-B group split occurred less than 100 mya.

The lack of CI is consistent with intermediate wMau infection frequencies

Across 671 genes (682,494 bases), the wMau genomes were identical and differed from wNo by

only 0.068%. Across the coding regions we analyzed, we found few SNVs and no CNVs among

wMau variants. Our analyses did identify four large deletions shared by all wMau genomes,

relative to wNo. Despite the close relationship between wMau and wNo, wNo causes CI while

wMau does not (Giordano et al. 1995; Merçot et al. 1995; Rousset and Solignac 1995, our data).

We searched for all pairs of loci known to cause CI and found only homologs to the *cinA-*

cinB$^{w\mathrm{No}}$ pair in wMau genomes. All wMau variants share a one-base-pair deletion in the wMau

region homologous to *cinB*$^{w\mathrm{No}}$. This mutation introduces a frameshift and more than ten stop

codons. Future functional work will help determine if disruption of this predicted-toxin locus

underlies the lack of CI in wMau. Regardless, the lack of CI is consistent with the prediction that

selection within host lineages does not directly act on the intensity of CI (Prout 1994; Turelli

1994). We predict that analysis of additional non-CI-causing strains will reveal additional

examples of genomic remnants of CI loci. Among non-CI *Wolbachia*, the relative frequency of

those with non-functional CI loci, versus no CI loci, is unknown.

Irrespective of whether CI was lost or never gained, non-CI *Wolbachia* have lower expected

equilibrium infection frequencies than do CI-causing variants (Kriesner et al. 2016). The wMau

infection frequency of approximately 0.34 on Mauritius (Giordano et al. 1995; our data) is

consistent with this prediction. Additional sampling of Mauritius, preferably over decades, will

determine whether intermediate *w*Mau frequencies are temporally stable. Such temporal stability

depends greatly on values of *F* and *μ* through time suggesting additional field-based estimates of

these parameters will be useful.

*w*Mau co-occurs with essentially the same mitochondrial haplotype as *w*Ma that infects *D.*

simulans on Madagascar and elsewhere in Africa and the South Pacific (Rousset and Solignac

1995; Merçot and Poinsot 1998; Ballard 2000a; James and Ballard 2000; James et al. 2002;

Ballard 2004), suggesting that *w*Mau and *w*Ma may be the same strain infecting different host

species following introgressive *Wolbachia* transfer (see below). *w*Mau and *w*Ma phenotypes are

also more similar to one another than to *w*No, with only certain crosses between *w*Ma-infected

D. simulans males and uninfected *D. simulans* females inducing CI (James and Ballard 2000).

Polymorphism in the strength of CI induced by *w*Ma could result from host modification of

Wolbachia-induced CI (Reynolds and Hoffmann 2002; Cooper et al. 2017), or from *Wolbachia*

titer variation that influences the strength of CI and/or the strength of CI rescue by infected

females. Alternatively, the single-base-pair deletion in the $cinB^{wNo}$ homolog or other mutations

that influence CI strength, could be polymorphic in *w*Ma. *w*Ma infection frequencies in *D.*

simulans are intermediate on Madagascar (infection frequency = 0.25, binomial confidence

intervals: 0.14, 0.40; James and Ballard 2000), consistent with no CI, suggesting replication of

rarely observed *w*Ma CI is needed. Including *D. simulans* from the island of Réunion in this

infection-frequency further supports the conjecture that *w*Ma causes little or no CI (infection

frequency = 0.31, binomial confidence intervals: 0.20, 0.45; James and Ballard 2000).

Unfortunately, no Madagascar *D. simulans* stocks available at the National *Drosophila* Species

Stock Center were *w*Ma infected, precluding detailed analysis of this strain.

Our genomic data indicate that *w*Mau may maintain an ability to rescue CI, as the $cinA^{wNo}$

homolog is intact in *w*Mau genomes with only one nonsynonymous substitution relative to

$cinA^{wNo}$; *cidA* in *w*Mel was recently shown to underlie transgenic-CI rescue (Shropshire et al.

2018). *w*Ma seems to sometimes rescue CI, but conflicting patterns have been found, and

additional experiments are needed to resolve this (Rousset and Solignac 1995; Bourtzis et al.

1998; Merçot and Poinsot 1998; James and Ballard 2000; Merçot and Poinsot 2003; Zabalou et

al. 2008). Future work that tests for CI rescue by *w*Mau and *w*Ma-infected females crossed with

males infected with *w*No or other CI-causing strains, combined with genomic analysis of CI loci

in *w*Ma, will be useful.

_w_Mau does not influence _D. mauritiana_ fecundity

While selection does not directly act on the level of CI (Prout 1994; Turelli 1994; Haygood and

Turelli 2009), it does act to increase the product *Wolbachia*-infected host fitness and the

efficiency of maternal transmission (Turelli 1994). Understanding the *Wolbachia* effects that

lead to spread from low frequencies and the persistence of non-CI causing *Wolbachia* at

intermediate frequencies is crucial to explaining *Wolbachia* prevalence among insects and other

arthropods. The four-fold fecundity effect of *w*Mau reported by Fast et al. (2011) in *D.*

mauritiana is inconsistent with our experiments and with the intermediate infection frequencies

observed in nature. We find no *w*Mau effects on host fecundity, regardless of host background,

with our estimates of *F* having BCa intervals that include 1. Small increases in *F* could allow the

deterministic spread of *w*Mau from low frequencies, although detecting very small increases in *F*

is difficult (Supplemental Figure 1B). Our results are consistent with an earlier analysis that

assessed egg lay of a single genotype and found no effect of *w*Mau on host fecundity (Giordano

et al. 1995). When combined with the low observed infection frequencies, our fecundity data are

also consistent with our mathematical analyses indicating that *Wolbachia* can increase host

fitness by at most about 50% for reasonable estimates of μ. Because fecundity is one of many

fitness components, analysis of other candidate phenotypes for aiding the spread of low-

frequency *Wolbachia* is needed.

Introgressive _Wolbachia_ transfer likely predominates in the _D. simulans_ clade

Hybridization and introgression in the *D. simulans* clade may have led to introgressive transfer

of *Wolbachia* among host species (Rousset and Solignac 1995). This has been observed in other

Drosophila (Turelli et al. 2018; Cooper et al. 2019) and *Nasonia* wasps (Raychoudhury et al.

2009). The number of *Wolbachia* strains in the *D. simulans* clade, and the diversity of

mitochondria they co-occur with, is complex. Figure 5A shows host relationships and Figure 5B

shows mitochondrial relationships, with co-occurring *Wolbachia* variants in parentheses. While

D. mauritiana is singly infected by *w*Mau, *D. simulans* is infected by several strains, including

CI-causing *w*Ha and *w*No that often co-occur as double infections within individuals (O'Neill

and Karr 1990; Merçot et al. 1995; Rousset and Solignac 1995). *w*Ha and *w*No are similar to

*w*Sh and *w*Sn, respectively, that infect *D. sechellia* (Giordano et al. 1995; Rousset and Solignac

1995). *w*Ha and *w*Sh also occur as single infections in *D. simulans* and in *D. sechellia*,

respectively (Rousset and Solignac 1995). In contrast, *w*No almost always co-occurs with *w*Ha in

doubly infected *D. simulans* individuals (James et al. 2002), and *w*Sn seems to occur only with

*w*Sh (Rousset and Solignac 1995). *D. simulans* has three distinct mitochondrial haplotypes (*si*I,

*si*II, *si*III) associated with *w*Au/*w*Ri (*si*II), *w*Ha/*w*No (*si*I), and *w*Ma (*si*III). The *si*I haplotype is

closely related to the *se* haplotype found with *w*Sh and *w*Sn in *D. sechellia* (Ballard 2000b). *w*Ma

co-occurs with the *si*III haplotype, which differs over its 13 protein-coding genes by only a

single-base pair from the *ma*I mitochondrial haplotype carried by *w*Mau-infected *D. mauritiana*.

A second haplotype (*ma*II) is carried by only uninfected *D. mauritiana* (Ballard 2000a; James

and Ballard 2000).

Figure 5. A) A nuclear relative chronogram. **B)** A mitochondrial relative chronogram with co-
occurring *Wolbachia* strains listed in parentheses. See the text for an interpretation of the results,
including the artifactual resolution of the phylogeny of the *D. simulans* clade.

The lack of whole *w*Ma genome data precludes us from confidently resolving the mode of

*w*Mau acquisition in *D. mauritiana*. However, mitochondrial relationships support the proposal

of Ballard (2000b) that *D. mauritiana* acquired *w*Mau and the *ma*I mitochondrial haplotype via

introgression from *w*Ma-infected *D. simulans* carrying *si*III. *D. mauritiana* mitochondria are

paraphyletic relative to *D. sechellia* and *D. simulans* mitochondria (Solignac and Monnerot

1986; Satta and Takahata 1990; Ballard 2000a, 2000b), with *ma*I sister to *si*III and *ma*II outgroup

to all other *D. simulans*-clade haplotypes (see Figure 5). Of the nine genomes we assessed, all

but one (uninfected-*R44*) carry the *ma*I haplotype, and genotypes carrying *ma*I are both *w*Mau-

infected ($N = 5$) and uninfected ($N = 3$). While *w*Ma-infected *D. simulans* carry *si*III, *w*No-

infected *D. simulans* carry *si*I. We estimate that *w*Mau and *w*No diverged about 55,000 years

ago, with only 0.068% sequence divergence over 682,494 bp. Nevertheless, it seems implausible

that *w*No (versus *w*Ma) was transferred directly to *D. mauritiana* as this requires horizontal or

paternal transmission of *w*No into a *D. mauritiana* background already carrying the *ma*I

mitochondrial haplotype. Although our nuclear result suggests a confident phylogenetic

resolution of the *D. simulans* clade (Figure 5A), this is an artifact of the bifurcation structure

imposed by the phylogenetic analysis. Population genetic analyses show a complex history of

introgression and probable shared ancestral polymorphisms (Kliman et al. 2000) among these

three species. Consistent with this, of the 20 nuclear loci we examined, 6 (*aconitase, aldolase,*

bicoid, ebony, enolase, ninaE) supported *D. mauritiana* as the outgroup within the *D. simulans*

clade, 7 (*glyp, pepck, pgm, pic, ptc, transaldolase, wingless*) supported *D. sechellia* as the

outgroup, and 7 (*esc, g6pdh, glys, pgi, tpi, white, yellow*) supported *D. simulans*. With successive

invasions of the islands and purely allopatric speciation, we expect the outgroup to be the island

endemic that diverged first. Figure 5B indicates that the *ma*II haplotype diverged from the other

mtDNA haplotypes roughly when the clade diverged, with the other haplotypes subject to a

complex history of introgression and *Wolbachia*-associated sweeps, as described by Ballard

(2000b).

Ballard (2000b) estimated that *si*III-*ma*I diverged about 4,500 years ago, which presumably

approximates the date of the acquisition of *w*Mau (and *si*III, which became *ma*I) by *D.*

mauritiana. This is surely many thousands of generations given previous estimates that consider

the temperature dependence of *Drosophila* development (Cooper et al. 2014; Cooper et al. 2018).

As shown by our mathematical analyses (Eq. 3), the apparent persistence of the *ma*II mtDNA

among *Wolbachia*-uninfected *D. mauritiana*—without its occurrence among infected

individuals—is unexpected. More extensive sampling of natural *D. mauritiana* populations is

needed to see if this unexpected pattern persists. The persistence of this haplotype is inconsistent

with simple null models, possibly indicating interesting fitness effects.

While paternal transmission has been observed in *D. simulans* (Hoffmann and Turelli 1988;

Turelli and Hoffmann 1995), it seems to be very rare (Richardson et al. 2012; Turelli et al.

2018). *w*No almost always occurs in *D. simulans* individuals also infected with *w*Ha,

complicating this scenario further. It is possible that horizontal or paternal transmission of *w*Ma

or *w*No between *D. simulans* backgrounds carrying different mitochondrial haplotypes underlies

the similarities of these strains within *D. simulans*, despite their co-occurrence with distinct

mitochondria. Given the diversity of *Wolbachia* that infect *D. simulans*-clade hosts, and known

patterns of hybridization and introgression among hosts (Garrigan et al. 2012; Brand et al. 2013;

Garrigan et al. 2014; Matute and Ayroles 2014; Schrider et al. 2018), determining relationships

among these *Wolbachia* and how *D. mauritiana* acquired *w*Mau will require detailed

phylogenomic analysis of nuclear, mitochondrial, and *Wolbachia* genomes in the *D. simulans*

clade.

AUTHOR CONTRIBUTIONS

MM performed the molecular and phenotypic work, participated in the design of the study, and

contributed to the writing; WC performed the phylogenetic and genomic analyses and

contributed to the writing; SR contributed to the molecular and phenotypic analyses and to the

writing; JB performed the library preparation and contributed to the writing; MT contributed to

the analyses, data interpretation, and writing; BSC designed and coordinated the study,

contributed to the analyses and data interpretation, and drafted the manuscript. All authors gave

final approval for publication.

ACKNOWLEDGMENTS

We thank Margarita Womack for sampling the *D. mauritiana* used in this study and Daniel

Matute for sharing them. We thank Michael Gerth for sharing *Nomada* genomic data. Isaac

Humble, Maria Kirby, and Tim Wheeler assisted with data collection. Michael Gerth, Michael

Hague, and Amelia Lindsey provided comments that improved earlier drafts of this manuscript.

Computational resources were provided by the University of Montana Genomics Core. Research

reported in this publication was supported by the National Institute Of General Medical Sciences

of the National Institutes of Health (NIH) under Award Number R35GM124701 to B.S.C. The

content is solely the responsibility of the authors and does not necessarily represent the official

views of the NIH. The authors declare no conflicts of interest.

LITERATURE CITED

Baldo, L., N. A. Ayoub, C. Y. Hayashi, J. A. Russell, J. K. Stahlhut, and J. H. Werren. 2008
   Insight into the routes of *Wolbachia* invasion: high levels of horizontal transfer in the
   spider genus *Agelenopsis* revealed by *Wolbachia* strain and mitochondrial DNA
   diversity. Molecular Ecology 17**:**557–569.

Ballard, J. W. O. 2000a. When one is not enough: introgression of mitochondrial DNA in
   Drosophila. Molecular Biology and Evolution 17**:**1126–1130.

Ballard, J. W. 2000b. Comparative genomics of mitochondrial DNA in *Drosophila simulans*.
   Journal of Molecular Evolution 51:64–75.

Ballard, J. W. O. 2004. Sequential evolution of a symbioint inferred from the host: *Wolbachia*
   and *Drosophila simulans*. Molecular Biology and Evolution 21:428–442.

Ballard, J. W., and R. G. Melvin. 2007. Tetracycline treatment influences mitochondrial
   metabolism and mtDNA density two generations after treatment in *Drosophila*. Insect
   Molecular Biology 16**:**799–802.

Barton, N. H., and M. Turelli. 2011. Spatial waves of advance with bistable dynamics:
   cytoplasmic and genetic analogues of allee effects. American Naturalist 178**:**E48–E75.

Beckmann, J. F., and A. M. Fallon. 2013. Detection of the *Wolbachia* protein WPIP0282 in
   mosquito spermathecae: implications for cytoplasmic incompatibility. Insect
   Biochemistry and Molecular Biology 43**:**867–878.

Beckmann, J. F., J. A. Ronau and M. Hochstrasser. 2017. A *Wolbachia* deubiquitylating enzyme
   induces cytoplasmic incompatibility. Nature Microbiology 2**:**17007.

Beckmann, J. F., M. Bonneau, H. Chen, M. Hochstrasser, D. Poinsot, H. Merçot, M. Weill, M.
   Sicard, and S. Charlat. 2019. The toxin-antidote model of cytoplasmic incompatibility:
   genetics and evolutionary implications. Trends in Genetics in press.

Boeva, V., T. Popova, K. Bleakley, P. Chiche, J. Cappo, G. Schleiermacher, I. Janoueix-Lerosey,
   O. Delattre, E. Barillot. 2012. Control-FREEC: a tool for assessing copy number and
   allelic content using next-generation sequencing data. Bioinformatics 28**:**423–425.

Bourtzis, K., S. L. Dobson, H. R. Braig and S. L. O'Neill. 1998. Rescuing *Wolbachia* have been
   overlooked. Nature 391**:**852–853.

Brand, C. L., S. B. Kingan, L. Wu and D. Garrigan. 2013. A Selective Sweep across Species
   Boundaries in *Drosophila*. Molecular Biology and Evolution 30**:**2177–2186.

Brownlie, J. C., B. N. Cass, M. Riegler, J. J. Witsenburg, I. Iturbe-Ormaetxe, E. A. McGraw, S.
   L. O'Neill. 2009. Evidence for Metabolic Provisioning by a Common Invertebrate
   Endosymbiont, *Wolbachia pipientis*, during Periods of Nutritional Stress. PLoS
   Pathogens 5**:**e1000368.

Carrington, L. B., J. R. Lipkowitz, A. A. Hoffmann, and M. Turelli. 2011. A re-examination of
   Wolbachia-induced cytoplasmic incompatibility in California *Drosophila simulans*. PLoS
   ONE 6:e22565

Caspari, E., and G. S. Watson. 1959. On the evolutionary importance of cytoplasmic sterility in
   mosquitoes. Evolution 13:568–570.

Cattel, J., K. Nikolouli, T. Andrieux, J. Martinez, F. Jiggins, S. Charlat, F. Vavre, D. Lejon, P.
   Gibert, and L. Mouton. 2018. Back and forth *Wolbachia* transfers reveal efficient strains
   to control spotted wing *Drosophila* populations. Journal of Applied Ecology 55:2408-
   2418.

Conner, W. R., M. L. Blaxter, G. Anfora, L. Ometto, O. Rota-Stabelli, and M. Turelli. 2017.
Genome comparisons indicate recent transfer of *w*Ri-like *Wolbachia* between sister
species *Drosophila suzukii* and *D. subpulchrella*. Ecology and Evolution 7:9391–9404.
Cooper, B. S., L. A. Hammad, and K. L. Montooth. 2014. Thermal adaptation of cellular
membranes in natural populations of *Drosophila melanogaster*. Functional Ecology.
28:886-894.
Cooper, B. S., P. S. Ginsberg, M. Turelli and D. R. Matute. 2017. *Wolbachia* in the *Drosophila*
yakuba complex: pervasive frequency variation and weak cytoplasmic incompatibility,
but no apparent effect on reproductive isolation. Genetics 205**:**333–351.
Cooper, B. S., A. Sedghifar, W. T. Nash, A. A. Comeault, D. R. Matute. 2018. A maladaptive
combination of traits contributes to the maintenance of a *Drosophila* hybrid zone. Current
Biology. 28:2940-2947.
Cooper, B. S., D. Vanderpool, W. R. Conner, D. R. Matute, and M. Turelli. 2019. Introgressive
and horizontal acquisition of *Wolbachia* by *Drosophila yakuba*-clade hosts and horizontal
transfer of incompatibility loci between distantly related *Wolbachia*. bioRxiv.
doi.org/10.1101/551036.
David, J., S. McEvey, M. Solignac and L. Tsacas. 1989. *Drosophila* communities on Mauritius
and the ecological niche of *D. mauritiana* (Diptera, Drosophilidae). Revue De Zoologie
Africaine – Journal of African Zoology 103:107–116.
Dean, M. D., and J. W. Ballard. 2004. Linking phylogenetics with population genetics to
reconstruct the geographic origin of a species. Molecular Phylogenetics and Evolution
32**:**998–1009.
Efron, B., and R. Tibshirani. 1993. *An Introduction to the Bootstrap*. Chapman & Hall, New
York, NY.
Ellegaard, K. M., L. Klasson, K. Näslund, K. Bourtzis and S. G. E. Andersson. 2013.
Comparative Genomics of *Wolbachia* and the Bacterial Species Concept. PLoS Genetics
9**:**e1003381.
Fast, E. M., M. E. Toomey, K. Panaram, D. Desjardins, E. D. Kolaczyk, H. M. Frydman. 2011.
Wolbachia enhance *Drosophila* stem cell proliferation and target the germline stem cell
niche. Science 334**:**990–992.
Fisher, R. A. 1937. The wave of advance of advantageous genes. Annals of Eugenics 7**:**355–369.
Garrigan, D., S. B. Kingan, A. J. Geneva, P. Andolfatto, A. G. Clark, K. R. Thornton, D. C.
Presgraves. 2012. Genome sequencing reveals complex speciation in the *Drosophila*
simulans clade. Genome Research 22**:**1499–1511.
Garrigan, D., S. B. Kingan, A. J. Geneva, J. P. Vedanayagam and D. C. Presgraves. 2014.
Genome diversity and divergence in *Drosophila mauritiana*: multiple signatures of faster
X evolution. Genome Biology and Evolution 6**:**2444–2458.
Gerth, M., and C. Bleidorn. 2016. Comparative genomics provides a timeframe for *Wolbachia*
evolution and exposes a recent biotin synthesis operon transfer. Nature Microbiology
2**:**16241.
Giordano, R., S. L. O'Neill and H. M. Robertson. 1995. *Wolbachia* infections and the expression
of cytoplasmic incompatibility in *Drosophila sechellia* and *D. mauritiana*. Genetics
140**:**1307–1317.
Gliddon, C. and C. Strobeck. 1975. Necessary and sufficient conditions for multiple-niche
polymorphism in haploids. American Naturalist 109:233–235.

Gloor, G. B., C. R. Preston, D. M. Johnsonschlitz, N. A. Nassif, R. W. Phillis, W. K. Benz, H.
848   M. Robertson, W. R. Engels. 1993. Type-1 repressors of P-element mobility. Genetics
849   135**:**81–95.

850   Gupta, V., R. B. Vasanthakrishnan, J. Siva-Jothy, K. M. Monteith, S. P. Brown, and P. F. Vale.
851   2017. The route of infection determines *Wolbachia* antibacterial protection in
852   *Drosophila*. Proceedings of the Royal Society B: Biological Sciences 284:20170809.

853   Hamm, C. A., D. J. Begun, A. Vo, C. C. R. Smith, P. Saelao, A. O. Shaver, J. Jaenike, M.
854   Turelli. 2014. *Wolbachia* do not live by reproductive manipulation alone: infection
855   polymorphism in *Drosophila suzukii* and *D. subpulchrella*. Molecular Ecology 23:4871–
856   4885.

857   Haygood, R., and M. Turelli. 2009. Evolution of incompatibility-inducing microbes in
858   subdivided host populations. Evolution 63:432–447

859   Hedges, L. M., J. C. Brownlie, S. L. O'Neill and K. N. Johnson. 2008. *Wolbachia* and virus
860   protection in insects. Science 322:702.

861   Hilgenboecker, K., P. Hammerstein, P. Schlattmann, A. Telschow and J. H. Werren. 2008. How
862   many species are infected with *Wolbachia*? A statistical analysis of current data. FEMS
863   Microbiology Letters 281**:**215–220.

864   Hoffmann, A. A., M. Turelli, and G. M. Simmons. 1986. Unidirectional incompatibility between
865   populations of *Drosophila simulans*. Evolution 40:692-701.

866   Hoffmann, A. A., and M. Turelli. 1988. Unidirectional incompatibility in *Drosophila simulans*:
867   Inheritance, geographic variation and fitness effects. Genetics 119:435–444.

868   Hoffmann, A. A., and M. Turelli. 1997. Cytoplasmic incompatibility in insects. In *Influential*
869   *passengers: inherited microorganisms and arthropod reproduction*, edited by S. L.
870   O'Neill, J. H. Werren, and A. A. Hoffmann. Oxford University Press, pp. 42–80.

871   Hoffmann, A. A., D. Clancy and J. Duncan. 1996. Naturally-occurring *Wolbachia* infection in
872   *Drosophila simulans* that does not cause cytoplasmic incompatibility. Heredity 76**:**1–8.

873   Hoffmann, A. A., M. Hercus and H. Dagher. 1998. Population dynamics of the *Wolbachia*
874   infection causing cytoplasmic incompatibility in *Drosophila melanogaster*. Genetics
875   148:221–231.

876   Hoffmann, A. A., M. Turelli, and L. G. Harshman. 1990. Factors affecting the distribution of
877   cytoplasmic incompatibility in *Drosophila simulans*. Genetics 126**:**933–948.

878   Hoffmann, A. A., B. L. Montgomery, J. Popovici, I. Iturbe-Ormaetxe, P. H. Johnson, F. Muzzi,
879   M. Greenfield, M. Durkan, Y. S. Leong, Y. Dong, H. Cook, J. Axford, A. G. Callahan, N.
880   Kenny, C. Omodei, E. A. McGraw, P. A. Ryan, S. A. Ritchie, M. Turelli, and S. L.
881   O'Neill. 2011. Successful establishment of *Wolbachia* in *Aedes* populations to suppress
882   dengue transmission. Nature 476:454–457.

883   Höhna, S., M. J. Landis, T. A. Heath, B. Boussau, N. Lartillot, B. R. Moore, J. P. Huelsenbeck,
884   F. Ronquist. 2016. RevBayes: Bayesian Phylogenetic Inference Using Graphical Models
885   and an Interactive Model-Specification Language. Systematic Biology. 65:726–736.

886   Hornett, E. A., S. Charlat, A. M. R. Duplouy, N. Davies, G. K. Roderick, N. Wedell, and G. D.
887   D. Hurst. 2006. Evolution of male-killer suppression in a natural population. PLoS
888   Biology 4:e283.

889   Huigens, M. E., R. P. de Almeida, P. A. Boons, R. F. Luck and R. Stouthamer. 2004. Natural
890   interspecific and intraspecific horizontal transfer of parthenogenesis-inducing *Wolbachia*
891   in *Trichogramma* wasps. Proceedings of the Royal Society B: Biological Sciences
892   271**:**509–515.

Hurst, G. D. D. and F. M. Jiggins. 2000. Male-killing bacteria in insects: mechanisms, incidence,
and implications. Emerging Infectious Diseases 6:329-336.
Jackman, S. D., B. P. Vandervalk, H. Mohamadi, J. Chu, S. Yeo, S. A. Hammond, G. Jahesh, H.
Khan, L. Coombe, R. L. Warren, I. Birol. 2017. ABySS 2.0: resource-efficient assembly
of large genomes using a Bloom filter. Genome Research 27**:**768–777.
James, A. C., and J. W. O. Ballard. 2000. Expression of cytoplasmic incompatibility in
Drosophila simulans and its impact on infection frequencies and distribution of
Wolbachia pipientis. Evolution 54**:**1661–1672.
James, A. C., M. D. Dean, M. E. McMahon and J. W. Ballard. 2002. Dynamics of double and
single *Wolbachia* infections in *Drosophila simulans* from New Caledonia. Heredity
88**:**182–189.
Jeyaprakash, A., and M. A. Hoy. 2000. Long PCR improves *Wolbachia* DNA amplification: wsp
sequences found in 76% of sixty-three arthropod species. Insect Molecular Biology
9**:**393–405.
Joshi, N. A., and J. N. Fass. 2011. Sickle: A sliding-window, adaptive, quality-based trimming
tool for FastQ files (Version 1.33). Available at https://github.com/najoshi/sickle.
Katoh, K., and D. M. Standley. 2013. MAFFT Multiple Sequence Alignment Software Version
7: Improvements in Performance and Usability. Molecular Biology and Evolution
30**:**772–780.
Kent, B. N., L. Salichos, J. G. Gibbons, A. Rokas, I. L. Newton, M. E. Clark, S. R. Bordenstein.
2011. Complete bacteriophage transfer in a bacterial endosymbiont (*Wolbachia*)
determined by targeted genome capture. Genome Biology and Evolution 3**:**209–218.
Klasson, L., T. Walker, M. Sebaihia, M. J. Sanders, M. A. Quail, A. Lord, S. Sanders, J. Earl, S.
L. O'Neill, N. Thomson, S. P. Sinkins, J. Parkhill. 2008. Genome evolution of *Wolbachia*
strain *w*Pip from the *Culex pipiens* group. Molecular Biology and Evolution 25**:**1877–
1887.
Klasson, L., J. Westberg, P. Sapountzis, K. Näslund, Y. Lutnaes, A. C. Darby, Z. Veneti, L.
Chen, H. R. Braig, and R. Garrett. 2009. The mosaic genome structure of the *Wolbachia*
*w*Ri strain infecting *Drosophila simulans*. Proceedings of the National Academy of
Sciences 106**:**5725–5730.
Kliman, R. M., P. Andolfatto, J. A. Coyne, F. Depaulis, M. Kreitman, A. J. Berry, J. McCarter, J.
Wakeley, and J. Hey. 2000. The population genetics of the origin and divergence of the
Drosophila simulans species complex. Genetics 156:1913–1931.
Kriesner, P., and A. A. Hoffmann. 2018. Rapid spread of a *Wolbachia* infection that does not
affect host reproduction in *Drosophila simulans* cage populations. Evolution 72:1475–
1487.
Kriesner, P., W. R. Conner, A. R. Weeks, M. Turelli, and A. A. Hoffmann. 2016. Persistence of
a *Wolbachia* infection frequency cline in *Drosophila melanogaster* and the possible role
of reproductive dormancy. Evolution 70:979–997.
Kriesner, P., A. A. Hoffmann, S. F., Lee, M. Turelli, and A. R. Weeks. 2013. Rapid sequential
spread of two *Wolbachia* variants in *Drosophila simulans*. PLoS Pathogens 9:e1003607.
Lachaise, D., J. R. David, F. Lemeunier, L. Tsacas and M. Ashburner. 1986. The reproductive
relationships of *Drosophila sechellia* with *D. mauritiana*, *D. simulans*, and *D.*
melanogaster from the Afrotropical region. Evolution 40:262–271.
Laven, H. 1951. Crossing Experiments with *Culex* Strains. Evolution 5:370-375.

Legrand, D., T. Chenel, C. Campagne, D. Lachaise and M. L. Cariou. 2011. Inter-island
divergence within *Drosophila mauritiana*, a species of the *D. simulans* complex: Past
history and/or speciation in progress? Molecular Ecology 20**:**2787–2804.
LePage, D. P., J. A. Metcalf, S. R. Bordenstein, J. On, J. I. Perlmutter, J. D. Shropshire, E. M.
Layton, L. J. Funkhouser-Jones, J. F. Beckmann, S. R. Bordenstein. 2017. Prophage WO
genes recapitulate and enhance *Wolbachia*-induced cytoplasmic incompatibility. Nature
543**:**243–247.
Li, H., and R. Durbin. 2009. Fast and accurate short read alignment with Burrows-Wheeler
transform. Bioinformatics 25**:**1754–1760.
Lindsey, A. R. I., D. W. Rice, S. R. Bordenstein, A. W. Brooks, S. R. Bordenstein, A. W.
Brooks, S. R. Bordenstein, I. L. Newton. 2018. Evolutionary Genetics of Cytoplasmic
Incompatibility Genes *cifA* and *cifB* in Prophage WO of *Wolbachia*. Genome Biology
and Evolution 10**:**434–451.
Lindsey, A. R. I., J. H. Werren, S. Richards and R. Stouthamer. 2016. Comparative Genomics of
a Parthenogenesis-Inducing *Wolbachia* Symbiont. Genes|Genomes|Genetics. 6:2113–
2123.
Martinez, J., B. Longdon, S. Bauer, Y.-S. Chan, W. J. Miller, K. Bourtzia, L. Teixeira, F. M.
Jiggins. 2014. Symbionts commonly provide broad spectrum resistance to viruses in
insects: A comparative analysis of *Wolbachia* strains. PLoS Pathogens 10:e1004369.
Matute, D. R., and J. F. Ayroles. 2014. Hybridization occurs between *Drosophila simulans* and
D. sechellia in the Seychelles archipelago. Journal of Evolutionary Biology 27**:**1057–
1068.
Mavingui, P., C. Valiente Moro, V. Tran-Van, F. Wisniewski-Dye, V. Raquin, G. Minard, F. H.
Tran, D. Voronin, Z. Rouy, P. Bustos, L. Lozano, V. Barbe, V. Gonzalez. 2012. Whole-
genome sequence of *Wolbachia* strain *w*AlbB, an endosymbiont of tiger mosquito vector
Aedes albopictus. Journal of Bacteriology 194**:**1840.
McDermott, S. R., and R. M. Kliman. 2008. Estimation of isolation times of the island species in
the *Drosophila simulans* complex from multilocus DNA sequence data. PLoS One
3**:**e2442.
Merçot, H., and D. Poinsot. 1998. …and discovered on Mount Kilimanjaro. Nature 391**:**853.
Merçot, H., and D. Poinsot. 2003. *Wolbachia* transmission in a naturally bi-infected *Drosophila*
simulans strain from New-Caledonia. Entomologia Experimentalis et Applicata 86**:**97–
103.
Merçot, H., B. Llorente, M. Jacques, A. Atlan and C. Montchamp-Moreau. 1995. Variability
within the Seychelles cytoplasmic incompatibility system in *Drosophila simulans*.
Genetics 141**:**1015–1023.
Nunes, M. D., P. O. Wengel, M. Kreissl and C. Schlotterer. 2010. Multiple hybridization events
between *Drosophila simulans* and *Drosophila mauritiana* are supported by mtDNA
introgression. Molecular Ecology 19**:**4695–4707.
Ochman, H., and A. C. Wilson. 1987. Evolution in bacteria: evidence for a universal substitution
rate in cellular organisms. Journal of Molecular Evolution 26:74–86.
O'Neill, S. L., R. Giordano, A. M. Colbert, T. L. Karr and H. M. Robertson. 1992. 16S rRNA
phylogenetic analysis of the bacterial endosymbionts associated with cytoplasmic
incompatibility in insects. Proceedings of the National Academy of Sciences 89**:**2699–
2702.

O'Neill, S. L., and T. L. Karr. 1990. Bidirectional incompatibility between conspecific
984   populations of *Drosophila simulans*. Nature 348**:**178–180.
985   Pinto, S. B., K. Stainton, S. Harris, Z. Kambris, E. R. Sutton, M. B. Bonsall, J. Parkhill, S. P.
986   Sinkins. 2013. Transcriptional regulation of *Culex pipiens* mosquitoes by *Wolbachia*
987   influences cytoplasmic incompatibility. PLoS Pathogens 9:e1003647.
988   Poinsot, D., K. Bourtzis, G, Markakis, C. Savakis, and H. Merçot. 1998. Wolbachia transfer from
989   *Drosophila melanogaster* into *D. simulans*: host effect and cytoplasmic incompatibility
990   relationships. Genetics 150:227–237.
991   Prout, T. 1994. Some evolutionary possibilities for a microbe that causes incompatibility in its
992   host. Evolution 48**:**909–911.
993   Raychoudhury, R., L. Baldo, D. C. S. G. Oliveira, and J. H. Werren. 2009. Modes of acquisition
994   of *Wolbachia*: horizontal transfer, hybrid introgression, and codivergence in the *Nasonia*
995   species complex. Evolution 63:165–183.
996   Reynolds, K. T., and A. A. Hoffmann. 2002. Male age, host effects and the weak expression or
997   non-expression of cytoplasmic incompatibility in *Drosophila* strains infected by
998   maternally transmitted *Wolbachia*. Genetics Research 80**:**79–87.
Richardson, M. F., L. A. Weinert, J. J. Welch, R. S. Linheiro, M. M. Magwire, F. M. Jiggins, C.
M. Bergman. 2012. Population genomics of the *Wolbachia* endosymbiont in *Drosophila*
melanogaster. PLoS Genetics 8**:**e1003129.
Ritchie S. A. 2018. *Wolbachia* and the near cessation of dengue outbreaks in Northern Australia
despite continued dengue importations via travellers. Journal of Travel
Medicine. 25:tay084.)
Rousset, F., and M. Solignac. 1995. Evolution of single and double *Wolbachia* symbioses during
speciation in the *Drosophila simulans* complex. Proceedings of the National Academy of
Sciences 92**:**6389–6393.
Satta, Y. and N. Takahata. 1990. Evolution of *Drosophila* mitochondrial DNA and the history of
the *melanogaster* subgroup. Proceedings of the National Academy of Sciences of the
United States of America 87:9558–9562.
Schmidt, T. L., N. H. Barton, G. Rašić, A. P. Turley, B. L. Montgomery, I. Iturbe-Ormaetxe, P.
E. Cook, P. A. Ryan, S. A. Ritchie, A. A. Hoffmann, S. L. O'Neill, and M. Turelli. 2017.
Local introduction and heterogeneous spatial spread of dengue-suppressing *Wolbachia*
through an urban population of *Aedes aegypti*. PLoS Biology 15:e2001894.
Schrider, D. R., J. Ayroles, D. R. Matute and A. D. Kern. 2018. Supervised machine learning
reveals introgressed loci in the genomes of *Drosophila simulans* and *D. sechellia*. PLoS
Genetics 14**:**e1007341.
Schuler, H., K. Köppler, S. Daxböck-Horvath, B. Rasool, S. Krumböck, D. Schwarz, T. S.
Hoffmeister, B. C. Schlick-Steiner, F. M. Steiner, A. Telschow, C. Stauffler, W.
Arthofer, M. Riegler. 2016. The hitchhiker's guide to Europe: the infection dynamics of
an ongoing *Wolbachia* invasion and mitochondrial selective sweep in *Rhagoletis cerasi*.
Molecular Ecology 25**:**1595–1609.
Seemann, T. 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics 30**:**2068-2069.
Shi, M., V. L. White, T. Schlub, J.-S. Eden, A. A. Hoffmann, E. C. Holmes. 2018. No detectable
effect of *Wolbachia w*Mel on the prevalence and abundance of the RNA virome of
Drosophila melanogaster. Proceedings of the Royal Society B: Biological Sciences
285:20181165.

Shropshire, J. D., J. On, E. M. Layton, H. Zhou and S. R. Bordenstein. 2018. One prophage WO gene rescues cytoplasmic incompatibility in *Drosophila melanogaster*. Proceedings of the National Academy of Sciences. 115:4987–4991.

Simao, F. A., R. M. Waterhouse, P. Ioannidis, E. V. Kriventseva and E. M. Zdobnov. 2015. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. Bioinformatics 31**:**3210–3212.

Siozios, S., A. Cestaro, R. Kaur, I. Pertot, O. Rota-Stabelli, and G. Anfora. 2013. Draft genome sequence of the *Wolbachia* endosymbiont of *Drosophila suzukii*. Genome Announcements. 1, e00032-13.

Solignac, M. and M. Monnerot. 1986. Race formation, speciation, and introgression within *Drosophila simulans*, *D. mauritiana*, and *D. schellia* inferred from mitochondrial DNA analysis. Evolution 40:531–539.

Sutton, E. R., S. R. Harris, J. Parkhill and S. P. Sinkins. 2014. Comparative genome analysis of *Wolbachia* strain *w*Au. BMC Genomics 15**:**928.

Taylor, M. J., D. Voronin, K. L. Johnston and L. Ford. 2013. *Wolbachia* filarial interactions. Cell Microbiology 15**:** 520–526.

Team, R. C. 2015. R: A language and environment for statistical computing, *R Foundation for Statistical Computing*, Vienna, Austria.

Teixeira, L., A. Ferreira and M. Ashburner. 2008. The bacterial symbiont *Wolbachia* induces resistance to RNA viral infections in *Drosophila melanogaster*. PLoS Biology 6**:**e1000002.

Toomey, M. E., K. Panaram, E. M. Fast, C. Beatty and H. M. Frydman. 2013. Evolutionarily conserved *Wolbachia*-encoded factors control pattern of stem-cell niche tropism in *Drosophila* ovaries and favor infection. Proceedings of the National Academy of Sciences 110**:**10788–10793.

Turelli, M. 1994. Evolution of incompatibility-inducing microbes and their hosts. Evolution 48:1500–1513.

Turelli, M. and N. H. Barton. 2017. Deploying dengue-suppressing *Wolbachia*: robust models predict slow but effective spatial spread in *Aedes aegypti*. Theoretical Population Biology 115:45–60.

Turelli, M. and A. A. Hoffmann. 1995. Cytoplasmic incompatibility in *Drosophila simulans*: dynamics and parameter estimates from natural populations. Genetics 140:1319–1338.

Turelli, M., A. A. Hoffmann, and S. W. McKechnie. 1992. Dynamics of cytoplasmic incompatibility and mtDNA variation in natural *Drosophila simulans* populations. Genetics 132:713–723.

Turelli, M., B. S. Cooper, K. M. Richardson, P. S. Ginsberg, B. Peckenpaugh, C. X. Antelope, K. J. Kim, M. R. May, A. Abrieux, D. A. Wilson, M. J. Bronski, B. R. Moore, J. J. Gao, M. B. Eisen, J. C. Chiu, W. R. Conner, A. A. Hoffmann. 2018. Rapid Global Spread of *w*Ri-like *Wolbachia* across Multiple *Drosophila*. Current Biology 28**:**963–971.

Vanthournout, B. and F. Hendrickx. 2016. Hidden suppression of sex ratio distortion suggests Red queen dynamics between *Wolbachia* and its dwarf spider host. Journal of Evolutionaly Biology 29:1488-1494.

Webster, C. L., F. M. Waldron, S. Robertson, D. Crowson, G. Ferrari, J. F. Quintana, J.-M. Brouqui, E. H. Bayne, B. Longdon, A. H. Buck, B. P. Lazzaro, J. Akorli, P. R. Haddrill, and D. J. Obbard. 2015. The Discovery, Distribution, and Evolution of Viruses Associated with *Drosophila melanogaster*. PLoS Biology 13:e1002210.

Weeks, A. R., M. Turelli, W. R. Harcombe, K. T. Reynolds, and A. A. Hoffmann. 2007. From
parasite to mutualist: rapid evolution of *Wolbachia* in natural populations of *Drosophila*.
PLoS Biology 5**:**e114.
Weinert, L. A., E. V. Araujo-Jnr, M. Z. Ahmed and J. J. Welch. 2015. The incidence of bacterial
endosymbionts in terrestrial arthropods. Proceedings of the Royal Society of London B:
Biological Sciences 282**:**20150249.
Werren, J. H., and D. M. Windsor. 2000. *Wolbachia* infection frequencies in insects: evidence of
a global equilibrium? Proceedings of the Royal Society B: Biological Sciences
267**:**1277–1285.
Werren, J. H., W. Zhang and L. R. Guo. 1995. Evolution and phylogeny of *Wolbachia*:
reproductive parasites of arthropods. Proceedings of the Royal Society B-Biological
Sciences 261**:**55–63.
Wu, M., L. V. Sun, J. Vamathevan, M. Riegler, R. Deboy, J. C. Brownlie, E. A. McGraw, W.
Martin, C. Esser, N. Ahmadinejad. 2004. Phylogenomics of the reproductive parasite
*Wolbachia pipientis w*Mel: a streamlined genome overrun by mobile genetic elements.
PLoS Biology 2**:**e69.
Yen, J. H. and A. R. Barr. 1971. New hypothesis of the cause of cytoplasmic incompatibility in
Culex pipiens L. Nature 232:657-658.
Zabalou, S., A. Apostolaki, S. Pattas, Z. Veneti, C. Paraskevopoulos, I. Livadaras, G. Markakis,
T. Brissac, H. Merçot, K. Bourtzis. 2008. Multiple rescue factors within a *Wolbachia*
strain. Genetics 178**:**2145–2160.
Zug, R., and P. Hammerstein. 2012. Still a host of hosts for *Wolbachia*: Analysis of recent data
suggests that 40% of terrestrial arthropod species are infected. PLoS ONE 7:e38544.

1097

1098

1099

1100

1101

1102

1103

1104

1105

1106

1107

1108

1109

1110

1111

Appendix 1. Mathematical analyses of mtDNA and *Wolbachia* dynamics

Our analysis follows the framework developed in Turelli et al. (1992), but is simplified by the

lack of CI. We suppose that introgression introduces a cytoplasm carrying *Wolbachia* and a

novel mtDNA haplotype, denoted B. Before *Wolbachia* introduction, we assume the population

is monomorphic for mtDNA haplotype A. With imperfect maternal *Wolbachia* transmission,

uninfected individuals will be produced with mtDNA haplotype B. Without horizontal or

paternal transmission (which are very rare, Turelli et al. 2018), all *Wolbachia*-infected

individuals will carry mtDNA haplotype B. Once *Wolbachia* is introduced, uninfected

individuals can have mtDNA haplotype A or B. We assume that these three cytoplasmic types

("cytotypes") differ only in fecundity, and denote their respective fecundities F_I, F_A and F_B.

Denote the frequencies of the three cytotypes among adults in generation t by $p_{I,t}$, $p_{A,t}$ and $p_{B,t}$,

with $p_{I,t} + p_{A,t} + p_{B,t} = 1$. Without CI, the frequency dynamics are

1124

1125 $$p_{I,t+1} = \frac{p_{I,t}F_I(1-\mu)}{\overline{F}}, \ p_{A,t+1} = \frac{p_{A,t}F_A}{\overline{F}}, \ \text{and} \ p_{B,t+1} = \frac{p_{B,t}F_B + p_{I,t}\mu F_I}{\overline{F}}, \text{with} \qquad (A1)$$

1126

1127 $$\overline{F} = F_I p_{I,t} + F_A p_{A,t} + F_B p_{B,t}. \qquad (A2)$$

1128

1129 If the uninfected population is initially monomorphic for mtDNA haplotype A, the *Wolbachia*

1130 infection frequency will increase when rare if and only if

1131

1132 $$F_I(1-\mu) > F_A. \qquad (A3)$$

1133

Turelli et al. (1992) showed that if a CI-causing *Wolbachia* is introduced with a cytoplasm

that contains a novel mtDNA haplotype B, which has no effect on fitness, *Wolbachia*-uninfected

individuals will eventually all carry haplotype B. This follows because eventually all uninfected

individuals have *Wolbachia*-infected maternal ancestors. This remains true for non-CI-causing

Wolbachia that satisfy (A3). However, we conjectured that if the introduced B mtDNA is

deleterious in the new host nuclear background, i.e., $F_A > F_B$, a stable polymorphism might be

maintained for the alternative mtDNA haplotypes. The motivation was that imperfect maternal

transmission seemed analogous to migration introducing a deleterious allele into an "island" of

uninfected individuals. To refute this conjecture, consider the equilibria of (A1) with

1143

1144 $$F_I > F_A \geq F_B. \tag{A4}$$

1145

1146 If all three cytotypes are to be stably maintained, we expect each to increase in frequency when

1147 rare. In particular, we expect the fitness-enhancing mtDNA haplotype A to increase when the

1148 population contains only infected individuals and uninfected individuals carrying the deleterious

1149 *Wolbachia*-associated mtDNA haplotype B. From (A1), $p_{A,t}$ increases when rare if and only if

1150

1151 $$F_A > \overline{F} = F_I p_{I,t} + F_B (1 - p_{I,t}). \tag{A5}$$

1152

1153 In the absence of haplotype A, we expect p_I to be at equilibrium between selection and imperfect

1154 maternal transmission, i.e.,

1155

1156 $$p_I = 1 - \frac{\mu F}{F - 1}, \tag{A6}$$

1157

1158 with $F = F_I/F_B$ (Hoffmann and Turelli 1997). Substituting (A6) into (A5) and simplifying, the

1159 condition for $p_{A,t}$ to increase when rare is

1160

1161 $$F_A(F_I - F_B) > F_I(1 - \mu)(F_I - F_B). \tag{A7}$$

1162

By assumption (A4), $F_I > F_B$; hence (A7) contradicts condition (A3), required for initial

Wolbachia invasion. Thus, simple selection on *Wolbachia*-uninfected mtDNA haplotypes cannot

stably maintain an mtDNA polymorphism. The "ancestral" mtDNA haplotype A is expected to

be replaced by the less-fit *Wolbachia*-associated haplotype B.

To understand the time scale over which this replacement occurs, let r_t denote the frequency

of haplotype A among *Wolbachia*-uninfected individuals, i.e., $r_t = p_{A,t}/(p_{A,t} + p_{B,t})$. From (A1),

1169

1170 $$r_{t+1} = \frac{r_t F_A}{r_t F_A + (1 - r_t)F_B + \mu F_I[p_{I,t}/(1 - p_{I,t})]} . \tag{A8}$$

1171

1172   If we assume that the mtDNA haplotypes do not affect fitness, i.e., $F_A = F_B$, and that the

1173   *Wolbachia* infection frequency has reached the equilibrium described by (A6), (A8) reduces to

1174

1175 $r_{t+1} = r_t/[F(1 - \mu)],$ (A9)

1176

1177   with $F = F_I/F_B$.

1178

1179

1180

1181

1182

1183

1184

1185

1186

1187

1188

1189

1190

1191

1192

1193

1194

1195

1196

1197

1198

1199

1200

1201

1202

 SUPPLEMENTAL INFORMATION

 Supplemental Tables

Supplemental Table 1. *w*Mau assembly statistics.

Genotype	Scaffold count	N50 (bp)	Longest scaffold (bp)	Total length (bp)
R9	36	60,027	169,305	1,266,004
R29	36	61,106	169,295	1,277,467
R31	39	63,676	169,381	1,272,847
R41	42	61,106	170,537	1,303,156
R60	38	63,156	221,751	1,282,564

Supplemental Table 2. Near-universal, single-copy proteobacteria genes (out of 221) found using BUSCO v. 3.0.0.

Genome	Complete	Duplicated	Fragment	Absent
*w*Ri	179	1	2	39
*w*Mel	179	1	2	39
*w*Au	180	1	2	38
*w*Ha	178	1	3	39
*w*No	180	1	4	36
R9	180	1	4	36
R29	180	1	4	36
R31	180	1	4	36
R41	180	1	4	36
R60	180	1	4	36

Supplemental Table 3. Copy number variants in *w*Mau relative to sister *w*No. Genomic positions are based on the *w*No reference. There were no CNVs among *w*Mau variants.

Start	End	Change	Wilcoxon Rank Sum Test	Kolmogorov-Smirnov Test
115,000	142,000	1 → 0	< 0.0001	< 0.0001
590,000	593,000	1 → 0	0.0027	0.0050
982,000	991,000	1 → 0	< 0.0001	< 0.0001
1,195,000	1,199,000	1 → 0	0.0005	0.0007

1211
1212
1213

Supplemental Table 4: Genes present in regions deleted in *w*Mau relative to *w*No. Genes predicted to be pseudogenized in *w*No are shaded grey.

Accession number	Name
Deletion 1 (115,000-142,000):	
wNO_RS00550	Hypothetical protein
wNO_RS06015	Ankyrin repeat domain protein
wNO_RS00560	Pseudo IS256 family transposase, frameshifted
wNO_RS00565	Recombinase family protein
wNO_RS00570	DUF2924 domain-containing protein
wNO_RS00575	Ankyrin repeat domain-containing protein
wNO_RS00580	Ankyrin repeat domain-containing protein
wNO_RS00585	Phage tail protein
wNO_RS00590	Baseplate assembly protein GpJ
wNO_RS00595	Pseudo baseplate assembly protein W, frameshifted
wNO_RS00600	Hypothetical protein
wNO_RS00605	Phage baseplate assembly protein V
wNO_RS00610	Hypothetical protein
wNO_RS00615	Putative minor tail protein Z
wNO_RS00620	Hypothetical protein
wNO_RS00625	Minor capsid protein E
wNO_RS00630	Head decoration protein
wNO_RS00635	S49 family peptidase

wNO_RS00640	Pseudo phage portal protein, frameshifted
wNO_RS00645	Phage head stabilizing protein GpW
wNO_RS00650	Phage terminase large subunit family protein
wNO_RS00655	Ankyrin repeat domain-containing protein
wNO_RS00660	Hypothetical protein
wNO_RS00665	Hypothetical protein
wNO_RS00670	Sigma-70 family RNA polymerase sigma factor
wNO_RS00675	ATP-binding protein
wNO_RS00680	IS110 family transposase
Deletion 2 (590,000-593,000):	
wNO_RS02645	XRE family transcriptional regulator
wNO_RS02650	Hypothetical protein
wNO_RS02655	Hypothetical protein
wNO_RS02660	XRE family transcriptional regulator
Deletion 3 (982,000-991,000):	
wNO_RS04690	Group II intron reverse transcriptase/maturase
wNO_RS06355	Hypothetical protein
wNO_RS04695	Pseudo hypothetical protein, partial
wNO_RS04700	Pseudo cell filamentation protein Fic, partial
wNO_RS04705	Hypothetical protein
wNO_RS04710	DNA methylase
wNO_RS04715	Hypothetical protein
wNO_RS04720	Ankyrin repeat domain-containing protein
wNO_RS04725	Phage terminase large subunit family protein
wNO_RS04730	Phage head stabilizing protein GpW
wNO_RS04735	DUF1016 domain-containing protein
Deletion 4 (1,195,000-1,199,000):	
wNO_RS06095	Ankyrin repeat domain protein
wNO_RS05590	Rpn family recombination-promoting nuclease/putative transposase

1214
1215

Supplemental Table 5. *w*Mau does not significantly affect *D. mauritiana* fecundity in comparisons of paired infected (I) and uninfected (U) strains sharing host nuclear backgrounds. *N* is the number of females that produced the means and SDs. *P* values are for two-tailed Wilcoxon tests. Cytoplasm sources are denoted with superscripts for introgressed strains.

Strain	Mean eggs laid/5 days	SD	N	P value
R31I	210.97	55.22	30	0.901
R31U	213.50	58.43	28	
R41I	265.43	31.47	30	0.355
R41U	239.69	73.43	29	
R31^{R41}I	257.83	58.15	30	0.762
R31^{R41}U	254.67	48.72	30	
R41^{R31}I	248.77	45.54	30	0.433
R41^{R31}U	250.67	76.35	30	

1216

1217

1218

1219 **Supplemental Figures**

1220
1221 **Supplemental Figure 1.** A) Our proposed upper bound, $1/(1 - 2\mu)$, on plausible relative
1222 fitness, *F*, as a function of μ ranging from perfect transmission ($\mu = 0$) to a level of
1223 imperfect transmission ($\mu = 0.2$) that has not been observed in nature. B) Our powers to
1224 detect values of *F* with $\alpha = 0.05$.

1225