Prenatal exposure to particulate matter (PM$_{2.5}$) from biomass fuel and low birth weight in a Sri Lankan birth cohort

Meghan Tipre, BDS, MSPH, DrPH[1]
Rajitha Wickremesinghe, MD, PhD[2]
Sumal Nandasena, MD[3]
Anuradhini Kasturiratne, MD[2]
Rodney Larson, PhD[4]
Sreelatha Meleth, PhD[5]
Udaya Wimalasiri BS[2]
Claudiu Lungu, PhD[1]
Tamika Smith, PhD[1]
Nalini Sathiakumar, MD, DrPH[1]

[1]University of Alabama at Birmingham, United States (US)
[2]University of Kelaniya, Faculty of Medicine, Sri Lanka
[3]National Institutes of Health Sciences, Sri Lanka
[4]Larson International. Inc
[5]Research Triangle Institute, Atlanta, US

Corresponding author:

Meghan Tipre, BDS, MSPH, DrPH
Research Scientist I
Ryals Building, 430
School of Public Health
University of Alabama at Birmingham
Birmingham Alabama, 35294
Email: meghan24@uab.edu

ABSTRACT

Background.

Prenatal exposure to particulate matter of size 2.5 µm ($PM_{2.5}$) emissions from biomass fuel may be associated with low birth weight (LBW) (<2.5 kg). We present results examining the association between $PM_{2.5}$ and LBW from a birth cohort of 545 pregnant women followed from first trimester until delivery, in Sri Lanka.

Methods.

Exposure to household air pollution (HAP) from biomass smoke was assessed using a detailed questionnaire; two-hour measurements of kitchen $PM_{2.5}$ were collected in a subset of households (n, 303, 56%). Data from questionnaire and measured $PM_{2.5}$ were used to estimate two-hour kitchen $PM_{2.5}$ concentrations in unmeasured households. Fetal growth including weight and gestational age were assessed at birth. Linear and logistic regression were used to evaluate the association between HAP and birth weight.

Results.

About 78% of the households used wood (n, 423). In linear regression models, we found an inverse association between a 10-unit increase in $PM_{2.5}$ and birth weight (β±SE, -0.03±0.02; p, 0.02) adjusted for covariates. Similarly, households primarily using wood (>50%) were negatively associated with birth weight as compared to LPG users (β, -0.13±0.06; p, 0.03). In logistic regression models, a 10-unit increase in $PM_{2.5}$ was associated with increased odds for LBW (OR, 1.26; 95%CI, 1.02-1.55; p, 0.04). The odds for LBW were highest among >50% wood users (OR, 2.82; 95%CI, 1.18-6.73; p, 0.01), compared to those using >50% LPG with wood (OR, 1.44; 95%CI, 0.57-3.63) and 100% LPG users (referent). The association between HAP exposure and birth weight/LBW were consistent among term-births (n,486).

Discussion.

The finding of a significant association between prenatal $PM_{2.5}$ exposure and LBW in a low-middle income country (LMIC) setting where competing risk factors are minimal fills a gap in the body of evidence linking HAP to LBW and underscore need to prevent and reduce of HAP exposure in LMICs.

Key words

Household air pollution, Low birth weight, Sri Lanka, Birth cohort

INTRODUCTION AND AIMS

Household air pollution (HAP) from combustion of solid fuels such as wood, dung and coal, used for cooking and heating, is ranked 10^{th} as a contributor to the global burden of disease (1,2) and the third leading risk factor for global mortality and disability-adjusted life years (DALYs)(3). Smoke emissions from solid fuel combustion in inefficient cookstoves and inadequate ventilation release relatively high concentrations of $PM_{2.5}$, carbon monoxide (CO) and several other organic and inorganic compounds (4). Long-term exposure to fine particulate matter ($PM_{2.5}$) of size 2.5 μm and smaller contributes to 4.2 million deaths and a loss of 103.1 million DALYs; majority of this burden is borne by low middle income countries (LMICs) (5)(6). More than three billion people accounting for 43% of the world's population use solid fuels (coal and biomass) for cooking and heating; about 80% of these people live in LMICs (7,8). In most LMICs, women of child-bearing age typically spend several hours a day in cooking chores. Consequently, they are exposed to high levels of HAP for long periods and are at high risk for adverse health effects from HAP. In Sri Lanka, a LMIC in South Asia with a population of 20 million, about 74% of the population uses some form of solid fuel (unprocessed wood) for cooking.(8,9) Recent reports from the Sri Lankan National Demographic Surveys indicate that the prevalence of solid fuel use remains mostly unchanged with only slight decrease from 78% to 74% in the last 15 years. (10–12)

Studies on exposure to $PM_{2.5}$ from either ambient or indoor sources have found positive associations with adverse pregnancy outcomes including stillbirths, low birth weight (LBW), small for gestational age (SGA), preterm births and birth defects (13–15). Results from a meta-analysis of ambient air pollution and birth weight found that birth weight in grams (g) was negatively associated with 10 μg/m^3 increase in ambient PM_{10} and $PM_{2.5}$ exposure during the

prenatal period, adjusted for maternal smoking (16). Studies that have evaluated the role of HAP from solid fuel combustion and LBW indicate a stronger association, than ambient air exposure studies (17–20). A meta-analysis found that exposure to HAP from solid fuels based mostly on self-reports, resulted in an 86.4 g reduction in birth weight, and a 35% and 29% increased risk of LBW and stillbirth, respectively (21). Several of the pollutants found in ambient and indoor air share similar toxic properties as those found in environmental tobacco smoke (ETS) (22). Evidence from ETS and animal studies indicate that these air pollutants are absorbed by maternal blood, and then may cross the placental barrier and cause adverse effects on the fetus (23,24).

Previous studies of HAP and adverse birth outcomes including birth weight and LBW have several limitations. Most studies are either cross-sectional or case-control. Many of these studies are also subject to exposure misclassification. The labor-intensive and high costs associated with conducting air quality monitoring in all households, limit several HAP studies to qualitative exposure assessment resulting in exposure misclassification, especially in households with stacking of fuel (25). Stacking of fuel is where the household supplements the main fuel with secondary fuel for cooking certain dishes such as rice or beans or for special occasion such as festivals or holidays.

Most studies to date have been conducted in LMICs including India, Guatemala, Pakistan, and Ghana, countries where other competing risk factors for adverse birth outcomes such as malnutrition, poverty, infections and poor access to health care, are highly prevalent (21,26). Further, in-home deliveries and limited access to prenatal care in rural regions may often result in gaps in information related to fetal growth and pregnancy-related complications making it difficult to control for residual confounding factors to estimate the true effect of HAP on fetal

health. Only a few were birth cohort or experimental in design but except for a study conducted by Balakrishna et. al., (2018), most had smaller sample size (27) .

To address some of these challenges, we designed a prospective birth-cohort study in Sri Lanka. The goal of the study was to evaluate the association between prenatal exposure to HAP from biomass smoke emissions and adverse birth outcomes. Our choice of the study setting is unique and offered several advantages. Sri Lanka has a literacy rate of 92% and maternal-child health indicators comparable to most high-income countries (28); thus, confounding/competing risk factors are minimized. Next, Sri Lanka's primary care infrastructure includes routine follow-up of almost all (>95%) pregnant women and their children until five years of age facilitating assembling a cohort of women early in their pregnancy, following them through childbirth, and to subsequently follow their children until five years of age.

MATERIALS AND METHODS

Study setting

Sri Lanka is divided into approximately 334 Medical Officers of Health (MoH) units areas under the management of a Ministry of Health; each MoH serves a population of 60,000 to 100,000 (29–31). The MoH area is further divided into several Public Health Inspector (PHI) areas and, within each PHI areas are Public Health Midwife (PHM) areas, the smallest administrative unit of the government health system in Sri Lanka. Typically, a PHI area serves a population of 10,000-15,000 which is subdivided into 3-4 PHM areas. The PHM is the female grass roots level family health worker primarily responsible for maternal-child health services and serving a population of 3000-5000 people (31,32).

The PHM registers newly pregnant women, typically before 10 weeks of gestation. Thereafter, women are followed at the field clinic and in their homes every four weeks until the

28th week of gestation, every two weeks from the 28th to the 36th week of gestation and weekly thereafter until delivery. The PHM is also responsible for mothers' immunizations and health education. Almost 99% of births in Sri Lanka occur in hospitals. After delivery, the PHM visits the mother at home (a minimum of three visits within the first 10 days) to follow-up and identify early post-partum complications of the mother, if any, and newborn problems including feeding complications.

Study area and recruitment

The study was conducted in the Ragama, Medical Officer of Health (MOH) which comprises of 16 PHM areas. Two research assistants (RAs) worked with the PHMs to identify eligible study participants. A total of 721 pregnant women ($n = 721$) were screened between July 2011 and June 2012 from the Ragama MOH area. Eligibility criteria included pregnant women in their first trimester; between 18 and 40 years of age; who planned to continue prenatal care and deliver at the Colombo North Teaching Hospital, Ragama; and permanently resided in the Ragama MoH area. Written informed consent in Sinhala was obtained from all women and the study procedures were explained prior to enrollment. Of the 721, 545 women were followed until delivery. Loss of participants were due to miscarriages (n, 60); twin births (n, 5); stillbirth (n, 9); delivered in different hospitals (n,13); lost to contact due to changed residence (n, 15) or refused to participate (n, 74).

Data collection

Women were interviewed at enrollment (mean, 11±2.5 weeks of gestation) to collect baseline information on demographic factors, residential history, medical history, maternal and paternal occupational history, history of active and passive tobacco smoking; and household characteristics. A section of the questionnaire collected detailed exposure information on factors

related to HAP including fuel use, kitchen characteristics and cooking practices. Information was also collected on other sources of indoor and outdoor pollution contributing to HAP. Maternal pregnancy history and infant anthropometric measurement at birth were obtained from hospital records. All interviews were conducted by trained bilingual study interviewers.

Prenatal exposure assessment

Qualitative information on exposure to HAP was obtained from the baseline questionnaire administered to all study participants. The baseline questionnaire elicited detailed information on cooking practices such as types of fuels used for cooking including primary and secondary cooking fuels; percent use of each fuel during cooking; daily average hours spent cooking; type of stove; presence of a chimney and its condition; ventilation in the kitchen during cooking including number and size of open doors and windows; and the type of kitchen, its dimensions and relative position with other rooms. Information was also collected on other sources of HAP such as tobacco smoking habits of household members, burning of incense and mosquito coils, and proximity of sources of ambient air pollution such as vehicular traffic, industry or garbage disposal sites.

Quantitative exposure to HAP was ascertained from two-hour $PM_{2.5}$ measurements in the kitchen during the cooking period in a subset of households (n, 304 out of 545). Based on baseline data, households were categorized into 'high' and 'low' exposure categories if the women used biomass fuel or clean fuel as primary cooking fuel, respectively. Using a stratified random sampling method, we selected 304 households (high exposure group, n, 158; low exposure group, n, 145) for conducting two-hour air quality measurements in the kitchen. At least one measurement per household was scheduled during the pregnancy period; in 106 households (36%), two sets of measurements were conducted in two of the three trimesters and

in 26 households, three measurements were conducted, once every trimester; all others had one measurement (n,197, 65%). All measurements were conducted during a two-hour cooking session while preparing a typical lunch. Using this data, $PM_{2.5}$ concentrations were estimated for unmeasured household (described below). $PM_{2.5}$ concentrations were measured using a TSI Dusttrak II Aerosol Monitor 8530. Inlets of monitors were fixed at 145 cm above the floor, 100 cm from the cook stove and at least 150 cm away from an open window or door.

Birth outcome assessment

All births occurred in the North Colombo Teaching Hospital. RAs collected data on birth outcomes from hospital records. Information abstracted included: birth weight, length and head circumference measured within 24 hours of birth, prenatal and delivery medical history. Gestational age was estimated based on the date of the last menstrual period and substantiated with ultrasound in the first trimester to classify infants as preterm vs. term births; pre-term births were defined as <37 weeks of gestation. LBW was defined as birth weight <2500 grams at birth. Term LBW were defined as full-term babies weighing <2500 grams at birth, as per standard definitions by the World Health Organization (33,34) .

Covariates and confounding variables

Data on covariates obtained from baseline questionnaire included: (i) demographic and SES factors such as maternal age, education, occupation, household income, type of housing, toilets, drinking water supply, and ownership of house; (ii) maternal lifestyle factors such as use of alcohol and tobacco; (iii) maternal exposure to toxic chemicals such as ETS, pesticides and lead; (iv) parity; (v) past and present obstetric history of pregnancy induced hypertension, gestational diabetes, other pregnancy-related complications, miscarriages and still births, iron deficiency, and calcium intake; (vi) and child's gender.

Ethical review

All study protocols were approved by the Ethics Review Committee of the Faculty of Medicine, University of Kelaniya and the Institutional Review Board of the University of Alabama at Birmingham (UAB).

Statistical analysis

    (i)   Exposure estimation.

<u>***Categorical HAP exposure.***</u> In the baseline questionnaire, information was collected on the primary fuel and supplementary fuels used in the households. A Likert scale was used to quantify the amount of fuels used typically during an average week. The scale ranged from 1 to 4, 1 being 0 to 25% of fuel use and 4 being 75% to 100% of fuel used. Most of the households (n, 423, 78%) in the study used a second fuel in varying quantities to supplement their primary fuel use. Combining the above information, we created two exposure variables. The first was a dichotomous variable classifying household into two categories: 'only LPG' and 'any wood use'. The second variable classified the households into three categories with gradients of wood use: (i) those using >50% to 100% of wood; (ii) those using >50-100% of LPG or kerosene and supplementing it with <50% wood; and (iii) those using 100% LPG. Since, only three households used 100% kerosene supplemented occasionally by LPG, these households were included in the third fuel category of 100% LPG.

<u>***Continuous PM$_{2.5}$.***</u> Next, using regression models, we examined the influence of several physical characteristics on concentration of PM$_{2.5}$ levels in the kitchen fuel types (>50% wood, >50% LPG + <50% wood, 100% LPG), kitchen type (indoor, temporary hut or outdoor); cookstoves (traditional cookstove (TCS), improved cookstove (ICS), kerosene or LPG stove); presence or absence of functional chimney, number of doors and windows open at the time of

cooking as a surrogate for kitchen ventilation, and average daily cooking time categorized into three groups (0 to 60 minutes; 60 to 90 minutes and >90 minutes).

Prediction model. Data from 275 out of 303 households with air quality measurements were used to develop the exposure model to estimate kitchen $PM_{2.5}$ airborne concentrations. Twenty-six samples were excluded as the comparison between fuel use and $PM_{2.5}$ values suggested potential misclassification of fuel use or were identified as possible outliers. For example, a household using >75% of LPG had measured value of $PM_{2.5}$ as high as $1000\mu g/m^3$ significantly greater than the concentration that would be expected. The final regression model was then used to quantify kitchen for $PM_{2.5}$ concentrations in kitchens without air quality measurements (n, 270; 50%); data on percentage of fuel use was missing for one subject and thus, had missing value for $PM_{2.5}$.

Data (n, 275) was partitioned randomly into training (n, 192) and test (n, 83) datasets using a 7:3 ratio respectively. The training dataset was used to identify factors significantly associated with log-transformed $PM_{2.5}$. Linear regression was used to model log-transformed $PM_{2.5}$ as a function of significant risk factors. In univariate statistics, geometric means for $PM_{2.5}$ were compared across physical variables. Factors significantly associated with log-transformed $PM_{2.5}$ at $\alpha < 0.05$ were included in a multivariable linear regression model to derive the final model. Adjusted R^2 was used to evaluate the model fit. Results identified three independent risk factors associated with log-transformed $PM_{2.5}$ including multilevel fuel categories, kitchen type, and functional chimney at $p < 0.05$ (Supplementary table 1-2). The model R^2 was 0.57 (supplementary table 2). The regression model equation is described below.

$$Ln(y) = \alpha + \beta_{fuel\_category} + \beta_{kitchen\,type} + \beta_{functional\,chimney}$$

where: α, intercept = 4.31; β_{wood} = 2.10, estimate for wood use >50%; $\beta_{LPG+Wood}$ = 0.42, estimate for LPG use 50% to 100% supplemented by wood; β_{LPG} = 0, estimate for 100% LPG; β_{K_ind} = -0.18, estimate for indoor kitchen; $\beta_{K\_temphut}$ = 1.17, estimate for temporary outdoor hut kitchen; $\beta_{K\_outdoor}$ = 0, estimate for outdoor open kitchen; and $\beta_{chimney}$ = 0.06, estimate for functional chimney.

Validation of the model. In the test dataset, the beta (β) coefficients from training model were used to predict $PM_{2.5}$ values. The difference between the observed and the predicted $PM_{2.5}$ values were calculated and residuals were plotted to evaluate the robustness of the predictive model (Supplementary table 2).

(ii)   HAP exposure and outcomes

Initial analyses compared the mean distribution of birth weight and prevalence of LBW with exposure variables and covariates using t-test, ANOVA, and the chi-square test (supplementary table 1). We restricted the latter comparisons and further analyses to those potential risk factors, to which at least 10 subjects were exposed.

HAP exposure was analyzed as continuous (log-transformed $PM_{2.5}$) and categorical variables (dichotomous and three-level fuel categories). We used linear regression to evaluate the association between HAP exposure and birth weight. Covariates significantly associated with exposure and/or birth weight in bivariate statistics at $p<0.05$ were included in the multivariable linear regression to compute adjusted beta-estimate and standard error, and adjusted means for birth weight.

Logistic regression was used to evaluate the association between HAP exposure and LBW. Covariates significantly associated with exposure and/or LBW in bivariate analysis at $p<0.05$ were included in the multivariable model. The odds ratio (OR) and 95% confidence intervals

(CIs) were computed for LBW in relation to HAP exposure, modeled as continuous or categorical variable, adjusting for other potential risk factors.

We further stratified the analysis by term and preterm births The association between HAP exposure and birth weight were further examined stratified by term and preterm births. All analyses were conducted using SAS version 9.4 (SAS Institute NC).

RESULTS

Of the total mothers (n, 545) followed-up from 1st trimester until delivery, 423 (78%) used wood in varying quantities, while 122 (22%) women used only LPG for cooking. The average age of mothers enrolled was 29 years (SD, 4.9; median, 29); all women were married. Significant differences were noted between wood users and LPG users for maternal education, spouse's education, main household water supply, ownership of the household, type of house (semi-permanent vs. permanent), income, kitchen type, functional chimney, and sources of HAP other than biomass fuel including ETS, burning of candles or incense (p<0.05). No remarkable differences were noted for mother's age at the time of enrollment, past and current obstetric history, parity or child's gender (Table 1).

Among households with air quality measurements, the geometric mean ± standard error for $PM_{2.5}$ among wood users (n, 157; 737± 80µg/m^3) was about four times higher than households using >50% of LPG with wood (n, 78; 193±29 µg/m^3) and almost 7 times higher than households using 100% LPG (n,68;100 ±10µg/m^3).

HAP exposure and birth weight

Of the total 545 newborns, 52% were boys. The mean birthweight of infants in the cohort was 2.95 (±0.45) kg. The proportion of LBW children in the cohort was 13% (n, 70 of 545) and the preterm births were 10% (n, 57). The total number of term LBW births were 46 (10%) out of

486 term births. About 65% of the mothers had vaginal delivery including forceps (n,6) and vacuum deliveries (n, 5).

Table 2 presents distribution of mean birth weight across all covariates. Birth weight was significantly associated with continuous $PM_{2.5}$, three-level fuel categories, type of house, pregnancy induced hypertension and preterm births at p <0.05. In multivariable linear regression models (Table 3), continuous log-transformed $PM_{2.5}$ was inversely associated with birth weight (β, -0.06; SE, 0.02; p, 0.03), adjusted for covariates. All multivariable models were adjusted for mother's education, father's education, income, permanent or semi-permanent house, ownership of the house, and pregnancy induced hypertension. In addition, models with categorical HAP exposure were also adjusted for type of kitchen and functional chimney.

Use of wood fuel as compared to LPG use was associated with lower birth weight but the association was non-significant (β,-0.07; SE, 0.05; p, 0.20). In model with three-level HAP exposure, with >50% wood use (β, -0.13; SE, 0.06; p, 0.03) was significant and inversely associated with birth weight as compared to 100% LPG use (as referent). No association was observed between >50% LPG with wood use and birth weight (β, 0.002; SE, 0.06; p, 0.53).

HAP exposure and LBW

The prevalence of LBW was significantly associated with continuous $PM_{2.5}$; a 10 unit (in $\mu g/m^3$) increase in $PM_{2.5}$ was associated with 1.26 times increased odds of LBW (95% CI, 1.01-1.55; p, 0.04). No association was observed between dichotomous fuel use variable and LBW (OR, 1.78; 95% CI, 0.78-4.09; p, 0.17). For three-level HAP exposure variable, the odds of LBW were significantly higher for >50% wood users (OR, 2.27; 95% CI, 0.94-5.12; p, 0.08) compared to 100% LPG users; no difference was noted between >50% LPG with wood users and 100% LPG users (Table 4).

Term births.

In stratified models by term birth, the relationship between exposure variables and birth weight and LBW were consistent as observed in the full models but effect estimates were lower. A moderately significant association was found between 10-unit increase in $PM_{2.5}$ and decrements in birth weight (β, -0.04; SE, 0.02; p, 0.03) and LBW (OR, 1.27; 95% CI, 0.98-1.65, p, 0.07). Similarly, >50% wood use was associated with decrease in birth weight (β, -0.12; SE, 0.06; p, 0.05) and prevalence of LBW (OR, 1.76; 95% CI, 0.61-5.03; p, 0.08), but the associations were not significant. We did not run models on preterm birth due to small number of observations (n, 55).

Discussion

In the present study, we investigated the relationship between prenatal exposure to HAP from biomass fuel use for cooking and birth weight, in a birth cohort study of 545 mother-child pairs in the Western Province of Sri Lanka. Our study modeled exposure to HAP as continuous and categorical variables. We found a significant association between two-hour kitchen concentrations of $PM_{2.5}$ and birth weight and prevalence of LBW, after adjusting for significant confounding factors. Similarly, households using most wood were associated with decrease in birth weight and increased prevalence of LBW as compared to households using clean fuels. The results are consistent with previous air pollution studies including ambient and HAP and LBW (15,16,35). In our study, a combination of qualitative and quantitative data from household with measured $PM_{2.5}$ were used to quantify kitchen concentrations of $PM_{2.5}$ in unmeasured households. To our knowledge, our study is the first to use both measured and estimated $PM_{2.5}$ measurements to evaluate the association between $PM_{2.5}$ and birth weight. The exposure

estimation method, effect estimates for birth weight and LBW in context of previous literature and the strengths and limitations of the study are discussed below.

HAP, birth weight, LBW and previous studies

The results of our birth-cohort evaluating the relationship between HAP and LBW were consistent with the largest birth-cohort study to date as well as several meta-analyses published in the last decade. Balakrishnan et al. (2017) reported a 4 grams (95% CI, 1.08-6.76) decrease in birthweight and 2% increase in prevalence of low birthweight (OR, 1.02; 95% CI, 1.01-1.04) with a 10-μg/m^3 increase in PM$_{2.5}$. Results from other meta-analyses also found increased prevalence of LBW between 10% to 45%, associated with HAP exposure (17,19,21,36–40). Similarly, a recent meta-analysis conducted by Sun et al. (2016), found significant association between 10μg/m^3 increment of ambient PM$_{2.5}$ and increased odds of LBW (OR,1.09; 95% CI, 1.03-1.15) (35). The effect estimates in our study are closer to the higher end of central estimates reported by individual studies and meta-analyses. In our study, the average two-hour kitchen PM$_{2.5}$ in kitchens using wood was almost 12 times higher than households using LPG (1649 vs. 143). Balakrishnan et al., (2018) reported 24-h average kitchen concentrations in households which were much lower than those reported in our study (wood use: 229±233 μg/m^3 vs. 1649±166 μg/m^3; LPG/electricity: 59±46 μg/m^3 vs. 143±12 μg/m^3), respectively. The higher concentration of kitchen PM$_{2.5}$ found in our study may account for the higher effect estimate reported in our study as compared to the results reported by Balakrishnan et.al. (2016). However, without additional data on 24-hour average kitchen concentration from our study, no firm conclusions can be drawn.

The increased odds for term LBW (OR = 1.69; 95% CI = 1.03, 2.02) associated with 10-unit increase in PM$_{2.5}$ were moderately higher than those reported by Tielsch et al., (2009) (OR =

1.21, 95% CI= 1.11, 1.31)(41) and Mavalankar et al., (1992) (OR=1.23; 95% CI : 1.01, 1.49)
(42).

Exposure estimation

In our study, information on stacking of fuel during an average week was used, to categorize HAP exposure into three categories. We developed a predicted model using information on fuel stacking, type of kitchen, and chimney to predict kitchen $PM_{2.5}$ concentrations. This novel approach was used to evaluate the feasibility of using questionnaire responses to predict $PM_{2.5}$ concentration in kitchen during cooking period, for households that did not have air pollution measurement data. Cross-validation approach evaluated the robustness of the model with a R^2 of 57%. Personal monitoring of exposure to HAP by wearing portable devices for a period of 24 to 48 hours can substantially reduce exposure misclassification and improve the power of the study to detect relationships between exposure to HAP and adverse health outcomes (43). However, conducting area or personal air quality monitoring in all study subjects significantly increases study costs and respondent burden.

Previous HAP studies have mostly relied on questionnaire responses as surrogate for exposure. In more recent studies, questionnaire data is supplemented with area measurements for short time periods (snapshots) at multiple time points to generate time-integrated exposure metric to characterize chronic exposure (25). In these studies, households are categorized based on their primary fuel use as households using wood, kerosene or LPG and electricity. Few studies have considered multiple types of fuel use or the proportion of fuels used, resulting in uncertainty in nature and magnitude of exposure variability and misclassification of exposure in epidemiologic studies (25).

Balakrishnan et al.(2013), conducted one of the first studies that included 24-hour $PM_{2.5}$ measurements in all study households (n, 474). They fitted a regression analysis to model 24-hour data to predict household $PM_{2.5}$ concentrations (44). They identified fuel type, kitchen type, ventilation, geographical location and cooking duration, to be significant predictors of $PM_{2.5}$ concentrations (R^2= 0.33) with a fair degree of correlation (r=0.56) between modeled and measured values.(45) Another study conducted by Baxter et al. (2007), used questionnaire information combined with ambient air quality measurements to predict indoor concentrations of nitrogen dioxide, $PM_{2.5}$, and elemental carbon in lower socio-economic status homes in urban Boston, Massachusetts (46). Their study found that outdoor concentrations, cooking time and occupant density were significant predictors of indoor $PM_{2.5}$ concentration (adjusted R^2=0.36). The inclusion of open windows as an effect modifier for ambient air increased the R^2 to 0.40. A recent study conducted by Zhou et al. (2018) in Shanghai, China, used outdoor air measurements and household characteristics to predict outdoor $PM_{2.5}$ infiltration of indoor air (47). The models stratified by seasons and including information on air conditioning use, meteorological factors, floor of residence and building age, predicted 60.0%-68.4% of the variance in 7-day averages of indoor $PM_{2.5}$ infiltration by outdoor air.

The R^2 for the exposure model developed in our study (57%) was lower than the R^2 (66%) reported by Zhou et al (47); and 1.6 times greater than the R^2 (0.36) reported by Balakrishnan et al. (2013) (45). The correlation between estimated and measured $PM_{2.5}$ (r, 0.74 vs. r, 0.56) was also higher in our study compared to that reported by Balakrishnan (2013) study.

Our results indicate moderate feasibility of using detailed questionnaire information to quantify kitchen $PM_{2.5}$ concentrations. Several other factors can affect personal exposure including the high degree of variability from stacking of fuels, different stove types and kitchen

designs, and time-activity pattern of the subjects i.e., the amount of time spent in the kitchen and inside the homes as compared to outside. Furthermore, changes in patterns of fuel use over the course of pregnancy could change personal exposure. Increased frequency, longer duration of air quality monitoring and time activity diary of time spent indoor could improve the accuracy of our predictive model.

Our study has several strengths. HAP exposure was analyzed in several different ways with consistent results. The retention rate in the study was almost 84% (545/656) after accounting for miscarriages, stillbirths and twin pregnancies. The study participants were identified through routine prenatal checkups at the antenatal clinics and were representative of the study population. All births occurred in hospitals and birth weight was recorded within 24-hours. Gestational age was determined by ultrasound examination in 1st trimester if there was doubt about the last menstrual period, consequently, reducing information bias for LBW. In Sri Lanka, government-funded public health system provides prenatal and antenatal care to almost 99% of the pregnant mothers. Thus, self-reported maternal medical history was verified with maternal pregnancy records obtained from the antenatal clinics. Maternal lifestyle, such as smoking cigarettes, and use of alcohol, are known confounding factors for adverse birth outcomes. However, none of the mothers in our study smoked or consumed alcohol. Information on ETS was self-reported but no significant differences were noted for LBW and normal children. We were able to collect and adjust for known confounding factors associated with LBW including pregnancy induced hypertension, gestational diabetes, and other complications during pregnancy such as essential hypertension, high head, hyperemesis, hyperthyroidism, hypothyroidism, and threatened miscarriage.

The major limitation of our study was the lack of 24-hour air quality measurements for indoor and ambient air quality. Data on indoor air quality was available for 60% of households in the study and were used to quantify indoor air quality for remaining 40%, which could have resulted in some exposure misclassification. However, internal consistency in results suggest minimum bias.

Conclusion.

Our study evaluating HAP and birth outcomes in Sri Lanka adds to the growing evidence linking HAP to LBW and term LBW. Although, air quality measurements were conducted in a subset of households, both qualitative and quantitative data from measured households were used to quantify HAP exposure in households without measured data. Results of this study are important as almost 74% of the population in Sri Lanka continue to use wood for cooking. Since almost one-third of world's population is exposed to HAP from solid fuel, even modest increase in the risk of adverse pregnancy outcomes has large-scale implications on the overall health of the population.

Acknowledgements

The Prenatal Exposure to Biomass Smoke and Infant Neurodevelopment in Sri Lanka, study was supported by the National Institutes of Environmental Sciences (NIEHS: grant R21ES018730).

Conflict of Interest

The authors declare that they have no competing financial interests.

References

1. Cohen AJ, Brauer M, Burnett R, Anderson HR, Frostad J, Estep K, et al. Estimates and
    25-year trends of the global burden of disease attributable to ambient air pollution: an
    analysis of data from the Global Burden of Diseases Study 2015. Lancet (London,
    England) [Internet]. 2017 May 13 [cited 2017 Sep 22];389(10082):1907–18. Available
    from: http://www.ncbi.nlm.nih.gov/pubmed/28408086

2. GBD 2013 Mortality and Causes of Death Collaborators G 2013 M and C of D. Global,
    regional, and national age-sex specific all-cause and cause-specific mortality for 240
    causes of death, 1990-2013: a systematic analysis for the Global Burden of Disease Study
    2013. Lancet (London, England). 2015;385(9963):117–71.

3. Collaborators G 2013 Daly and H, Bernabe E. Global, regional, and national disability-
    adjusted life years (DALYs) for 306 diseases and injuries and healthy life expectancy
    (HALE) for 188 countries, 1990–2013: quantifying the epidemiological transition. Lancet
    [Internet]. 2015 Nov 28 [cited 2016 Jan 19];386(1009). Available from:
    https://kclpure.kcl.ac.uk/portal/en/publications/global-regional-and-national-
    disabilityadjusted-life-years-dalys-for-306-diseases-and-injuries-and-healthy-life-
    expectancy-hale-for-188-countries-19902013-quantifying-the-epidemiological-
    transition(35529eb4-ddf4-40b7-b12d-b67c06e8f85d)/export.html

4. Naeher LP, Brauer M, Lipsett M, Zelikoff JT, Simpson CD, Koenig JQ, et al. Woodsmoke
    health effects: a review. Inhal Toxicol [Internet]. 2007 Jan [cited 2014 Oct 15];19(1):67–
    106. Available from: http://www.ncbi.nlm.nih.gov/pubmed/17127644

5.   Cohen AJ, Brauer M, Burnett R, Anderson HR, Frostad J, Estep K, et al. Estimates and 25-year trends of the global burden of disease attributable to ambient air pollution: an analysis of data from the Global Burden of Diseases Study 2015. Lancet [Internet]. 2017 May 13 [cited 2018 Feb 14];389(10082):1907–18. Available from: http://www.ncbi.nlm.nih.gov/pubmed/28408086

6.   World Health Organization. WHO | Household air pollution: Disease impacts [Internet]. WHO. World Health Organization; 2017 [cited 2018 Jul 11]. Available from: http://www.who.int/airpollution/household/health-impacts/diseases/en/

7.   Amegah AK, Jaakkola JJ. Household air pollution and the sustainable development goals. Bull World Health Organ [Internet]. 2016 Mar 1 [cited 2018 Oct 27];94(3):215–21. Available from: http://www.who.int/entity/bulletin/volumes/94/3/15-155812.pdf

8.   Elledge MF, Phillips MJ, Thornburg VE, Everett KH, Nandasena S. A profile of biomass stove use in Sri Lanka. Int J Environ Res Public Health [Internet]. 2012 [cited 2017 Nov 1];9(4):1097–110. Available from: http://www.ncbi.nlm.nih.gov/pubmed/22690185

9.   Census.lk. Department of Census and Statistics-Sri Lanka [Internet]. Department of Census and Statistics. 2014 [cited 2014 Sep 7]. Available from: http://www.statistics.gov.lk/

10.   Elledge MF, Phillips MJ, Thornburg VE, Everett KH, Nandasena S. A profile of biomass stove use in Sri Lanka. Int J Environ Res Public Health [Internet]. 2012 Apr [cited 2014 Nov 14];9(4):1097–110. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3366602&tool=pmcentrez&rendertype=abstract

11. Nandasena S, Wickremasinghe AR, Sathiakumar N. Biomass fuel use for cooking in Sri Lanka: analysis of data from national demographic health surveys. Am J Ind Med [Internet]. 2012 Dec [cited 2014 Nov 14];55(12):1122–8. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3319003&tool=pmcentrez&rendertype=abstract

12. WHO. WHO | Sri Lanka [Internet]. WHO. World Health Organization; 2015 [cited 2018 Oct 24]. Available from: http://www.who.int/gho/countries/lka/en/

13. Zhang K, Lu Y, Zhao H, Guo J, Gehendra M, Qiu H, et al. Association between atmospheric particulate matter and adverse pregnancy outcomes in the population [Internet]. Vol. 9, Int J Clin Exp Med. 2016 [cited 2018 Oct 31]. Available from: www.ijcem.com/

14. Nieuwenhuijsen MJ, Dadvand P, Grellier J, Martinez D, Vrijheid M. Environmental risk factors of pregnancy outcomes: a summary of recent meta-analyses of epidemiological studies. Environ Heal [Internet]. 2013 Dec 15 [cited 2018 Oct 31];12(1):6. Available from: http://ehjournal.biomedcentral.com/articles/10.1186/1476-069X-12-6

15. Srám RJ, Binková B, Dejmek J, Bobak M. Ambient air pollution and pregnancy outcomes: a review of the literature. Environ Health Perspect [Internet]. 2005 Apr [cited 2018 Oct 31];113(4):375–82. Available from: http://www.ncbi.nlm.nih.gov/pubmed/15811825

16. Amegah AK, Quansah R, Jaakkola JJK. Household air pollution from solid fuel use and risk of adverse pregnancy outcomes: a systematic review and meta-analysis of the empirical evidence. Nawrot TS, editor. PLoS One [Internet]. 2014 Dec 2 [cited 2018 Oct

22];9(12):e113920. Available from: https://dx.doi.org/10.1371/journal.pone.0113920

17.   Lamichhane DK, Leem J-H, Lee J-Y, Kim H-C. A meta-analysis of exposure to
      particulate matter and adverse birth outcomes. Environ Health Toxicol [Internet]. 2015
      [cited 2018 Jul 11];30:e2015011. Available from:
      http://www.ncbi.nlm.nih.gov/pubmed/26796890

18.   Shah PS, Balkhair T, Knowledge Synthesis Group on Determinants of Preterm/LBW
      births. Air pollution and birth outcomes: A systematic review. Environ Int [Internet]. 2011
      Feb [cited 2018 Jul 11];37(2):498–516. Available from:
      http://www.ncbi.nlm.nih.gov/pubmed/21112090

19.   Yucra S, Tapia V, Steenland K, Naeher LP, Gonzales GF. Maternal exposure to biomass
      smoke and carbon monoxide in relation to adverse pregnancy outcome in two high
      altitude cities of Peru. Environ Res [Internet]. 2014 Apr [cited 2015 Jan 9];130:29–33.
      Available from: http://www.sciencedirect.com/science/article/pii/S0013935114000127

20.   Smith KR, Pillarisetti A. Household Air Pollution from Solid Cookfuels and Its Effects on
      Health [Internet]. Injury Prevention and Environmental Health. The International Bank for
      Reconstruction and Development / The World Bank; 2017 [cited 2019 Feb 7]. Available
      from: http://www.ncbi.nlm.nih.gov/pubmed/30212117

21.   Amegah AK, Quansah R, Jaakkola JJK. Household Air Pollution from Solid Fuel Use and
      Risk of Adverse Pregnancy Outcomes: A Systematic Review and Meta-Analysis of the
      Empirical Evidence. Nawrot TS, editor. PLoS One [Internet]. 2014 Dec 2 [cited 2018 Jul
      11];9(12):e113920. Available from: http://dx.plos.org/10.1371/journal.pone.0113920

22.   Mansi G, Raimondi F, Pichini S, Capasso L, Sarno M, Zuccaro P, et al. Neonatal urinary

cotinine correlates with behavioral alterations in newborns prenatally exposed to tobacco smoke. Pediatr Res [Internet]. 2007 Feb [cited 2015 Mar 9];61(2):257–61. Available from: http://www.ncbi.nlm.nih.gov/pubmed/17237732

23. El Marroun H, Schmidt MN, Franken IHA, Jaddoe VW V, Hofman A, van der Lugt A, et al. Prenatal tobacco exposure and brain morphology: a prospective study in young children. Neuropsychopharmacology [Internet]. 2014 Mar [cited 2018 Oct 27];39(4):792–800. Available from: http://www.ncbi.nlm.nih.gov/pubmed/24096296

24. Lambers DS, Clark KE. The maternal and fetal physiologic effects of nicotine. Semin Perinatol [Internet]. 1996 Apr [cited 2018 Oct 27];20(2):115–26. Available from: http://www.ncbi.nlm.nih.gov/pubmed/8857697

25. Clark ML, Peel JL, Balakrishnan K, Breysse PN, Chillrud SN, Naeher LP, et al. Health and household air pollution from solid fuel use: the need for improved exposure assessment. Environ Health Perspect [Internet]. 2013 Oct [cited 2014 Oct 30];121(10):1120–8. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3801460&tool=pmcentrez&rendertype=abstract

26. Carter E, Norris C, Dionisio KL, Balakrishnan K, Checkley W, Clark ML, et al. Assessing Exposure to Household Air Pollution: A Systematic Review and Pooled Analysis of Carbon Monoxide as a Surrogate Measure of Particulate Matter. [cited 2017 Nov 4]; Available from: https://doi.org/

27. Balakrishnan K, Ghosh S, Thangavel G, Sambandam S, Mukhopadhyay K, Puttaswamy N, et al. Exposures to fine particulate matter (PM2.5) and birthweight in a rural-urban,

mother-child cohort in Tamil Nadu, India. Environ Res [Internet]. 2018 Feb [cited 2019 Mar 24];161:524–31. Available from: http://www.ncbi.nlm.nih.gov/pubmed/29227900

28. Statistics | Sri Lanka | UNICEF [Internet]. [cited 2018 Oct 27]. Available from: https://www.unicef.org/infobycountry/sri_lanka_statistics.html#0

29. World Health Organization. WHO | The contribution of Public Health Midwives to better health in rural communities in Sri Lanka [Internet]. WHO. World Health Organization; 2011 [cited 2019 Jan 31]. Available from: https://www.who.int/workforcealliance/forum/2011/hrhawardscs28/en/

30. Gunathunga W, Fernando DN. Assessment of community maternal care performance of public health midwives of a province in Sri Lanka: a multi-method approach. Southeast Asian J Trop Med Public Health [Internet]. 2000 Jun [cited 2019 Jan 31];31(2):310–8. Available from: http://www.ncbi.nlm.nih.gov/pubmed/11127332

31. Fernando D. Health care systems in transition III. Sri Lanka, Part I. An overview of Sri Lanka's health care system [Internet]. [cited 2019 Mar 24]. Available from: https://watermark.silverchair.com/220014.pdf?token=AQECAHi208BE49Ooan9kkhW_E rcy7Dm3ZL_9Cf3qfKAc485ysgAAAm0wggJpBgkqhkiG9w0BBwagggJaMIICVgIBAD CCAk8GCSqGSIb3DQEHATAeBglghkgBZQMEAS4wEQQMPRcwlIF_cdFEXV90AgE QgIICILrjIprbO7H-
0K_RIzvRvruxyJkBhytoSOxfDY4wsKRKGeemYLeR2IxKKCZCp0fstFne-c42-
aPcfQhGxF8QWjNzjOS2tN1TUSdil1gTCyDTO9JxFtDX-
ovDy1Q5uqwJMfvsMklSzPkoexXFyKZCWwRM5GwPsJP9ogAG0ptS4XqA0FW3S8Yp
1JDDozw6dRY9TVY55icxofeW-BwzvCOQjlM7vOoFQENPgPiLkxxpPd-G00o-

Vv2aS6XXtwQ22kGlKPWns63OV-qlekNQ-

e7FI0VYid51u07F5rJIZM7MN7BLd2RylOpRkyrUn_ph6Wow_SAkxEyhdX69KVy87Jf

LabHUdMuQyMJHcWPI1u05LzAoIINd-AUZbzWBC1_07qL_PQxCOds-

uX2WDysl_qCL-GPWdLxFZY_-

sgi26_TCHOPwrEAKesrhUeqhlqWVF8GARDBQ_5JJyRCSOWTgrHjmnQzyIxkegoMsz

FERhK0GkaLp9dXXTt_wReRfQUgjT98zXeYeSXCU59beF4jH_xOy00MVGGhzte3Od

WZvniYFhGLoMk0OKR__tBd9-

mmZXiYDm2ZnUxkWbA_PhOljVcDKR_sUB9NNJKBc40fir2Zk9pEUe2pmZp2-

KowQfx1w5agvhLRs6VBonf_I83KOWfxZozvlQrdLtt8S9yxAnz0JT1OkHEyG4TaPPOY

L2jhGtx3BKwaTgQ2J47FRzEhvyHM8TwE

32. Case study from Sri Lanka PRIMARY HEALTH CARE SYSTEMS (PRIMASYS)
[Internet]. 2017 [cited 2019 Mar 24]. Available from: http://apps.who.int/bookorders.

33. Cutland CL, Lackritz EM, Mallett-Moore T, Bardají A, Chandrasekaran R, Lahariya C, et
al. Low birth weight: Case definition & guidelines for data collection, analysis, and
presentation of maternal immunization safety data. Vaccine [Internet]. 2017 [cited 2019
Feb 8];35(48 Pt A):6492–500. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/29150054

34. World Health Organization. Preterm birth [Internet]. World Health Organization. 2018
[cited 2019 Feb 8]. Available from: https://www.who.int/news-room/fact-
sheets/detail/preterm-birth

35. Sun X, Luo X, Zhao C, Zhang B, Tao J, Yang Z, et al. The associations between birth
weight and exposure to fine particulate matter (PM2.5) and its chemical constituents

during pregnancy: A meta-analysis. Environ Pollut [Internet]. 2016 Apr 1 [cited 2018 Oct 22];211:38–47. Available from:

https://www.sciencedirect.com/science/article/pii/S0269749115302360

36.   Wylie BJ, Coull BA, Hamer DH, Singh MP, Jack D, Yeboah-Antwi K, et al. Impact of biomass fuels on pregnancy outcomes in central East India. Environ Health [Internet]. 2014 Jan [cited 2015 Feb 11];13(1):1. Available from:

http://www.ehjournal.net/content/13/1/1

37.   Sreeramareddy CT, Shidhaye RR, Sathiakumar N. Association between biomass fuel use and maternal report of child size at birth--an analysis of 2005-06 India Demographic Health Survey data. BMC Public Health [Internet]. 2011 Jan [cited 2015 Feb 11];11:403. Available from:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3125371&tool=pmcentrez&rendertype=abstract

38.   Gilliland F, Avol E, Kinney P, Jerrett M, Dvonch T, Lurmann F, et al. Air pollution exposure assessment for epidemiologic studies of pregnant women and children: lessons learned from the Centers for Children's Environmental Health and Disease Prevention Research. Environ Health Perspect [Internet]. 2005 Oct [cited 2018 Oct 18];113(10):1447–54. Available from: http://www.ncbi.nlm.nih.gov/pubmed/16203261

39.   Bruce NG, Dherani MK, Das JK, Balakrishnan K, Adair-Rohani H, Bhutta ZA, et al. Control of household air pollution for child survival: estimates for intervention impacts. BMC Public Health [Internet]. 2013 Jan [cited 2015 Feb 11];13 Suppl 3:S8. Available from:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3847681&tool=pmcentrez&re
ndertype=abstract

40.   Misra P, Srivastava R, Krishnan A, Sreenivaas V, Pandav CS. Indoor air pollution-related
      acute lower respiratory infections and low birthweight: a systematic review. J Trop Pediatr
      [Internet]. 2012 Dec 1 [cited 2018 Oct 22];58(6):457–66. Available from:
      https://academic.oup.com/tropej/article-lookup/doi/10.1093/tropej/fms017

41.   Tielsch JM, Katz J, Thulasiraj RD, Coles CL, Sheeladevi S, Yanik EL, et al. Exposure to
      indoor biomass fuel and tobacco smoke and risk of adverse reproductive outcomes,
      mortality, respiratory morbidity and growth among newborn infants in south India. Int J
      Epidemiol [Internet]. 2009 Oct 1 [cited 2015 Feb 10];38(5):1351–63. Available from:
      http://ije.oxfordjournals.org/content/38/5/1351.abstract?ijkey=0fb09a0ddfb109306303f59
      967b5f56811fd4e65&keytype2=tf_ipsecsha

42.   MAVALANKAR D V, GRAY RH, TRIVEDI CR. Risk Factors for Preterm and Term
      Low Birthweight in Ahmedabad, India. Int J Epidemiol [Internet]. 1992 Apr 1 [cited 2015
      Feb 11];21(2):263–72. Available from:
      http://ije.oxfordjournals.org/content/21/2/263.abstract?ijkey=ce4e912ab7e2639100ad7f0d
      eb0b6a267787d9ee&keytype2=tf_ipsecsha

43.   Charles McCoy D, Waldman M, Field Team C, Fink G. Measuring early childhood
      development at a global scale: Evidence from the Caregiver-Reported Early Development
      Instruments. Early Child Res Q [Internet]. 2018 [cited 2019 Feb 1];45:58–68. Available
      from: https://doi.org/10.1016/j.ecresq.2018.05.002

44.   Balakrishnan K, Ghosh S, Thangavel G, Sambandam S, Mukhopadhyay K, Puttaswamy

N, et al. Exposures to fine particulate matter (PM2.5) and birthweight in a rural-urban, mother-child cohort in Tamil Nadu, India. Environ Res [Internet]. 2018 Feb 1 [cited 2018 Nov 4];161:524–31. Available from: https://www.sciencedirect.com/science/article/pii/S0013935117312276

45.   Balakrishnan K, Ghosh S, Ganguli B, Sambandam S, Bruce N, Barnes DF, et al. State and national household concentrations of PM2.5 from solid cookfuel use: results from measurements and modeling in India for estimation of the global burden of disease. Environ Health [Internet]. 2013 Sep 11 [cited 2018 Oct 19];12(1):77. Available from: http://www.ncbi.nlm.nih.gov/pubmed/24020494

46.   Baxter LK, Clougherty JE, Paciorek CJ, Wright RJ, Levy JI. Predicting residential indoor concentrations of nitrogen dioxide, fine particulate matter, and elemental carbon using questionnaire and geographic information system based data. Atmos Environ (1994) [Internet]. 2007 Oct [cited 2018 Oct 19];41(31):6561–71. Available from: http://www.ncbi.nlm.nih.gov/pubmed/19830252

47.   Zhou X, Cai J, zhao Y, Chen R, Wang C, Zhao A, et al. Estimation of residential fine particulate matter infiltration in Shanghai, China. Environ Pollut [Internet]. 2018 Feb [cited 2018 Oct 19];233:494–500. Available from: http://www.ncbi.nlm.nih.gov/pubmed/29102879

Table 1. Descriptive statistics comparing variables between ever wood users and LPG users

	Total N	Wood N(%)	LPG N(%)	P-value
Total subjects	545	423	122	
Demographic factors				
				0.3020
Mother's age		423	122	
Mean (SD)	29(4.9)	29(4.9)	28(4.9)	
Median (IQR)	29(7)	29(7)	28(7.5)	
Missing	18			
Mother's education				0.0054
0-5 years	74	58(78)	16(22)	
5-10 years	314	258(82)	56(18)	
>10 years	140	96(69)	44(31)	
Missing	17	11(65)	6(35)	
Father's education				**0.0003**
0-5 years	68	57(84)	11(16)	
>5-10 years	332	271(82)	61(18)	
>10 years	126	82(65)	44(35)	
Missing	19	13(68)	6(32)	
Water source				
Piped water	133	103(77)	30(23)	0.3987
Other source of water (well, tube well)	394	308(78)	86(22)	
Missing	19	12(67)	6(33)	
Toilet				
Private flushed toilet	448	346(77)	102(23)	0.3987
Other (shared flushed, private pit, compositing, bucket)	76	62(82)	14(18)	
Missing	21	15(71)	6(29)	
Owning a house				**0.0033**
Yes	437	56(66)	29(34)	
No	85	351(80)	86(20)	
Missing	23	16(70)	7(30)	
Type of house				**0.0002**
Semi-permanent or improvised	349	122(69)	56(31)	
Permanent	178	289(83)	60(17)	
Missing	18	12(67)	6(33)	
Income				**0.0006**
< 15000	96	84(88)	12(13)	
15000-20000	93	82(88)	11(12)	
20001 - 25000	93	68(74)	24(26)	
>25000	242	173(71)	69(29)	
Missing	22	16(73)	6(27)	
Parity		136(74)	48(26)	0.0892
0	184	170(78)	49(22)	

Table 1. Descriptive statistics comparing variables between ever wood users and LPG users

	Total N	Wood N(%)	LPG N(%)	P-value
1	219	108(84)	20(16)	
≥2	128	9(64)	5(36)	
Missing	14			
Past miscarriage				0.3586
Yes	28	24(86)	4(14)	
No	517	399(77)	118(23)	
Pregnancy hypertension				0.2664
Yes	26	19(73)	7(27)	
No	519	404(78)	115(22)	
Gestational diabetes				0.5695
Yes	19	16(84)	3(16)	
No	526	407(77)	119(23)	
Kitchen characteristics				
Continuous $PM_{2.5}$ (in µg/m^3)				**<.0001**
0-35	10	3(30)	7(70)	
35-75	36	18(50)	18(50)	
75-28245	498	401(81)	97(19)	
Kitchen type				0.4826
Indoor	465	353(76)	112(24)	
Temporary hut	47	46(98)	1(2)	
Outdoor	15	12(80)	3(20)	
Missing	18	12(67)	6(33)	
Functional chimney				**<.0001**
No	264	229(87)	35(13)	
Yes	281	194(69)	87(31)	
Daily cooking time (in minutes)				
≤ 60	148	111(75)	37(25)	
>60 - ≤ 90	218	166(76)	52(24)	0.2474
>90	154	129(84)	25(16)	
Missing	25	17(68)	8(32)	
Open windows				0.3730
0	258	205(80)	51(20)	
1	262	198(76)	63(24)	
2	28	20(71)	8(29)	
Other source of indoor pollution (smoking, incense)				0.3763
1	95	77(81)	18(19)	
0	450	346(77)	104(23)	
Outdoor sources of pollution				0.1221
1	40	34(85)	6(15)	
0	508	389(77)	116(23)	

Table 1. Descriptive statistics comparing variables between ever wood users and LPG users

	Total N	Wood N(%)	LPG N(%)	P-value
Environmental risk				0.2444
Yes	259	203(78)	56(22)	
No	286	220(77)	66(23)	
Birth outcomes				
Child's gender				0.1942
Male	285	208(80)	52(20)	
Female	261	214(75)	70(25)	
Missing	1	1(100)	0(0)	
Preterm births				0.4394
Yes	57	42(74)	15(26)	
No	486	380(78)	106(22)	
Missing	2	1(50)	1(50)	

Significant at p-value <0.05

Table 2. Distribution of mean birth weight and prevalence of LBW across demographic, maternal and household characteristics

	Total	Birth weight	LBW
	N	Mean (SD)	N(%)
Total subjects	545		70(100)
Demographic factors			
Mother's age			
Mean (SD)	29(4.9)		28(4.9)
Median (IQR)	29(7)		28(5.5)
Missing	18		
Mother's education			
0-5 years	74	2.9(0.5)	9(12)
5-10 years	314	2.9(0.5)	41(13)
>10 years	140	3(0.5)	18(13)
Missing	17		2(13)
Father's education			
0-5 years	68	2.9(0.5)	12(18)
>5-10 years	332	2.9(0.5)	45(14)
>10 years	126	3(0.5)	11(8.7)
Missing	19		2(11)
Water source			
Piped water	133	3(0.5)	12(9.0)
Other source of water (well, tube well)	394	2.9(0.5)	56(14)
Missing	19		2 (11)
Toilet			
Private flushed toilet	448	2.9(0.5)	60(13)
Other (shared flushed, private pit, compositing, bucket)	76	3.0(0.5)	8(11)
Missing	21		2 (10)
Owning a house			
Yes	437	3.0(0.5)	58(13)
No	85	2.9(0.4)	10(12)
Missing	23		2 (8.7)
Type of house			
Semi-permanent or improvised	349	**2.9(0.5)****	51(15)
Permanent	178	**3(0.4)**	17(10)
Missing	18		2(11)
Income			
< 15000	96	2.9(0.4)	11(11)
15000-20000	93	3(0.5)	10(11)
20001 - 25000	93	3(0.5)	16(17)
>25000	242	2.9(0.5)	30(12)
Missing	22		3(14)
Parity			
0	184	2.9(0.4)	23(13)

Table 2. Distribution of mean birth weight and prevalence of LBW across demographic, maternal and household characteristics

	Total	Birth weight	LBW
	N	Mean (SD)	N(%)
1	219	3(0.5)	29(13)
≥2	128	2.9(0.5)	18(14)
Missing	14		0
Past miscarriage			
Yes	28	2.9(0.6)	4(14)
No	517	3(0.5)	66(13)
Pregnancy hypertension			
Yes	26	**2.4(0.7)****	**13(50)****
No	519	**3(0.4)**	**57(11)**
Gestational diabetes			
Yes	19	2.9(0.5)	4(6)
No	526	3(0.5)	66(94)
Kitchen characteristics			
Dichotomous fuel use			
Wood	423	2.9(0.4)	58(14)
LPG	122	3.0(0.5)	12(10)
Fuel percentage			
50%-100% Wood	283	**2.9(0.5)****	43(15)
>50% LPG + wood	139	**2.9(0.5)**	15(11)
100% LPG	122	**3(0.5)**	12(10)
Missing	1		**1**
Continuous PM$_{2.5}$ (in µg/m^3)			
0-35	10	2.9(0.4)	1(10)
35-75	36	3.1(0.5)	4(11)
75-28245	498	2.97(0.5)	65(13)
Kitchen type			
Indoor	465	3(0.5)	58(12)
Temporary hut	47	2.9(0.5)	8(17)
Outdoor	15	2.9(0.6)	2(13)
Missing	18		2 (11)
Functional chimney			
No	264	2.9(0.5)	35(13)
Yes	281	3(0.5)	35(12)
Daily cooking time (in minutes)			
≤ 60	148	3.0(0.5)	18(12)
>60 - ≤ 90	218	2.94(0.5)	28(13)
>90	154	2.9(0.5)	21(14)
Missing	25	3.0(0.5)	3(0.6)
Open windows			
0	258	3(0.5)	32(13)
1	262	3(0.5)	36(14)

Table 2. Distribution of mean birth weight and prevalence of LBW across demographic, maternal and household characteristics

	Total	Birth weight	LBW
	N	Mean (SD)	N(%)
2	28	2.9(0.4)	2(7.1)
Other source of indoor pollution (smoking, incense)			
1	95	**2.8(0.5)***	**18(19)***
0	450	**3(0.5)**	**58(12)**
Outdoor sources of pollution			
1	40	2.9(0.5)	5(13)
0	508	3(0.5)	65(13)
Environmental risk			
Yes	259	2.9(0.5)	28(11)
No	286	3(0.5)	42(15)
Birth outcomes			
Child's gender			
Male	285	2.97(0.5)	38(13)
Female	261	2.9(0.4)	32(12)
Missing	1		0
Preterm births			
Yes	57	**2.5(0.5)****	**24(42)****
No	486	**3.0(0.4)**	**46(9)**
Missing	2		

** significant at p <0.05; * significant at p<0.1

Table 3. Relationship between HAP exposure and birth weight (in kgs)

	Exposure	Crude β (Standard error (SE))	Adjusted β(SE)	P-value	Adjusted R^2, sample size
Model[1]	Continuous $PM_{2.5}$	-0.06(-3.17)	**-0.03(0.02)**	**0.0339**	0.10
Model[2]	Wood	-0.07(0.05)	-0.07(0.05)	0.2008	0.09
	LPG	0			
Model[3]	>50% wood	-0.10 (0.05)	**-0.13(0.06)**	**0.0229**	0.11
	>50% LPG + wood	0.02 (0.06)	0.002(0.06)	0.9724	
	100% LPG	0			

[1]Adjusted for mother's education, father's education, type of house, ownership of the house, toilet, income, average daily cooking time, other sources of indoor pollution, pregnancy induced hypertension

[2,3]Adjusted for mother's education, father's education, type of house, ownership of the house, toilet, income, kitchen type, presence of functional chimney, average daily cooking time, other sources of indoor pollution, pregnancy induced hypertension

Table 4. Relationship between HAP exposure and LBW

	Exposure	Crude OR (95% CI)	Adjusted OR (95% CI)	P-value
Model[1]	Continuous $PM_{2.5}$	1.14(0.94-1.38)	**1.24(1.01-1.55)**	**0.0428**
Model[2]	Wood	1.46(0.76-2.81)	1.82 (0.81-4.11)	0.1500
	LPG	1.00	1.00	
Model[3]	>50% wood	1.64 (0.83-3.24)	**2.14(0.90-5.07)**	**0.0791**
	>50% LPG + wood	1.11 (0.50-2.47)	1.44(0.57-3.63)	0.6377
	100% LPG	1.00	1.00	

[1]Adjusted for mother's education, father's education, type of house, ownership of the house, toilet, income, average daily cooking time, other sources of indoor pollution, pregnancy induced hypertension

[2,3]Adjusted for mother's education, father's education, type of house, ownership of the house, toilet, income, kitchen type, presence of functional chimney, average daily cooking time, other sources of indoor pollution, pregnancy induced hypertension